

# SC223 - Linear Algebra

Aditya Tatu

Lecture 7



August 8, 2023

# LU Decomposition

$$A = \begin{bmatrix} 1 & -2 & -1 & -1 \\ 2 & 0 & 3 & 2 \\ -2 & 3 & -2 & 1 \\ 3 & -4 & 2 & 1 \end{bmatrix}$$

- $E_3 E_2 E_1 A = EA = U \Rightarrow A = E^{-1} U$ .
- By Theorem 2,  $E^{-1}$  is a lower triangular matrix. Define  $L := E^{-1}$ .  
Thus  $A = LU$ , known as the **LU decomposition**.
- For this example:

$$\underbrace{\begin{bmatrix} 1 & -2 & -1 & -1 \\ 2 & 0 & 3 & 2 \\ -2 & 3 & -2 & 1 \\ 3 & -4 & 2 & 1 \end{bmatrix}}_A = \underbrace{\begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ -2 & -1/4 & 1 & 0 \\ 3 & 1/2 & -10/11 & 1 \end{bmatrix}}_L \underbrace{\begin{bmatrix} 1 & -2 & -1 & -1 \\ 0 & 4 & 5 & 4 \\ 0 & 0 & -11/4 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}}_U$$

Is  $A = LU$  always possible?

$$\left[ \begin{array}{cccc|c} 1 & 2 & 1 & -2 & 3 \\ 2 & 4 & 0 & 1 & 4 \\ -2 & -3 & 0 & 5 & -4 \\ 0 & 1 & 2 & 1 & 2 \end{array} \right]$$

Is  $A = LU$  always possible?

$$\left[ \begin{array}{cccc|c} 1 & 2 & 1 & -2 & 3 \\ 2 & 4 & 0 & 1 & 4 \\ -2 & -3 & 0 & 5 & -4 \\ 0 & 1 & 2 & 1 & 2 \end{array} \right]$$

$$\bullet E_1 = \left[ \begin{array}{cccc} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right],$$

Is  $A = LU$  always possible?

$$\left[ \begin{array}{cccc|c} 1 & 2 & 1 & -2 & 3 \\ 2 & 4 & 0 & 1 & 4 \\ -2 & -3 & 0 & 5 & -4 \\ 0 & 1 & 2 & 1 & 2 \end{array} \right]$$

$$\bullet E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

Is  $A = LU$  always possible?

$$\left[ \begin{array}{cccc|c} 1 & 2 & 1 & -2 & 3 \\ 2 & 4 & 0 & 1 & 4 \\ -2 & -3 & 0 & 5 & -4 \\ 0 & 1 & 2 & 1 & 2 \end{array} \right]$$

$$\bullet E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$\bullet P_{23} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix},$$

Is  $A = LU$  always possible?

$$\left[ \begin{array}{cccc|c} 1 & 2 & 1 & -2 & 3 \\ 2 & 4 & 0 & 1 & 4 \\ -2 & -3 & 0 & 5 & -4 \\ 0 & 1 & 2 & 1 & 2 \end{array} \right]$$

$$\bullet E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$\bullet P_{23} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, P_{23} E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

Is  $A = LU$  always possible?

$$\left[ \begin{array}{cccc|c} 1 & 2 & 1 & -2 & 3 \\ 2 & 4 & 0 & 1 & 4 \\ -2 & -3 & 0 & 5 & -4 \\ 0 & 1 & 2 & 1 & 2 \end{array} \right]$$

$$\begin{aligned} \bullet E_1 &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \\ 0 & 1 & 2 & 1 \end{bmatrix} \\ \bullet P_{23} &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, P_{23} E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \end{bmatrix} \\ \bullet E_2 &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2 & 0 & 1 \end{bmatrix}, \end{aligned}$$

-1



Is  $A = LU$  always possible?

$$\left[ \begin{array}{cccc|c} 1 & 2 & 1 & -2 & 3 \\ 2 & 4 & 0 & 1 & 4 \\ -2 & -3 & 0 & 5 & -4 \\ 0 & 1 & 2 & 1 & 2 \end{array} \right]$$

$$\bullet E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$\bullet P_{23} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, P_{23} E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$\bullet E_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2 & 0 & 1 \end{bmatrix}, E_2 P_{23} E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Is  $A = LU$  always possible?

$$\overline{E_3} \overline{E_2} \overline{E_1} \underbrace{P_{23} A}_{\overline{A}} = U$$

$$\left[ \begin{array}{cccc|c} 1 & 2 & 1 & -2 & 3 \\ 2 & 4 & 0 & 1 & 4 \\ -2 & -3 & 0 & 5 & -4 \\ 0 & 1 & 2 & 1 & 2 \end{array} \right]$$

$$\bullet E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$\bullet P_{23} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, P_{23} E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$\bullet E_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2 & 0 & 1 \end{bmatrix}, E_2 P_{23} E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -2 & 5 \\ 0 & 0 & 0 & 0 \end{bmatrix} = U$$

~~-1~~

Is  $A = LU$  always possible?

$$\left[ \begin{array}{cccc|c} 1 & 2 & 1 & -2 & 3 \\ 2 & 4 & 0 & 1 & 4 \\ -2 & -3 & 0 & 5 & -4 \\ 0 & 1 & 2 & 1 & 2 \end{array} \right]$$

$$\bullet E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$\bullet P_{23} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, P_{23} E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$\bullet E_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2 & 0 & 1 \end{bmatrix}, E_2 P_{23} E_1 A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = U$$

$$\bullet E_2 P_{23} E_1 A = U.$$

$$A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 2 & 4 & 0 & 1 \\ -2 & -3 & 0 & 5 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$E_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2 & 0 & 1 \end{bmatrix}, P_{23} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$P_{23}A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ -2 & -3 & 0 & 5 \\ 2 & 4 & 0 & 1 \\ 0 & 1 & 2 & 1 \end{bmatrix} \quad \bar{E}_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ -2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad \bar{E}_1 P_{23}A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -2 & 5 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$\bar{E}_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{bmatrix}, \quad \bar{E}_2 \bar{E}_1 P_{23}A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -2 & 5 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \bar{U}$$

$$A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 2 & 4 & 0 & 1 \\ -2 & -3 & 0 & 5 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$E_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2 & 0 & 1 \end{bmatrix}, P_{23} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\overline{E}_2 \overline{E}_1 P_{23} A = \overline{U}$$

$$P_{23} A = (\overline{E}_1)^{-1} (\overline{E}_2)^{-1} \overline{U}$$

$$\boxed{P_{23} A = \overline{L} \overline{U}}$$

$$A = \begin{bmatrix} 1 & 2 & 1 & -2 \\ 2 & 4 & 0 & 1 \\ -2 & -3 & 0 & 5 \\ 0 & 1 & 2 & 1 \end{bmatrix}$$

$$E_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2 & 0 & 1 \end{bmatrix}, P_{23} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\bar{E}_3 P_{14} \bar{E}_2 P_{23} \bar{E}_1 A = U$$

$$E'_3 E'_2 E'_1 (P_{14} \cdot P_{23}) A = U$$

$$\left( \begin{matrix} P_{1 \leftrightarrow 4} \\ 2 \leftrightarrow 3 \end{matrix} \right) A = \underbrace{(E'_1)^{-1} (E'_2)^{-1} (E'_3)^{-1}}_L U$$

# Why bother with $LU$ decomposition?

- Solve  $Ax = b$ .

## Why bother with $LU$ decomposition?

- Solve  $Ax = b$ .
- Given the  $LU$  decomposition of  $A$ , solve:  $LUx = b$ .



# Why bother with $LU$ decomposition?

- Solve  $Ax = b$ .
- Given the  $LU$  decomposition of  $A$ , solve:  $LUx = b$ .
- Letting  $y := Ux$  *reduces* the original problem to two *simpler* linear systems:

# Why bother with $LU$ decomposition?

- Solve  $Ax = b$ .
- Given the  $LU$  decomposition of  $A$ , solve:  $LUx = b$ .
- Letting  $y := Ux$  *reduces* the original problem to two *simpler* linear systems:
  - ▶ Solve for  $y$  in  $Ly = b$ .  $\sim O(n^2)$
  - ▶ Solve for  $x$  in  $Ux = y$ .  $\sim O(n^2)$

# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .

# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .
- $Ax = x_1 \cdot a_{*1} + x_2 \cdot a_{*2} + \dots + x_n \cdot a_{*n} = b \in \mathbb{R}^m$ .

# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .
- $Ax = x_1 \cdot a_{*1} + x_2 \cdot a_{*2} + \dots + x_n \cdot a_{*n} = b \in \mathbb{R}^m$ .
- When does a solution to  $Ax = b$  exist?

# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .
- $Ax = x_1 \cdot a_{*1} + x_2 \cdot a_{*2} + \dots + x_n \cdot a_{*n} = b \in \mathbb{R}^m$ .
- When does a solution to  $Ax = b$  exist?
- ▶ **A solution to  $Ax = b$  exists if and only if  $b$  belongs to the set of all possible linear combinations of columns of  $A$ .**

# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .
- $Ax = x_1 \cdot a_{*1} + x_2 \cdot a_{*2} + \dots + x_n \cdot a_{*n} = b \in \mathbb{R}^m$ .
- When does a solution to  $Ax = b$  exist?
  - ▶ **A solution to  $Ax = b$  exists if and only if  $b$  belongs to the set of all possible linear combinations of columns of  $A$ .**
  - ▶ **Column Space:** The set of all possible linear combinations of columns of  $A$  is called the Column space of matrix  $A$ , and is denoted by  $C(A)$ .

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .
- $Ax = x_1 \cdot a_{*1} + x_2 \cdot a_{*2} + \dots + x_n \cdot a_{*n} = b \in \mathbb{R}^m$ .
- When does a solution to  $Ax = b$  exist?
  - **A solution to  $Ax = b$  exists if and only if  $b$  belongs to the set of all possible linear combinations of columns of  $A$ .**
  - **Column Space:** The set of all possible linear combinations of columns of  $A$  is called the Column space of matrix  $A$ , and is denoted by  $C(A)$ .

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

- **Properties:**



# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .
- $Ax = x_1 \cdot a_{*1} + x_2 \cdot a_{*2} + \dots + x_n \cdot a_{*n} = b \in \mathbb{R}^m$ .
- When does a solution to  $Ax = b$  exist?
  - **A solution to  $Ax = b$  exists if and only if  $b$  belongs to the set of all possible linear combinations of columns of  $A$ .**
  - **Column Space:** The set of all possible linear combinations of columns of  $A$  is called the Column space of matrix  $A$ , and is denoted by  $C(A)$ .

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

► **Properties:**

► Let  $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$ .

# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .
- $Ax = x_1 \cdot a_{*1} + x_2 \cdot a_{*2} + \dots + x_n \cdot a_{*n} = b \in \mathbb{R}^m$ .
- When does a solution to  $Ax = b$  exist?
  - ▶ **A solution to  $Ax = b$  exists if and only if  $b$  belongs to the set of all possible linear combinations of columns of  $A$ .**
  - ▶ **Column Space:** The set of all possible linear combinations of columns of  $A$  is called the Column space of matrix  $A$ , and is denoted by  $C(A)$ .

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

▶ **Properties:**

▶ Let  $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$ . Does  $\mathbf{0}_m \in C(A)$ ?

# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .
- $Ax = x_1 \cdot a_{*1} + x_2 \cdot a_{*2} + \dots + x_n \cdot a_{*n} = b \in \mathbb{R}^m$ .
- When does a solution to  $Ax = b$  exist?
  - **A solution to  $Ax = b$  exists if and only if  $b$  belongs to the set of all possible linear combinations of columns of  $A$ .**
  - **Column Space:** The set of all possible linear combinations of columns of  $A$  is called the Column space of matrix  $A$ , and is denoted by  $C(A)$ .

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

► **Properties:**

► Let  $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$ .  $\mathbf{0}_m \in C(A)$  for any matrix  $A$

# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .
- $Ax = x_1 \cdot a_{*1} + x_2 \cdot a_{*2} + \dots + x_n \cdot a_{*n} = b \in \mathbb{R}^m$ .
- When does a solution to  $Ax = b$  exist?
  - ▶ **A solution to  $Ax = b$  exists if and only if  $b$  belongs to the set of all possible linear combinations of columns of  $A$ .**
  - ▶ **Column Space:** The set of all possible linear combinations of columns of  $A$  is called the Column space of matrix  $A$ , and is denoted by  $C(A)$ .

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

▶ **Properties:**

- ▶ Let  $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$ .  $\mathbf{0}_m \in C(A)$  for any matrix  $A$
- ▶ If  $b_1, b_2 \in C(A)$ ,  $\forall p, q \in \mathbb{R}$ ,  $p \cdot b_1 + q \cdot b_2$

# General Picture

- Let  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ . Solve  $Ax = b$ .
- $Ax = x_1 \cdot a_{*1} + x_2 \cdot a_{*2} + \dots + x_n \cdot a_{*n} = b \in \mathbb{R}^m$ .
- When does a solution to  $Ax = b$  exist?
  - ▶ **A solution to  $Ax = b$  exists if and only if  $b$  belongs to the set of all possible linear combinations of columns of  $A$ .**
  - ▶ **Column Space:** The set of all possible linear combinations of columns of  $A$  is called the Column space of matrix  $A$ , and is denoted by  $C(A)$ .

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

▶ **Properties:**

- ▶ Let  $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$ .  $\mathbf{0}_m \in C(A)$  for any matrix  $A$
- ▶ If  $b_1, b_2 \in C(A)$ ,  $\forall p, q \in \mathbb{R}$ ,  $p \cdot b_1 + q \cdot b_2 \in C(A)$ .

# What about multiple solutions?

- Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .

## What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) =$

## What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .



## What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  
 $\forall k \in \mathbb{R}, A(x + k \cdot z) =$

## What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  
 $\forall k \in \mathbb{R}, A(x + k \cdot z) = b \Rightarrow$  Multiple Solutions!

# What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  $\forall k \in \mathbb{R}, A(x + k \cdot z) = b \Rightarrow$  Multiple Solutions!
- ▶ **Nullspace:** For a matrix  $A \in \mathbb{R}^{m \times n}$ , the *Nullspace* is the set of vectors that get mapped to  $\mathbf{0}_m$ , and is denoted by  $N(A)$ .

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

# What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  $\forall k \in \mathbb{R}, A(x + k \cdot z) = b \Rightarrow$  Multiple Solutions!
- ▶ **Nullspace:** For a matrix  $A \in \mathbb{R}^{m \times n}$ , the *Nullspace* is the set of vectors that get mapped to  $\mathbf{0}_m$ , and is denoted by  $N(A)$ .

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶  $Ax = \mathbf{0}_m$  are also called **Homogeneous equations**.

# What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  $\forall k \in \mathbb{R}, A(x + k \cdot z) = b \Rightarrow$  Multiple Solutions!
- ▶ **Nullspace:** For a matrix  $A \in \mathbb{R}^{m \times n}$ , the *Nullspace* is the set of vectors that get mapped to  $\mathbf{0}_m$ , and is denoted by  $N(A)$ .

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶  $Ax = \mathbf{0}_m$  are also called **Homogeneous equations**.
- ▶ **Properties:**

# What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  $\forall k \in \mathbb{R}, A(x + k \cdot z) = b \Rightarrow$  Multiple Solutions!
- ▶ **Nullspace:** For a matrix  $A \in \mathbb{R}^{m \times n}$ , the *Nullspace* is the set of vectors that get mapped to  $\mathbf{0}_m$ , and is denoted by  $N(A)$ .

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶  $Ax = \mathbf{0}_m$  are also called **Homogeneous equations**.
- ▶ **Properties:**
- ▶  $\mathbf{0}_n$

# What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  $\forall k \in \mathbb{R}, A(x + k \cdot z) = b \Rightarrow$  Multiple Solutions!
- ▶ **Nullspace:** For a matrix  $A \in \mathbb{R}^{m \times n}$ , the *Nullspace* is the set of vectors that get mapped to  $\mathbf{0}_m$ , and is denoted by  $N(A)$ .

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶  $Ax = \mathbf{0}_m$  are also called **Homogeneous equations**.
- ▶ **Properties:**
- ▶  $\mathbf{0}_n \in N(A)$ .

# What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  $\forall k \in \mathbb{R}, A(x + k \cdot z) = b \Rightarrow$  Multiple Solutions!
- ▶ **Nullspace:** For a matrix  $A \in \mathbb{R}^{m \times n}$ , the *Nullspace* is the set of vectors that get mapped to  $\mathbf{0}_m$ , and is denoted by  $N(A)$ .

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶  $Ax = \mathbf{0}_m$  are also called **Homogeneous equations**.
- ▶ **Properties:**
- ▶  $\mathbf{0}_n \in N(A)$ .
- ▶ If  $x, y \in N(A)$ ,  $\forall p, q \in \mathbb{R}, p \cdot x + q \cdot y$



# What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  $\forall k \in \mathbb{R}, A(x + k \cdot z) = b \Rightarrow$  Multiple Solutions!
- ▶ **Nullspace:** For a matrix  $A \in \mathbb{R}^{m \times n}$ , the *Nullspace* is the set of vectors that get mapped to  $\mathbf{0}_m$ , and is denoted by  $N(A)$ .

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶  $Ax = \mathbf{0}_m$  are also called **Homogeneous equations**.
- ▶ **Properties:**
- ▶  $\mathbf{0}_n \in N(A)$ .
- ▶ If  $x, y \in N(A)$ ,  $\forall p, q \in \mathbb{R}, p \cdot x + q \cdot y \in N(A)$ .

# What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  $\forall k \in \mathbb{R}, A(x + k \cdot z) = b \Rightarrow$  Multiple Solutions!
- ▶ **Nullspace:** For a matrix  $A \in \mathbb{R}^{m \times n}$ , the *Nullspace* is the set of vectors that get mapped to  $\mathbf{0}_m$ , and is denoted by  $N(A)$ .

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶  $Ax = \mathbf{0}_m$  are also called **Homogeneous equations**.
- ▶ **Properties:**
- ▶  $\mathbf{0}_n \in N(A)$ .
- ▶ If  $x, y \in N(A)$ ,  $\forall p, q \in \mathbb{R}, p \cdot x + q \cdot y \in N(A)$ .
- **Summary:** If  $\exists z \in N(A), z \neq \mathbf{0}_n$ , then  $Ax = b$  will have

# What about multiple solutions?

- ▶ Let  $Ax = b$  and  $Ay = b$ , with  $x \neq y$ .
- ▶ Then,  $A(x - y) = \mathbf{0}_m$ .
- ▶ Similarly, let  $z \in \mathbb{R}^n$  be such that  $Az = \mathbf{0}_m$ . Then, if  $Ax = b$ ,  $\forall k \in \mathbb{R}, A(x + k \cdot z) = b \Rightarrow$  Multiple Solutions!
- ▶ **Nullspace:** For a matrix  $A \in \mathbb{R}^{m \times n}$ , the *Nullspace* is the set of vectors that get mapped to  $\mathbf{0}_m$ , and is denoted by  $N(A)$ .

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶  $Ax = \mathbf{0}_m$  are also called **Homogeneous equations**.
- ▶ **Properties:**
- ▶  $\mathbf{0}_n \in N(A)$ .
- ▶ If  $x, y \in N(A)$ ,  $\forall p, q \in \mathbb{R}, p \cdot x + q \cdot y \in N(A)$ .
- **Summary:** If  $\exists z \in N(A), z \neq \mathbf{0}_n$ , then  $Ax = b$  will have infinitely many solutions, if one exists!

- If there are multiple solutions to  $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$  such that  $Az = \mathbf{0}_m$ .

- If there are multiple solutions to  $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$  such that  $Az = \mathbf{0}_m$ .
- Re-writing, we get  $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$ , with at least one non-zero entry in  $z$ , say  $z_k$ .

- If there are multiple solutions to  $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$  such that  $Az = \mathbf{0}_m$ .
- Re-writing, we get  $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$ , with at least one non-zero entry in  $z$ , say  $z_k$ .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{-z_i}{z_k} a_{*i}$ . Thus we can write column  $k$  as a *linear combination* of other columns.

- If there are multiple solutions to  $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$  such that  $Az = \mathbf{0}_m$ .
- Re-writing, we get  $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$ , with at least one non-zero entry in  $z$ , say  $z_k$ .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{-z_i}{z_k} a_{*i}$ . Thus we can write column  $k$  as a *linear combination* of other columns. We say that the vectors  $\{a_{*i}, i = 1, \dots, n\}$  is a **linearly dependent** set of vectors.

- If there are multiple solutions to  $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$  such that  $Az = \mathbf{0}_m$ .
- Re-writing, we get  $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$ , with at least one non-zero entry in  $z$ , say  $z_k$ .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{-z_i}{z_k} a_{*i}$ . Thus we can write column  $k$  as a *linear combination* of other columns. We say that the vectors  $\{a_{*i}, i = 1, \dots, n\}$  is a **linearly dependent** set of vectors.
- **Linear Independent set of vectors:** A set of vectors  $\{a_{*i}, i = 1, \dots, n\}$  is said to be linearly independent if none of the vectors can be written as a linear combination of the remaining vectors.



- If there are multiple solutions to  $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$  such that  $Az = \mathbf{0}_m$ .
- Re-writing, we get  $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$ , with at least one non-zero entry in  $z$ , say  $z_k$ .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{-z_i}{z_k} a_{*i}$ . Thus we can write column  $k$  as a *linear combination* of other columns. We say that the vectors  $\{a_{*i}, i = 1, \dots, n\}$  is a **linearly dependent** set of vectors.
- **Linear Independent set of vectors:** A set of vectors  $\{a_{*i}, i = 1, \dots, n\}$  is said to be linearly independent if none of the vectors can be written as a linear combination of the remaining vectors.
- A set of vectors  $\{a_{*i}, i = 1, \dots, n\}$  is said to be linearly independent if

$$\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m \Rightarrow$$

- If there are multiple solutions to  $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$  such that  $Az = \mathbf{0}_m$ .
- Re-writing, we get  $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$ , with at least one non-zero entry in  $z$ , say  $z_k$ .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{-z_i}{z_k} a_{*i}$ . Thus we can write column  $k$  as a *linear combination* of other columns. We say that the vectors  $\{a_{*i}, i = 1, \dots, n\}$  is a **linearly dependent** set of vectors.
- **Linear Independent set of vectors:** A set of vectors  $\{a_{*i}, i = 1, \dots, n\}$  is said to be linearly independent if none of the vectors can be written as a linear combination of the remaining vectors.
- A set of vectors  $\{a_{*i}, i = 1, \dots, n\}$  is said to be linearly independent if

$$\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m \Rightarrow z_1 = z_2 = \dots = z_n = 0$$

- If there are multiple solutions to  $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$  such that  $Az = \mathbf{0}_m$ .
- Re-writing, we get  $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$ , with at least one non-zero entry in  $z$ , say  $z_k$ .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{-z_i}{z_k} a_{*i}$ . Thus we can write column  $k$  as a *linear combination* of other columns. We say that the vectors  $\{a_{*i}, i = 1, \dots, n\}$  is a **linearly dependent** set of vectors.
- **Linear Independent set of vectors:** A set of vectors  $\{a_{*i}, i = 1, \dots, n\}$  is said to be linearly independent if none of the vectors can be written as a linear combination of the remaining vectors.
- A set of vectors  $\{a_{*i}, i = 1, \dots, n\}$  is said to be linearly independent if

$$\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m \Rightarrow z_1 = z_2 = \dots = z_n = 0$$

- **Column Rank of a Matrix:** The number of linearly independent columns of a matrix is called the Column Rank.

- If there are multiple solutions to  $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$  such that  $Az = \mathbf{0}_m$ .
- Re-writing, we get  $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$ , with at least one non-zero entry in  $z$ , say  $z_k$ .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{-z_i}{z_k} a_{*i}$ . Thus we can write column  $k$  as a *linear combination* of other columns. We say that the vectors  $\{a_{*i}, i = 1, \dots, n\}$  is a **linearly dependent** set of vectors.
- **Linear Independent set of vectors:** A set of vectors  $\{a_{*i}, i = 1, \dots, n\}$  is said to be linearly independent if none of the vectors can be written as a linear combination of the remaining vectors.
- A set of vectors  $\{a_{*i}, i = 1, \dots, n\}$  is said to be linearly independent if

$$\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m \Rightarrow z_1 = z_2 = \dots = z_n = 0$$

- **Column Rank of a Matrix:** The number of linearly independent columns of a matrix is called the Column Rank.

- **Matrix Transpose:** For  $A \in \mathbb{R}^{m \times n}$  given by

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix},$$

the matrix

$$\begin{bmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & \dots & a_{m2} \\ \vdots & \vdots & \vdots & \vdots \\ \underbrace{a_{1n}}_{a_{1*}} & \underbrace{a_{2n}}_{a_{2*}} & \dots & \underbrace{a_{mn}}_{a_{m*}} \end{bmatrix} \in \mathbb{R}^{n \times m},$$

is called the *transpose of A*, and is denoted by  $A^T$ .

- **Matrix Transpose:** For  $A \in \mathbb{R}^{m \times n}$  given by

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix},$$

the matrix

$$\begin{bmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & \dots & a_{m2} \\ \vdots & \vdots & \vdots & \vdots \\ \underbrace{a_{1n}}_{a_{1*}} & \underbrace{a_{2n}}_{a_{2*}} & \dots & \underbrace{a_{mn}}_{a_{m*}} \end{bmatrix} \in \mathbb{R}^{n \times m},$$

is called the *transpose* of  $A$ , and is denoted by  $A^T$ .

- **Beware of the notation:**  $a_{i*}$  denotes the  $i^{th}$  row of  $A$  written as a column matrix.

- All linear combination of columns gives the column space, how about linear combination of rows?

- All linear combination of columns gives the column space, how about linear combination of rows?
- **Rowspace:** The set of all linear combinations of rows of the matrix is called the Row space.



- All linear combination of columns gives the column space, how about linear combination of rows?
- **Rowspace:** The set of all linear combinations of rows of the matrix is called the Row space.
- Since rows of  $A$  are columns of  $A^T$ , notation for row space is  $C(A^T)$ , and can be written as

$$C(A^T) = \{A^T y \mid \forall y \in \mathbb{R}^m\} \subseteq \mathbb{R}^n$$

- All linear combination of columns gives the column space, how about linear combination of rows?
- **Rowspace:** The set of all linear combinations of rows of the matrix is called the Row space.
- Since rows of  $A$  are columns of  $A^T$ , notation for row space is  $C(A^T)$ , and can be written as

$$C(A^T) = \{A^T y \mid \forall y \in \mathbb{R}^m\} \subseteq \mathbb{R}^n$$

- **Properties:**

- All linear combination of columns gives the column space, how about linear combination of rows?
- **Rowspace:** The set of all linear combinations of rows of the matrix is called the Row space.
- Since rows of  $A$  are columns of  $A^T$ , notation for row space is  $C(A^T)$ , and can be written as

$$C(A^T) = \{A^T y \mid \forall y \in \mathbb{R}^m\} \subseteq \mathbb{R}^n$$

- **Properties:**

- ▶  $\mathbf{0}_n \in C(A^T)$
- ▶ If  $x_1, x_2 \in C(A^T)$ , then  $\forall p, q \in \mathbb{R}, p \cdot x_1 + q \cdot x_2 \in C(A^T)$ .

- All linear combination of columns gives the column space, how about linear combination of rows?
- **Rowspace:** The set of all linear combinations of rows of the matrix is called the Row space.
- Since rows of  $A$  are columns of  $A^T$ , notation for row space is  $C(A^T)$ , and can be written as

$$C(A^T) = \{A^T y \mid \forall y \in \mathbb{R}^m\} \subseteq \mathbb{R}^n$$

- **Properties:**

- ▶  $\mathbf{0}_n \in C(A^T)$
- ▶ If  $x_1, x_2 \in C(A^T)$ , then  $\forall p, q \in \mathbb{R}, p \cdot x_1 + q \cdot x_2 \in C(A^T)$ .
- **Row Rank of a Matrix:** The number of linearly independent rows of a matrix is called the Column Rank.

- **Left Nullspace:** The set of solutions in  $\mathbb{R}^m$  to  $A^T y = \mathbf{0}_n$  is called the *Left Nullspace* of matrix  $A$ .

- **Left Nullspace:** The set of solutions in  $\mathbb{R}^m$  to  $A^T y = \mathbf{0}_n$  is called the *Left Nullspace* of matrix  $A$ .
- Since this is the same as Nullspace of the matrix  $A^T$ , left nullspace is denoted by  $N(A^T)$ .

$$N(A^T) := \{y \in \mathbb{R}^m \mid A^T y = \mathbf{0}_n\} \subseteq \mathbb{R}^m$$

- **Properties:**

- ▶  $\mathbf{0}_m \in N(A^T)$
- ▶ If  $y_1, y_2 \in N(A^T)$ , then  $\forall p, q \in \mathbb{R}, p \cdot y_1 + q \cdot y_2 \in N(A^T)$ .

- **Left Nullspace:** The set of solutions in  $\mathbb{R}^m$  to  $A^T y = \mathbf{0}_n$  is called the *Left Nullspace* of matrix  $A$ .
- Since this is the same as Nullspace of the matrix  $A^T$ , left nullspace is denoted by  $N(A^T)$ .

$$N(A^T) := \{y \in \mathbb{R}^m \mid A^T y = \mathbf{0}_n\} \subseteq \mathbb{R}^m$$

- **Properties:**

- ▶  $\mathbf{0}_m \in N(A^T)$
- ▶ If  $y_1, y_2 \in N(A^T)$ , then  $\forall p, q \in \mathbb{R}, p \cdot y_1 + q \cdot y_2 \in N(A^T)$ .

- **The Column Space, Row Space, Null Space and Left Nullspace are called the Four Fundamental Subspaces associated with a matrix.**