

SC223 - Linear Algebra

Aditya Tatu

Lecture 37



November 8, 2023

Summary of Lecture 36

- Norm: For a vector space $(V, +, \cdot)$ over \mathbb{R} or \mathbb{C} , **norm** is a function $\|\cdot\| : V \rightarrow \mathbb{R}$ that satisfies:
 - ▶ Positive definiteness: $\forall x \in V, \|x\| \geq 0, \|x\| = 0 \Leftrightarrow x = \theta$
 - ▶ Absolute Homogeneity: $\forall x \in V, \forall a \in \mathbb{F}, \|a \cdot x\| = |a| \cdot \|x\|$
 - ▶ Triangular Inequality: $\forall x, y \in V, \|x + y\| \leq \|x\| + \|y\|$.
- A vector space $(V, +, \cdot)$ with a norm $\|\cdot\|$ is called a **Normed Vector space** (NVS).

Inner Product

● **Definition:** (Inner Product) Given a vector space $(V, +, \cdot)$ over \mathbb{F} (either \mathbb{R} or \mathbb{C}), an **inner product** is any mapping $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{F}$ such that it satisfies the following properties:

1. Positive definite: $\forall x \in V, \langle x, x \rangle \geq 0, = 0 \Leftrightarrow x = \theta$.

Inner Product

● **Definition:** (Inner Product) Given a vector space $(V, +, \cdot)$ over \mathbb{F} (either \mathbb{R} or \mathbb{C}), an **inner product** is any mapping $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{F}$ such that it satisfies the following properties:

1. Positive definite: $\forall x \in V, \langle x, x \rangle \geq 0, = 0 \Leftrightarrow x = \theta$.

2. Linear in first argument:

$$\forall x, y, z \in V, \forall a, b \in \mathbb{F}, \langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$$

Inner Product

● **Definition:** (Inner Product) Given a vector space $(V, +, \cdot)$ over \mathbb{F} (either \mathbb{R} or \mathbb{C}), an **inner product** is any mapping $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{F}$ such that it satisfies the following properties:

1. Positive definite: $\forall x \in V, \langle x, x \rangle \geq 0, = 0 \Leftrightarrow x = \theta$.
2. Linear in first argument:
 $\forall x, y, z \in V, \forall a, b \in \mathbb{F}, \langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$
3. Conjugate symmetry: $\langle y, x \rangle = \overline{\langle x, y \rangle}$

$$\begin{aligned}\langle x, ay + bz \rangle &= \overline{\langle ay + bz, x \rangle} \\ &= \overline{a\langle y, x \rangle + b\langle z, x \rangle} \\ &= \overline{a} \overline{\langle y, x \rangle} + \overline{b} \overline{\langle z, x \rangle} \\ &= \overline{a} \langle x, y \rangle + \overline{b} \langle x, z \rangle\end{aligned}$$

Inner Product

● **Definition:** (Inner Product) Given a vector space $(V, +, \cdot)$ over \mathbb{F} (either \mathbb{R} or \mathbb{C}), an **inner product** is any mapping $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{F}$ such that it satisfies the following properties:

1. Positive definite: $\forall x \in V, \langle x, x \rangle \geq 0, = 0 \Leftrightarrow x = \theta$.
2. Linear in first argument:
 $\forall x, y, z \in V, \forall a, b \in \mathbb{F}, \langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$
3. Conjugate symmetry: $\langle y, x \rangle = \overline{\langle x, y \rangle}$

● **Definition:** (Inner Product Space) A vector space V with an inner product is called an **Inner Product space**(IPS) and is denoted by $(V, \langle \cdot, \cdot \rangle)$.

Example 1: $V = \mathbb{R}^n$, $\langle x, y \rangle := \sum_{i=1}^n x_i y_i$

$$\begin{aligned}\langle ax + by, z \rangle &= \sum_{i=1}^n (ax_i + by_i) z_i \\ &= a \sum_{i=1}^n x_i z_i + b \sum_{i=1}^n y_i z_i \\ &= a \langle x, z \rangle + b \langle y, z \rangle\end{aligned}$$

Examples of IPS

- \mathbb{R}^n : Euclidean inner product - $x, y \in \mathbb{R}^n$, $\langle x, y \rangle = \sum_{i=1}^n x_i y_i = x^T y$, where x and y are written as column vectors.

Examples of IPS

- \mathbb{R}^n : Euclidean inner product - $x, y \in \mathbb{R}^n$, $\langle x, y \rangle = \sum_{i=1}^n x_i y_i = x^T y$, where x and y are written as column vectors.
- \mathbb{C}^n : Euclidean inner product - $x, y \in \mathbb{C}^n$, $\langle x, y \rangle = \sum_{i=1}^n x_i \bar{y}_i = y^* x$, where x and y are written as column vectors, and y^* denotes the *conjugate transpose* of y .

Examples of IPS

- \mathbb{R}^n : Euclidean inner product - $x, y \in \mathbb{R}^n$, $\langle x, y \rangle = \sum_{i=1}^n x_i y_i = x^T y$, where x and y are written as column vectors.
- \mathbb{C}^n : Euclidean inner product - $x, y \in \mathbb{C}^n$, $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i} = y^* x$, where x and y are written as column vectors, and y^* denotes the *conjugate transpose* of y .
- $\mathcal{P}_n([0, 1])$: L_2 inner product -
 $\forall p, q \in \mathcal{P}_n([0, 1]), \langle p, q \rangle = \int_0^1 p(t) q(t) dt.$

$$\langle p, q \rangle := \int_0^1 p(t) \overline{q(t)} dt.$$

Examples of IPS

● \mathbb{R}^n : Euclidean inner product - $x, y \in \mathbb{R}^n$, $\langle x, y \rangle = \sum_{i=1}^n x_i y_i = x^T y$, where x and y are written as column vectors.

● \mathbb{C}^n : Euclidean inner product - $x, y \in \mathbb{C}^n$, $\langle x, y \rangle = \sum_{i=1}^n x_i \bar{y}_i = y^* x$, where x and y are written as column vectors, and y^* denotes the *conjugate transpose* of y .

● $\mathcal{P}_n([0, 1])$: L_2 inner product -

$\forall p, q \in \mathcal{P}_n([0, 1])$, $\langle p, q \rangle = \int_0^1 p(t)q(t) dt$.

● Let $G \in \mathbb{R}^{n \times n}$ be such that $G = G^T$ and $x^T G x > 0, \forall x \in \mathbb{R}^n, \neq \mathbf{0}_n$. Such a matrix G is said to be *Symmetric Positive-Definite* (SPD). Then, on $\mathbb{R}^n, \forall x, y \in \mathbb{R}^n$, $\langle x, y \rangle = x^T G y$ is a valid inner product.

$$\forall x, y, \quad \langle x, y \rangle = x^T G y = \langle y, x \rangle = y^T G x.$$
$$\Rightarrow \boxed{G = G^T}$$
$$x^T G y = x^T G^T y.$$

$$\langle x, x \rangle = x^T G x \geq 0 \quad \& \text{ if } x \neq \vec{0}, \quad x^T G x > 0$$

Examples of IPS

- \mathbb{R}^n : Euclidean inner product - $x, y \in \mathbb{R}^n$, $\langle x, y \rangle = \sum_{i=1}^n x_i y_i = x^T y$, where x and y are written as column vectors.
- \mathbb{C}^n : Euclidean inner product - $x, y \in \mathbb{C}^n$, $\langle x, y \rangle = \sum_{i=1}^n x_i \bar{y}_i = y^* x$, where x and y are written as column vectors, and y^* denotes the *conjugate transpose* of y .
- $\mathcal{P}_n([0, 1])$: L_2 inner product -
 $\forall p, q \in \mathcal{P}_n([0, 1]), \langle p, q \rangle = \int_0^1 p(t)q(t) dt$.
- Let $G \in \mathbb{R}^{n \times n}$ be such that $G = G^T$, and $x^T G x > 0, \forall x \in \mathbb{R}^n, \neq \mathbf{0}_n$. Such a matrix G is said to be *Symmetric Positive-Definite* (SPD). Then, on \mathbb{R}^n , $\forall x, y \in \mathbb{R}^n$, $\langle x, y \rangle = x^T G y$ is a valid inner product.




Figure: Orthogonal vectors with inner product: (left) $\langle x, y \rangle = x^T y$, (right)

$$\langle x, y \rangle = x^T G y, G = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$

Definitions

$$|\langle x, y \rangle| \leq \sqrt{\langle x, x \rangle} \sqrt{\langle y, y \rangle}$$

- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, $\forall x \in V$, $\|x\| = \sqrt{\langle x, x \rangle}$ is a valid norm, called the **induced norm**.

$$\|x\| := \sqrt{\langle x, x \rangle}$$

$$\begin{aligned} \|a \cdot x\| &= \sqrt{\langle ax, ax \rangle} = \sqrt{a \langle x, ax \rangle} \\ &= \sqrt{a \cdot \bar{a} \langle x, x \rangle} = \sqrt{|a|^2 \langle x, x \rangle} \\ &= |a| \cdot \sqrt{\langle x, x \rangle} = |a| \cdot \|x\|. \end{aligned}$$

$$\begin{aligned} \|x+y\|^2 &= \langle x, x+y \rangle + \langle y, x+y \rangle \\ &= \langle x, x \rangle + \langle y, y \rangle + 2\operatorname{Re}\{\langle x, y \rangle\} \\ &\leq \|x\|^2 + \|y\|^2 + 2|\langle x, y \rangle| \end{aligned}$$

Definitions

- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, $\forall x \in V$, $\|x\| = \sqrt{\langle x, x \rangle}$ is a valid norm, called the **induced norm**.
- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, two vectors $x, y \in V$ are said to be **orthogonal** if $\langle x, y \rangle = 0$

Definitions

- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, $\forall x \in V$, $\|x\| = \sqrt{\langle x, x \rangle}$ is a valid norm, called the **induced norm**.
- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, two vectors $x, y \in V$ are said to be **orthogonal** if $\langle x, y \rangle = 0$, and are said to be **orthonormal** if $\langle x, y \rangle = 0$, $\|x\| = \|y\| = 1$.

Definitions

- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, $\forall x \in V$, $\|x\| = \sqrt{\langle x, x \rangle}$ is a valid norm, called the **induced norm**.
- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, two vectors $x, y \in V$ are said to be **orthogonal** if $\langle x, y \rangle = 0$, and are said to be **orthonormal** if $\langle x, y \rangle = 0, \|x\| = \|y\| = 1$.
- A set of vectors $\{v_1, \dots, v_n\}$ is said to be **orthogonal** if $\langle v_i, v_j \rangle = 0, \forall i \neq j$

Definitions

- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, $\forall x \in V$, $\|x\| = \sqrt{\langle x, x \rangle}$ is a valid norm, called the **induced norm**.
- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, two vectors $x, y \in V$ are said to be **orthogonal** if $\langle x, y \rangle = 0$, and are said to be **orthonormal** if $\langle x, y \rangle = 0$, $\|x\| = \|y\| = 1$.
- A set of vectors $\{v_1, \dots, v_n\}$ is said to be **orthogonal** if $\langle v_i, v_j \rangle = 0, \forall i \neq j$, and is said to be **orthonormal** if $\langle v_i, v_j \rangle = 0, \forall i \neq j$, $\|v_i\| = 1, \forall i$.

Definitions

- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, $\forall x \in V$, $\|x\| = \sqrt{\langle x, x \rangle}$ is a valid norm, called the **induced norm**.
- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, two vectors $x, y \in V$ are said to be **orthogonal** if $\langle x, y \rangle = 0$, and are said to be **orthonormal** if $\langle x, y \rangle = 0$, $\|x\| = \|y\| = 1$.
- A set of vectors $\{v_1, \dots, v_n\}$ is said to be **orthogonal** if $\langle v_i, v_j \rangle = 0, \forall i \neq j$, and is said to be **orthonormal** if $\langle v_i, v_j \rangle = 0, \forall i \neq j$, $\|v_i\| = 1, \forall i$.
- A set of orthonormal vectors that also forms a basis of the given vector space is called an **Orthonormal basis**.

Definitions

- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, $\forall x \in V$, $\|x\| = \sqrt{\langle x, x \rangle}$ is a valid norm, called the **induced norm**.
- Given an IPS $(V, \langle \cdot, \cdot \rangle)$, two vectors $x, y \in V$ are said to be **orthogonal** if $\langle x, y \rangle = 0$, and are said to be **orthonormal** if $\langle x, y \rangle = 0$, $\|x\| = \|y\| = 1$.
- A set of vectors $\{v_1, \dots, v_n\}$ is said to be **orthogonal** if $\langle v_i, v_j \rangle = 0, \forall i \neq j$, and is said to be **orthonormal** if $\langle v_i, v_j \rangle = 0, \forall i \neq j$, $\|v_i\| = 1, \forall i$.
- A set of orthonormal vectors that also forms a basis of the given vector space is called an **Orthonormal basis**.
- A matrix $A \in \mathbb{R}^{n \times n}$ or $\mathbb{C}^{n \times n}$ is said to be an **orthogonal matrix** if all its n columns are orthonormal, i.e., $A^* A = I$, where A^* denotes the conjugate transpose of A . In this case, $A^{-1} = A^*$.

————— END OF CLASS —————

Properties

- **Proposition 23** (Pythagoras Theorem): In an IPS $(V, \langle \cdot, \cdot \rangle)$, if $\langle x, y \rangle = 0$, $\|x + y\|^2 = \|x\|^2 + \|y\|^2$.

Properties

- **Proposition 23** (Pythagoras Theorem): In an IPS $(V, \langle \cdot, \cdot \rangle)$, if $\langle x, y \rangle = 0$, $\|x + y\|^2 = \|x\|^2 + \|y\|^2$.
- **Orthogonal Decomposition:** Let $x, y \neq \theta \in V$. Find $w \in V$ such that $x = a \cdot y + w$, with $a \in \mathbb{F}$, $\langle w, y \rangle = 0$.

Properties

- **Proposition 23** (Pythagoras Theorem): In an IPS $(V, \langle \cdot, \cdot \rangle)$, if $\langle x, y \rangle = 0$, $\|x + y\|^2 = \|x\|^2 + \|y\|^2$.
- **Orthogonal Decomposition:** Let $x, y \neq \theta \in V$. Find $w \in V$ such that $x = a \cdot y + w$, with $a \in \mathbb{F}$, $\langle w, y \rangle = 0$.

$$\langle w, y \rangle = 0$$

$$\langle x - a \cdot y, y \rangle = \langle x, y \rangle - a \langle y, y \rangle = 0$$

$$a = \frac{\langle x, y \rangle}{\langle y, y \rangle}$$

$$\text{Thus, } x = \frac{\langle x, y \rangle}{\langle y, y \rangle} \cdot y + \left(x - \frac{\langle x, y \rangle}{\langle y, y \rangle} \cdot y \right)$$

Properties

- **Proposition 24** (Cauchy-Schwartz inequality): In an IPS $(V, \langle \cdot, \cdot \rangle)$,
 $|\langle x, y \rangle| \leq \|x\| \|y\|$.

Properties

- **Proposition 24** (Cauchy-Schwartz inequality): In an IPS $(V, \langle \cdot, \cdot \rangle)$, $|\langle x, y \rangle| \leq \|x\| \|y\|$.
- **Proof:** If $x = \theta$, or $y = \theta$, both sides are equal to zero. So let us assume $x, y \neq \theta$.

Properties

● **Proposition 24** (Cauchy-Schwartz inequality): In an IPS $(V, \langle \cdot, \cdot \rangle)$,
 $|\langle x, y \rangle| \leq \|x\| \|y\|$.

● Proof: If $x = \theta$, or $y = \theta$, both sides are equal to zero. So let us assume $x, y \neq \theta$.

● From previous proposition,

$$x = \frac{\langle x, y \rangle}{\langle y, y \rangle} \cdot y + \left(x - \frac{\langle x, y \rangle}{\langle y, y \rangle} \cdot y \right) = \frac{\langle x, y \rangle}{\langle y, y \rangle} \cdot y + w, \text{ with } w \perp y.$$

Properties

● **Proposition 24** (Cauchy-Schwartz inequality): In an IPS $(V, \langle \cdot, \cdot \rangle)$,
 $|\langle x, y \rangle| \leq \|x\| \|y\|$.

● Proof: If $x = \theta$, or $y = \theta$, both sides are equal to zero. So let us assume $x, y \neq \theta$.

● From previous proposition,

$$x = \frac{\langle x, y \rangle}{\langle y, y \rangle} \cdot y + \left(x - \frac{\langle x, y \rangle}{\langle y, y \rangle} \cdot y \right) = \frac{\langle x, y \rangle}{\langle y, y \rangle} \cdot y + w, \text{ with } w \perp y.$$

Gram-Schmidt Procedure

- **Proposition 25** (Gram-Schmidt Procedure): Let $\{v_1, \dots, v_m\}$ be a list of linearly independent vectors. Then there exists a list of orthonormal vectors $\{e_1, \dots, e_m\}$ such that $\text{span}(\{v_1, \dots, v_j\}) = \text{span}(\{e_1, \dots, e_j\}), \forall j = 1, \dots, m$.

Gram-Schmidt Procedure

- **Proposition 25** (Gram-Schmidt Procedure): Let $\{v_1, \dots, v_m\}$ be a list of linearly independent vectors. Then there exists a list of orthonormal vectors $\{e_1, \dots, e_m\}$ such that $\text{span}(\{v_1, \dots, v_j\}) = \text{span}(\{e_1, \dots, e_j\}), \forall j = 1, \dots, m$.
- Let $e_1 = \frac{v_1}{\|v_1\|}$. Define $e_2 = \frac{v_2 - \langle v_2, e_1 \rangle e_1}{\|v_2 - \langle v_2, e_1 \rangle e_1\|}$.

Gram-Schmidt Procedure

- **Proposition 25** (Gram-Schmidt Procedure): Let $\{v_1, \dots, v_m\}$ be a list of linearly independent vectors. Then there exists a list of orthonormal vectors $\{e_1, \dots, e_m\}$ such that $\text{span}(\{v_1, \dots, v_j\}) = \text{span}(\{e_1, \dots, e_j\})$, $\forall j = 1, \dots, m$.
- Let $e_1 = \frac{v_1}{\|v_1\|}$. Define $e_2 = \frac{v_2 - \langle v_2, e_1 \rangle e_1}{\|v_2 - \langle v_2, e_1 \rangle e_1\|}$.
- Similarly, $e_3 = \frac{v_3 - (\langle v_3, e_1 \rangle e_1 + \langle v_3, e_2 \rangle e_2)}{\|v_3 - (\langle v_3, e_1 \rangle e_1 + \langle v_3, e_2 \rangle e_2)\|}$, and $e_k = \frac{v_k - \sum_{i=1}^{k-1} \langle v_k, e_i \rangle e_i}{\|v_k - \sum_{i=1}^{k-1} \langle v_k, e_i \rangle e_i\|}$.

Gram-Schmidt Procedure

- **Proposition 25** (Gram-Schmidt Procedure): Let $\{v_1, \dots, v_m\}$ be a list of linearly independent vectors. Then there exists a list of orthonormal vectors $\{e_1, \dots, e_m\}$ such that $\text{span}(\{v_1, \dots, v_j\}) = \text{span}(\{e_1, \dots, e_j\})$, $\forall j = 1, \dots, m$.
- Let $e_1 = \frac{v_1}{\|v_1\|}$. Define $e_2 = \frac{v_2 - \langle v_2, e_1 \rangle e_1}{\|v_2 - \langle v_2, e_1 \rangle e_1\|}$.
- Similarly, $e_3 = \frac{v_3 - (\langle v_3, e_1 \rangle e_1 + \langle v_3, e_2 \rangle e_2)}{\|v_3 - (\langle v_3, e_1 \rangle e_1 + \langle v_3, e_2 \rangle e_2)\|}$, and $e_k = \frac{v_k - \sum_{i=1}^{k-1} \langle v_k, e_i \rangle e_i}{\|v_k - \sum_{i=1}^{k-1} \langle v_k, e_i \rangle e_i\|}$.
- Observe that $\text{span}(\{v_1, \dots, v_j\}) = \text{span}(\{e_1, \dots, e_j\})$.

Gram-Schmidt Procedure

- **Proposition 25** (Gram-Schmidt Procedure): Let $\{v_1, \dots, v_m\}$ be a list of linearly independent vectors. Then there exists a list of orthonormal vectors $\{e_1, \dots, e_m\}$ such that $\text{span}(\{v_1, \dots, v_j\}) = \text{span}(\{e_1, \dots, e_j\})$, $\forall j = 1, \dots, m$.
- Let $e_1 = \frac{v_1}{\|v_1\|}$. Define $e_2 = \frac{v_2 - \langle v_2, e_1 \rangle e_1}{\|v_2 - \langle v_2, e_1 \rangle e_1\|}$.
- Similarly, $e_3 = \frac{v_3 - (\langle v_3, e_1 \rangle e_1 + \langle v_3, e_2 \rangle e_2)}{\|v_3 - (\langle v_3, e_1 \rangle e_1 + \langle v_3, e_2 \rangle e_2)\|}$, and $e_k = \frac{v_k - \sum_{i=1}^{k-1} \langle v_k, e_i \rangle e_i}{\|v_k - \sum_{i=1}^{k-1} \langle v_k, e_i \rangle e_i\|}$.
- Observe that $\text{span}(\{v_1, \dots, v_j\}) = \text{span}(\{e_1, \dots, e_j\})$.
- It is easy to see that $e_1 \perp e_2$. Assume that $\{e_1, \dots, e_j\}$ are orthonormal.

Gram-Schmidt Procedure

- **Proposition 25** (Gram-Schmidt Procedure): Let $\{v_1, \dots, v_m\}$ be a list of linearly independent vectors. Then there exists a list of orthonormal vectors $\{e_1, \dots, e_m\}$ such that $\text{span}(\{v_1, \dots, v_j\}) = \text{span}(\{e_1, \dots, e_j\})$, $\forall j = 1, \dots, m$.
- Let $e_1 = \frac{v_1}{\|v_1\|}$. Define $e_2 = \frac{v_2 - \langle v_2, e_1 \rangle e_1}{\|v_2 - \langle v_2, e_1 \rangle e_1\|}$.
- Similarly, $e_3 = \frac{v_3 - (\langle v_3, e_1 \rangle e_1 + \langle v_3, e_2 \rangle e_2)}{\|v_3 - (\langle v_3, e_1 \rangle e_1 + \langle v_3, e_2 \rangle e_2)\|}$, and $e_k = \frac{v_k - \sum_{i=1}^{k-1} \langle v_k, e_i \rangle e_i}{\|v_k - \sum_{i=1}^{k-1} \langle v_k, e_i \rangle e_i\|}$.
- Observe that $\text{span}(\{v_1, \dots, v_j\}) = \text{span}(\{e_1, \dots, e_j\})$.
- It is easy to see that $e_1 \perp e_2$. Assume that $\{e_1, \dots, e_j\}$ are orthonormal.
- Then $\forall l = 1, \dots, j$, with $e_{j+1}^\sim = v_{j+1} - \sum_{i=1}^j \langle v_{j+1}, e_i \rangle e_i$

$$\begin{aligned}\langle e_{j+1}, e_l \rangle &= \frac{1}{\|e_{j+1}^\sim\|} \left(\langle v_{j+1}, e_l \rangle - \sum_{i=1}^j \langle v_{j+1}, e_i \rangle \langle e_i, e_l \rangle \right) \\ &= \frac{1}{\|e_{j+1}^\sim\|} (\langle v_{j+1}, e_l \rangle - \langle v_{j+1}, e_l \rangle) = 0\end{aligned}$$

Orthogonal Complement

- Let V be a FD IPS and let U be a subset of V . The **Orthogonal Complement** of U is defined as

$$U^\perp = \{v \in V \mid \langle v, u \rangle = 0, \forall u \in U\}$$

Orthogonal Complement

- Let V be a FD IPS and let U be a subset of V . The **Orthogonal Complement** of U is defined as

$$U^\perp = \{v \in V \mid \langle v, u \rangle = 0, \forall u \in U\}$$

- **Proposition 26:** Irrespective of whether U is a subspace of V or not, U^\perp is a subspace.

Orthogonal Projection

- Let U be a subspace of FD IPS V , and $V = U \oplus U^\perp$.

Orthogonal Projection

- Let U be a subspace of FD IPS V , and $V = U \oplus U^\perp$.
- Define $P_U \in \mathcal{L}(V)$ as $\forall v \in V$, if $v = u + w, u \in U, w \in U^\perp$, $P_U(v) = u$.

Orthogonal Projection

- Let U be a subspace of FD IPS V , and $V = U \oplus U^\perp$.
- Define $P_U \in \mathcal{L}(V)$ as $\forall v \in V$, if $v = u + w, u \in U, w \in U^\perp$, $P_U(v) = u$.
- P_U is said to be the *Orthogonal Projection Operator on U* .

Orthogonal Projection

- Let U be a subspace of FD IPS V , and $V = U \oplus U^\perp$.
- Define $P_U \in \mathcal{L}(V)$ as $\forall v \in V$, if $v = u + w, u \in U, w \in U^\perp$, $P_U(v) = u$.
- P_U is said to be the *Orthogonal Projection Operator on U* .
- It has the following properties:
 1. $\text{Range}(P_U) =$

Orthogonal Projection

- Let U be a subspace of FD IPS V , and $V = U \oplus U^\perp$.
- Define $P_U \in \mathcal{L}(V)$ as $\forall v \in V$, if $v = u + w, u \in U, w \in U^\perp$, $P_U(v) = u$.
- P_U is said to be the *Orthogonal Projection Operator on U* .
- It has the following properties:
 1. $\text{Range}(P_U) = U$
 2. $\text{Null}(P_U) =$

Orthogonal Projection

- Let U be a subspace of FD IPS V , and $V = U \oplus U^\perp$.
- Define $P_U \in \mathcal{L}(V)$ as $\forall v \in V$, if $v = u + w, u \in U, w \in U^\perp$, $P_U(v) = u$.
- P_U is said to be the *Orthogonal Projection Operator on U* .
- It has the following properties:
 1. $\text{Range}(P_U) = U$
 2. $\text{Null}(P_U) = U^\perp$
 3. Idempotent: $(P_U)^2 =$

Orthogonal Projection

- Let U be a subspace of FD IPS V , and $V = U \oplus U^\perp$.
- Define $P_U \in \mathcal{L}(V)$ as $\forall v \in V$, if $v = u + w, u \in U, w \in U^\perp$, $P_U(v) = u$.
- P_U is said to be the *Orthogonal Projection Operator on U* .
- It has the following properties:
 1. $\text{Range}(P_U) = U$
 2. $\text{Null}(P_U) = U^\perp$
 3. Idempotent: $(P_U)^2 = P_U$
 4. (Conjugate) Symmetric: If $U = \mathbb{R}^n$ (or \mathbb{C}^n), $P_U^T =$

Orthogonal Projection

- Let U be a subspace of FD IPS V , and $V = U \oplus U^\perp$.
- Define $P_U \in \mathcal{L}(V)$ as $\forall v \in V$, if $v = u + w, u \in U, w \in U^\perp$, $P_U(v) = u$.
- P_U is said to be the *Orthogonal Projection Operator on U* .
- It has the following properties:
 1. $\text{Range}(P_U) = U$
 2. $\text{Null}(P_U) = U^\perp$
 3. Idempotent: $(P_U)^2 = P_U$
 4. (Conjugate) Symmetric: If $U = \mathbb{R}^n$ (or \mathbb{C}^n), $P_U^T = P_U$ ($P_U^* = P_U$).

Orthogonal Projection

- Let U be a subspace of FD IPS V , and $V = U \oplus U^\perp$.
- Define $P_U \in \mathcal{L}(V)$ as $\forall v \in V$, if $v = u + w, u \in U, w \in U^\perp$, $P_U(v) = u$.
- P_U is said to be the *Orthogonal Projection Operator on U* .
- It has the following properties:
 1. $\text{Range}(P_U) = U$
 2. $\text{Null}(P_U) = U^\perp$
 3. Idempotent: $(P_U)^2 = P_U$
 4. (Conjugate) Symmetric: If $U = \mathbb{R}^n$ (or \mathbb{C}^n), $P_U^T = P_U$ ($P_U^* = P_U$).
 5. $\forall v \in V, P_U(v) = \arg \min_{u \in U} \|u - v\|^2$.