

FORMULAS AND CONCEPTS YOU MIGHT FIND USEFUL

- **Sine waves and complex exponentials:**

$$\cos(\theta) = \frac{e^{j\theta} + e^{-j\theta}}{2}$$

and

$$\sin(\theta) = \frac{e^{j\theta} - e^{-j\theta}}{2j}$$

- **Geometric series:** For any $\alpha \neq 1$ (complex or real),

$$\sum_{n=0}^N \alpha^n = \frac{1 - \alpha^{N+1}}{1 - \alpha}.$$

If $|\alpha| < 1$, then

$$\sum_{n=0}^{\infty} \alpha^n = \frac{1}{1 - \alpha}.$$

- **LTI properties:** An LTI system with impulse response $h[n]$ is

- **memoryless** if $h[n] = c\delta[n]$ for some constant c .
- **causal** if $h[n] = 0$ for all $n < 0$.
- **stable** if $\sum_{n=-\infty}^{\infty} |h[n]| < \infty$.
- **invertible** if there is $g[n]$ such that $g[n] * h[n] = \delta[n]$.

- **DTFS:** For a signal with period N , and $\omega_0 = \frac{2\pi}{N}$

$$x[n] = \sum_{k=0}^{N-1} a_k e^{j\omega_0 kn}$$

where

$$a_k = \frac{1}{N} \sum_{n=0}^{N-1} x[n] e^{-j\omega_0 kn}.$$

- **DTFT:** For any signal,

$$x[n] = \frac{1}{2\pi} \int_0^{2\pi} X(e^{j\omega}) e^{j\omega n} d\omega$$

where

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}.$$

- **Properties of DTFT:**

- **Linearity:**

$$\begin{aligned} x[n] &\rightarrow X(e^{j\omega}) \\ y[n] &\rightarrow Y(e^{j\omega}) \end{aligned} \implies \alpha x[n] + \beta y[n] \rightarrow \alpha X(e^{j\omega}) + \beta Y(e^{j\omega})$$

- **Time reversal:**

$$x[n] \rightarrow X(e^{j\omega}) \implies x[-n] \rightarrow X(e^{-j\omega})$$

- **Time shifting:**

$$x[n] \rightarrow X(e^{j\omega}) \implies x[n - n_0] \rightarrow X(e^{j\omega}) e^{-j\omega n_0}$$

- **Frequency shifting:**

$$x[n] \rightarrow X(e^{j\omega}) \implies x[n] e^{j\omega_0 n} \rightarrow X(e^{j(\omega - \omega_0)})$$

- **Convolution:**

$$\begin{aligned} x[n] &\rightarrow X(e^{j\omega}) \\ y[n] &\rightarrow Y(e^{j\omega}) \end{aligned} \implies x[n] * y[n] \rightarrow X(e^{j\omega}) Y(e^{j\omega})$$

- **Multiplication:**

$$\begin{aligned} x[n] &\rightarrow X(e^{j\omega}) \\ y[n] &\rightarrow Y(e^{j\omega}) \end{aligned} \implies x[n] y[n] \rightarrow \frac{1}{2\pi} \left[X(e^{j\omega}) * Y(e^{j\omega}) \right]$$

- **Differentiation in frequency domain:**

$$x[n] \rightarrow X(e^{j\omega}) \implies nx[n] \rightarrow j \frac{dX(e^{j\omega})}{d\omega}$$

- **z-Transform:**

$$X(z) = \sum_{n=-\infty}^{\infty} x[n] z^{-n}.$$

Region of convergence (ROC) is the set of z for which the above sum converges.

- **Properties of the z-Transform:**

- If $X(z) = \frac{(z-z_0)(z-z_1)\cdots(z-z_M)}{(z-p_0)(z-p_1)\cdots(z-p_K)}$, then z_m are *zeros* and p_k are *poles*. In addition, there are $K-M$ zeros at ∞ if $K > M$, $M-K$ poles at ∞ if $M > K$.
- The ROC can only take one of three possible forms:
 - * $|z| < a$ when $x[n]$ is left-sided,
 - * $a < |z| < b$ when $x[n]$ is two-sided,
 - * $b < |z|$ when $x[n]$ is right-sided.

In addition, the ROC cannot contain any poles.

- An LTI system that has a transfer function $H(z)$ with ROC \mathcal{R} is
 - * **causal** if \mathcal{R} is of the form $b < |z|$ and $H(z)$ has no pole at $z = \infty$.
 - * **stable** if \mathcal{R} contains the unit circle $|z| = 1$.
- The DTFT exists (converges) if the ROC contains $|z| = 1$.
- **Linearity:** If $x[n] \rightarrow X(z)$ with ROC \mathcal{R}_1 and $y[n] \rightarrow Y(z)$ with ROC \mathcal{R}_2 , then

$$\alpha x[n] + \beta y[n] \rightarrow \alpha X(z) + \beta Y(z)$$

with an ROC containing $\mathcal{R}_1 \cap \mathcal{R}_2$. There might be a zero-pole cancellation resulting in a larger ROC.

- **Time reversal:** If $x[n] \rightarrow X(z)$ with ROC \mathcal{R} , then

$$x[-n] \rightarrow X(z^{-1})$$

with ROC = $\mathcal{R}^{-1} \triangleq \{z : z^{-1} \in \mathcal{R}\}$

- **Time shifting:** If $x[n] \rightarrow X(z)$ with ROC \mathcal{R} , then

$$x[n - n_0] \rightarrow X(z) z^{-n_0}$$

with ROC = \mathcal{R} (possibly excluding $z = 0$ or $z = \infty$).

- **Frequency shifting:** If $x[n] \rightarrow X(z)$ with ROC \mathcal{R} , then

$$x[n] z_0^n \rightarrow X(z/z_0)$$

with ROC = $|z_0| \mathcal{R} \triangleq \{z : z/z_0 \in \mathcal{R}\}$.

- **Convolution:** If $x[n] \rightarrow X(z)$ with ROC \mathcal{R}_1 and $y[n] \rightarrow Y(z)$ with ROC \mathcal{R}_2 , then

$$x[n] * y[n] \rightarrow X(z) Y(z)$$

with an ROC containing $\mathcal{R}_1 \cap \mathcal{R}_2$. There might be a zero-pole cancellation resulting in a larger ROC.

- **Differentiation in z-domain:** If $x[n] \rightarrow X(z)$ with ROC \mathcal{R} , then

$$nx[n] \rightarrow -z \frac{dX(z)}{dz}$$

with ROC = \mathcal{R} (possibly excluding $z = 0$ or $z = \infty$).

• **Some known signal/z-Transform pairs:**

- If $x[n] = a^n u[n]$, then

$$X(z) = \frac{1}{1 - az^{-1}}$$

with ROC = $\{z : |z| > a\}$.

- If $x[n] = -a^n u[-n - 1]$, then

$$X(z) = \frac{1}{1 - az^{-1}}$$

with ROC = $\{z : |z| < a\}$.

- If $x[n] = na^n u[n]$, then

$$X(z) = \frac{az^{-1}}{(1 - az^{-1})^2}$$

with ROC = $\{z : |z| > a\}$.

- If $x[n] = -na^n u[-n - 1]$, then

$$X(z) = \frac{az^{-1}}{(1 - az^{-1})^2}$$

with ROC = $\{z : |z| < a\}$.

- If $x[n] = \delta[n - n_0]$, then

$$X(z) = z^{-n_0}$$

with ROC = $\{z : |z| > 0\}$ if $n_0 > 0$. If $n_0 < 0$, then ROC is the entire complex plane with the exception of ∞ .

- **Continuous time Fourier Transform (CTFT):** For any signal $x(t)$,

$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\Omega) e^{j\Omega t} d\Omega$$

where

$$X(j\Omega) = \int_{-\infty}^{\infty} x(t) e^{-j\Omega t} dt.$$

• **Properties of CTFT:**

- **Linearity:**

$$\begin{aligned} x(t) &\rightarrow X(j\Omega) \\ y(t) &\rightarrow Y(j\Omega) \end{aligned} \implies \alpha x(t) + \beta y(t) \rightarrow \alpha X(j\Omega) + \beta Y(j\Omega)$$

- **Time reversal:**

$$x(t) \rightarrow X(j\Omega) \implies x(-t) \rightarrow X(-j\Omega)$$

- **Time shifting:**

$$x(t) \rightarrow X(j\Omega) \implies x(t - t_0) \rightarrow X(j\Omega) e^{-j\Omega t_0}$$

- **Frequency shifting:**

$$x(t) \rightarrow X(j\Omega) \implies x(t) e^{j\Omega_0 t} \rightarrow X(j(\Omega - \Omega_0))$$

- **Convolution:**

$$\begin{aligned} x(t) &\rightarrow X(j\Omega) \\ y(t) &\rightarrow Y(j\Omega) \end{aligned} \implies x(t) * y(t) \rightarrow X(j\Omega) Y(j\Omega)$$

- **Multiplication:**

$$\begin{aligned} x(t) &\rightarrow X(j\Omega) \\ y(t) &\rightarrow Y(j\Omega) \end{aligned} \implies x(t)y(t) \rightarrow \frac{1}{2\pi} [X(j\Omega) * Y(j\Omega)]$$

- **Differentiation in time domain:**

$$x(t) \rightarrow X(j\Omega) \implies \frac{dx(t)}{dt} \rightarrow j\Omega X(j\Omega)$$

- **Differentiation in frequency domain:**

$$x(t) \rightarrow X(j\Omega) \implies -jtx(t) \rightarrow \frac{dX(j\Omega)}{d\Omega}$$

- **Some known signal-CTFT pairs:**

- If $x(t) = \delta(t - t_0)$, then

$$X(j\Omega) = e^{-j\Omega t_0}.$$

- If $x(t) = e^{j\Omega_0 t}$, then

$$X(j\Omega) = 2\pi \delta(\Omega - \Omega_0).$$

- If $x(t) = \frac{\sin(At)}{t}$, then

$$X(j\Omega) = \begin{cases} \pi & -A \leq \Omega \leq A \\ 0 & \text{otherwise} \end{cases}$$

- If $x(t) = \begin{cases} 1/2 & -B \leq t \leq B \\ 0 & \text{otherwise} \end{cases}$, then

$$X(j\Omega) = \frac{\sin(B\Omega)}{\Omega}.$$

- **Sampling and reconstruction:** If $x_d[n] = x_c(nT)$, x_d is said to be the sampled version of the continuous-time signal x_c . The sampling period is T , and the sampling frequency is $\Omega_s = \frac{2\pi}{T}$.

- Defining the intermediate signal $x_s(t)$ as

$$x_s(t) = x_c(t) \left(\sum_{k=-\infty}^{\infty} \delta(t - kT) \right),$$

we have the relation

$$X_s(j\Omega) = \frac{1}{T} \sum_{k=-\infty}^{\infty} X_c(j(\Omega - k\Omega_s)).$$

- The relation between $X_s(j\Omega)$ and $X_d(e^{j\omega})$ is

$$X_d(e^{j\omega}) = X_s \left(j \left(\frac{\omega}{T} \right) \right)$$

or

$$X_s(j\Omega) = X_d(e^{j\Omega T}).$$

- Therefore,

$$X_d(e^{j\omega}) = \frac{1}{T} \sum_{k=-\infty}^{\infty} X_c \left(j \left(\frac{\omega - k2\pi}{T} \right) \right).$$

- These imply that in order for us to reconstruct $x_c(t)$ perfectly, we need

$$\Omega_s = \frac{2\pi}{T} \geq 2\Omega_M$$

where Ω_M is the bandwidth, i.e., maximum frequency where $X_c(j\Omega) \neq 0$.

- Reconstruction is accomplished by eliminating the replicas of $X_c(j\Omega)$ from $X_s(j\Omega)$. Ideally, this is done by a filter

$$H(j\Omega) = \begin{cases} T & -\frac{\pi}{T} \leq \Omega \leq \frac{\pi}{T} \\ 0 & \text{otherwise} \end{cases}$$

which, in the time domain becomes $h(t) = \frac{\sin(\frac{\pi}{T}t)}{\frac{\pi}{T}t}$.