

# CS663 - Assignment 5 - Question 2

Kavan Vavadiya  
Roll No: 210100166

Kushal Agarwal  
Roll No: 210100087

Anshika Raman  
Roll No: 210050014

November 2024

## Analyzing Blurred Images with Fourier Transform Techniques

The objective is to reconstruct two images,  $f_1$  and  $f_2$ , from two blurred images  $g_1$  and  $g_2$ , using known blur kernels  $h_1$  and  $h_2$ . The images  $f_1$  and  $f_2$  represent an external scene and its reflection, respectively. The relationships between the images and the kernels are given by:

$$g_1 = f_1 + h_2 * f_2$$

$$g_2 = f_2 + h_1 * f_1$$

Applying the Fourier Transform to both equations, we obtain:

$$\mathcal{F}(g_1) = \mathcal{F}(f_1) + \mathcal{F}(h_2 * f_2) \Rightarrow G_1 = F_1 + H_2 F_2$$

$$\mathcal{F}(g_2) = \mathcal{F}(f_2) + \mathcal{F}(h_1 * f_1) \Rightarrow G_2 = F_2 + H_1 F_1$$

To isolate  $F_1$  and  $F_2$ , we manipulate these equations:

$$F_1 = \frac{G_1 - H_2 G_2}{1 - H_1 H_2}, \quad F_2 = \frac{G_2 - H_1 G_1}{1 - H_1 H_2}$$

Finally, the inverse Fourier Transforms of these expressions yield the deblurred images:

$$f_1 = \mathcal{F}^{-1} \left( \frac{G_1 - H_2 G_2}{1 - H_1 H_2} \right)$$

$$f_2 = \mathcal{F}^{-1} \left( \frac{G_2 - H_1 G_1}{1 - H_1 H_2} \right)$$

We note that the division by  $1 - H_1 H_2$  can lead to numerical instability if this term is close to zero, indicating potential problems in specific frequency domains where the product  $H_1 H_2$  approaches unity. This necessitates careful examination of the system's frequency response to ensure accurate image recovery.