කෘෂිකර්ම දෙපාර්තමේන්තුව
விவசாயத் திணைக்களம்
Department of Agriculture

වී පර්යේෂණ හා සංවර්ධන ආයතනය - බතලගොඩ
நெல் ஆராய்ச்சி அபிவிருத்தி நிறுவனம் - பதலகொட
Rice Research and Development Institute (RRDI) - Bathalagoda

Nursery systems - Dapog method

Introduction

Dapog method.

- is easy to establish as it can be located anywhere on a flat surface
- require small nursery area
- is the fastest method to raise paddy seedlings
- It takes 12 - 15 days to transplant
- requires carefully controlled irrigation

Steps of Establishing Dapog method

This nursery can be established on any flat surface

- First, construct a raised earthened bed
- Cover the bed with green banana leaves or plastic sheeting
- Surface of the bed should be kept as level as possible
- Pre-germinated seeds should be to sow
- After sowing the seeds, from the 3rd day seed bed should be kept continuously flooded
- Seedlings for one hectare can be produced in a 100m

Advantages

- Ability raises seedlings in relatively short period of time

- Require a small area to raise seedlings
- Can establish at any place
- Excellent for replacing small quantities of seedlings destroyed by pest
- Minimal transplanting shock (Recovery of seedlings once transplanted is fast)

Disadvantages

- Transplanting can be relatively difficult because seedlings are small

Dapog nurseries are more suitable for short aged varieties. However, before transplanting, the field should be leveled well and free of water since the seedlings are very small.