

බස්නාහිර පළාත් අධ්‍යාපන දෙපාර්තමේන්තුව

Western Province Educational Department

අධ්‍යයන පොදු සහතික පත්‍ර (උසස් පෙළ) විභාගය, 2023 (2024)

General Certificate of Education (Adv. Level) Examination, 2023 (2024)

සංයුක්ත ගණිතය

II

Combined Mathematics

II

10

E

II

2023.12.15/ 08.30 - 11.40

පැය තුනයි

Three hours

අමතර කියවීම් කාලය

- මිනිත්තු 10 යි

Additional Reading Time

- 10 minutes

additional reading time to go through the question paper, select the questions and decide on the questions that you give priority in answering.

Index Number

Instructions:

- * This question paper consists of two parts.

Part A (Questions 1 - 10) and **Part B** (Questions 11 - 17)

- * **Part A:**

Answer **all** questions. Write your answers to each question in the space provided. You may use additional sheets if more space is needed.

- * **Part B:**

Answer **five** questions only. Write your answers on the sheets provided.

- * At the end of the time allotted, tie the answer scripts of the two parts together so that **Part A** is on top of **Part B** and hand them over to the supervisor.

- * You are permitted to remove only **Part B** of the question paper from the Examination Hall.

For Examiners' Use only

(10) Combined Mathematics II		
Part	Question No.	Marks
A	1	
	2	
	3	
	4	
	5	
	6	
	7	
	8	
	9	
	10	
B	11	
	12	
	13	
	14	
	15	
	16	
	17	
	Total	
	Percentage	

Final Marks


In Numbers	
In Words	

Code Numbers


Marking Examiner	
Checked by:	1
	2
Supervised by:	

Part A


01. Two particles A and B of equal size and mass $2m$ and m are at rest on a smooth horizontal on a straight line, perpendicular to the wall as shown in the figure. The particle B is projected horizontally with velocity u towards the wall. Show that, if $e > \frac{1}{2}$, then B collides with the wall again after the collision between A and B in the subsequent motion, where e is the coefficient of restitution between B and the wall and between A and B .



02. A staircase of height a and breadth a at a distance $2a$ from a fixed point O on the horizontal ground. A particle is projected with speed u at an angle θ to the horizontal from O and it just reaches point A , at the end of the third step. Show that $u^2 \sin 2\theta = 8ag$ and further, by considering the vertical motion from O to A , $\sin^2 \theta = \frac{6ga}{u^2}$. Hence deduce $\tan \theta = \frac{3}{2}$.




03. Two particles of masses m and $2m$ placed on the same inclined face of inclination 30° to horizontal of a smooth wedge of mass $3m$ are attached to the two ends of a light inextensible string passing over a smooth pulley attached to the wedge. When the system is released from rest, the wedge moves with an acceleration F relative to the Earth. Show that the magnitude of the acceleration of the particles **relative to the wedge** is $4\sqrt{3}F$.




04. A car of mass m kg and maximum power H kW moves up a straight level road inclined at an angle α to the horizontal with a maximum speed u ms^{-1} against a constant resistance. Find the resistance to the motion. When the car moves with an acceleration a ms^{-2} on a level road against that resistance, then show that the speed of the car is $\frac{1000Hu}{mua - mg \sin \alpha + 1000H}$.

05. A particle of mass m traces out a circle of radius r at a height h inside a smooth cone fixed vertically with the vertex downwards as shown in the diagram. Show that the speed of the particle is $u = \sqrt{gh}$.




06. In the usual notation, $-\mathbf{i} + 2\mathbf{j}$ and $2\alpha\mathbf{i} + \alpha\mathbf{j}$ are the position vectors of two points A and B , respectively, with respect to a fixed point O . where $\alpha(>0)$ is a constant. Using the **scalar product**, show that $\angle AOB = \frac{\pi}{2}$. Let C be a point such that $OACB$ is a rectangle. If the vector \overrightarrow{OC} lies on the y -axis, find the value of α .

07. A uniform rod of length $4a$ and weight w is in equilibrium in a fixed smooth hemispherical bowl of radius $\sqrt{3}a$ such that a part of the rod of length a is at the outside as shown in the figure. Find the inclination of the rod to the horizontal and show that the reactions of the bowl on the rod are equal to $\frac{w}{\sqrt{3}}$.



08. A particle of weight w is kept in equilibrium on a rough inclined plane with an inclination 30° to the horizontal, by a light elastic string attached to the particle and a point on the plane. If the coefficient of friction between the plane and the particle is $\frac{\sqrt{3}}{4}$ and the tension in the elastic string is T , show that $\frac{w}{8} \leq T \leq \frac{7w}{8}$.



09. A and B be two events in the event space corresponding to the sample space Ω . Given that

$$P(A') = \frac{1}{4}, P(B) = \frac{1}{3} \text{ and } P(A \cup B) = \frac{5}{6}. \text{ Show that } A \text{ and } B' \text{ are independent events.}$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10. The mean and standard deviation of grade 13 students in General English are 50 is 5 respectively. These scores should be adjusted using a linear transformation $y_i = ax_i + b$ such that the mean is 66 and the standard deviation is 6. Here $a, b > 0$. Find, the values of a and b . Also, find the **transformed mark** for the original mark 55.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

බස්නාහිර පළාත් අධ්‍යාපන දෙපාර්තමේන්තුව

Western Province Educational Department

අධ්‍යයන පොදු සහතික පත්‍ර (උසස් පෙළ) විභාගය, 2023 (2024)
General Certificate of Education (Adv. Level) Examination, 2023 (2024)

සංයුක්ත ගණිතය

II

Combined Mathematics

II

10

E

II

2023.12.15/ 08.30 - 11.40

Part B

■ Answer only **five** questions.

11. (a) Santa Claus travels through the sky at a speed v at a horizontal height h above ground level, riding on a cart with reindeer. At a certain instant, Santa sees a small boy named P standing at a point B on the ground $2d$ horizontal distance ahead of a point A vertically below the cart, and instantly drops a gift parcel named Q . At the same instant the boy P starts running towards A along BA with constant acceleration a from rest, parallel to the direction of motion of the cart. Child P just catches the parcel Q at ground level.

Sketch the velocity-time graphs for the horizontal motion of Q and the motion of child P in the same diagram. Also, draw the velocity-time graph for the vertical motion of Q in a separate diagram. **Hence**, show that the time taken to caught the parcel Q by P is $\frac{2gd - ah}{g^2v}$ from the time it is released, and the distance travelled by the boy at the instant is $\frac{ah}{g}$.

- (b) A ship S is in breakdown at a point L in still sea d km East of a point O on a straight coast extending North and travels along LM with a constant speed u kmh^{-1} sounding the hazard horn to reach a port M at d km North of O . At the same instant, a fishing boat B_1 starts moving from point N which is at d km South of O with a speed of $2u$ kmh^{-1} in the North-East direction and a relief boat B_2 starts from the point O to intercept the ship S with a speed u kmh^{-1} relative to the Earth.


Draw the velocity triangles on the same figure to determine the velocities of the boats relative to the ship. Find the shortest distance between S and B_1 and show that if the hazard horn is heard

at a distance of R km $\left(\sqrt{\frac{2}{5}}d < R < \sqrt{2}d\right)$ then B_1 will hear the horn for a time $\frac{2\sqrt{5R^2 - 2d^2}}{5u}$.

Also find the time taken by relief boat B_2 to intercept S .

12. (a) In the figure ABC is a triangle with $\hat{ACB} = \alpha$, $\hat{ABC} = \frac{\pi}{2}$


is the vertical cross-section through the center of gravity of a smooth uniform wedge of mass $5m$ placed on a smooth horizontal ground. AB is a line of the maximum slope of the face containing it. D is a fixed point in ABC plane such that AD is horizontal.



A particle P of mass $4m$ and a movable pulley Q of mass $2m$ are attached to both ends of a light elastic string passing over two small smooth pulleys fixed at A and D .

Two particles R and S of masses and respectively are attached to the two ends of a light inelastic string passing over the moving pulley Q . When the system is released from rest with all the strings taut, then **write the equations** sufficient to determine the acceleration of the particles and the tension in the strings.

- (b) A rough thin horizontal tube BC is attached to the end B of a thin smooth thin semicircular tube AB of radius $2a$ and a smooth thin tube CD of radius a is attached as shown in the figure. (The radius passing through D makes an angle of $\frac{\pi}{3}$ with the vertical.) A particle P of mass m is placed at A and projected horizontally into the tube with speed u .



When the particle P is inside the tube AB , show that the radius through which the particle makes an angle θ with the upward vertical then $\dot{\theta}^2 = \frac{1}{4a^2}(u^2 + 4ga(1 - \cos\theta))$ and also find the reaction on the particle by the tube.

Now, let $u = \sqrt{ga}$. Find the velocity of the particle at B . In the subsequent motion, if the particle P just reached to the point D , show that the energy loss in the tube BC is $3mga$. Hence, show that the velocity of the particle at C is $\sqrt{3ga}$.

13. Two springs AP and PB of natural lengths $2l$ and l are attached to a particle P of mass m and the other ends of the springs are fixed to two points A and B at a distant $6l$ apart on a smooth horizontal table. The modulus of elasticity of the spring AP is $2mg$ and the modulus of elasticity of the spring

PB is mg . Show that the particle is in equilibrium at a point C , at a distance $\frac{7}{2}l$ from A .

Now the particle is displaced to the point M on AB , where $AM = l$ and released from rest. If

$AP = x$, then show that $\ddot{x} + \frac{2g}{l}\left(x - \frac{7}{2}l\right) = 0$ for $l < x < 2l$. Let Q and R be two points on AB

such that $AQ = 2l$ and $AR = 5l$. By using the formulae $\dot{x}^2 = \frac{2g}{l}(c^2 - x^2)$, show that the velocity of

P at Q is $2\sqrt{2gl}$. Here c is the amplitude. Show that the time taken to move M to Q is

$$\sqrt{\frac{l}{2g}} \cos^{-1} \frac{3}{5}.$$


Now, let $CP = y$. Show that $\ddot{y} + \frac{2g}{l}y = 0$ for $-\frac{3l}{2} \leq y \leq \frac{3l}{2}$. Assuming that the solution of the

above equation is of the form $y = \alpha \cos \omega t + \beta \sin \omega t$, and $t = 0$ at Q , then find the values of the constants α, β and ω . Hence, find the center and amplitude of the simple harmonic motion performed by the particle from Q to R .

14. (a) The position vectors of two points A and B with respect to O are \mathbf{a} and \mathbf{b} respectively. C is a point on the line that passes through O parallel to AB . Find the position vector of C such that $OABC$ is a parallelogram. Let D be a point on produced line OC and E be a point on OB such that $OE : EB = 3 : 1$. Find the position vector D such that A, E, D are collinear. Find the

ratio $AE : ED$. If $AB = \sqrt{5}$ units and $2OA = OB$ then show that $\hat{BOA} = \cos^{-1} \left(\frac{5(|\mathbf{a}|^2 - 1)}{4|\mathbf{a}|^2} \right)$


- (b) In the hexagon $OABCDE$, the sides OA and OD lie on the X and Y axis respectively. $OA = DC = 2$ m, $OD = AC = 4$ m, $\hat{EOD} = \hat{BAC} = \frac{\pi}{3}$ and $\hat{DEO} = \hat{ABC} = \frac{\pi}{2}$.



The forces $5N$, $2\sqrt{3}N$, $2N$, $3N$, $6N$ and $4\sqrt{3}N$ act along the sides AO , AB , BC , CD , ED and EO respectively. Find the magnitude, direction and the equation of the line of action of the resultant. Hence, find the coordinates of the point where the line of action of the resultant meets OX .


When a couple of magnitude M act in the OXY plane, the resultant passes through A . Find the magnitude and direction of M .

15. (a) Four rods AB, BC, CD, DA each of length $2a$ and weight w are smoothly jointed at ends A, B, C and D . The system is at equilibrium in a vertical plane by a light rod of length $2a$, joining the points B and C with the midpoints BC of CD and touching two pegs in the same horizontal level as shown in the figure. Find the reactions of the pegs on rods BC and CD and the reactions at joints A and C . Also, show that the




thrust of the light rod is $\frac{2w}{\sqrt{3}}$.

- (b) Framework shown in the figure consists of five light rods AB, BC, CD, BD and AD smoothly jointed at their ends. Here, it is given that $AB = BC$, $AD = DC$ and $\hat{ADB} = \hat{BDC} = 30^\circ$. It is freely hinged at A and a load $3W$ is suspended at the joint C . The framework is kept in equilibrium in a vertical plane with AD vertical and AB horizontal by a force R at the joint D in the direction shown in the figure. Draw a stress diagram, using Bow's notation. Hence, find the magnitude of R , and the reaction at the hinge A . Also find the stresses in the five rods, stating whether they are tensions or thrusts.



16. By using the integration, show that the center of mass of


- (i) a uniform **solid** hemisphere of radius r is at a distance $\frac{3r}{8}$ from its center.
- (ii) a uniform **hollow** hemispherical frustum of radius $2r$ and height $\sqrt{3}r$ is at a distance $\frac{\sqrt{3}}{2}r$ from its center.



A flower vase is made by attaching a solid hemisphere of radius $2r$ and a hollow cylinder of radius r and height h to a uniform hollow hemispherical frustum of radius $2r$ and height $\sqrt{3}r$ as shown in the diagram.

The mass per unit area of the cylinder and hollow hemispherical frustum is ρ and the mass per unit volume of the

solid hemisphere is $\frac{k\rho}{r}$. Show that the center of mass of the vase is located at a distance $\frac{3[2\sqrt{3}rh + h^2 + 6r^2 - 4kr^2]}{2(3h + 6\sqrt{3}r + 8kr)}$ above OA .



Now, let $h = 2r$. The hemispherical surface of the vase is placed on a rough horizontal ground so that its axis of symmetry is vertical. The vase is slightly displaced from its equilibrium position so that the axis of symmetry makes a small angle with the vertical. Show that the vase falls if

$k > \frac{5 + 2\sqrt{3}}{2}$. What happened when

i. $k = \frac{5 + 2\sqrt{3}}{2}$

ii. $k < \frac{5 + 2\sqrt{3}}{2}$

17. (a) An insurance company classifies all their policyholders into three categories as high risk, high risk and low risk. 40% of the policyholders belong to the high-risk category and 35% belong to the risk category. According to last year's statistics, the probability of someone in the high-risk, risk and low-risk categories being in an accident is 60%, 20% and 5%, respectively.

Find the probability that,

- i. A person selected at random has been faced in an accident in the last year.
- ii. An accident victim is randomly selected to be in the low-risk category,

- (b) The marks obtained by 100 students of a certain school for the subject of Composite Mathematics are given in the table below.

Class intervals	21 - 35	36 - 50	51 - 65	66 - 80	81 - 95
No. of students	8	31	40	15	6

Determine the mean, standard deviation and mode of the distribution given above.

Find the median, first and third quartiles and draw box and whisker plot for the distribution and draw the shape of the distribution.