



First Term Test – 2018 November

Combined Mathematics- II

Grade 13

3 hours

Name : .....

Instructions :

- ★ This question paper consists of two parts.  
**Part A** (Questions 1 – 10) and **Part B** (Questions 11 – 16)
- ★ **Part A**  
Answer all questions. Write your answer in the space provided.
- ★ **Part B**  
Answer only 5 questions.
- ★ At the end of the time allocated, time the answers of the two parts together so that **Part A** is on top of **Part B** before handing them over to the supervisor.
- ★ You are permitted to remove only **Part B** of the question paper from the Examination Hall.

Part	Question NO.	Marks Awarded
A	01	
	02	
	03	
	04	
	05	
	06	
	07	
	08	
	09	
	10	
B	11	
	12	
	13	
	14	
	15	
	16	

Final Mark

**Part A**

- 01). A particle moves on a straight line with a uniform acceleration and describes distances  $x$  and  $y$  in successive intervals of time  $t_1$  and  $t_2$  respectively. Prove that the acceleration is  $\frac{2(yt_1 - xt_2)}{t_1t_2(t_1 + t_2)}$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 02). A ship **A** is travelling due east at  $2u \text{ kmh}^{-1}$  and a second ship **B** is travelling to a direction  $30^\circ$  east of south, with a velocity  $u \text{ kmh}^{-1}$ . Initially if **A** is  $d \text{ km}$  due south to **B**, Find the velocity of ship **A** relative to ship **B** and the shortest distance occur between them.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

- 
- This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

- 
- This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

- 05). A vertical cliff is 63m high. A stone A is projected horizontally with a velocity  $28 \text{ ms}^{-1}$  while another stone B is projected from the bottom of the cliff with a velocity  $35 \text{ ms}^{-1}$  at an angle  $\alpha$  to horizontal. If both stones move under gravity and collide in air, Show that  $\cos \alpha = \frac{4}{5}$ .  
Find the time taken for the collision.


- 06). The figure shows a vertical cross section of a uniform rod AB, resting between two planes. The inclined plane is smooth while the vertical plane is rough with the coefficient of friction is  $\frac{1}{2}$ .  
If the rod is in equilibrium, find the angle  $\theta$ .



- 07). A body of mass  $M$  kg rests on a smooth plane. The hanging mass is  $m$  kg. If the hanging mass moves downwards,

show that the tension of the string is  $\frac{M m g}{M + m} (1 + \sin \alpha)$ .

Deduce for what values of  $\sin \alpha$ , does the hanging mass moves upward.



- 08). A one tone car tows a caravan of mass 800kg along a levelled horizontal road using a rigid tow bar. The resistance to the motion of the car is 150N and the caravan experience a resistance 250N. If the engine of the car is working at 13 KW and it is moving with velocity  $10 \text{ ms}^{-1}$ . Find the tension of the tow bar.

- 
- This image shows a full page of white paper with horizontal dotted lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- 
- This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

## Part – B

11. (a). A stone **S** is projected vertically upward from a point **O** , on a horizontal ground with a velocity ***u***. At the same instant a balloon **B** is projected from rest ,from a point which is at a vertical height ***h*** from the point **O**, where (  $h < \frac{u^2}{2g}$  ). The balloon moves with an acceleration  $\frac{g}{2}$ .

If **B** and **S** just touch in air, draw a velocity time graph for the motion of **B** relative to **S**.

Hence find the time taken by **S** to touch **B**.

Also show that  $u^2 = 3gh$ .

- (b). A particle is projected under gravity from a point **O** on a horizontal ground , with a velocity ***u*** inclined  $\theta$  to horizontal.

With respect to the OXY plane through **O**, if the particle just pass through a point **P**( ***x*** , ***y*** )

Show that the locus of the particle is given by  $y = x \tan \theta - \frac{gx^2}{2u^2} \sec^2 \theta$ .

Deduce that the horizontal range of the particle is  $\frac{u^2}{g} \sin 2\theta$ .

Find the angle  $\theta$ , (  $0 < \theta < \frac{\pi}{2}$  ), such that the horizontal range is maximum for a constant velocity ***u***.

Show that the maximum height attained by the particle at that moment is  $\frac{a}{4}$ , where ***a*** is the

maximum horizontal range. Further if this particle just pass through two walls of height  $\frac{a}{8}$ ,

at apart ***b*** distance each other , show that  $a = \sqrt{2}b$

12. (a). A plane can travel with velocity ***u*** in still air. When a wind blows with velocity ***v*** (  $< u$  ) from North, the plane travels from city **A** to city **B** and back to **A**, without stay at **B**. The city **B** is located at a distance ***d*** in a direction  $30^\circ$  West of North from the city **A**. In both trips the plane travels in a path making an angle  $\alpha$  with **AB**.


Show that  $\sin \alpha = \frac{v}{2u}$ , and the time taken for both trips is  $\frac{d \left( \sqrt{4u^2 - v^2} \right)}{2(u^2 - v^2)}$ .

- (b). An engine of mass **M** metric tons moves on a straight horizontal path at a constant speed  $v \text{ kmh}^{-1}$  against a constant resistance from the road. If the engine works with its maximum power ***H*** kW. Find the resistance from the road to the motion of the engine.
- Find the acceleration of the engine when it moves with a speed  $u \text{ kmh}^{-1}$  along a straight road inclined at an angle  $\alpha$  to horizontal, against the same road resistance.

13. (a). Each of the weights A and B has a mass of  $m$  and the mass of C is  $km$ . All the pulleys are light and smooth. If the system is released from rest, find the acceleration of the moveable pulley.

Deduce that the scale pan will ascend if  $k > 4$ .

When the system is moving freely, find the tension of the string and the reaction between the weights A and B.



- (b). The figure shows the cross section ABC of

a wedge of  $\hat{ACB} = 90^\circ$  and  $\hat{CAB} = 60^\circ$  of mass  $m$  which is placed on a smooth horizontal table.


The inclined planes AC and BC are equally rough and  $\mu$  being the coefficient of friction between each particle and the planes.

The mass of the particles on AC and BC are  $m$  and  $2m$  respectively, and connected

by a light inextensible string pass through a smooth pulley at C as shown in the figure.

Write necessary equations and determine the accelerations of each particles  $m$  and  $2m$ .

when  $\mu = \frac{1}{2}$ .



14. (a). A system of forces consisting three forces with respect to OXY plane given below. Where  $\underline{i}$  and  $\underline{j}$  denote unit vectors along OX and OY axes respectively.

Point	Position vector	Force
A	$2\underline{i} + 3\underline{j}$	$3\underline{i} + 4\underline{j}$
B	$6\underline{i} + \underline{j}$	$-\underline{i} + 6\underline{j}$
C	$-3\underline{i} + 2\underline{j}$	$-3\underline{i} - 3\underline{j}$

Mark these forces in a single diagram. Find the magnitude and direction of the resultant force.

Show that the Cartesian equation of the line of action of this resultant force is  $7x + y = 51$ .

An additional force of  $\lambda\underline{i} + \mu\underline{j}$  is now introduced to the system at the origin with a couple G, such that the system to be in equilibrium. Find  $\lambda$ ,  $\mu$  and  $G$ .



- (b). A, B and C are three non collinear points. The position vectors of A and B with respect to the point C is  $\underline{a}$  and  $\underline{b}$  respectively. The point E is on AC such that  $AE : EC = 2 : 3$  and the point D is on CB such that  $CD : DB = 4 : 1$

show that  $\overrightarrow{BE} = \frac{1}{5}(3\underline{a} - 5\underline{b})$

Write a similar expression for  $\overrightarrow{AD}$ . The lines AD and BE intersect at X.

Write  $\overrightarrow{AX}$  using two different occasions and hence build up a suitable vector equation in the form of  $\alpha\underline{a} + \beta\underline{b} = \underline{0}$ , where  $\alpha, \beta \in \mathbb{R}$ .

Deduce the ratio of AX : XD

15. (a). Four equal uniform rods each of length  $2a$  and weight  $w$  are smoothly jointed together at their ends to form a rhombus ABCD. It is suspended from the vertex A, and is maintained in equilibrium with C below A and with  $\hat{DAB} = 2\theta$  by means of an inextensible string connected to the mid points of the rods AB and BC.

Show that the reaction at the joint C is  $\frac{w}{2}\sqrt{4 + \tan^2 \theta}$ .


Find the tension of the string.

- (b). A frame work is made up of five light rods freely jointed as shown in the figure.

A and B joints rest on smooth supports.

A load of 900 N acts at D.

Find the reactions at A and B.



Draw a stress diagram, using Bow's notation and hence find the stress in each rod.

Classify them in to tension or thrust.

16. (a). A circular wire is fixed in a vertical plane with two rings of weight  $w$  and  $\lambda w$  are attached to the extreme ends A and B of a light rod AB. The rod is in equilibrium in the upper part of the circular frame, such that the rod subtended  $2\alpha$  at the center and the rod make an angle  $\theta$  at A with the downward vertical.

Show that  $(1 - \lambda)\tan\theta = (1 + \lambda)\cot\alpha$

When  $\lambda = \frac{1}{2}$  and  $\theta = \frac{\pi}{3}$ , Find the reactions on each ring from the rod and the stress in the light rod using the triangle of forces.

(b). Two equal uniform rods  $AC$ ,  $CB$  each of weight  $w$ , are smoothly hinged together at  $C$ , to form a shape of equilateral triangle  $ABC$  while the ends  $A$  and  $B$  rest on a rough horizontal floor. The coefficient of friction of both ends  $A$  and  $B$  is  $\mu$ . A force  $\mathbf{P}$  is applied at the point  $D$  on the rod  $AC$ , such that  $AD = \frac{3}{4}AC$ , where the force  $\mathbf{P}$  is perpendicular to the rod  $AC$  and away from  $B$ .

Show that the ratio between the frictional force to the normal reaction at  $A$  is  $\frac{2w + 3P}{\sqrt{3}(4w + P)}$ .

Find the above ratio at  $B$  also.

Hence find the magnitude of force  $\mathbf{P}$ , such that both ends  $A$  and  $B$  slip together.

Further show that  $\mu = \frac{\sqrt{3}}{2}$ , for the above occasion.

