


Date : 11 August 2019

A CONSOLIDATED QUESTION PAPER-CUM-ANSWER BOOKLET



Keep Practising!  
K.

# MAINS TEST SERIES-2019

(JUNE-2019 to SEPT.-2019)

Under the guidance of K. Venkanna

## MATHEMATICS

PAPER - I : FULL SYLLABUS

TEST CODE: TEST-11: IAS(M)/11-AUG.-2019

Time: 3 Hours

Maximum Marks: 250

### INSTRUCTIONS

1. This question paper-cum-answer booklet has 50 pages and has 34 PART/SUBPART questions. Please ensure that the copy of the question paper-cum-answer booklet you have received contains all the questions.
2. Write your Name, Roll Number, Name of the Test Centre and Medium in the appropriate space provided on the right side.
3. A consolidated Question Paper-cum-Answer Booklet, having space below each part/sub part of a question shall be provided to them for writing the answers. Candidates shall be required to attempt answer to the part/sub-part of a question strictly within the pre-defined space. Any attempt outside the pre-defined space shall not be evaluated."
4. Answer must be written in the medium specified in the admission Certificate issued to you, which must be stated clearly on the right side. No marks will be given for the answers written in a medium other than that specified in the Admission Certificate.
5. Candidates should attempt Question Nos. 1 and 5, which are compulsory, and any THREE of the remaining questions selecting at least ONE question from each Section.
6. The number of marks carried by each question is indicated at the end of the question. Assume suitable data if considered necessary and indicate the same clearly.
7. Symbols/notations carry their usual meanings, unless otherwise indicated.
8. All questions carry equal marks.
9. All answers must be written in blue/black ink only. Sketch pen, pencil or ink of any other colour should not be used.
10. All rough work should be done in the space provided and scored out finally.
11. The candidate should respect the instructions given by the invigilator.
12. The question paper-cum-answer booklet must be returned in its entirety to the invigilator before leaving the examination hall. Do not remove any

READ INSTRUCTIONS ON THE  
LEFT SIDE OF THIS PAGE  
CAREFULLY

Name SWIT SHANKAR

Roll No. —

Test Centre ORM DELHI

Medium ENGLISH

Do not write your Roll Number or Name anywhere else in this Question Paper-cum-Answer Booklet.

I have read all the instructions and shall abide by them

Swit shankar  
Signature of the Candidate

I have verified the information filled by the candidate above

Signature of the invigilator

### IMPORTANT NOTE:

Whenever a question is being attempted, all its parts/ sub-parts must be attempted contiguously. This means that before moving on to the next question to be attempted, candidates must finish attempting all parts/sub-parts of the previous question attempted. This is to be strictly followed. Pages left blank in the answer-book are to be clearly struck out in ink. Any answers that follow pages left blank may not be given credit.

**DO NOT WRITE ON  
THIS SPACE**

## INDEX TABLE

QUESTION	No.	PAGE NO.	MAX. MARKS	MARKS OBTAINED
1	(a)			08
	(b)			09
	(c)			08
	(d)			09
	(e)			08
2	(a)			14
	(b)			10
	(c)			13
	(d)			
3	(a)			
	(b)			
	(c)			
	(d)			
4	(a)			
	(b)			
	(c)			
	(d)			
5	(a)			09
	(b)			09
	(c)			08
	(d)			09
	(e)			08
6	(a)			12
	(b)			09
	(c)			11
	(d)			11
7	(a)			
	(b)			
	(c)			
	(d)			
8	(a)			11
	(b)			08
	(c)			12
	(d)			14
Total Marks				

210  
250

BAHAT X 2000

OBRAZET ZEPLA	ZERAVI XGM	OKRES	AK	ADMINTO
1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25
26	27	28	29	30
31	32	33	34	35
36	37	38	39	40
41	42	43	44	45
46	47	48	49	50
51	52	53	54	55
56	57	58	59	60
61	62	63	64	65
66	67	68	69	70
71	72	73	74	75
76	77	78	79	80
81	82	83	84	85
86	87	88	89	90
91	92	93	94	95
96	97	98	99	100
101	102	103	104	105
106	107	108	109	110
111	112	113	114	115
116	117	118	119	120
121	122	123	124	125
126	127	128	129	130
131	132	133	134	135
136	137	138	139	140
141	142	143	144	145
146	147	148	149	150
151	152	153	154	155
156	157	158	159	160
161	162	163	164	165
166	167	168	169	170
171	172	173	174	175
176	177	178	179	180
181	182	183	184	185
186	187	188	189	190
191	192	193	194	195
196	197	198	199	200
201	202	203	204	205
206	207	208	209	210
211	212	213	214	215
216	217	218	219	220
221	222	223	224	225
226	227	228	229	230
231	232	233	234	235
236	237	238	239	240
241	242	243	244	245
246	247	248	249	250
251	252	253	254	255
256	257	258	259	260
261	262	263	264	265
266	267	268	269	270
271	272	273	274	275
276	277	278	279	280
281	282	283	284	285
286	287	288	289	290
291	292	293	294	295
296	297	298	299	300
301	302	303	304	305
306	307	308	309	310
311	312	313	314	315
316	317	318	319	320
321	322	323	324	325
326	327	328	329	330
331	332	333	334	335
336	337	338	339	340
341	342	343	344	345
346	347	348	349	350
351	352	353	354	355
356	357	358	359	360
361	362	363	364	365
366	367	368	369	370
371	372	373	374	375
376	377	378	379	380
381	382	383	384	385
386	387	388	389	390
391	392	393	394	395
396	397	398	399	400
401	402	403	404	405
406	407	408	409	410
411	412	413	414	415
416	417	418	419	420
421	422	423	424	425
426	427	428	429	430
431	432	433	434	435
436	437	438	439	440
441	442	443	444	445
446	447	448	449	450
451	452	453	454	455
456	457	458	459	460
461	462	463	464	465
466	467	468	469	470
471	472	473	474	475
476	477	478	479	480
481	482	483	484	485
486	487	488	489	490
491	492	493	494	495
496	497	498	499	500
501	502	503	504	505
506	507	508	509	510
511	512	513	514	515
516	517	518	519	520
521	522	523	524	525
526	527	528	529	530
531	532	533	534	535
536	537	538	539	540
541	542	543	544	545
546	547	548	549	550
551	552	553	554	555
556	557	558	559	560
561	562	563	564	565
566	567	568	569	570
571	572	573	574	575
576	577	578	579	580
581	582	583	584	585
586	587	588	589	590
591	592	593	594	595
596	597	598	599	600
601	602	603	604	605
606	607	608	609	610
611	612	613	614	615
616	617	618	619	620
621	622	623	624	625
626	627	628	629	630
631	632	633	634	635
636	637	638	639	640
641	642	643	644	645
646	647	648	649	650
651	652	653	654	655
656	657	658	659	660
661	662	663	664	665
666	667	668	669	670
671	672	673	674	675
676	677	678	679	680
681	682	683	684	685
686	687	688	689	690
691	692	693	694	695
696	697	698	699	700
701	702	703	704	705
706	707	708	709	710
711	712	713	714	715
716	717	718	719	720
721	722	723	724	725
726	727	728	729	730
731	732	733	734	735
736	737	738	739	740
741	742	743	744	745
746	747	748	749	750
751	752	753	754	755
756	757	758	759	760
761	762	763	764	765
766	767	768	769	770
771	772	773	774	775
776	777	778	779	780
781	782	783	784	785
786	787	788	789	790
791	792	793	794	795
796	797	798	799	800
801	802	803	804	805
806	807	808	809	810
811	812	813	814	815
816	817	818	819	820
821	822	823	824	825
826	827	828	829	830
831	832	833	834	835
836	837	838	839	840
841	842	843	844	845
846	847	848	849	850
851	852	853	854	855
856	857	858	859	860
861	862	863	864	865
866	867	868	869	870
871	872	873	874	875
876	877	878	879	880
881	882	883	884	885
886	887	888	889	890
891	892	893	894	895
896	897	898	899	900
901	902	903	904	905
906	907	908	909	910
911	912	913	914	915
916	917	918	919	920
921	922	923	924	925
926	927	928	929	930
931	932	933	934	935
936	937	938	939	940
941	942	943	944	945
946	947	948	949	950
951	952	953	954	955
956	957	958	959	960
961	962	963	964	965
966	967	968	969	970
971	972	973	974	975
976	977	978	979	980
981	982	983	984	985
986	987	988	989	990
991	992	993	994	995
996	997	998	999	1000



**SECTION - A**

1. (a) Let  $T$  be the linear operator on  $C^2$  which is represented in the standard ordered basis by the matrix.

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}. \text{ Find the minimal polynomial for } T.$$

[10]

Finding out eigen values of  $A$

$$|A - \lambda I| = 0$$

$$\Rightarrow \begin{vmatrix} -\lambda & -1 \\ 1 & -\lambda \end{vmatrix} = 0$$

$$\Rightarrow \lambda^2 + 1 = 0$$

$$\Rightarrow \lambda = \pm i$$

Since  $T$  is linear operator on  $C^2$  then  $\lambda = i, -i$  are two distinct values

and hence minimal polynomial is given by

$$(x+i)(x-i) = x^2 + 1$$

88-

1. (b) If  $A$  be an  $n$ -rowed non-singular matrix,  $\mathbf{X}$  be an  $n \times 1$  matrix,  $\mathbf{B}$  be an  $n \times 1$  matrix, the system of equations  $\mathbf{AX} = \mathbf{B}$  has a unique solution. [10]

Given that  $A$  is  $n$ -rowed non-singular matrix.

This implies  $A$  is invertible matrix.

Let  $x_1, x_2$  are two solutions that satisfies  $\mathbf{AX} = \mathbf{B}$ . Then we need to prove that  $x_1 = x_2$  to show that solution is unique.

Now, we have

$$\mathbf{AX}_1 = \mathbf{B} \quad -(1)$$

$$\mathbf{AX}_2 = \mathbf{B} \quad -(2)$$

Since  $A$  is non-singular that means  $A^{-1}$  exists  
Multiplying post with  $A^{-1}$  in both the equation  
we obtain

$$(A^{-1}A)x_1 = A^{-1}\mathbf{B} \Rightarrow x_1 = A^{-1}\mathbf{B} \quad -(3)$$

$$(A^{-1}A)x_2 = A^{-1}\mathbf{B} \Rightarrow x_2 = A^{-1}\mathbf{B} \quad -(4)$$

From (3) and (4),

$$x_1 = x_2$$

Q9

Hence the system of equations  $\mathbf{AX} = \mathbf{B}$  has a unique solution.

1. (c) Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be defined by setting

$$f(x, y) = \begin{cases} \frac{x^2 y}{x^4 + y^2}, & \text{if } (x, y) \neq (0, 0), \\ 0, & \text{if } (x, y) = (0, 0). \end{cases}$$

Show that  $f$  possesses partial derivatives at  $(0, 0)$  but is not differentiable at the point. [10]

Partial derivative of  $f$  at  $(0, 0)$

$$f_x(0, 0) = \lim_{h \rightarrow 0} \frac{f(0+h, 0) - f(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{0-0}{h} = 0$$

$$f_y(0, 0) = \lim_{k \rightarrow 0} \frac{f(0, 0+k) - f(0, 0)}{k} = \lim_{k \rightarrow 0} \frac{0-0}{k} = 0$$

Thus,  $f_x(0, 0)$  and  $f_y(0, 0)$  exist

Now Assume that  $f$  is differentiable at  $(0, 0)$

then

$$f(0+h, 0+k) - f(0, 0) = Ah + BK + \sqrt{h^2+k^2} \phi(h, k)$$

where  $\lim_{(h, k) \rightarrow (0, 0)} \phi(h, k) \rightarrow 0$ .

$$\Rightarrow \frac{f(h, k)}{\sqrt{h^2+k^2}} = \phi(h, k)$$

$$\Rightarrow \lim_{\substack{(h, k) \rightarrow (0, 0)}} \phi(h, k) = \lim_{(h, k) \rightarrow (0, 0)} \frac{h^2 k}{(h^4+k^2)(h^2+k^2)^{1/2}}$$

$$= \lim_{h \rightarrow 0} \frac{h^2 (mh)}{(h^4+m^2 h^2)(h^2+m^2 h^2)^{1/2}} \quad [\text{Put } K=mh]$$

$$= \lim_{h \rightarrow 0} \frac{m}{(1+m^2 h^2)(1+m^2 h^2)^{1/2}}$$

$$= m$$

Since limit depends upon variable  $m$  and therefore  $f$  is not differentiable at  $(0, 0)$

1. (d) Show that the height of an open cylinder of given surface and greatest volume is equal to the radius of its base. [10]

Let the height of cylinder be  $h$  and radius be  $r$ .


Now

$$\text{Surface area, } S \text{ (say)} = 2\pi rh + \pi r^2$$

$$\Rightarrow dS = 2\pi r dh + 2\pi h dr + 2\pi r dr$$

$$\Rightarrow 0 = r dh + h dr + r dr$$

$$\Rightarrow \frac{dh}{dr} = - \left[ \frac{h+r}{r} \right] \quad \textcircled{1}$$



$$\text{and Volume (V say)} = \pi r^2 h$$

$$\frac{dV}{dr} = 2\pi rh + \pi r^2 \frac{dh}{dr} \quad \text{--- (2)}$$

Now for volume to be maximum/minimum

$$\frac{dr}{dr} = 0$$

$$\Rightarrow 2\pi rh + \pi r^2 \left[ -\frac{h+r}{r} \right] = 0 \quad [\text{Using } \textcircled{1}]$$

$$\Rightarrow rh - r^2 = 0$$

$$\Rightarrow r(h-r) = 0$$

$$\Rightarrow r \neq 0 \quad h = r \quad \checkmark$$

Now

$$\begin{aligned} \frac{d^2V}{dr^2} &= 2\pi h + 2\pi r \frac{dh}{dr} + 2\pi \frac{dh}{dr} + \pi r^2 \frac{d^2h}{dr^2} \\ &= 2\pi \left[ 1 + r [-2] + r [-2] + \frac{\pi r^2}{2} \right] < 0 \end{aligned}$$

Hence Volume is maximum when radius is equal to height for a given surface area.

1. (e) Show that the straight lines whose direction cosines are given by

$$al + bm + cn = 0, fm + gn + hm = 0.$$

are perpendicular if

$$f/a + g/b + h/c = 0$$

are parallel if

$$\sqrt{af} \pm \sqrt{bg} \pm \sqrt{ch} = 0.$$

[10]

Given  $al + bm + cn = 0 \quad \text{--- (1)}$

$$fm + gn + hm = 0 \quad \text{--- (2)}$$

Putting the value of  $l$  from (1) into (2)  
we get

$$fm + (gn + hm) \frac{(-bm - cn)}{a} = 0$$

$$\Rightarrow (af - bg - hc)m n - bh m^2 - cg n^2 = 0$$

$$\Rightarrow bh \left( \frac{m}{n} \right)^2 + (bg + hc - af) \frac{m}{n} + cg = 0 \quad \text{--- (3)}$$

$$\Rightarrow \frac{m_1 m_2}{n_1 n_2} = \frac{cg}{bh}$$

Similarly  $\frac{l_1 l_2}{n_1 n_2} = \frac{cf}{ah}$

For 1st lines,  $m_1 m_2 + n_1 n_2 + l_1 l_2 = 0$

$$\Rightarrow \frac{cg}{bh} + 1 + \frac{cf}{ah} = 0$$

$$\Rightarrow \boxed{\frac{f}{a} + \frac{g}{b} + \frac{h}{c} = 0} \quad [\text{on multiplying by } \frac{n}{c}]$$

shown

For II lines, direction ratio of lines would be equal.

$$a = 0 \quad \text{of (3)}$$

$$\Rightarrow (bg + hc - af)^2 - 4bh(g - c) = 0$$

$$\Rightarrow bg + hc - af = \pm 2\sqrt{bg} \sqrt{hc}$$

$$\Rightarrow bg + hc \pm 2\sqrt{bg} \sqrt{hc} = af$$

$$\Rightarrow (\sqrt{bg} \pm \sqrt{hc})^2 = (\pm \sqrt{af})^2$$

$$\Rightarrow \boxed{\pm \sqrt{bg} \pm \sqrt{hc} + \sqrt{af} = 0} \quad \text{--- shown}$$

2. (a) (i) Define the linear span,  $L(S)$ , of a subset  $S$  of a vector space  $V(F)$ . What is  $L(S)$  if (i)  $S = \{0\}$ , if (ii)  $S = \emptyset$ . Let  $V = \mathbb{R}^3$  ( $\mathbb{R}$ ),  $S = \{\alpha_1 = (1, 1, 0), \alpha_2 = (0, -1, 1), \alpha_3 = (1, 0, 1)\}$ . Prove that  $(a, b, c) \in L(S)$  if and only if  $a = b + c$ .

- (ii) Let  $W$  be the subspace of  $\mathbb{R}^4$  generated by vectors  $(1, -2, 5, -3), (2, 3, 1, -4)$  and  $(3, 8, -3, -5)$ . Find a basis and dimension of  $W$ . Extend this basis of  $W$  to a basis of  $\mathbb{R}^4$ . [17]

2 (a)  $L(S)$  is set of vectors formed by  
 (i) these vectors are independent  
 (ii) these vectors form all the vector of set  $S$ .  
 $L(S)$  if  $S = \{0\}$  is  $0$ .  
 $L(S)$  if  $S = \emptyset$  is  $\emptyset$

Now let  $(a, b, c) \in L(S)$  and let  $k_1, k_2, k_3 \in F$

$$\text{then } (a, b, c) = k_1(1, 1, 0) + k_2(0, -1, 1) + k_3(1, 0, 1)$$

$$\Rightarrow a = k_1 + k_3, b = k_1 - k_2, c = k_2 + k_3$$

$$\Rightarrow a = b + c$$

(iii) writing given vectors into matrix form

$$\left[ \begin{array}{cccc} 1 & -2 & 5 & -3 \\ 2 & 3 & 1 & -4 \\ 3 & 8 & -3 & -5 \end{array} \right] \xrightarrow{\substack{R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 3R_1}} \left[ \begin{array}{cccc} 1 & -2 & 5 & -3 \\ 0 & 7 & -9 & 2 \\ 0 & 14 & -18 & 4 \end{array} \right]$$

$$\text{R}_3 \rightarrow R_3 - 2R_2$$

$$\sim \left[ \begin{array}{cccc} 1 & -2 & 5 & -3 \\ 0 & 7 & -9 & 2 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Basis of  $W = \{(1, -2, 5, -3), (0, 7, -9, 2)\}$

and dimension of  $W$  is 2.

Extending the basis to a basis of  $\mathbb{R}^4$  gives

$$\{(1, -2, 5, -3), (0, 7, -9, 2), (0, 0, 1, 0), (0, 0, 0, 1)\}$$

$\sim (4)$

2. (b) (i) Evaluate:  $\iint_D x \sin(x+y) dx dy$ , where D is the region bounded by  $0 \leq x \leq \pi$  and  $0 \leq y \leq \frac{\pi}{2}$ .

$$0 \leq y \leq \frac{\pi}{2}.$$

(ii) If  $w = f[xy / (x^2 + y^2)]$  is a differentiable function of  $u = f[xy / (x^2 + y^2)]$  show that  $x \left( \frac{\partial w}{\partial x} \right) + y \left( \frac{\partial w}{\partial y} \right) = 0$ . [18]

(i) Put  $x+y=2v$ ,  $x=uv$ . gives  $y=u(1-v)$

$$\therefore \frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} v & u \\ 1-v & -u \end{vmatrix} = -u$$

and  $0 \leq uv \leq \pi$  and  $0 \leq u(1-v) \leq \frac{1}{2}$

$$\iint_D x [\sin(x+y)] dx dy = \int \int uv \sin u \cdot u du dv$$

??

(ii)  $w = f \left( \frac{xy}{x^2+y^2} \right)$

$$\frac{\partial w}{\partial x} = f' \left( \frac{xy}{x^2+y^2} \right) \cdot \frac{(x^2+y^2)y - xy(2x)}{(x^2+y^2)^2} = -\frac{x^2y + y^3}{(x^2+y^2)^2} f'$$

$$\frac{\partial w}{\partial y} = f' \left( \frac{xy}{x^2+y^2} \right) \cdot \frac{(x^2+y^2)x - xy(2y)}{(x^2+y^2)^2} = -\frac{xy^2 + x^3}{(x^2+y^2)^2} f'$$

NOW,  $x \frac{\partial w}{\partial x} + y \frac{\partial w}{\partial y} = f' \left[ \frac{-x^3y + y^3x - xy^3 + x^3y}{(x^2+y^2)^2} \right] = 0$

Hence

$$x \frac{\partial v}{\partial x} + y \frac{\partial v}{\partial y} = 0$$

shown

10'

2. (c) Show that the equation of the plane containing the line

$$\frac{y}{b} + \frac{z}{c} = 1, x = 0$$

and parallel to the line  $\frac{x}{a} - \frac{z}{c} = 1, y = 0$

$$\text{is } \frac{x}{a} - \frac{y}{b} - \frac{z}{c} + 1 = 0$$

and if  $2d$  is the S.D. show that  $d^2 = a^2 + b^2 + c^2$ .

[15]

Equation of plane containing the lines

$$\frac{y}{b} + \frac{z}{c} = 1, n=0 \text{ is}$$

$$\lambda x + \frac{y}{b} + \frac{z}{c} - 1 = 0 \quad \text{--- (1)}$$

Now plane (1) is parallel to line

$$\frac{x-a}{a} = \frac{y}{b} = \frac{z}{c} \quad \text{--- (2)}$$

then,

$$\lambda \cdot a + 0 \cdot y_b + c \cdot \cancel{\frac{z}{c}} = 0$$

$$\Rightarrow \lambda a = -1$$

$$\Rightarrow \lambda = -1/a$$

Hence equation of plane would be

$$\frac{x}{a} - \frac{y}{b} - \frac{z}{c} + 1 = 0$$

Now, the shortest distance between the line and plane is  $2d$

then

$$2d = \sqrt{\left| \frac{a}{a} - \frac{0}{b} - \frac{0}{c} + 1 \right|^2}$$

$$\sqrt{\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}}$$

$$\Rightarrow d^2 = \frac{1}{a^2 + b^2 + c^2}$$

$$\rightarrow d^{-2} = a^{-2} + b^{-2} + c^{-2}$$

shown

13


3. (a) (i) Show that the matrix

$$A = \begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

satisfies Cayley-Hamilton theorem. Hence or otherwise obtain the value of  $A^{-1}$  and  $A^{-2}$ .

- (ii) Consider the linear transformation  $T : \mathbf{R}^3 \rightarrow \mathbf{R}^2$ , defined by  $T(x, y, z) = (x + y, 2z)$ . Find the matrix of  $T$  with respect to the bases  $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$  and  $\{\mathbf{u}'_1, \mathbf{u}'_2\}$  of  $\mathbf{R}^3$  and  $\mathbf{R}^2$ , where

$\mathbf{u}_1 = (1, 1, 0)$ ,  $\mathbf{u}_2 = (0, 1, 4)$ ,  $\mathbf{u}_3 = (1, 2, 3)$  and  $\mathbf{u}'_1 = (1, 0)$ ,  $\mathbf{u}'_2 = (0, 2)$ . Use this matrix to find the image of the vector  $\mathbf{u} = (2, 3, 5)$ . [20]



## SECTION - B

5. (a) Solve  $(xy^2 - x^2) dx + (3x^2y^2 + x^2y - 2x^3 + y^2) dy = 0.$

[10]

This is of the form  $M dx + N dy = 0$

where  $M = xy^2 - x^2$  and  $N = 3x^2y^2 + x^2y - 2x^3 + y^2$

$$\text{Here } \frac{1}{M} \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) = \frac{1}{xy^2 - x^2} [6xy^2 + 2xy - 6x^2 - 2xy] = 6$$

$$\text{thus I.F.} = e^{\int 6 dy} = e^{6y}$$


Thus, multiplying by I.F.

$$\int e^{6y} (xy^2 - x^2) dx + \int e^{6y} (y^2) dy = C$$

~~keeping y constant. term not containing x~~ where C is integration constant

$$\Rightarrow e^{6y} \left( \frac{x^2y^2}{2} - \frac{x^3}{3} \right) + \frac{y^2 e^{6y}}{6} - \frac{1}{6} \int 2ye^{6y} = C$$

$$\Rightarrow e^{6y} \left( \frac{x^2y^2}{2} - \frac{x^3}{3} \right) + \frac{e^{6y} y^2}{6} - \frac{1}{18} ye^{6y} + \frac{1}{108} e^{6y} = C \quad \text{Ans}$$



5. (b) Find the orthogonal trajectories of the family of curves  $r = c(\cos \theta + \sin \theta)$ , where  $c$  is the parameter. [10]

$$\text{Given } r = c(\cos \theta + \sin \theta)$$

Taking log both sides

$$\log r = \log c + \log(\cos \theta + \sin \theta)$$

Differentiating the equation w.r.t.  $\theta$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{-\sin \theta + \cos \theta}{\cos \theta + \sin \theta}$$

For orthogonal trajectory, replacing  $\frac{dr}{d\theta}$  by  $-\frac{1}{r^2} \frac{d\theta}{dr}$

$$\Rightarrow \frac{1}{r} \left( -r^2 \frac{d\theta}{dr} \right) = \frac{-\sin \theta + \cos \theta}{\cos \theta + \sin \theta}$$

$$\Rightarrow \frac{\cos \theta + \sin \theta}{\sin \theta - \cos \theta} d\theta = \frac{dr}{r}$$

Integrating,

$$\log(\sin\theta - \cos\theta) = \log r + \log C \quad \text{where } \log C \text{ is any arbitrary constant}$$

$$\Rightarrow r = C(\sin\theta - \cos\theta)$$


This is required orthogonal trajectory.

-09-

5. (c) A particle of mass  $m$ , is falling under the influence of gravity through a medium whose resistance equals  $\mu$  times the velocity. If the particle were released from rest, determine the distance fallen through in time  $t$ . [10]

According to question,

$$\begin{aligned} \frac{d^2x}{dt^2} &= mg - \mu v & \dots (1) \\ \Rightarrow v \frac{dy}{dx} &= mg - \mu v \quad [\because \frac{dv}{dt} = \frac{dy}{dx} \cdot \frac{dx}{dt}] \\ \Rightarrow \frac{v \frac{dy}{dx}}{mg - \mu v} &= dx \end{aligned}$$



$$\Rightarrow \frac{1}{\mu} \left( \frac{mg}{mg - \mu v} - 1 \right) dy = dx$$

Integrating

$$\Rightarrow -\frac{mg}{\mu^2} \log |mg - \mu v| - \frac{v}{\mu} = x + C \quad \dots (2)$$

Now at  $v=0, x=0$

$$(2) \text{ gives } -\frac{mg}{\mu^2} \log mg = C \quad \dots (3)$$

Using (3) in (2)

$$x = \frac{mg}{\mu^2} \log \frac{mg}{mg-\mu v} - \frac{v}{\mu} \quad (4)$$

(i) Can be written as

$$\frac{dv}{dt} = mg - \mu v \Rightarrow \frac{dv}{mg - \mu v} = dt$$

Integrating,  $-\frac{1}{\mu} \log(mg - \mu v) = t + C_1$  where  $C_1$  is const  
— (5)

Now at  $t=0, v=0$

$$C_1 = -\frac{1}{\mu} \log mg$$

Putting in (5)

$$\frac{1}{\mu} \log \frac{mg}{mg - \mu v} = t \Rightarrow v = \frac{mg}{\mu} (1 - e^{-\mu t}) \quad (6)$$

Using (6) in (4)

~~$$x = \frac{mg}{\mu} t - \frac{mg}{\mu^2} (1 - e^{-\mu t})$$~~

~~$$r = \frac{mg}{\mu} [ \mu t - 1 + e^{-\mu t} ]$$~~

5. (d) A particle moves so that its position vector is given by  $\mathbf{r} = \cos \omega t \mathbf{i} + \sin \omega t \mathbf{j}$  where  $\omega$  is a constant; show that (i) the velocity of the particle is perpendicular to  $\mathbf{r}$ , (ii) the acceleration is directed towards the origin and has magnitude proportional to the distance from the origin, (iii)  $\mathbf{r} \times \frac{d\mathbf{r}}{dt}$  is a constant vector. [10]

i)  $\mathbf{r} = \cos \omega t \mathbf{i} + \sin \omega t \mathbf{j}$

$$\mathbf{v} = \dot{\mathbf{r}} = \omega(-\sin \omega t \mathbf{i} + \cos \omega t \mathbf{j})$$

Now,  $\mathbf{v} \cdot \mathbf{r} = -\cos \omega t \sin \omega t + \cos \omega t \sin \omega t = 0$

$\Rightarrow \mathbf{v}$  and  $\mathbf{r}$  are perpendicular

ii)  $\mathbf{a} = \ddot{\mathbf{r}} = \omega(-\cos \omega t \mathbf{i} - \sin \omega t \mathbf{j}) = -\omega^2 \mathbf{r}$

clearly acceleration is directed towards origin and mag. proportional to distance from origin

(iii)

$$\mathbf{r} \times \frac{d\mathbf{r}}{dt} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \cos\omega t & \sin\omega t & 0 \\ -\sin\omega t & \cos\omega t & 0 \end{vmatrix}$$

$$= \hat{i}(0-0) + \hat{j}(0-0) + \hat{k}(\omega \cos^2\omega t + \omega^2 \sin\omega t)$$

$$= \underline{\omega} \text{ (a constant vector)}$$

- 89 -

5. (e) (i) Prove that  $\mathbf{r}^n \mathbf{r}$  is an irrotational vector for any value of n but is solenoidal only if  $n+3=0$ .  
(ii) If  $\mathbf{a}$  is constant vector, prove that  
 $\operatorname{div}\{\mathbf{r}^n(\mathbf{a} \times \mathbf{r})\} = 0$ .

[10]

(i) For irrotational vector  $\nabla \times \mathbf{F} = 0$  where  
 $\mathbf{F}$  is any vector

$$\begin{aligned} \nabla \times (\mathbf{r}^n \vec{r}) &= \cancel{\frac{\partial}{\partial r}} \nabla \mathbf{r}^n \times \vec{r} + \mathbf{r}^n (\nabla \times \vec{r}) \quad [\nabla \times (\phi \mathbf{F}) \\ &= n \mathbf{r}^{n-1} \cancel{\frac{\partial}{\partial r}} \vec{r} \times \vec{r} + \mathbf{r}^n (0) \quad = \nabla \phi \times \mathbf{F} + \\ &= n \mathbf{r}^{n-2} (0) + \mathbf{r}^n (0) \quad \phi (\nabla \times \mathbf{F}) \\ &= 0 \end{aligned}$$

Hence  $\mathbf{r}^n \vec{r}$  is ~~irrotational~~ for any value of  $n$

NOW for solenoidal,

$$\nabla \cdot \mathbf{v} = 0$$

$$\begin{aligned} \text{Now, } \nabla \cdot (r^n \hat{r}) &= \nabla \cdot r^n \hat{r} + r^n \nabla \cdot \hat{r} \quad [\because \nabla \cdot (\phi F) \\ &= n r^{n-1} \frac{\hat{r}}{r} + r^n (3) \quad = \nabla \phi \cdot F + \phi \nabla \cdot F] \\ &= n r^{n-2+2} + r^n (3) \\ &= (n+3) r^n \end{aligned}$$

Now, it is clear that  $r^n \hat{r}$  is solenoidal when

$$n+3=0$$

$$\begin{aligned} (\text{iii}) \operatorname{div} \{ r^n (axr) \} &= \nabla \cdot r^n \cdot (axr) + r^n \nabla \cdot (axr) \quad [\because \nabla \cdot (\phi F) = \nabla \phi \cdot F] \\ &= n r^{n-1} \frac{\hat{r}}{r} [axr] + r^n \nabla \cdot \left( \sum i (a_2 z - a_3 y) \right) \quad [\because a = a_1 + a_2 j + a_3 k] \\ &= 0 + r^n (0) \end{aligned}$$

✓ 08

$$\text{Hence } \operatorname{div} [r^n (axr)] = 0$$

6. (a) Solve the differential equation  $y = x - 2ap + ap^2$ . Find the singular solution and interpret it geometrically. [13]

$$\text{Given } y = x - 2ap + ap^2 \quad \text{--- (1)}$$

$$\frac{dy}{dx} = 1 - 2a \frac{dp}{dx} + 2ap \frac{d^2p}{dx^2}$$

$$\Rightarrow p = 1 - 2a(1-p) \frac{dp}{dx}$$

$$\Rightarrow 0 = (1-p) \left[ 1 - 2a \frac{dp}{dx} \right]$$

$$\Rightarrow 1 - 2a \frac{dp}{dx} = 0$$

$$\Rightarrow x - 2ap = c \quad \text{where } c \text{ is any constant}$$

$$\Rightarrow p = \frac{x-c}{2a} \quad \text{--- (2)}$$

Put it in (1),

$$y = x - 2a \frac{x-c}{2a} + \left( \frac{x-c}{2a} \right)^2 - \frac{c^2}{4a} \quad \text{--- (3)} \quad y = x - (x+c) + \frac{(x+c)^2}{4a}$$

$$\Rightarrow y = -c + \frac{(x+c)^2}{4a} - \frac{c^2}{4a} \quad \text{--- (3)}$$

Now p-discriminant from  $\alpha$

$$\text{is } 4a^2 - 4a(x-y) = 0 \Rightarrow a-x+y=0$$

and c-discriminant gives

$$a-x+y=0$$

and it also satisfies  $\alpha$

Hence  $a=x-y \neq 0$  is unique solution

Geometric interpretation, the line  $(a-x+y=0)$   
touches the family of curves given by  
forementioned differential equations,

12

5. (e) (i) Prove that  $r^n \mathbf{r}$  is an irrotational vector for any value of n but is solenoidal only if  $n+3=0$ .  
(ii) If  $\mathbf{a}$  is constant vector, prove that  
 $\operatorname{div} \{r^n(\mathbf{a} \times \mathbf{r})\} = 0$ .

[10]



6. (b) Solve  $(d^4y/dx^4) + 6(d^3y/dx^3) + 11(d^2y/dx^2) + 6(dy/dx) = 20 e^{-2x} \sin x$ .

[10]

Given

$$(D^4 + 6D^3 + 11D^2 + 6D) y = 20e^{-2x} \sin x \quad \text{--- (1)}$$

C.F. of (1) is given by

$$D(D^3 + 6D^2 + 11D + 6) = 0$$

$$\Rightarrow D=0, -1, -2, -3$$

$$\text{C.F.} = C_1 e^0 + C_2 e^{-x} + C_3 e^{-2x} + C_4 e^{-3x} \quad \text{where } C_1, C_2, C_3, C_4 \text{ are constants}$$

and

$$\text{P.I.} = 20 \frac{1}{D(D+1)(D+2)(D+3)} e^{-2x} \sin x$$

$$= 20e^{-2x} \frac{1}{(D-2)(D-1)D(D+1)} \sin x$$

$$= 20e^{-2x} \frac{1}{(D-2)(D-1)D(D+1)} \sin x$$

$$= 20e^{-2x} \frac{1}{D^4 - 2D^3 - D^2 + 2D} \sin x$$

$$= 20e^{-2x} \frac{1}{(-1)^2 - 2D(-1) + 1 + 2D} \sin x$$

$$= 20e^{-2x} \frac{1}{2 + 4D} \sin x$$

$$= 10e^{-2x} \frac{(1+2D)}{1-4D^2} \sin x$$

$$= \frac{10e^{-2x}}{5} [ \sin x - 2 \cos x ] = 2e^{-2x} [ \sin x - 2 \cos x ]$$

$$y = (\text{C.F.} + \text{P.I.}) = C_1 e^0 + C_2 e^{-x} + C_3 e^{-2x} + C_4 e^{-3x}$$

$$+ 2e^{-2x} [ \sin x - 2 \cos x ] \text{ due.}$$

6. (b) Solve  $(d^4y/dx^4) + 6(d^3y/dx^3) + 11(d^2y/dx^2) + 6(dy/dx) = 20 e^{-2x} \sin x.$  [10]

6. (c) Solve  $y'' + 3y' + 2y = x + \cos x$  by the method of variation of parameters.

[12]

Given  $y'' + 3y' + 2y = x + \cos x \quad \text{--- (1)}$

Taking C.F. of (1)

$$(D^2 + 3D + 2)y = 0$$

$$\Rightarrow (D+2)(D+1)y = 0$$

$$\text{C.F. of (1)} = C_1 e^{-2x} + C_2 e^{-x} \quad \text{where } C_1 \text{ and } C_2 \text{ are constants.}$$

let  $u = e^{-2x}$  and  $v = e^{-x}$

P.I. of (1) would be given by.

$$u f(x) + v g(x)$$

where

$$\begin{aligned} f(x) &= \int -\frac{vR}{w} dx = \int \frac{-e^{-x}(x + \cos x)}{uv_1 - u_1 v} dx \\ &= \int \frac{-e^{-x}(x + \cos x)}{e^{-3x}} dx \\ &= \int -e^{2x}(x + \cos x) dx \\ &= \int -xe^{2x} dx - \int e^{2x} \cos x dx \\ &= -\frac{x e^{2x}}{2} + \frac{e^{2x}}{4} - \frac{e^{2x}}{5}(2 \cos x + 8 \sin x) \end{aligned}$$

and  $g(x) = \int \frac{uR}{w} dx = \int \frac{e^{-2x}(x + \cos x)}{e^{-3x}} dx$

$$= \int xe^{2x} dx + \int e^{2x} \cos x dx$$

$$= \frac{xe^{2x}}{2} - \frac{e^{2x}}{2} + \frac{e^{2x}}{2}(2 \cos x + 8 \sin x)$$

thus,

$$y = C_1 e^{-2x} + C_2 e^{-x} + \left[ -\frac{x}{2} + \frac{1}{2} - \frac{1}{5}(2 \cos x + 8 \sin x) \right]$$

$$+ \left[ x - 1 + \frac{1}{2}(2 \cos x + 8 \sin x) \right]$$

$$\therefore y = C_1 e^{-2x} + C_2 e^{-x} + \frac{x}{2} - \frac{3}{10} + \frac{1}{10} \cos x + \frac{3}{10} \sin x$$

6. (d) (i) Show that  $\int_0^\infty \frac{\sin t}{t} dt = \frac{\pi}{2}$  by using Laplace transform method.

(ii) By using Laplace Transform Method solve the initial value problem  $(D^3 - D^2 - D + 1)y = 8t e^{-t}$

$y = D^2 y = 0$ ,  $Dy = 1$  when  $t = 0$ .

[15]

$$(i) \int_0^\infty e^{-st} \frac{\sin t}{t} dt = \frac{1}{s^2+1} = \mathcal{L}\{\sin t\}$$

then  $\mathcal{L}\left(\frac{\sin t}{t}\right) = \int_s^\infty \frac{1}{u^2+1} du = \left[\tan^{-1} u\right]_s^\infty = \frac{\pi}{2} - \tan^{-1}s$

now, put  $s=0$

$$\int_0^\infty e^{(0)t} \frac{\sin t}{t} dt = \frac{\pi}{2} - \tan^0$$

$$\Rightarrow \boxed{\int_0^\infty \frac{\sin t}{t} dt = \frac{\pi}{2}}$$

shown

(ii) Given

$$(s^3 - s^2 - s + 1) y = 8te^{-t}$$

Taking Laplace transform

$$\mathcal{L}\{s^3 y\} - \mathcal{L}\{s^2 y\} - \mathcal{L}\{sy\} + \mathcal{L}\{y\} = \mathcal{L}\{8te^{-t}\}$$

$$\Rightarrow [s^3 F(s) - s^2 f(0) - [s^2 F(s) - sf(0) - f'(0)] - [s F(s) - f(0)] + f(s) = \frac{8}{(s+1)^2}$$

$$\Rightarrow -sf'(0) - f''(0)$$

$$\Rightarrow [s^3 - s^2 - s + 1] F(s) - s + 1 = \frac{8}{(s+1)^2} \quad [\text{let } \mathcal{L}\{y\} = f(s)]$$

$$\Rightarrow (s^3 - s^2 - s + 1) F(s) = s - 1 + \frac{8}{(s+1)^2}$$

$$\Rightarrow F(s) = \frac{1}{(s^2 - 1)} + \frac{8}{(s+1)^2 (s-1)(s^2 - 1)}$$

$$\Rightarrow F(s) = \frac{1}{s^2 - 1} + \frac{8}{(s+1)^3 (s-1)^2}$$

Taking Laplace inverse

$$\Rightarrow y(t) = \mathcal{L}^{-1} \left\{ -\frac{1}{2(s+1)} + \frac{1}{2(s-1)} \right\} + \mathcal{L}^{-1} \left\{ \frac{8}{(s+1)^3 (s-1)^2} \right\}$$

$$\Rightarrow y(t) = -\frac{1}{2} e^{-t} + \frac{1}{2} e^t + ?$$

Please  
complete  
it.

11

7. (a) A uniform beam of length  $2a$  rests with its ends on two smooth planes which intersect in a horizontal line. If the inclinations of the planes to the horizontal are  $\alpha$  and  $\beta$  ( $\alpha > \beta$ ), show that the inclination  $\theta$  of the beam to the horizontal in one of the equilibrium positions is given by

$$\tan \theta = \frac{1}{2}(\cot \beta - \cot \alpha)$$

and show that the beam is unstable in this position.

[15]

8. (a) Find the curvature and torsion of the circular helix  $x = a \cos \theta$ ,  $y = a \sin \theta$ ,  $z = a\theta \cot \alpha$ . [12]

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\Rightarrow \vec{r} = a \cos \theta \hat{i} + a \sin \theta \hat{j} + a\theta \cot \alpha \hat{k}$$

$$\Rightarrow \dot{\vec{r}} (= v) = -a \sin \theta \hat{i} + a \cos \theta \hat{j} + a \cot \alpha \hat{k}$$

$$\Rightarrow \ddot{\vec{r}} (= a) = -a \cos \theta \hat{i} - a \sin \theta \hat{j}$$

$$\Rightarrow \dddot{\vec{r}} (= a') = +a \sin \theta \hat{i} - a \cos \theta \hat{j}$$

and  $\dot{\vec{r}} \times \ddot{\vec{r}} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -a \sin \theta & a \cos \theta & a \cot \alpha \\ a \cos \theta & -a \sin \theta & 0 \end{vmatrix} = \hat{i}(a^2 \sin \theta \cot \alpha) + \hat{j}(-a^2 \cos^2 \theta) + \hat{k}(a^2)$

Now,

$$K (\text{Curvature}) = \frac{|\dot{\vec{r}} \times \ddot{\vec{r}}|}{|\dot{\vec{r}}|^3}$$

$$= \frac{\sqrt{(a^2 \sin \theta \cot \alpha)^2 + (a^2 \cos^2 \theta)^2 + (a)^2}}{(a^2 \sin^2 \theta + a^2 \cos^2 \theta + a^2 \cot^2 \alpha)^{3/2}}$$

$$= \frac{\sqrt{a^4 \cot^2 \alpha + a^4}}{(a^2 + a^2 \cot^2 \alpha)^{3/2}}$$

$$= \frac{1}{a} \frac{\cosec \alpha}{\cosec^3 \alpha}$$

$$K = \frac{\sin^2 \alpha}{a}$$

and  $T (\text{Torsion}) = \frac{(\dot{\vec{r}} \times \ddot{\vec{r}}) \cdot \ddot{\vec{r}}}{|\dot{\vec{r}} \times \ddot{\vec{r}}|^2}$

$$= \frac{a^3 \sin^2 \theta \cot \alpha + a^3 \cos^2 \theta \cot \alpha}{a^4 \cosec^2 \alpha}$$

$$= \frac{a^3 \cot \alpha}{a^4 \cosec^2 \alpha}$$

$$T = \frac{\cosec \alpha \sin \alpha}{a} = \frac{\sin^2 \alpha}{2a} \text{ rad.}$$

8. (b) Evaluate  $\oint_C (x^2 + y^2) dx + 3xy^2 dy$

where C, a circle of radius two with centre at the origin of the xy plane, is traversed in the positive sense. [10]

$$C = x^2 + y^2 = 4$$

$\oint_C (x^2 + y^2) dx + 3xy^2 dy$  - This is of the form  $M dx + N dy$

Applying Green's theorem

$$\oint_C (x^2 + y^2) dx + 3xy^2 dy = \iint_D (3y^2 - 2y) dxdy$$

Put  $x = r\cos\theta, y = r\sin\theta$

$$\begin{aligned}
 &= \iint_{D} 3r^2 \cos^2\theta \cdot r \cdot r^2 \sin^2\theta \, drd\theta \quad [\iint_D y \, dxdy = 0 \text{ (odd)}] \\
 &= \frac{3}{4} \left[ r^4 \right]_{0}^{2} \int_{0}^{2\pi} \cos^2\theta \, d\theta \quad \text{if } 1 \text{ is odd} \\
 &= \frac{3}{4} \cdot [16] \cdot \frac{1}{2} \cdot 2\pi
 \end{aligned}$$

8. (d) Verify Stoke's theorem for the vector

$A = 3y\mathbf{i} - xz\mathbf{j} + yz^2\mathbf{k}$ , where S is the surface of the paraboloid  $2z = x^2 + y^2$  bounded by  $z = 2$  and C is its boundary. [15]

Stoke's theorem states

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S \text{curl } \mathbf{A} \cdot \mathbf{n} dS$$

$$\oint_C \mathbf{A} \cdot d\mathbf{r} = \iint_S (3y\mathbf{i} - xz\mathbf{j} + yz^2\mathbf{k}) \cdot (dx\mathbf{i} + dy\mathbf{j} + dz\mathbf{k})$$

Now  $C \equiv 4 = x^2 + y^2$

Use  $x = 2\cos\theta, y = 2\sin\theta, z=2$

$$= - \int_0^{2\pi} 6\sin\theta (-2\sin\theta) - (4\cos\theta)(2\cos\theta) d\theta$$

$$= - \int_0^{2\pi} -12\sin^2\theta d\theta - 8\cos^2\theta d\theta.$$

$$= -4 \left[ - \int_0^{1/2} 12\sin^2\theta d\theta - 8 \int_0^{1/2} \cos^2\theta d\theta \right]$$

$$= -4 \left[ -12 \cdot \frac{\pi}{4} - 8 \cdot \frac{\pi}{4} \right]$$

$$= +20\pi$$

[Integration would be clockwise]

Now,

$$\text{curl } \mathbf{A} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 3y & -xz & yz^2 \end{vmatrix}$$

$$= (z^2 + x)\mathbf{i} + (0 - 0)\mathbf{j} + (-z - 3)\mathbf{k}$$

$$= (z^2 + x)\mathbf{i} - (z + 3)\mathbf{k}$$

$$\text{Now, } \text{div}(\text{curl } \mathbf{A}) = 1 - 1 = 0$$

By Gauss theorem,

$$\iint_S \operatorname{curl} A \cdot \hat{n} dS + \iint_{C'} \operatorname{curl} A \cdot \hat{n} dS = 0$$

$$\Rightarrow \iint_S \operatorname{curl} A \cdot \hat{n} dS = - \iint_{C'} (\operatorname{curl} A) \cdot \hat{r} dS$$

$$= + \iint_{C'} (z+3) dx dy$$

$$= S \iint_{C'} dz dy$$

$$= S \cdot (\pi \cdot \frac{R^2}{2})$$

$$= 20\pi$$

Hence  $\oint_C F dr = \iint_S \operatorname{curl} A \cdot \hat{n} dS = 20\pi$

verified

## ROUGH SPACE