

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

1(a) Let  $H$  be a subgroup of group  $G$ . Then  $W = \bigcap_{g \in G} gHg^{-1}$  is a normal subgroup of  $G$ .

Sol'n: Let  $a = ghg^{-1}$  and  $b = gh_1g^{-1}$  be two elements of  $K$ . Then

$$\begin{aligned} ab^{-1} &= ghg^{-1}(gh_1g^{-1})^{-1} \\ &= ghg^{-1}((g^{-1})^{-1}h_1^{-1}g^{-1}) \\ &= ghg^{-1}gh_1^{-1}g^{-1} \\ &= gh_1^{-1}g^{-1} \end{aligned}$$

$\left[ gHg^{-1} = \{ghg^{-1} / h \in H\} \right]$

Now  $h, h_1 \in H$  and  $H$  is a subgroup of  $G$ . Hence  $hh_1^{-1} \in H$ . Then from (\*) above

$$ab^{-1} = g(hh_1^{-1})g^{-1} \in gHg^{-1}$$

Hence  $gHg^{-1}$  is a subgroup of  $G$  for all  $g \in G$ .

Since the intersection of subgroups is a subgroup,  $W$  is a subgroup of  $G$ . Let  $x \in G, w \in W$ . Then  $w \in gHg^{-1} \forall g \in G$ . We have to show that  $xw^{-1} \in gHg^{-1} \forall g \in G$ , which in turn will yield that  $xw^{-1} \in W$ .

Let  $g \in G$  and let us suppose that  $xw^{-1} \notin gHg^{-1}$ .

Then  $xw^{-1} = ghg^{-1}$  for some  $h \in H$ .

thus  $g^{-1}xw^{-1}g = h \in H$ .

$$\Rightarrow (g^{-1}x)w(g^{-1}x)^{-1} \in H$$

Put  $y = x^{-1}g$ . Then  $g = xy$ . Hence in order to show that

$xw^{-1} \in gHg^{-1}$  for a given  $g \in G$ ,

first we need to find  $y \in G$  such that  $g = xy$ .

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

Since  $g = x(x^{-1}g)$ ,  
 we can choose  $y = x^{-1}g$ .  
 So there exists  $y \in G$  such that  $g = xy$ .  
 Since  $y \in G$ , we have  $w \in yHy^{-1}$  and  $w = yhy^{-1}$  for some  $h \in H$ .  
 Hence  $xwx^{-1} = x(yhy^{-1})x^{-1}$   
 $= xyhy^{-1}x^{-1}$   
 $= (xy)h(xy)^{-1}$   
 $= ghg^{-1} \in gHg^{-1}$

Since  $g \in G$  was arbitrary,  
 $xwx^{-1} \in gHg^{-1}$  for all  $g \in G$ .

Thus,  $H$  is a normal subgroup of  $G$ .

---

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

1.(b)

Let  $R = \left\{ \begin{bmatrix} a & b \\ b & a \end{bmatrix} \mid a, b \in \mathbb{Z} \right\}$  and let  $\phi$  be the mapping that takes  $\begin{bmatrix} a & b \\ b & a \end{bmatrix}$  to  $a-b$ .

- Show that  $\phi$  is a homomorphism.
- Determine the kernel of  $\phi$ .
- Show that  $R/\ker\phi$  is isomorphic to  $\mathbb{Z}$ .

Soln: Let  $\phi: R \rightarrow \mathbb{Z}$  such that

$$\phi \begin{pmatrix} a & b \\ b & a \end{pmatrix} = a-b.$$

we have

$$\begin{aligned} \phi \left( \begin{bmatrix} a & b \\ b & a \end{bmatrix} + \begin{bmatrix} a_1 & b_1 \\ b_1 & a_1 \end{bmatrix} \right) &= \phi \begin{bmatrix} a+a_1 & b+b_1 \\ b+b_1 & a+a_1 \end{bmatrix} \\ &= (a+a_1) - (b+b_1) \\ &= (a-b) + (a_1 - b_1) \\ &= \phi \begin{pmatrix} a & b \\ b & a \end{pmatrix} + \phi \begin{pmatrix} a_1 & b_1 \\ b_1 & a_1 \end{pmatrix} \end{aligned}$$

$\therefore \phi$  is homomorphism.

Now  $\ker\phi = \left\{ \begin{bmatrix} a & b \\ b & a \end{bmatrix} \mid \phi \left( \begin{bmatrix} a & b \\ b & a \end{bmatrix} \right) = 0 \right\}$

since  $\phi \begin{pmatrix} a & b \\ b & a \end{pmatrix} = a-b$

$$\begin{aligned} \Rightarrow a-b &= 0 \\ \Rightarrow a &= b \end{aligned}$$

$\therefore \ker\phi = \left\{ \begin{pmatrix} a & a \\ a & a \end{pmatrix} \mid a \in \mathbb{Z} \right\}$ .

for each  $a \in \mathbb{Z} \exists A \in R$  such that-

$$\phi(A) = a$$

$\therefore \phi$  is onto.

$\therefore$  By Fundamental theorem of Homomorphism.

$$\frac{R}{\ker\phi} \cong \mathbb{Z}$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

1(c) Let  $f(x) = \frac{n}{n+2}$  if  $\frac{1}{n+2} \leq x \leq \frac{1}{n}$ , where  $n=1, 2, 3, \dots$ .

and  $f(0)=0$ . Prove that  $f$  is Riemann integrable in  $[0, 1]$

$$\underline{\text{Sol'n}} : f(x) = \begin{cases} \frac{n}{n+2}, & \frac{1}{n+2} \leq x \leq \frac{1}{n}, n=1, 2, \dots \\ 0, & x=0 \end{cases}$$

$$= \frac{1}{3}; \frac{1}{3} \leq x \leq 1$$

$$= \frac{1}{4}; \frac{1}{4} \leq x \leq \frac{1}{2}$$

$$= \frac{1}{5}; \frac{1}{5} \leq x \leq \frac{1}{3}$$

$$= \frac{1}{6}; \frac{1}{6} \leq x \leq \frac{1}{4}$$

⋮

$$= \frac{n-1}{n+1}; \frac{1}{n+1} \leq x \leq \frac{1}{n-1}$$

$$= 0, x=0.$$

$\Rightarrow f(x)$  is bounded and continuous on  $[0, 1]$

except at the points  $\frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \dots$

The set of points of discontinuity of  $f$  on  $[0, 1]$   
 is  $\{\frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \dots\}$  which has one limit point '0'.

The set of points of discontinuity of  $f$  on  
 $[0, 1]$  has a finite no. of limit points.

$\therefore f$  is integrable on  $[0, 1]$



**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

1.(c)

Use Cauchy's theorem and/or Cauchy integral formula to evaluate the following integrals

(i)  $\int_{|z|=4} \frac{z^4}{(z-1)^3} dz$       (ii)  $\int_{|z-1|=4} \frac{z^4}{z-1} dz.$

Soln: (i) Given that  $\int_{|z|=4} \frac{z^4}{(z-1)^3} dz.$

Here  $f(z) = z^4$  is analytic in  $|z| = 4$ .  
 Comparing the given integral with  $\int_{|z|=4} \frac{f(z)}{(z-z_0)^n} dz$   
 since  $f(z) = z^4$  is analytic in  $|z| = 4$ .  
 $\therefore$  we can apply the Cauchy's integral formula

$$\int_{|z|=4} \frac{f(z)}{(z-z_0)^3} dz = \frac{2\pi i}{2!} f''(z_0)$$

$$f(z) = z^4 \Rightarrow f'(z) = 4z^3 - \\ f''(z) = 12z^2$$

$$\int_{|z|=4} \frac{z^4}{(z-1)^3} dz = \frac{2\pi i}{2!} f''(z_0) \\ = \frac{2\pi i}{2!} (12)(3)! \quad (\because z_0 = 1) \\ = -12\pi i.$$

—————

(ii) Given

$$\int_{|z-1-i|=4} \frac{z^2}{z-1} dz.$$

We have  $|z-1-i| = 4$  is a circle with centre at  $z = 1+i$  and radius  $4/4 = 1$ .

i.e., at the centre  $(1, 1)$ .

$$\text{Also } z-1 = 0$$

$$\Rightarrow z = 1$$

$$\Rightarrow z = (1, 0) \quad z = 1 \text{ lies inside } C.$$



Let  $f(z) = z^2$ , which is clearly analytic at every point within  $C$  and on  $C$ .

$$\begin{aligned} \therefore \int \frac{f(z)}{z-1} dz &= \int \frac{f(z)}{z-z_0} dz, \text{ where } z_0 = 1 \\ &= 2\pi i f(z_0) \quad (\text{By Cauchy's integral formula}) \\ &= 2\pi i (3)^2 \\ &= 18\pi i. \end{aligned}$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

1(e) Give the dual of the LP problem:  $\min z = 2x_1 + 3x_2 + 5x_3$   
 Subject to the constraints:  $2x_1 + 3x_2 + 5x_3 \geq 2$ ,  
 $3x_1 + x_2 + 7x_3 = 3$ ,  $x_1 + 4x_2 + 6x_3 \leq 5$ ,  $x_1, x_2 \geq 0$  and  
 $x_3$  is unrestricted.

Soln: Since the variable  $x_3$  is unrestricted in sign the given LP problem can be transformed into standard primal form by substituting  $x_3 = x_3' - x_3''$ , where  $x_3' \geq 0$ ,  $x_3'' \geq 0$ . Therefore standard primal becomes:

$$\text{Max } z' = -2x_1 - 3x_2 - 4(x_3' - x_3'')$$

Subjected to constraints:

$$-2x_1 - 3x_2 - 5(x_3' - x_3'') \leq -2$$

$$3x_1 + x_2 + 7(x_3' - x_3'') \leq 3$$

$$-3x_1 - x_2 - 7(x_3' - x_3'') \leq -3$$

$$x_1 + 4x_2 + 6(x_3' - x_3'') \leq 5$$

$$x_1, x_2, x_3', x_3'' \geq 0.$$

The dual of the given standard primal is,

$$\min z' = -2w_1 + 3(w_2' - w_2'') + 5w_3$$

Subject to the constraints:

$$-2w_1 + 3(w_2' - w_2'') + w_3 \geq -2$$

$$-3w_1 + (w_2' - w_2'') + 4w_3 \geq -3$$

$$-5w_1 + 7(w_2' - w_2'') + 6w_3 \geq -4$$

$$5w_1 - 7(w_2' - w_2'') - 6w_3 \geq 4$$

$$w_1, w_2', w_2'', w_3 \geq 0$$

Again we may write

$$\min z' = -2w_1 + 3w_2 + 5w_3$$

$$-2w_1 + 3w_2 + w_3 \geq -2$$

$$-3w_1 + w_2 + 4w_3 \geq -3$$

$$5w_1 - 7w_2 - 6w_3 = 4$$

$$w_1, w_3 \geq 0 \text{ and}$$

$w_2$  is unrestricted.

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

2. (a) (i) Let  $M$  be the set of all  $3 \times 3$  matrices of the following form:

$$\begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ b & c & a \end{bmatrix}$$

where  $a, b, c \in \mathbb{Z}_2$ . Show that with standard matrix addition and multiplication (over  $\mathbb{Z}_2$ ),  $M$  is a commutative ring. Find all the idempotent elements of  $M$ .

Solution:- Given Matrix  $M = \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ b & c & a \end{bmatrix}; a, b, c \in \mathbb{Z}_2$

Let  $A_1 = \begin{bmatrix} a_1 & 0 & 0 \\ 0 & a_1 & 0 \\ b_1 & c_1 & a_1 \end{bmatrix}$  and  $A_2 = \begin{bmatrix} a_2 & 0 & 0 \\ 0 & a_2 & 0 \\ b_2 & c_2 & a_2 \end{bmatrix}$   $a_1, b_1, c_1 \in \mathbb{Z}_2$   $a_2, b_2, c_2 \in \mathbb{Z}_2$

For Matrix Addition

$$A_1 + A_2 = \begin{bmatrix} a_1 & 0 & 0 \\ 0 & a_1 & 0 \\ b_1 & c_1 & a_1 \end{bmatrix} + \begin{bmatrix} a_2 & 0 & 0 \\ 0 & a_2 & 0 \\ b_2 & c_2 & a_2 \end{bmatrix} = \begin{bmatrix} a_1 + a_2 & 0 & 0 \\ 0 & a_1 + a_2 & 0 \\ b_1 + b_2 & c_1 + c_2 & a_1 + a_2 \end{bmatrix}$$

$$A_2 + A_1 = \begin{bmatrix} a_2 & 0 & 0 \\ 0 & a_2 & 0 \\ b_2 & c_2 & a_2 \end{bmatrix} + \begin{bmatrix} a_1 & 0 & 0 \\ 0 & a_1 & 0 \\ b_1 & c_1 & a_1 \end{bmatrix} = \begin{bmatrix} a_1 + a_2 & 0 & 0 \\ 0 & a_1 + a_2 & 0 \\ b_1 + b_2 & c_1 + c_2 & a_1 + a_2 \end{bmatrix}$$

Since  $a_1 + a_2 = a_2 + a_1, b_1 + b_2 = b_2 + b_1, c_1 + c_2 = c_2 + c_1$   
as  $a_1, a_2, b_1, b_2, c_1, c_2 \in \mathbb{Z}_2$

and addition of two is always commutative.  
in  $\mathbb{Z}_2$ .

Hence,  $M$  is commutative in standard matrix addition.

For Matrix Multiplication

$$A_1 \cdot A_2 = \begin{bmatrix} a_1 & 0 & 0 \\ 0 & a_1 & 0 \\ b_1 & c_1 & a_1 \end{bmatrix} \begin{bmatrix} a_2 & 0 & 0 \\ 0 & a_2 & 0 \\ b_2 & c_2 & a_2 \end{bmatrix} = \begin{bmatrix} a_1 a_2 & 0 & 0 \\ 0 & a_1 a_2 & 0 \\ b_1 a_2 + a_1 b_2 & c_1 a_2 + a_1 c_2 & a_1 a_2 \end{bmatrix}$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

M is closed under matrix Multiplication.

$$\begin{aligned}
 A_2 \cdot A_1 &= \begin{bmatrix} a_2 & 0 & 0 \\ 0 & a_2 & 0 \\ b_2 & c_2 & a_2 \end{bmatrix} \begin{bmatrix} a_1 & 0 & 0 \\ 0 & a_1 & 0 \\ b_1 & c_1 & a_1 \end{bmatrix} = \begin{bmatrix} a_2 a_1 & 0 & 0 \\ 0 & a_2 a_1 & 0 \\ b_2 a_1 + a_2 b_1 & c_2 a_1 + a_2 c_1 & a_2 a_1 \end{bmatrix} \\
 &= \begin{bmatrix} a_1 a_2 & 0 & 0 \\ 0 & a_1 a_2 & 0 \\ b_1 b_2 + b_1 a_2 & a_1 c_2 + c_1 a_2 & c_1 a_2 \end{bmatrix}
 \end{aligned}$$

as multiplication is always commutative in  $\mathbb{Z}_2$ .

therefore ;  $A_1 A_2 = A_2 A_1$

Hence, M is commutative

Idempotent means  $\Rightarrow M^2 = I$

$$M^2 = \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ b & c & a \end{bmatrix} \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \\ b & c & a \end{bmatrix} = \begin{bmatrix} a^2 & 0 & 0 \\ 0 & a^2 & 0 \\ 2ab & 2ac^2 & a^2 \end{bmatrix}$$

$$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

i.e  $a^2 = 1$ ,  $2ab = 0 = 2ac$  i.e  $b=c=0$   
 $a = \pm 1$ .

Hence; Idempotent elements of M are.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ & } \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

2.(c)(i): The function  $f$  defined by

$$f(z) = u + iv = \begin{cases} \frac{\operatorname{Im}(z^2)}{z} & \text{if } z \neq 0 \\ 0 & \text{if } z=0 \end{cases}$$

Satisfies the C-R equations at the origin, yet it is not differentiable there.

Sol'n: The function  $f$  defined by

$$f(z) = u + iv = \begin{cases} \frac{\operatorname{Im}(z^2)}{z} & ; z \neq 0 \\ 0 & ; z=0 \end{cases}$$

$$= \begin{cases} \frac{2xy}{x-iy} & ; z \neq 0 \\ 0 & ; z=0 \end{cases} \quad \begin{aligned} z^2 &= (x+iy)^2 \\ &= x^2 - y^2 + 2ixy \\ \operatorname{Im}(z^2) &= 2xy \end{aligned}$$

$$= \begin{cases} \frac{2xy(x-iy)}{x^2+y^2} & ; z \neq 0 \\ 0 & ; z=0 \end{cases}$$

Here  $u = \frac{2xy}{x^2+y^2}$  ;  $v = \frac{2xy^2}{x^2+y^2}$  where  $x \neq 0, y \neq 0$ .

To show that Cauchy - Riemann equations are satisfied at  $z=0$ :

Since  $f(0)=0$

$$\Rightarrow u(0,0) + iv(0,0) = 0$$

$$\Rightarrow u(0,0) = v(0,0) = 0$$

$$\text{Now } \left(\frac{\partial u}{\partial x}\right)_{(0,0)} = \lim_{y \rightarrow 0} \frac{u(x,0) - u(0,0)}{x} = \lim_{x \rightarrow 0} \frac{0-0}{x} = 0$$

$$\left(\frac{\partial u}{\partial y}\right)_{(0,0)} = \lim_{y \rightarrow 0} \frac{u(0,y) - u(0,0)}{y} = \lim_{y \rightarrow 0} \frac{0-0}{y} = 0$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

Similarly  $\left(\frac{\partial v}{\partial x}\right)_{(0,0)} = 0$  and  $\left(\frac{\partial u}{\partial y}\right)_{(0,0)} = 0$

$$\therefore \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \& \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \text{ at } z=0$$

$\therefore$  Cauchy-Riemann equations are satisfied.

To Prove that  $f(z)$  does not differentiable at  $(0,0)$ .

$$\begin{aligned} f'(0) &= \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z} \\ &= \lim_{(x,y) \rightarrow 0} \frac{2xy(x+iy)}{(x^2+y^2)(x+iy)} \end{aligned}$$

Let  $(x,y) \rightarrow (0,0)$  along  $y=mx$

$$\begin{aligned} f'(0) &= \lim_{x \rightarrow 0} \frac{2x^2m}{x^2+m^2x^2} \\ &= \lim_{x \rightarrow 0} \frac{2m}{1+m^2} \end{aligned}$$

Clearly which depends on  $m$ .

$\therefore f'(0)$  does not exist.

$\therefore f(z)$  does not differentiable at  $(0,0)$

$\therefore$  the given statement is true i.e. the given function  $f$  satisfies C-R equations although it is not differentiable at  $(0,0)$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

Q(C)(ii), The integral function  $f(z)$  satisfies everywhere the inequality  $|f(z)| \leq A|z|^k$  where  $A$  and  $k$  are positive constants. Prove that  $f(z)$  is a polynomial of degree not exceeding  $k$ .

Sol'n: Since  $f(z)$  is analytic in the finite part of the plane, therefore by Taylor's theorem.

$$f(z) = \sum_{n=0}^{\infty} a_n z^n, \text{ where } |z| < R$$

Now, if  $\max |f(z)| = M(r)$  on the circle  $|z|=r$  ( $r < R$ ), Then by Cauchy's inequality, we have

$$\begin{aligned} |a_n| &\leq \frac{M(r)}{r^n} \text{ for all } n \\ &= \frac{A(r)^k}{r^n} \quad \text{since } M(r) = |f(r)| = A|r|^k \\ &\qquad \qquad \qquad \text{when } |z| \rightarrow \infty. \end{aligned}$$

$$= \frac{A r^k}{r^n} = A r^{k-n}$$

Hence as  $r \rightarrow \infty$ , the right hand side tends to zero.

Since  $n > k$ .

i.e.  $a_n = 0$  for  $n > k$ .

i.e. all the coefficients  $a_n$  for which  $n > k$  becomes zero.

$$\therefore f(z) = a_0 + a_1 z + a_2 z^2 + \dots + a_k z^k$$

which is a polynomial of degree  $k$ .

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

3(a) (i) If in a ring  $R$ , with unity  $(xy)^2 = x^2y^2$  for all  $x, y \in R$  then show that  $R$  is commutative.

(ii) Show that the ring  $R$  of real valued continuous functions on  $[0,1]$  has zero divisors.

Sol'n: (i) we have  $(xy)^2 = x^2y^2 \forall x, y \in R$  — ①

Replacing  $y$  by  $y+1 \in R$  in ①, we get

$$[x(y+1)]^2 = x^2(y+1)^2$$

$$\Rightarrow (xy+x)^2 = x^2(y^2+2y+1)$$

$$\Rightarrow (xy+x)(xy+x) = x^2(y^2+2y+1)$$

$$\Rightarrow (xy)^2 + (xy)x + x(xy) + x^2 = x^2y^2 + 2x^2y + x^2 \quad \text{— ②}$$

$$\Rightarrow (xy)^2 + (xy)x + x(xy) + x^2 = (xy)^2 + 2x^2y + x^2$$

$$\Rightarrow (xy)x + x(xy) = 2x^2y \quad (\because \text{LCL \& RCL in } (R, +))$$

$$\Rightarrow xyx + x^2y = 2x^2y$$

$$\Rightarrow xyx = x^2y \quad \forall x, y \in R \quad (\text{RCL}) \quad \text{— ③}$$

Replacing  $x$  by  $x+1 \in R$  in ③,

$$(x+1)y(x+1) = (x+1)^2y$$

$$\Rightarrow (x+1)(yx+y) = (x+1)(xy+y)$$

$$\Rightarrow xyx + xy + yx + y = x^2y + xy + xy + y$$

$$\Rightarrow yx = xy \quad \forall x, y \in R \quad (\because \text{LCL \& RCL in } (R, +))$$

$\therefore R$  is a commutative ring.

(ii) Consider the functions  $f$  and  $g$  defined on  $[0,1]$

by

$$f(x) = \frac{1}{2} - x, \quad 0 \leq x \leq \frac{1}{2}$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

$$= 0, \frac{1}{2} \leq x \leq 1$$

$$\text{and } g(x) = 0, 0 \leq x \leq \frac{1}{2}$$

$$= x - \frac{1}{2}, \frac{1}{2} \leq x \leq 1$$

then  $f$  and  $g$  are continuous functions and  
 $f \neq 0, g \neq 0.$

whereas  $gf(x) = g(x)f(x) = 0 \cdot (\frac{1}{2} - x) \text{ if } 0 \leq x \leq \frac{1}{2}$

$$= (x - \frac{1}{2}) \cdot 0 = 0 \text{ if } \frac{1}{2} \leq x \leq 1$$

i.e.  $gf(x) = 0$  for all  $x$

i.e.  $gf = 0$  but  $f \neq 0, g \neq 0.$

                  .

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

3(b) Let  $f_n(x) = \frac{x}{1+nx^2}$  for all real  $x$ . Show that  $f_n$  converges uniformly to a function  $f$ . What is  $f$ ? Show that for  $x \neq 0$ ,  $f'_n(x) \rightarrow f'(x)$  but  $f'_n(0)$  does not converge to  $f'(0)$ . Show that the maximum value  $|f_n(x)|$  can take is  $\frac{1}{2\sqrt{n}}$ .

$$\text{Sol'n: Here } f(x) = \lim_{n \rightarrow \infty} f_n(x) = \lim_{n \rightarrow \infty} \frac{x}{1+nx^2} = 0 \quad \forall x \in \mathbb{R}$$

$$|f_n(x) - f(x)| = \left| \frac{x}{1+nx^2} - 0 \right| = \left| \frac{x}{1+nx^2} \right|$$

$$\text{Let } y = \frac{x}{1+nx^2}$$

$$\text{then } \frac{dy}{dx} = \frac{(1+nx^2)(1-x) \cdot 2nx}{(1+nx^2)^2} = \frac{1-nx^2}{(1+nx^2)^2}$$

$$\text{For max. (or) min } \frac{dy}{dx} = 0$$

$$\Rightarrow 1-nx^2 = 0 \Rightarrow x = \frac{1}{\sqrt{n}}$$

$$\text{Also } \frac{d^2y}{dx^2} = \frac{(1+nx^2)^2(-2nx) - (1-nx^2) \cdot 2(1+nx^2) \cdot 2nx}{(1+nx^2)^4}$$

$$= \frac{-2nx(1+nx^2) - 4nx(1-nx^2)}{(1+nx^2)^3}$$

$$\left. \frac{d^2y}{dx^2} \right|_{x=\frac{1}{\sqrt{n}}} = \frac{-2\sqrt{n}(1+1)}{(1+1)^3} = -\frac{\sqrt{n}}{2} < 0.$$

$\Rightarrow y$  is maximum when  $x = \frac{1}{\sqrt{n}}$  and maximum

value of  $y$

$$= \frac{\frac{1}{\sqrt{n}}}{1+1} = \frac{1}{2\sqrt{n}}$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

$$\therefore M_n = \max_{x \in [a,b]} |f_n(x) - f(x)| = \max_{x \in [a,b]} \left| \frac{x}{1+nx^2} \right|$$

$$= \frac{1}{2\sqrt{n}} \rightarrow 0 \text{ as } n \rightarrow \infty$$

Hence  $\{f_n\}$  converges uniformly to  $f$  on  $[a,b]$ .

$$\Rightarrow f(x) = 0 \quad \forall x \in \mathbb{R}$$

$$\Rightarrow f'(x) = 0 \quad \forall x \in \mathbb{R}$$

when  $x \neq 0$ ,

$$f_n'(x) = \frac{(1+nx^2) \cdot (1-x) \cdot 2nx}{(1+nx^2)^2} = \frac{1-nx^2}{(1+nx^2)^2}$$

$$\lim_{n \rightarrow \infty} f_n'(x) = \lim_{n \rightarrow \infty} \frac{1-nx^2}{(1+nx^2)^2} \quad | \text{ form } \frac{\infty}{\infty}$$

$$= \lim_{n \rightarrow \infty} \frac{-x^2}{2(1+nx^2) \cdot x^2} = 0 = f'(x)$$

So that if  $x \neq 0$ , the formula  $\lim_{n \rightarrow \infty} f_n'(x) = f'(x)$  is true.

At  $x=0$ ,

$$f_n'(0) = \lim_{h \rightarrow 0} \frac{f_n(0+h) - f_n(0)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{h}{1+nh^2} - 0}{h} = \lim_{h \rightarrow 0} \frac{1}{1+nh^2} = 1$$

So that  $\lim_{n \rightarrow \infty} f_n'(0) = 1 \neq f'(0)$ .

Hence at  $x=0$ , the formula  $\lim_{n \rightarrow \infty} f_n'(x) = f'(x)$  is false.

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

3(c) Using the simplex method solve the LPP problem:

$$\text{minimize } Z = 2x_1 + 3x_2.$$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 + 7x_2 \geq 7$$

$$x_1, x_2 \geq 0.$$

Sol? The objective function of the given LPP is of minimization type.

So, we convert it into maximization type.

$$\text{Max } Z' = \text{Min}(-Z)$$

$$= -x_1 - 3x_2$$

Now we write the given LPP in the standard form

$$\text{Max } Z' = -x_1 - x_2 + 0s_1 + 0s_2 - MA_1 - MA_2$$

Subject to

$$2x_1 + x_2 - s_1 + A_1 = 4$$

$$x_1 + 7x_2 - s_2 + A_2 = 7$$

$$A_1, A_2, x_1, x_2, s_1, s_2 \geq 0.$$

where  $s_1, s_2$  are the surplus variables

$A_1, A_2$  are the artificial variables.

Now the initial BFS is

$$s_1 = s_2 = x_1 = x_2 = 0 \quad (\text{non-basic})$$

$$A_1 = 4, A_2 = 7 \quad (\text{Basic})$$

Thus the initial simplex table is

CB Basis	$C_j$		$-C_j$				b	0
	$x_1$	$x_2$	$s_1$	$s_2$	$A_1$	$A_2$		
$-M A_1$	2	1	-1	0	1	0	4	4
$-M A_2$	1	(7)	0	-1	0	1	7	1
$Z_j = \sum C_{Bij} x_j$	-3M	-8M	M	M	-M	-M	-11M	
$C_j = Z_j - Z_B$	-13M	-18M	-M	-M	0	0		

From the above table,

the variable  $x_2$  is entering variable,  $A_2$  is the outgoing variable and omit column for the variable in the next simplex table. Here (7) is the key element and convert it into unity and all other elements in this column to zero.

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

Then the new simplex table is:

$C_j$	-1	-1	0	0	-M		
CB Basis	$x_1$	$x_2$	$s_1$	$s_2$	$A_1$	b	$\theta$
-M	$A_1$	( $\frac{13}{7}$ )	0	-1	$y_7$	1	$\frac{21}{13} \rightarrow$
-1	$x_2$	$y_7$	1	0	- $y_7$	0	1 $\neq$

$$Z_j = \sum a_{ij} C_B - \frac{13M}{7} - \frac{1}{7} - 1 M - \frac{M}{7} + \frac{1}{7} - M - 3M - 1$$

$$C_j = C_j - Z_j \quad \frac{13M - 6}{7} \quad 0 \quad -M \quad \frac{M}{7} - \frac{1}{7} \quad 0$$

↑  
From the above table,

$x_2$  is the entering variable,  $A_1$  is the outgoing variable and omit its column in the next simplex table. Here ( $\frac{13}{7}$ ) is the key element and make it unity and all other elements in its column equal to zero. Then the revised simplex table is

$C_j$	-1	-1	0	0			
CB Basis	$x_1$	$x_2$	$s_1$	$s_2$	$A_1$	b	$\theta$
-1	$x_1$	1	0	$-\frac{1}{13}$	$\frac{1}{13}$	$\frac{21}{13}$	
-1	$x_2$	0	1	$y_{13}$	$-\frac{2}{13}$	$10/13$	
$Z_j = \sum a_{ij} C_B$		-1	-1	$\frac{6}{13}$	$\frac{4}{13}$	$-\frac{31}{13}$	
$C_j = C_j - Z_j$		0	0	$-\frac{6}{13}$	$-\frac{4}{13}$		

From the above table, all  $C_j \leq 0$ .

there remains no artificial variable in the basis.

∴ The solution is an optimal BFS to the problem and is given by

$$x_1 = \frac{21}{13}, \quad x_2 = \frac{10}{13}$$

$$\therefore \text{Max } Z' = -\frac{31}{13}$$

Hence the optimal value of the objective function is  $\text{Min } Z = \text{Max } Z' = \frac{31}{13}$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

4(a)

Let  $H$  be a subgroup of a group  $G$  such that  $(G:H) = 2$ . Then prove that  $H$  is a normal subgroup of  $G$ .

Is converse true? Justify your answer.

Sol Let  $(G, \cdot)$  be a group and  $H \trianglelefteq G$  such that  $(G:H) = 2$ .  
 To prove  $H \trianglelefteq G$ .

$$\therefore (G:H) = 2$$

There are two distinct left & right cosets of  $H$  in  $G$ .

Let  $H$  &  $Ha$ ;  $h \in H$  be two distinct right cosets & left cosets of  $H$  in  $G$ !

$$\text{Then } G = H \cup Ha = \overbrace{H \cup cH}^{\text{AFG}} \quad (1)$$

$$\therefore a \in G \Rightarrow a \in H \text{ or } a \notin H$$

Case(i) When  $a \in H$

$$\begin{aligned} \therefore Ha &= H = aH \\ &\Rightarrow Ha = cH. \end{aligned}$$

Case(ii) When  $a \notin H$

$$\therefore Ha \neq H \text{ & } cH \neq H$$

$$\therefore G = H \cup Ha \quad \text{&} \quad G = H \cup cH$$

$$\therefore H \cup Ha = H \cup cH$$

$$\Rightarrow Ha = cH$$

$$\therefore Ha = aH \nrightarrow a \in G.$$

$$\therefore H \trianglelefteq G.$$

The converse of the above need not be true.

i.e. If  $H \trianglelefteq G$  then  $(G:H) \neq 2$

for example:

Let  $G = \{1, -1, i, -i, j, -j, k, -k\}$  be a group

Let  $H = \{1, -1\}$

Clearly  $H \trianglelefteq G$

$$\text{But } (G:H) = 8$$

$$\neq 2.$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

4(b) →

Discuss the convergence of the sequence  $\{x_n\}$

$$\text{where } x_n = \frac{\sin\left(\frac{n\pi}{2}\right)}{8}$$

Soln: let  $x_n = \frac{\sin\left(\frac{n\pi}{2}\right)}{8} \quad \forall n \in \mathbb{N}$

Since  $\lim_{n \rightarrow \infty} x_n = \begin{cases} 0 & \text{if } n \text{ is even} \\ \frac{1}{8} & \text{if } n = 4m+1 \\ -\frac{1}{8} & \text{if } n = 4m+3 \end{cases}$

does not exist

$\therefore (x_n)$  does not converge.



**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

4.(C) →

Use the method of contour integration to prove that

$$\int_0^{2\pi} \frac{d\theta}{(a+b\cos\theta+c\sin\theta)^2} = \frac{2\pi a}{\sqrt[3]{a^2-b^2-c^2}}, \quad a^2 > b^2 + c^2.$$

Sol'n:  $I = \int_0^{2\pi} \frac{d\theta}{(a+b\cos\theta+c\sin\theta)^2}$

$$= \int_0^{2\pi} \frac{d\theta}{\left[a + \frac{1}{2}(e^{i\theta} + e^{-i\theta}) + \frac{c}{2i}(e^{i\theta} - e^{-i\theta})\right]^2}$$

write  $e^{i\theta} = z \Rightarrow d\theta = \frac{dz}{iz}$

$$I = \int_C \frac{dz/iz}{\left[a + \frac{b}{2}(z + \frac{1}{z}) + \frac{c}{2i}(z - \frac{1}{z})\right]^2}$$

$$= \int_C \frac{4iz dz}{\left[2ia + bi(z+1) + c(z-1)\right]^2}$$

$$= \int_C \frac{4iz dz}{\left[\frac{2}{z}(bi+c) + 2iz + (bi-c)\right]^2}$$

$$= \frac{4i}{(bi+c)^2} \int_C \frac{z dz}{\left[z + \frac{2iaz}{bi+c} + \frac{bi-c}{bi+c}\right]^2} = \frac{4i}{(bi+c)} \int_C f(z) dz \quad \text{--- (1)}$$

where  $C$  is the unit circle.

$f(z)$  has poles of order 2 given by

$$z + \frac{2ia}{bi+c} z + \frac{bi-c}{bi+c} = 0.$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

$$\Rightarrow z = \frac{-\frac{2ic}{bi+c} \pm \sqrt{\frac{iac}{(bi+c)^2} - \frac{4(bi+c)}{bi+c}}}{2}$$

$$\Rightarrow z = \frac{1}{2} \left[ \frac{-2ic}{bi+c} \pm \frac{2i \sqrt{a^2 - b^2 - c^2}}{bi+c} \right]$$

$$\Rightarrow z = \frac{i}{bi+c} \left[ -a \pm \sqrt{a^2 - b^2 - c^2} \right]$$

$$z = \frac{i}{bi+c} (-a + \sqrt{a^2 - b^2 - c^2}) \quad & z = \frac{i}{(bi+c)} (-a - \sqrt{a^2 - b^2 - c^2}) \\ = \alpha \text{ (say)} \quad & \quad = \beta \text{ (say)}$$

only  $z = \alpha$  lies inside  $C$  of order 2.

Residue at  $z = \alpha$  is

$$\begin{aligned} & \underset{z \rightarrow \alpha}{\lim} \frac{d}{dz} \left\{ (z-\alpha)^2 \frac{z}{(z-\alpha)^2 (z-\beta)^2} \right\} \\ &= \underset{z \rightarrow \alpha}{\lim} \frac{d}{dz} \left[ \frac{z}{(z-\beta)^2} \right] \\ &= \underset{z \rightarrow \alpha}{\lim} \frac{-(z+\beta)}{(z-\beta)^3} \\ &= -\frac{(z+\beta)}{(z-\beta)^2} \\ &= -\frac{i}{(bi+c)} \frac{(-2a)}{\left(\frac{2i}{bi+c}\right)^3 \left(\sqrt{a^2 - b^2 - c^2}\right)^3} = \frac{a}{(bi+c)^2 \left(\sqrt{a^2 - b^2 - c^2}\right)^2} \end{aligned}$$

Then from ① . .

$$\begin{aligned} I &= \frac{4i}{(bi+c)^2} [2\pi i] (\text{Residue at } z = \alpha) \\ &= \frac{ui}{(bi+c)^2} (2\pi i) \frac{-a}{\left(\frac{4}{bi+c}\right)^2 \left(\sqrt{a^2 - b^2 - c^2}\right)^3} \\ &= \frac{2\pi a}{\left(a^2 - b^2 - c^2\right)^{3/2}} \\ \therefore \int_{C} \frac{dz}{z^2 + bz + c^2} &= \frac{2\pi a}{\sqrt{a^2 - b^2 - c^2}} \end{aligned}$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

4.(d), Make a graphical representation of the set of constraints of the following LPP. Find the extreme points of the feasible region. Finally, solve the problem graphically.

$$\text{Max. } Z = 2x_1 + x_2$$

$$\text{Subject to } x_1 + x_2 \geq 5$$

$$2x_1 + 3x_2 \leq 20$$

$$4x_1 + 3x_2 \leq 25$$

$$x_1, x_2 \geq 0.$$


Sol'n: Let us consider the constraint  $x_1 + x_2 \geq 5$  as equality  $x_1 + x_2 = 5$  represents a line which passes through  $(0, 5)$  &  $(5, 0)$  in the  $XY$ -plane.

Similarly, we get

$$2x_1 + 3x_2 = 20 \Rightarrow (10, 0) \text{ & } (0, \frac{20}{3})$$

$$4x_1 + 3x_2 = 25 \Rightarrow (\frac{25}{4}, 0) \text{ & } (0, \frac{25}{3})$$

the graph representation is as shown below:



**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

The feasible region is given by ABCDE which satisfy all the constraints simultaneously. Coordinates of extreme points are

(Corner points)

$$A(5, 0), B\left(\frac{25}{4}, 0\right), C\left(\frac{5}{2}, 5\right), D\left(0, \frac{20}{3}\right), E(0, 5)$$

To maximize  $Z = 2x_1 + x_2$

The values of the objective function  $Z = 2x_1 + x_2$  at these extreme points are

$$Z(0, 5) = 5$$

$$Z(5, 0) = 10$$

$$Z\left(\frac{25}{4}, 0\right) = \frac{25}{2}$$

$$Z\left(\frac{5}{2}, 5\right) = 10$$

$$Z\left(0, \frac{20}{3}\right) = \frac{20}{3}$$

∴ The maximum value of  $Z$  is at the point

$$\left(\frac{25}{4}, 0\right)$$

and the maximum value is

$$Z = \frac{25}{2}$$

∴ The optimal solution is

$$x_1 = \frac{25}{4}, x_2 = 0 \text{ and } Z = \frac{25}{2}.$$

=====

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

5.(a), solve  $(D_x^3 - 7D_x D_y^2 - 6D_y^3)z = \sin(x+2y) + e^{3x+y}$ .

Soln. Here note that  $D_x$  and  $D_y$  stand for  $D$  and  $D'$  respectively.

∴ Auxiliary equation is  $m^3 - 7m^2 - 6m = 0$  so that  $m = -1, -2, 3$ .

∴ C.F. =  $\phi_1(y-x) + \phi_2(y-2x) + \phi_3(y+3x)$ ,  $\phi_1, \phi_2, \phi_3$  being arbitrary functions.

Now P.I. corresponding to  $\sin(x+2y)$

$$\begin{aligned} &= \frac{1}{D_x^3 - 7D_x D_y^2 - 6D_y^3} \sin(x+2y) \\ &= \frac{1}{1^3 - 7 \cdot 1 \cdot 2^2 - 6 \cdot 2^3} \iiint \sin v dv dv dv \\ &= \left(-\frac{1}{75}\right) \iiint (-\cos v) dv dv dv = \left(-\frac{1}{75}\right) \left(-\sin v\right) dv \\ &= \left(-\frac{1}{75}\right) \cos v = \left(-\frac{1}{75}\right) \cos(x+2y) \end{aligned}$$

and P.I. Corresponding to  $e^{3x+y}$

$$\begin{aligned} &= \frac{1}{D_x^3 - 7D_x D_y^2 - 6D_y^3} e^{3x+y} = \frac{1}{D_x - 3D_y} \left[ \frac{1}{(D_x + D_y)(D_x + 2D_y)} e^{3x+y} \right] \\ &= \frac{1}{D_x - 3D_y} \cdot \frac{1}{(3+1)(3+2)} \iint e^v dv dv, \text{ where } v = 3x+y \\ &= \frac{1}{20} \cdot \frac{1}{D_x - 3D_y} \int e^v dv = \frac{1}{20} \cdot \frac{1}{D_x - 3D_y} e^v \\ &= \frac{1}{20} \cdot \frac{1}{(D_x - 3D_y)^1} e^{3x+y} \\ &= \frac{1}{20} \cdot \frac{x}{1!1!} e^{3x+y} = \frac{x}{20} e^{3x+y} \end{aligned}$$

with  $a = 2, b = 1, m = 1$

Hence the required general

Solution is  $z = C.F. + P.I.$

$$z = \phi_1(y-x) + \phi_2(y-2x) + \phi_3(y+3x) - \frac{1}{75} \cos(x+2y) + \frac{x}{20} e^{3x+y}$$

using formula

$$\frac{1}{(bD - aD')^n} \phi(ax+by) = \frac{x^n}{b^n \cdot n!} \phi(ax+by)$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

5(b) Find a surface satisfying  $\sigma + s = 0$ , i.e.,  $(D^2 + DD')Z = 0$  and touching the elliptic paraboloid  $Z = 4x^2 + y^2$  along its section by the plane  $y = 2x + 1$ .

Sol'n: Given  $(D^2 + DD')Z = 0 \quad \dots \textcircled{1}$

Here auxiliary equation is  $m^2 + m = 0$  so that  $m = 0, -1$

$\therefore$  Solution of  $\textcircled{1}$  is  $Z = C.F = \phi_1(y) + \phi_2(y-x) \quad \dots \textcircled{2}$

Since  $\textcircled{2}$  touches the curve given by

$$Z = 4x^2 + y^2 \quad \dots \textcircled{3}$$

$$\text{and } y = 2x + 1 \quad \dots \textcircled{4}$$

values of  $P \left( \frac{\partial Z}{\partial x} \right)$  and  $Q \left( = \frac{\partial Z}{\partial y} \right)$  obtained from  $\textcircled{2}$  and  $\textcircled{3}$  must be equal for any point on  $\textcircled{4}$

$$-\phi'_2(y-x) = 8x \text{ for } y = 2x + 1$$

$$\Rightarrow \phi'_2(x+1) = -8x \quad \dots \textcircled{5}$$

$$\text{and } \phi'_1(y) + \phi'_2(y-x) = 2y \text{ for } y = 2x + 1$$

$$\Rightarrow \phi'_1(2x+1) + \phi'_2(x+1) = 4x+2 \quad \dots \textcircled{6}$$

$$\text{from } \textcircled{5} \quad \phi'_2(x) = 8-8x$$

$$\text{Integrating it, } \phi_2(x) = 8x - 4x^2 + C_1 \quad \dots \textcircled{7}$$

$$\text{Subtracting } \textcircled{6} - \textcircled{5} \Rightarrow \phi'_1(2x+1) = 12x+2 = 6(2x+1)-4$$

$$\text{so that } \phi'_1(x) = 6x-4$$

$$\text{Integrating it } \phi_1(x) = 3x^2 - 4x + C_2 \quad \dots \textcircled{8}$$

$$\text{from } \textcircled{8} \quad \phi_1(y) = 3y^2 - 4y + C_2$$

$$\text{and from } \textcircled{7} \quad \phi_2(y-x) = 8(y-x) - 4(y-x)^2 + C_1$$

putting the above values of  $\phi_1(y)$  and  $\phi_2(y-x)$

in  $\textcircled{2}$  we get

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

$$z = 3y^2 - 4y + c_2 + 8(y-x) - 4(y-x)^2 + c_1$$

$$\Rightarrow z = -y^2 + 4y - 8x - 4x^2 + 8xy + c_3 \text{ where } c_3 = c_1 + c_2 \quad \textcircled{9}$$

Equating the values of  $z$  from  $\textcircled{3}$  and  $\textcircled{9}$  we get

$$4x^2 + y^2 = -y^2 + 4y - 8x - 4x^2 + 8xy + c_3 \text{ where } y = 2x+1$$

$$c_3 = 8x^2 + 2y^2 - 4y + 8x - 8xy$$

$$= 8x^2 + 2(2x+1)^2 - 4(2x+1) + 8x - 8x(2x+1)$$

$$= -2$$

Hence from  $\textcircled{9}$  the required surface is

$$\underline{4x^2 - 8xy + y^2 + 8x - 4y + 2 + 2 = 0}.$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

5.(c) →

From the following table, estimate the number of students who obtained marks between 40 and 45 using Newton's forward interpolation formula.

Marks :            30-40    40-50    50-60    60-70    70-80

No. of Students :    31        42        51        35        31

Soln

First we prepare the cumulative frequency table, as follows:-

Marks less than ( $x$ ):    40    50    60    70    80

No. of students ( $y_x$ ):    31    73    124    159    190

Now the difference table is -

$x$	$y_x$	$\Delta y_x$	$\Delta^2 y_x$	$\Delta^3 y_x$	$\Delta^4 y_x$
40	31	42			
50	73	51	9	-25	
60	124	35	-16		37
70	159	31	-4	12	
80	190				

We shall find  $y_{45}$  i.e. number of students with marks less than 45. Taking  $x_0 = 40$ ,  $x = 45$ , we have

$$P = \frac{x - x_0}{h} = \frac{5}{10} = 0.5$$

∴ Using Newton's forward interpolation formula, we get,

$$y_{45} = y_{40} + P \Delta y_{40} + \frac{P(P-1)}{2!} \Delta^2 y_{40}$$

$$+ \frac{P(P-1)(P-2)}{3!} \Delta^3 y_{40}$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

$$\begin{aligned}
 &= 31 + 0.5 \times 42 + \frac{0.5(-0.5)}{2} \times 9 + \frac{0.5(-0.5)(-15)}{6} \\
 &\quad \times (-25) + \frac{0.5(+0.5)(-15)(-2.5)}{24} \times 37 \\
 &= 31 + 21 - 1.125 - 1.5625 - 1.4453 \\
 &= 47.87, \text{ on simplification.}
 \end{aligned}$$

The number of students with marks less than 45 is 47.87 ie. 48.

But the number of students with marks less than 40 is 31.

Hence the number of students getting marks between 40 and 45 = 48 - 31  
 $= 17.$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

5(d)

Using Gauss Seidel iterative method and the starting solution  $x_1 = x_2 = x_3 = 0$  determine the solution of the following system of equations in two iterations,  $10x_1 - x_2 - x_3 = 8$ ;  $x_1 + 10x_2 + x_3 = 12$ ,  $x_1 - x_2 + 10x_3 = 10$ .

Solution:- The given equation can be re-written as -

$$x_1 = \frac{8 + x_2 + x_3}{10} = 0.8 + 0.1x_2 + 0.1x_3$$

$$x_2 = \frac{12 - x_1 - x_3}{10} = 1.2 - 0.1x_1 - 0.1x_3$$

$$x_3 = \frac{10 - x_1 + x_2}{10} = 1 - 0.1x_1 + 0.1x_2$$

Also given that  $x_1 = x_2 = x_3 = 0$

and we need to find the solution of the system of equation in two iteration.

Iteration -1

Taking;  $x_1 = x_2 = x_3 = 0$

$$x_1 = 0.8 + 0.1 \times 0 + 0.1 \times 0 = 0.8$$

$$x_2 = 1.2 - 0.1 \times 0.8 - 0.1 \times 0 = 1.12.$$

$$x_3 = 1 - 0.1 \times 0.8 + 0.1 \times 1.12 = 1.032.$$

After first iteration;  $x_1 = 0.8$

$$x_2 = 1.12$$

$$x_3 = 1.032.$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

Iteration - II

$$x_1 = 0.8, x_2 = 1.12, x_3 = 1.032$$

$$x_1 = 0.8 + 0.1 \times 1.12 + 0.1 \times 1.032 = 1.0152$$

$$\begin{aligned}x_2 &= 1.2 - 0.1 \times 1.0152 - 0.1 \times 1.032 \\&= 1.2 - 0.10152 - 0.1032 = 0.9953\end{aligned}$$

$$\begin{aligned}x_3 &= 1 - 0.1 \times 1.0152 + 0.1 \times 0.9953 \\&= 1 - 0.10152 + 0.09953 = 0.99801.\end{aligned}$$

∴ After 2<sup>nd</sup> Iteration  $\Rightarrow$

$$\boxed{\begin{aligned}x_1 &= 1.0152 \\x_2 &= 0.9953 \\x_3 &= 0.99801\end{aligned}}$$

which is required solution

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

5(e)

Prove that the necessary and sufficient condition that vortex lines may be at right angles to the streamlines are  $u, v, w = \mu \left( \frac{\partial \phi}{\partial x}, \frac{\partial \phi}{\partial y}, \frac{\partial \phi}{\partial z} \right)$ , where  $\mu$  and  $\phi$  are functions of  $x, y, z, t$ .

Sol'n: The differential equations of streamlines and vortex lines are respectively.

$$\frac{dx}{u} = \frac{dy}{v} = \frac{dz}{w} \quad \dots \quad (1)$$

$$\text{and } \frac{dx}{\xi} = \frac{dy}{\eta} = \frac{dz}{\zeta} \quad \dots \quad (2)$$

$\therefore$  (1) and (2) will intersect orthogonally iff

$$u\xi + v\eta + w\zeta = 0$$

$$\Rightarrow u \left( \frac{\partial \omega}{\partial y} - \frac{\partial v}{\partial z} \right) + v \left( \frac{\partial u}{\partial z} - \frac{\partial w}{\partial x} \right) + w \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right) = 0$$

But this is the condition that-

$udx + vdy + wdz$  is perfect differential

$$\Rightarrow udx + vdy + wdz = \mu d\phi$$

$$= \mu \left( \frac{\partial \phi}{\partial x} dx + \frac{\partial \phi}{\partial y} dy + \frac{\partial \phi}{\partial z} dz \right)$$

$$\text{this} \Rightarrow u, v, w = \mu \left( \frac{\partial \phi}{\partial x}, \frac{\partial \phi}{\partial y}, \frac{\partial \phi}{\partial z} \right).$$

=====.

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

6(a) Find a partial differential equation by eliminating  
 a, b, c from  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ .

Soln: Given that  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1 \quad \text{--- (1)}$

Differentiating (1) w.r.t x and y, we get

$$\frac{\partial z}{\partial x} + \frac{2z}{c^2} \frac{\partial z}{\partial x} = 0 \Rightarrow c^2 z + a^2 z \frac{\partial z}{\partial x} = 0 \quad \text{--- (2)}$$

$$\text{and } \frac{\partial z}{\partial y} + \frac{2z}{c^2} \frac{\partial z}{\partial y} = 0 \Rightarrow c^2 z + b^2 z \frac{\partial z}{\partial y} = 0 \quad \text{--- (3)}$$

Differentiating (2) w.r.t x and (3) w.r.t y, we have

$$c^2 + a^2 \left( \frac{\partial z}{\partial x} \right)^2 + a^2 z \frac{\partial^2 z}{\partial x^2} = 0 \quad \text{--- (4)}$$

$$c^2 + b^2 \left( \frac{\partial z}{\partial y} \right)^2 + b^2 z \frac{\partial^2 z}{\partial y^2} = 0 \quad \text{--- (5)}$$

$$\text{from (2), } c^2 = -a^2 z \left( \frac{\partial z}{\partial x} \right).$$

putting this value of  $c^2$  in (4) and dividing by  $a^2$ , we obtain

$$-\frac{2}{a^2} \frac{\partial z}{\partial x} + \left( \frac{\partial z}{\partial x} \right)^2 + 2 \frac{\partial^2 z}{\partial x^2} = 0 \quad \text{--- (6)}$$

Similarly, from (3) & (5),

$$-b^2 \frac{\partial z}{\partial y} + \left( \frac{\partial z}{\partial y} \right)^2 + 2 \frac{\partial^2 z}{\partial y^2} = 0 \quad \text{--- (7)}$$

Differentiating (2) partially w.r.t y, we get

$$a^2 \left\{ \frac{\partial^2 z}{\partial y^2} \left( \frac{\partial z}{\partial x} \right)^2 + 2 \frac{\partial^2 z}{\partial x \partial y} \right\} = 0$$

$$\text{i.e. } \frac{\partial z}{\partial x} \frac{\partial^2 z}{\partial y^2} + 2 \frac{\partial^2 z}{\partial x \partial y} = 0 \quad \text{--- (8)}$$

$\therefore$  (6), (7) and (8) are three possible forms of the required partial differential equations.

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

6(b), Reduce the equation  $y_r + (x+y)s + xt = 0$  to canonical form and hence find its general solution.

Soln:

$$\text{Given } y_r + (x+y)s + xt = 0 \rightarrow ①$$

$$\text{Comparing } ① \text{ with } Rr + Ss + Tt + f(x, y, z, P, Q) = 0$$

here  $R = y, S = x+y$  and  $T = x$  So that

$$S^2 - 4RT = (x+y)^2 - 4xy = (x-y)^2 > 0 \text{ for } x \neq y.$$

ans so ① is hyperbolic. It's 1-quadratic equation

$$R\lambda^2 + S\lambda + T = 0 \text{ reduces to } y\lambda^2 + (x+y)\lambda + x = 0$$

$$(\text{or}) (y\lambda + x)(\lambda + 1) = 0$$

So that  $\lambda = -1, -\frac{x}{y}$ . Then the corresponding characteristic equations are given by

$$\frac{dy}{dx} - 1 = 0 \text{ and } \frac{dy}{dx} - \left(-\frac{x}{y}\right) = 0$$

$$\text{Integrating these } y - x = c_1 \text{ and } \frac{y^2}{2} - \frac{x^2}{2} = c_2$$

In order to reduce one ① to its canonical form, we choose.

$$u = y - x \text{ and } v = \frac{y^2}{2} - \frac{x^2}{2} \rightarrow ②$$

$$\begin{aligned} \therefore r = \frac{\partial t}{\partial x} &= \frac{\partial t}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial t}{\partial v} \frac{\partial v}{\partial x} \\ &= -\left(\frac{\partial t}{\partial u} + x \frac{\partial t}{\partial v}\right), \text{ using } ② \rightarrow ③ \end{aligned}$$

$$s = \frac{\partial t}{\partial y} = \frac{\partial t}{\partial u} \frac{\partial u}{\partial y} + \frac{\partial t}{\partial v} \frac{\partial v}{\partial y} = \frac{\partial t}{\partial u} + y \frac{\partial t}{\partial v}, \text{ using } ②$$

↪ ④

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

$$\begin{aligned}
 r &= \frac{\partial^2 t}{\partial u^2} = \frac{\partial}{\partial u} \left( \frac{\partial t}{\partial u} \right) = -\frac{\partial}{\partial u} \left( \frac{\partial t}{\partial u} \right) - \frac{\partial}{\partial u} \left( u \frac{\partial t}{\partial v} \right) \text{ using } ① \\
 &= -\frac{\partial}{\partial u} \left( \frac{\partial t}{\partial u} \right) - \left[ u \frac{\partial}{\partial u} \left( \frac{\partial t}{\partial v} \right) + \frac{\partial t}{\partial v} \right] = -\frac{\partial}{\partial u} \left( \frac{\partial t}{\partial u} \right) \\
 &\quad - u \frac{\partial}{\partial u} \left( \frac{\partial t}{\partial v} \right) - \frac{\partial t}{\partial v} \\
 \Rightarrow & - \left[ \frac{\partial}{\partial u} \left( \frac{\partial t}{\partial u} \right) \frac{\partial u}{\partial u} + \frac{\partial}{\partial v} \left( \frac{\partial t}{\partial u} \right) \frac{\partial v}{\partial u} \right] - \\
 &\quad u \left[ \frac{\partial}{\partial u} \left( \frac{\partial t}{\partial v} \right) \frac{\partial u}{\partial u} + \frac{\partial}{\partial v} \left( \frac{\partial t}{\partial v} \right) \frac{\partial v}{\partial u} \right] - \frac{\partial t}{\partial v} \\
 &= - \left( -\frac{\partial^2 t}{\partial u^2} - u \frac{\partial^2 t}{\partial u \partial v} \right) - u \left( -\frac{\partial^2 t}{\partial u \partial v} - u \frac{\partial^2 t}{\partial v^2} \right) - \frac{\partial t}{\partial v} \text{ using } ② \\
 r &= \frac{\partial^2 t}{\partial u^2} + u \frac{\partial^2 t}{\partial u \partial v} + u^2 \frac{\partial^2 t}{\partial v^2} - \frac{\partial t}{\partial v} \\
 \text{Now, } t &= \frac{\partial^2 t}{\partial y^2} = \frac{\partial}{\partial y} \left( \frac{\partial t}{\partial y} \right) = \frac{\partial}{\partial y} \left( \frac{\partial t}{\partial u} + y \frac{\partial t}{\partial v} \right) \\
 &= \frac{\partial}{\partial y} \left( \frac{\partial t}{\partial u} \right) + \frac{\partial}{\partial y} \left( y \frac{\partial t}{\partial v} \right) \text{ by } ④ \\
 &= \frac{\partial}{\partial y} \left( \frac{\partial t}{\partial u} \right) + y \frac{\partial}{\partial y} \left( \frac{\partial t}{\partial v} \right) + \frac{\partial t}{\partial v} \\
 &= \frac{\partial}{\partial u} \left( \frac{\partial t}{\partial u} \right) \frac{\partial u}{\partial y} + \frac{\partial}{\partial v} \left( \frac{\partial t}{\partial u} \right) \frac{\partial v}{\partial y} + y \left\{ \frac{\partial}{\partial u} \left( \frac{\partial t}{\partial v} \right) \frac{\partial u}{\partial y} + \right. \\
 &\quad \left. \frac{\partial}{\partial v} \left( \frac{\partial t}{\partial v} \right) \frac{\partial v}{\partial y} \right\} + \frac{\partial t}{\partial v} \\
 \Rightarrow & \frac{\partial^2 t}{\partial u^2} + y \frac{\partial^2 t}{\partial u \partial v} + y \left( \frac{\partial^2 t}{\partial u \partial v} + y \frac{\partial^2 t}{\partial v^2} \right) + \frac{\partial t}{\partial v}.
 \end{aligned}$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

$$\therefore t = \frac{\partial^2 t}{\partial u^2} + 2y \frac{\partial^2 t}{\partial u \partial v} + y^2 \frac{\partial^2 t}{\partial v^2} + \frac{\partial t}{\partial v}$$

Also  $S = \frac{\partial^2 t}{\partial x \partial y} = \frac{\partial}{\partial x} \left( \frac{\partial t}{\partial y} \right) = \frac{\partial}{\partial x} \left( \frac{\partial t}{\partial u} + y \frac{\partial t}{\partial v} \right)$  using (4)

$$= \frac{\partial}{\partial x} \left( \frac{\partial t}{\partial u} \right) + \frac{\partial}{\partial x} \left( y \frac{\partial t}{\partial v} \right) = \frac{\partial}{\partial x} \left( \frac{\partial t}{\partial u} \right) + y \frac{\partial}{\partial x} \left( \frac{\partial t}{\partial v} \right)$$

$$= \frac{\partial}{\partial u} \left( \frac{\partial t}{\partial u} \right) \frac{\partial u}{\partial x} + \frac{\partial}{\partial v} \left( \frac{\partial t}{\partial u} \right) \frac{\partial v}{\partial x} + y \left\{ \frac{\partial}{\partial u} \left( \frac{\partial t}{\partial v} \right) \frac{\partial u}{\partial x} \right. \\ \left. + \frac{\partial}{\partial v} \left( \frac{\partial t}{\partial v} \right) \frac{\partial v}{\partial x} \right\}$$

$$\Rightarrow -\frac{\partial^2 t}{\partial u^2} - x \frac{\partial^2 t}{\partial u \partial v} - y \frac{\partial^2 t}{\partial u \partial v} - xy \frac{\partial^2 t}{\partial v^2} \text{ using (2)}$$

$$\therefore S = -\frac{\partial^2 t}{\partial u^2} - (x+y) \frac{\partial^2 t}{\partial u \partial v} - xy \frac{\partial^2 t}{\partial v^2}$$

↳ (7)

using (5), (6) and (7) in (1) we get :

$$y \left( \frac{\partial^2 t}{\partial u^2} + 2x \frac{\partial^2 t}{\partial u \partial v} + y^2 \frac{\partial^2 t}{\partial v^2} - \frac{\partial t}{\partial v} \right) + \\ (x+y) \left\{ -\frac{\partial^2 t}{\partial u^2} - (x+y) \frac{\partial^2 t}{\partial u \partial v} - xy \frac{\partial^2 t}{\partial v^2} \right\} \\ + x \left\{ \frac{\partial^2 t}{\partial u^2} + 2y \frac{\partial^2 t}{\partial u \partial v} + y^2 \frac{\partial^2 t}{\partial v^2} + \frac{\partial t}{\partial v} \right\} = 0$$

$$(or) \left\{ 4xy - (x+y)^2 \right\} \frac{\partial^2 t}{\partial u \partial v} - y \frac{\partial t}{\partial v} + x \frac{\partial t}{\partial v} = 0$$

$$(or) (y-x)^2 \frac{\partial^2 t}{\partial u \partial v} + (y-x) \frac{\partial t}{\partial v} = 0$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

$$(or) u^2 \frac{\partial^2 z}{\partial u \partial v} + u \frac{\partial z}{\partial v} = 0 \quad (or) u \frac{\partial^2 z}{\partial v \partial u} + \frac{\partial z}{\partial u} = 0 \rightarrow (8)$$

[ $\because u \neq 0$  and  $y-u=u$ , by (2)]

(8) is the required Canonical form of (1).

Solution of (8) multiplying both sides of (8) by  $v$  we get

$$uv \left( \frac{\partial^2 z}{\partial u \partial v} \right) + v \left( \frac{\partial z}{\partial v} \right) = 0 \quad (or) (uv D D' + v D') z = 0 \rightarrow (9)$$

where  $D \equiv \frac{\partial}{\partial u}$  and  $D' \equiv \frac{\partial}{\partial v}$ . To reduce (9) into linear equation with constant coefficients, we take new variables  $x$  and  $y$  as follows

$$\text{let } u = e^x \text{ and } v = e^y \text{ so that } x = \log u, y = \log v \rightarrow (10)$$

Let  $D_1 \equiv \frac{\partial}{\partial x}$  and  $D'_1 \equiv \frac{\partial}{\partial y}$  Then (9) reduces to

$$(D_1 D'_1 + D'_1) z = 0 \quad (or) D'_1 (D_1 + 1) z = 0$$

Its general solution is

$$z = e^{-x} \phi_1(y) + \phi_2(x) = u^{-1} \phi_1(\log v) + \phi_2(\log u)$$

$$(or) z = u^{-1} \psi_1(v) + \psi_2(u) = (y-u)^{-1} \psi_1(y^2 - u^2) + \psi_2(y-u),$$

where  $\psi_1$  and  $\psi_2$  are arbitrary functions.



**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

6.(d). The points of trisection of a string are pulled aside through a distance  $h$  on opposite sides of the position of equilibrium, and the string is released from rest. Derive an expression for the string at any subsequent time and show that the middle point of the string always remains at rest.

Sol'n:

Let the length of the string is  $= 3l$ .

The coordinates of A, C and D are  $(l, h)$ ,  $(2l, -h)$ ,  $(3l, 0)$  respectively. Initial deflection is given by OABCD. Equation of OA is

$$u = \frac{h}{l} (x-0) \text{ i.e. } u = \frac{hx}{l}$$

Equations of AC and CD respectively


are given by

$$u-h = \frac{-h-h}{2l-l} (x-l)$$

$$\Rightarrow u = \frac{h(3l-2x)}{l}$$

$$\text{and } u-(-h) = \frac{0-(-h)}{3l-2l} (x-2l)$$

$$\Rightarrow u = \frac{h(x-3l)}{l}$$



The required deflection is given by

$$u(x, t) = \sum_{n=1}^{\infty} E_n \cos \frac{n\pi ct}{3l} \sin \frac{n\pi x}{3l} \quad \text{--- (I)}$$

$$\text{where } E_n = \frac{2}{3l} \cdot \int_0^{3l} f(x) \sin \frac{n\pi x}{3l} dx.$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

The displacement  $y(x,t)$  of any point of the string is given by

$$\text{Boundary Conditions } y(0,t) = y(3l,t) = 0 \quad \forall t \geq 0 \quad \textcircled{1}$$

$$\text{Initial Conditions } y(x,0) = \begin{cases} hx/l & \text{when } 0 \leq x \leq l \\ \frac{h(3l-2x)}{l} & \text{when } l \leq x \leq 2l \\ \frac{h(x-3l)}{l} & \text{when } 2l \leq x \leq 3l \end{cases} \quad \textcircled{2}$$

$$\text{and } \left(\frac{\partial y}{\partial t}\right)_{t=0} = 0 \quad \textcircled{3}$$

we have

$$y(x,t) = \sum_{n=1}^{\infty} \left\{ E_n \cos \frac{n\pi ct}{3l} + F_n \sin \frac{n\pi ct}{3l} \right\} \sin \frac{n\pi x}{3l} \quad \textcircled{4}$$

Differentiating  $\textcircled{3}$  partially w.r.t  $t$ , we get

$$\frac{\partial y}{\partial t} = \sum_{n=1}^{\infty} \left\{ -E_n \frac{n\pi c}{3l} \sin \frac{n\pi ct}{3l} + \frac{n\pi c}{3l} F_n \cos \frac{n\pi ct}{3l} \right\} \sin \frac{n\pi x}{3l} \quad \textcircled{5}$$

Putting  $t=0$  in  $\textcircled{4}$  and  $\textcircled{5}$

and using the I.C  $\textcircled{5}$  and  $\textcircled{3}$  we get

$$\textcircled{6} \equiv \left(\frac{\partial y}{\partial t}\right)_{t=0} = \sum_{n=1}^{\infty} F_n \frac{n\pi c}{3l} \sin \frac{n\pi x}{3l} = 0 \quad (\text{by } \textcircled{3})$$

$$\text{where } F_n = \frac{2}{n\pi c} \int_0^{3l} f(x) \sin \frac{n\pi x}{3l} dx = 0$$

$$\textcircled{7} \equiv y(x,0) = f(x) = \sum_{n=1}^{\infty} E_n \sin \frac{n\pi x}{3l}$$

$$\text{where } E_n = \frac{2}{3l} \int_0^{3l} f(x) \sin \frac{n\pi x}{3l} dx \quad \textcircled{8}$$

Now

$$E_n = \frac{2}{3l} \int_0^{3l} f(x) \sin \frac{n\pi x}{3l} dx$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

$$\begin{aligned}
 &= \frac{2}{3l} \left[ \int_0^l f(x) \sin \frac{n\pi x}{3l} dx + \int_l^{2l} f(x) \sin \frac{n\pi x}{3l} dx + \int_{2l}^{3l} f(x) \sin \frac{n\pi x}{3l} dx \right] \\
 &= \frac{2}{3l} \left[ \int_0^l \frac{h}{l} \sin \frac{n\pi x}{3l} dx + \int_l^{2l} \frac{h(3l-2x)}{l} \sin \frac{n\pi x}{3l} dx \right. \\
 &\quad \left. + \int_{2l}^{3l} \frac{h(x-3l)}{l} \sin \frac{n\pi x}{3l} dx \right]
 \end{aligned}$$

Continuing in this way we get

$$\begin{aligned}
 E_n &= \frac{18h}{n^2\pi^2} \left[ \sin \frac{n\pi}{3} - \left\{ \sin \left( n\pi - \frac{n\pi}{3} \right) \right\} \right] \\
 &= \frac{18h}{n^2\pi^2} \left[ \sin \frac{n\pi}{3} - \left( \sin n\pi \cos \frac{n\pi}{3} - \cos n\pi \sin \frac{n\pi}{3} \right) \right] \\
 &= \frac{18h}{n^2\pi^2} \left[ \sin \frac{n\pi}{3} - 0 + \cos n\pi \sin \frac{n\pi}{3} \right] \quad [\because \sin n\pi = 0] \\
 &= \frac{18h}{n^2\pi^2} [1 + \cos n\pi] \sin \frac{n\pi}{3} = \frac{18h}{n^2\pi^2} [1 + (-1)^n] \sin \frac{n\pi}{3}.
 \end{aligned}$$

Thus  $E_n = 0$  if  $n$  is odd.

Thus  $E_n = 0$  if  $n$  is even put  $n = 2m$ ,  $m = 1, 2, \dots$

$$E_n = \frac{36h}{m^2\pi^2} \sin \frac{m\pi}{3} \quad \text{if } n \text{ is even, } m = 1, 2, \dots$$

$$\text{i.e., } E_n = \frac{36h}{4m^2\pi^2} \sin \frac{2m\pi}{3}, \quad m = 1, 2, \dots$$

$$= \frac{9h}{m^2\pi^2} \sin \frac{2m\pi}{3}$$

Putting the value of  $E_n$  &  $F_n$  in ④ the required deflection

is given by  $y(x,t) = \sum_{m=1}^{\infty} \frac{9h}{m^2\pi^2} \sin \frac{2m\pi}{3} \sin \frac{n\pi x}{3l} \cos \frac{n\pi ct}{3l}$

$$\Rightarrow y(x,t) = \frac{9h}{\pi^2} \sum_{m=1}^{\infty} \sin \frac{2m\pi}{3} \cos \frac{n\pi ct}{3l} \sin \frac{n\pi x}{3l} \quad ⑦$$

Putting  $x = \frac{3l}{2}$  in ⑦, we find that the displacement of the midpoint of the string i.e.  $y\left(\frac{3l}{2}, t\right) = 0$ .

because  $\sin m\pi = 0$ , for all integral values of  $m$ .

This shows that the mid-point of the string always rests.

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** *by K. Venkanna*

---

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**


Ques: 7(b)(i) Draw the circuit diagram for  $\bar{F} = A\bar{B}C + \bar{C}B$  using NAND to NAND logic long.

Solution:-

$$\bar{F} = A\bar{B}C + \bar{C}B$$

$$\bar{F} = \overline{\overline{A}\overline{B}C} + \overline{\overline{C}B}$$

$$\boxed{\bar{F} = \overline{\overline{A}\overline{B}C \cdot \overline{B}\overline{C}}}$$



Ques: 7(b) ii) In a Boolean Algebra B, for any a and b prove that  $ab' + a'b = 0$  if and only if  $a=b$ .

Solution: Given  $ab' + a'b = 0$   
 $\Leftrightarrow ab' = (a'b)'$   $[(a'b)' = (a'b)']$

$$ab' = a'' + b'$$

$$ab' = a + b'$$

$$ab' - (b') = a$$

$$b'(a-1) = a$$

$$-(b') = a$$

$$(b')' = a \Rightarrow \boxed{b=a}$$
 Hence proved.

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

7.(b)

(iii) Design a logic circuit having three inputs A, B, C such that output is 1 when A=0 or whenever B=C=1. Also obtain logic circuit using only NAND gates?

Solution

A	B	C	output = y	$m_i$ (output).
0	0	0	1	$m_0 = \bar{A}\bar{B}\bar{C}$
0	0	1	1	$m_1 = \bar{A}\bar{B}C$
0	1	0	1	$m_2 = \bar{A}BC$
0	1	1	1	$m_3 = \bar{A}B\bar{C}$
1	0	0	0	$m_4 = A\bar{B}\bar{C}$
1	0	1	0	$m_5 = A\bar{B}C$
1	1	0	0	$m_6 = AB\bar{C}$
1	1	1	1	$m_7 = ABC$

$$y = 1 \quad (\text{output})$$

$$y = m_0 + m_1 + m_2 + m_3 + m_7$$

$$y = \bar{A}\bar{B}\bar{C} + \bar{A}\bar{B}C + \bar{A}B\bar{C} + \bar{A}BC + ABC$$


$$y = \bar{A} [\bar{B}\bar{C} + \bar{B}C + B\bar{C} + BC] + ABC$$

$$y = \bar{A} [\bar{B}[\bar{C}+C] + B[\bar{C}+C]] + ABC$$

$$[\because \bar{C}+C=1]$$

$$y = \bar{A} [\bar{B}+B] + ABC \quad [\because \bar{B}+B=1]$$

$y = \bar{A} + ABC$



**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

f(x)

Using Runge - Kutta method of fourth order, solve

$$\frac{dy}{dx} = \frac{y^2 - x^2}{y^2 + x^2} \quad \text{with } y(0) = 1 \text{ at } x = 0.2, 0.4.$$

Sol'n: we have  $f(x, y) = \frac{y^2 - x^2}{y^2 + x^2}$

To find  $y(0.2)$ :

Here  $x_0 = 0, y_0 = 1, h = 0.2$

$$K_1 = h f(x_0, y_0) = (0.2) f(0, 1) = 0.2$$

$$K_2 = h f\left(x_0 + \frac{h}{2}, y_0 + \frac{k_1}{2}\right) = (0.2) f(0.1, 1.1) = 0.19672$$

$$K_3 = h f\left(x_0 + \frac{h}{2}, y_0 + \frac{k_2}{2}\right) = (0.2) f(0.1, 1.09836) = 0.1967$$

$$K_4 = h f\left(x_0 + h, y_0 + k_3\right) = (0.2) f(0.2, 1.1967) = 0.1891$$

$$K = \frac{1}{6}(K_1 + 2K_2 + 2K_3 + K_4) = 0.19599$$

$$\therefore y(0.2) = y_0 + K = 1.196.$$

To find  $y(0.4)$ :

Here  $x_1 = 0.2, y_1 = 1.196, h = 0.2$

$$K_1 = h f(x_1, y_1) = 0.1891$$

$$K_2 = h f\left(x_1 + \frac{h}{2}, y_1 + \frac{k_1}{2}\right) = (0.2) f(0.3, 1.2906) = 0.1795$$

$$K_3 = h f\left(x_1 + \frac{h}{2}, y_1 + \frac{k_2}{2}\right) = (0.2) f(0.3, 1.2858) = 0.1793$$

$$K_4 = h f\left(x_1 + h, y_1 + k_3\right) = (0.2) f(0.4, 1.3753) = 0.1688$$

$$K = \frac{1}{6}(K_1 + 2K_2 + 2K_3 + K_4) = 0.1792$$

$$\therefore y(0.4) = y_1 + K = 1.196 + 0.1792 = 1.3752$$

—————>

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** *by K. Venkanna*

---

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

8(a), Two equal rods AB and BC, each of length  $l$  smoothly joined at B are suspended from A and oscillate in a vertical plane through A. Show that the periods of normal oscillations are  $\frac{2\pi}{n}$ , where  $n^2 = \left(3 \pm \frac{6}{\sqrt{7}}\right) \frac{g}{l}$ .

Sol'n: Let AB and BC be the rods of equal length  $l$  and mass  $M$ . At time  $t$ , let the two rods make angles  $\theta$  and  $\phi$  to the vertical respectively.

Referred to A as origin horizontal and vertical lines AX and AY as axes the coordinates of C.G.  $G_1$ , of rod AB and that of C.G.  $G_2$  of rod BC are given by

$$x_{G_1} = \frac{1}{2}l \sin \theta, \quad y_{G_1} = \frac{1}{2}l \cos \theta$$

$$x_{G_2} = l \sin \theta + \frac{1}{2}l \sin \phi,$$

$$y_{G_2} = l \cos \theta + \frac{1}{2}l \cos \phi.$$

$\therefore$  If  $v_{G_1}$  and  $v_{G_2}$  are velocities of  $G_1$  &  $G_2$ , then

$$\begin{aligned} v_{G_1}^2 &= \dot{x}_{G_1}^2 + \dot{y}_{G_1}^2 = \left(\frac{1}{2}l \cos \theta \dot{\theta}\right)^2 + \left(-\frac{1}{2}l \sin \theta \dot{\theta}\right)^2 \\ &= \frac{1}{4}l^2 \dot{\theta}^2 \end{aligned}$$

$$\begin{aligned} v_{G_2}^2 &= \dot{x}_{G_2}^2 + \dot{y}_{G_2}^2 = \left(l \cos \theta \dot{\theta} + \frac{1}{2}l \cos \phi \dot{\phi}\right)^2 + \left(-l \sin \theta \dot{\theta} - \frac{1}{2}l \sin \phi \dot{\phi}\right)^2 \\ &= l^2 \left[\dot{\theta}^2 + \frac{1}{4}\dot{\phi}^2 + \dot{\theta}\dot{\phi} \cos(\theta - \phi)\right] \\ &= l^2 \left[\dot{\theta}^2 + \frac{1}{4}\dot{\phi}^2 + \dot{\theta}\dot{\phi}\right] \quad (\because \theta, \phi \text{ are small}) \end{aligned}$$


If  $T$  be the total kinetic energy and  $W$  the work function of the system, then

$$T = \text{K.E. of rod AB} + \text{K.E. of rod BC}$$

$$= \left[\frac{1}{2}M \cdot \frac{1}{3} \left(\frac{1}{2}l\right)^2 \dot{\theta}^2 + \frac{1}{2}M \cdot v_{G_1}^2\right] + \left[\frac{1}{2}m \cdot \frac{1}{3} \left(\frac{1}{2}l\right)^2 \dot{\phi}^2 + \frac{1}{2}M \cdot v_{G_2}^2\right]$$

$$= \frac{1}{2}M \left[ \frac{1}{12}l^2 \dot{\theta}^2 + \frac{1}{4}l^2 \dot{\phi}^2 \right] + \frac{1}{2}M \left[ \frac{1}{12}l^2 \dot{\theta}^2 + l^2 (\dot{\theta}^2 + \frac{1}{4}\dot{\phi}^2 + 2\dot{\theta}\dot{\phi}) \right]$$

$$= \frac{1}{2}Ml^2 \left( \frac{4}{3}\dot{\theta}^2 + \frac{1}{3}\dot{\phi}^2 + 2\dot{\theta}\dot{\phi} \right)$$



**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

$$\begin{aligned} \text{and } W &= MgY_{G_1} + MgY_{G_2} + C \\ &= Mg \left[ \frac{1}{2}l \cos\theta + l \cos\theta + \frac{1}{2}l \cos\phi \right] + C \\ &= \frac{1}{2}Mgl (3 \cos\theta + \cos\phi). \end{aligned}$$

∴ Lagrange's  $\theta$ -equation is  $\frac{d}{dt} \left( \frac{\partial T}{\partial \dot{\theta}} \right) - \frac{\partial T}{\partial \theta} = \frac{\partial W}{\partial \theta}$

$$\text{i.e. } \frac{d}{dt} \left[ \frac{1}{2}Ml^2 (8/3 \dot{\theta} + \dot{\phi}) \right] - 0 = \frac{1}{2}Mgl (-3 \sin\theta) = -\frac{3}{2}Mgl\theta \quad (\because \theta \text{ is small})$$

$$\Rightarrow 8\ddot{\theta} + 3\ddot{\phi} = -9c\theta, \text{ (where } c = g/l) \quad \text{--- (1)}$$

Equations (1) and (2) can be written as

$$(8D^2 + 9c)\theta + 3D^2\phi = 0 \text{ and } 3D^2\theta + \theta + (2D^2 + 3c)\phi = 0$$

Eliminating  $\phi$  between these two equations, we get

$$\begin{aligned} [(2D^2 + 3c)(8D^2 + 9c) - 9D^4]\theta &= 0 \\ \Rightarrow (7D^4 + 42cD^2 + 27c^2)\theta &= 0 \end{aligned}$$

If the periods of normal oscillations are  $2\pi/n$ , then  
the solution of (3), must be

$$\theta = A \cos(nt + B)$$

$$\therefore D^2\theta = -n^2\theta \text{ and } D^4\theta = n^4\theta.$$

Substituting in (3), we get

$$\begin{aligned} (7n^4 - 42cn^2 + 27c^2)\theta &= 0 \\ \Rightarrow 7n^4 - 42cn^2 + 27c^2 &= 0 \quad \because \theta \neq 0. \end{aligned}$$

$$\therefore n^2 = \frac{42 \pm \sqrt{(42c)^2 - 4 \cdot 7 \cdot 27c^2}}{2 \cdot 7}$$

$$\Rightarrow n^2 = \left( 3 \pm \frac{6}{\sqrt{7}} \right) c = \left( 3 + \frac{6}{\sqrt{7}} \right) \frac{g}{l} \quad (\because c = g/l)$$

————— .

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

8(b) A Sphere of radius  $a$  and mass  $M$  rolls down a rough plane inclined at an angle  $\alpha$  to the horizontal. If  $x$  be the distance of the point of contact of the sphere from a fixed point on the plane, find the acceleration by using Hamilton's equation.

Sol'n: Let a sphere of radius  $a$  and mass  $M$  roll down a rough plane inclined at an angle  $\alpha$  starting initially from a fixed point  $O$  of the plane. In time  $t$ , let the sphere roll down a distance  $x$  and during time let it turn through an angle  $\theta$ .

Since there is no slipping

$$\therefore x = OA = \text{arc } AB = a\theta.$$

$$\text{so that } \dot{x} = a\dot{\theta}$$

If  $T$  and  $V$  are the kinetic and potential energies of the sphere, then  $T = \frac{1}{2}MK^2\dot{\theta}^2 + \frac{1}{2}M\dot{x}^2 = \frac{1}{2}M\frac{2}{5}a^2\dot{\theta}^2 + \frac{1}{2}M(a\dot{\theta})^2$

$$\Rightarrow T = \frac{7}{10}M\dot{x}^2$$

and  $V = -MgOL = -Mgx \sin\alpha$  (since the sphere moves down the plane)

$$\therefore L = T - V = \frac{7}{10}M\dot{x}^2 + Mgx \sin\alpha$$

Here  $x$  is the only generalised coordinate.


$$\therefore p_x = \frac{\partial L}{\partial \dot{x}} = \frac{7}{5}M\dot{x} \quad \text{--- (1)}$$

Since  $L$  does not contain  $t$  explicitly,

$$\therefore H = T + V = \frac{7}{10}M\dot{x}^2 - Mgx \sin\alpha$$

$$\Rightarrow H = \frac{7}{10}M \left( \frac{5}{7M}p_x \right)^2 - Mgx \sin\alpha$$

$$= \frac{5}{14M}p_x^2 - Mgx \sin\alpha \quad \text{from (1)}$$



**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

Hence the two Hamilton's equations are

$$\dot{p}_x = -\frac{\partial H}{\partial x} = Mg \sin \alpha - (H_1), \quad \dot{x} = \frac{\partial H}{\partial p_x} = \frac{5}{7M} p_x - (H_2)$$

Differentiating (H<sub>2</sub>) and using (H<sub>1</sub>), we get

$$\ddot{x} = \frac{5}{7M} \dot{p}_x = \frac{5}{7M} Mg \sin \alpha$$

$$\Rightarrow \boxed{\ddot{x} = \frac{5}{7} g \sin \alpha}$$

which gives the required acceleration.  
=====.

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

Ques: 8(c)) Show that  $\phi = x f(\lambda)$  is a possible form for the velocity potential for an incompressible fluid motion. If the fluid velocity  $\vec{q} \rightarrow 0$  as  $\lambda \rightarrow \infty$ , find the surfaces of constant speed.

Solution:- Given;  $\phi = x f(\lambda) \quad \dots \quad (1)$

$$q_r = -\nabla \phi = -\nabla [x f(\lambda)].$$

$$q_r = -[f(\lambda) \nabla x + x \nabla f(\lambda)] \quad \dots \quad (2)$$

$$\text{Now, } r^2 = x^2 + y^2 + z^2$$

$$\Rightarrow 2x \left( \frac{\partial \lambda}{\partial x} \right) = 2x \Rightarrow \frac{\partial \lambda}{\partial x} = \frac{x}{\lambda} \quad \dots \quad (3)$$

$$\text{Similarly, } \frac{\partial \lambda}{\partial y} = \frac{y}{\lambda} \text{ and } \frac{\partial \lambda}{\partial z} = \frac{z}{\lambda} \quad \dots \quad (4)$$

$$\text{Also; } \nabla x = \left[ i \left( \frac{\partial}{\partial x} \right) + j \left( \frac{\partial}{\partial y} \right) + k \left( \frac{\partial}{\partial z} \right) \right] x = i$$

$$\text{and } \nabla f(\lambda) = \left[ i \left( \frac{\partial}{\partial x} \right) + j \left( \frac{\partial}{\partial y} \right) + k \left( \frac{\partial}{\partial z} \right) \right] f(\lambda)$$

$$\nabla f(\lambda) = i f'(\lambda) \left( \frac{\partial \lambda}{\partial x} \right) + j f'(\lambda) \left( \frac{\partial \lambda}{\partial y} \right) + k f'(\lambda) \left( \frac{\partial \lambda}{\partial z} \right)$$

$$\nabla f(\lambda) = \hat{i} f'(\lambda) \left( \frac{x}{\lambda} \right) + \hat{j} f'(\lambda) \left( \frac{y}{\lambda} \right) + \hat{k} f'(\lambda) \left( \frac{z}{\lambda} \right)$$

$$\nabla f(\lambda) = \frac{1}{\lambda} f'(\lambda) (\hat{i} x + \hat{j} y + \hat{k} z) = \frac{1}{\lambda} f'(\lambda) \cdot \vec{\lambda}$$

$$\therefore (2) \Rightarrow q_r = -f(\lambda) i - \left( \frac{x}{\lambda} \right) f'(\lambda) \cdot \vec{\lambda} \quad \dots \quad (5)$$

For a possible motion of an incompressible fluid, we have —

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS** by K. Venkanna

$$\nabla \cdot \mathbf{q} = 0 \quad \text{or} \quad \nabla \cdot (-\nabla \phi) = 0$$

$$\Rightarrow \nabla^2 \phi = 0$$

$$\Rightarrow \left[ \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right] [x f(x)] = 0, \text{ (using (1))} \quad \text{--- (5)}$$

$$\begin{aligned} \text{Now, } \frac{\partial^2}{\partial x^2} [x f(x)] &= \frac{\partial}{\partial x} \left[ \frac{\partial}{\partial x} [x f(x)] \right] \\ &= \frac{\partial}{\partial x} \left[ f(x) + x \frac{\partial f(x)}{\partial x} \right] \end{aligned}$$

$$\therefore \frac{\partial^2}{\partial x^2} [x f(x)] = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial x} + x \frac{\partial^2 f(x)}{\partial x^2} = 2 \frac{\partial f}{\partial x} + x \frac{\partial^2 f(x)}{\partial x^2}.$$

$$\text{Also; } \frac{\partial^2}{\partial y^2} [x f(x)] = x \frac{\partial^2 f}{\partial y^2}$$

$$\frac{\partial^2}{\partial z^2} [x f(x)] = x \frac{\partial^2 f}{\partial z^2}$$

$\therefore$  (6) becomes

$$2 \frac{\partial f}{\partial x} + x \left( \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \right) = 0 \quad \text{--- (7)}$$

$$\text{Now, } \frac{\partial f}{\partial x} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial x} = f'(x/x) \quad \text{--- using (3)}$$

$$\text{and } \frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right) = \frac{\partial}{\partial x} \left( f'(x/x) \right) = \frac{f''}{x} + x \frac{\partial}{\partial x} \left( \frac{f'}{x} \right) \quad \text{--- (8)}$$

$$= \frac{f'}{x} + x \frac{\partial}{\partial x} \left( \frac{f'}{x} \right) \frac{\partial x}{\partial x} = \frac{f'}{x} + x \left( \frac{\partial f'' - f'}{x^2} \right) \frac{x}{x}$$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

$$\frac{\partial^2 f}{\partial x^2} = \frac{f'}{x} + \frac{x^2}{x^2} f'' - \frac{x^2}{x^3} f' \quad \dots \quad (9)$$

Similarly,

$$\frac{\partial^2 f}{\partial y^2} = \frac{f'}{x} + \frac{y^2}{x^2} f'' - \frac{y^2}{x^3} f' \quad \dots \quad (10)$$

$$\text{and } \frac{\partial^2 f}{\partial z^2} = \frac{f'}{x} + \frac{z^2}{x^2} f'' - \frac{z^2}{x^3} f' \quad \dots \quad (11)$$

Adding (9), (10) and (11), we get.

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = \frac{3f'}{x} + \left[ \frac{x^2 + y^2 + z^2}{x^2} \right] f'' - \left( \frac{x^2 + y^2 + z^2}{x^3} \right) f'$$

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = \frac{3f'}{x} + \frac{x^2}{x^2} \cdot f'' - \frac{1}{x} \cdot f'$$

$$\boxed{\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = \frac{3f'}{x} + f'' - \frac{1}{x} f' = \frac{2}{x} f' + f''} \quad \dots \quad (12)$$

Using (8) and (12), (7) reduces to

$$\frac{2f'(x)}{x} + x \left( \frac{2f'}{x} + f'' \right) = 0$$

$$\Rightarrow f'' + \frac{4f'}{x} = 0$$

$$\Rightarrow \boxed{\frac{f''}{f'} + \frac{4}{x} = 0}$$

Integrating  $\log f' + 4 \log x = \log c_1$

so that  $\Rightarrow \boxed{f' = c_1 x^{-4}} \quad \dots \quad (13)$

Integrating (13);  $f = -\left(\frac{c_1}{3}\right) x^{-3} + c_2$

**INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS**  
**MATHEMATICS by K. Venkanna**

$c_2$  being an arbitrary constant

Substituting the values of  $f'$  and  $f$  from

(13) & (14) in (5), we get

$$\vec{q} = -\left\{ \left( c_1/3x^3 \right) - c_2 \right\} \vec{i} - \left( c_1 x / x^5 \right) \vec{x} \quad (15)$$

Given, that  $\vec{q} \rightarrow 0$  as  $x \rightarrow \infty$ ,

hence (15) shows that  $c_2 = 0$

$$\therefore \text{from (15)} ; \boxed{\vec{q} = \frac{c_1}{3x^3} \left( \vec{i} - \frac{3x\vec{x}}{x^2} \right)} \quad 16$$

$$\text{Now}; \quad q^2 = \vec{q} \cdot \vec{q} = \frac{c_1^2}{9x^6} \left( \vec{i} - \frac{3x\vec{x}}{x^2} \right) \cdot \left( \vec{i} - \frac{3x\vec{x}}{x^2} \right)$$

$$= \frac{c_1^2}{9x^6} \left[ \vec{i} \cdot \vec{i} - \frac{6x \cdot \vec{x} \cdot \vec{i}}{x^2} + \frac{9x^2 \cdot x^2}{x^4} \right]$$

$$\text{as } \vec{x} \cdot \vec{x} = x^2 \\ \text{& } \vec{x} \cdot \vec{i} = x$$

$$= \frac{c_1^2}{9x^6} \left( 1 - \frac{6x^2}{x^2} + \frac{9x^2 \cdot x^2}{x^4} \right)$$

$$= \frac{c_1^2}{9x^6} \left( 1 + \frac{3x^2}{x^2} \right)$$

$$\boxed{q^2 = \frac{c_1^2}{9x^8} (x^2 + 3x^2)}$$

Hence, the required surfaces of constant speed are

$$q^2 = \text{constant} \Leftrightarrow \left( \frac{c_1^2}{9x^8} \right) (x^2 + 3x^2) = \text{constant}$$

$$\boxed{q^2 = (x^2 + 3x^2)x^{-8} = \text{constant}} \quad \text{required solution}$$