

11 Years
Solved Papers
2009-2019



Civil Services **Main Examination**

TOPICWISE PREVIOUS YEARS' SOLVED PAPERS

Mathematics
Paper-II



11 Years

Previous Years Solved Papers

Civil Services Main Examination

(2009-2019)

Mathematics Paper-II

Topicwise Presentation



MADE EASY
Publications

10.8.2020

Previous Years Solved Papers of
Civil Services Main Examination

Mathematics : Paper-II

CONTENTS

SI.	TOPIC	PAGE No.
Unit-1	Modern Algebra	1-37
	1. Groups.....	1
	2. Rings.....	18
	3. Fields.....	32
Unit-2	Real Analysis	38-81
	1. Real Number System	38
	2. Sequences	38
	3. Series	45
	4. Functions	55
	5. Riemann Integral.....	66
	6. Improper Integral.....	72
	7. Functions of Several (Two or Three) Variables.....	73
Unit-3	Complex Analysis.....	82-113
	1. Analytic Functions	82
	2. Taylor's Series	93
	3. Singularity	96
	4. Laurant's Series	96
	5. Cauchy's Residue Theorem.....	102
	6. Contour Integration.....	107

Unit-4	Linear Programming Problems.....	114-144
1.	Basic Feasible Solutions.....	114
2.	Optimal Solutions.....	115
3.	Graphical Method of Solution.....	117
4.	Simplex Method of Solution.....	123
5.	Duality.....	135
6.	Transportation Problems	137
7.	Assignment Problems	141
Unit-5	Partial Differential Equations	145-209
1.	Formulation of P.D.E.....	145
2.	Solution of Quasilinear PDE of 1st Order, Lagrange's Auxiliary Equations and Charpit's Auxiliary Equations	150
3.	Cauchy's Method of Characteristics	159
4.	Linear PDE of Second Order with Constant Coefficient.....	165
5.	Canonical Form.....	172
6.	Application of PDE.....	182
Unit-6	Numerical Analysis and Computer Programming	210-256
1.	Solution of Algebraic and Transcendental Equation of One Variable.....	210
2.	Solutions of System of Linear Equations.....	214
3.	Interpolation.....	221
4.	Numerical Integration	229
5.	Ordinary Differential Equation.....	237
6.	Computer Programming	241
Unit-7	Mechanics & Fluid Dynamics	257-313
1.	Generalised Coordinate.....	257
2.	Moment of Inertia.....	272
3.	Motion of Rigid Bodies in Two Dimensions.....	282
4.	Equation of Continuity.....	283
5.	Euler's Equation of Motion for Inviscid Flow.....	295
6.	Source and Sink	300
7.	Vortex Motion	303
8.	Navier-Stoke's Equation for Viscous Flow.....	310

3

Complex Analysis

1. Analytic Functions

1.1 Let $f(z) = \frac{a_0 + a_1 z + \dots + a_{n-1} z^{n-1}}{b_0 + b_1 z + \dots + b_n z^n}$, $b_n \neq 0$. Assume that the zeroes of the denominator are simple.

Show that the sum of the residues of $f(z)$ at its poles is equal to $\frac{a_n - 1}{b_n}$.

(2009 : 12 Marks)

Solution:

Proof : As the zeroes of denominator are simple, so $b_0 + b_1 z + \dots + b_n z^n$ can be written as

$$b_n(z - z_0)(z - z_1)(z - z_2) \dots (z - z_{n-1})$$

or

$$f(z) = \frac{a_0 + a_1 z + \dots + a_{n-1} z^{n-1}}{b_n(z - z_0)(z - z_1)(z - z_2) \dots (z - z_{n-1})} \quad \dots(1)$$

As maximum degree of 'z' in numerator is $n-1$ and degree of denominator is n (i.e., degree of denominator is greater than degree of numerator).

So, $f(z)$ can be written as

$$f(z) = \frac{A_0}{z - z_0} + \frac{A_1}{z - z_1} + \frac{A_2}{z - z_2} + \dots + \frac{A_{n-1}}{z - z_{n-1}} \quad \dots(2)$$

$$= \frac{A_0(z - z_1)(z - z_2) \dots (z - z_{n-1}) + A_1(z - z_0)(z - z_2) - (z - z_{n-1}) + \dots + A_{n-1}(z - z_0)(z - z_{n-2})}{(z - z_0)(z - z_1) \dots (z - z_{n-1})} \quad \dots(3)$$

Now, we know that residues of $f(z)$ are $A_0, A_1, A_2, \dots, A_{n-1}$.

According to problem, we need to calculate $A_0 + A_1 + A_2 + \dots + A_{n-1}$.

From (3), the coefficient of z^{n-1} in numerator will be calculated as :

$$A_0 + A_1 + A_2 + \dots + A_{n-1}$$

because on observing

$$\begin{aligned} & A_0(z - z_1)(z - z_2) - (z - z_{n-1}) \\ & + A_1(z - z_0)(z - z_2) - (z - z_{n-1}) \\ & + A_2(z - z_0)(z - z_1)(z - z_3) - (z - z_{n-1}) \\ & + \dots \\ & + A_{n-1}(z - z_0)(z - z_1) \dots (z - z_{n-2}) \end{aligned}$$

In each of terms respectively coefficient of z^{n-1} will be $A_0, A_1, A_2, \dots, A_{n-1}$. So, the overall coefficient of z^{n-1} in numerator will be $A_0 + A_1 + A_2 + \dots + A_{n-1}$.

Now, comparing this coefficient of z^{n-1} with (1) i.e.,

$$\frac{a_0 + a_1 z + a_2 z^2 + \dots + a_{n-1} z^{n-1}}{(z - z_0)(z - z_1) \dots (z - z_{n-1})} \quad \dots(4)$$

In this fraction, coefficient of z^{n-1} will be $\frac{a_{n-1}}{b_n}$.
 (as denominator $(z - z_0)(z - z_1) \dots (z - z_{n-1})$ is common (3) and (4))

$$\therefore A_0 + A_1 + A_2 \dots A_{n-1} = \frac{a_{n-1}}{b_n}$$

or Sum of residues = $\frac{a_{n-1}}{b_n}$. Thus Proved.

- 1.2 Show that $u(x, y) = 2x - x^3 + 3xy^2$ is a harmonic function. Find a harmonic conjugate of $u(x, y)$. Hence, find the analytic function f for which $u(x, y)$ is the real part.

(2010 : 12 Marks)

Solution:

Given

$$u(x, y) = 2x - x^3 + 3xy^2$$

$$\frac{\partial u}{\partial x} = 2 - 3x^2 + 3y^2, \quad \frac{\partial^2 u}{\partial x^2} = -6x$$

$$\frac{\partial u}{\partial y} = 6xy, \quad \frac{\partial^2 u}{\partial y^2} = 6x$$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = -6x + 6x = 0$$

$\therefore u(x, y)$ is a harmonic function.

Let v be its harmonic conjugate.

$$\therefore \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} = 2 - 3x^2 + 3y^2$$

$$\Rightarrow v = 2y - 3x^2y + y^3 + f(x) \quad \dots(1)$$

$$\text{Also, } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} = 6xy$$

$$\Rightarrow \frac{\partial v}{\partial x} = -6xy$$

$$\Rightarrow v = -3x^2y + g(y) \quad \dots(2) \quad (f(x) \text{ and } g(y) \text{ are some arbitrary functions})$$

Comparing (1) and (2), we get

$$v = 2y + y^3 - 3x^2y + c, \text{ where } c \text{ is a constant.}$$

$$\text{Now, let } f = u(x, y) + iV(x, y)$$

$$\Rightarrow f = (2x - x^3 + 3xy^2) + i(2y + y^3 - 3x^2y + c)$$

$$f(z) = \int (\phi_1(z, 0) - i\phi_2(z, 0)) dz$$

$$= \int (2 - 3z^2) dz = 2z - z^3 + A, \text{ where } A \text{ is a constant}$$

$$\Rightarrow f(z) = 2z - z^3 + A$$

- 1.3 (i) Evaluate the line integral $\int_C f(z) dz$ where $f(z) = z^2$, C is the boundary of the triangle with vertices $A(0, 0)$, $B(1, 0)$, $C(1, 2)$ in that order.

- (ii) Find the image of the finite vertical strip $R : x = 5 \text{ to } x = 9, -\pi \leq y \leq \pi$ of z -plane under exponential function.

(2010 : 15 Marks)

Solution:

(i) Given : $f(z) = z^2$

Now, $f(z) = z^2$ is analytic in whole given region.

Line integral of closed boundary curve of analytic function is always zero.

$\therefore \int_C f(z) dz$ in the given region is 0.

(ii) Given : x varies from 5 to 9 and y varies from $-\pi$ to π .

Now, $z = x + iy$

Image under exponential function will be


$$e^z = e^{x+iy} = e^x \cdot e^{iy}$$

As x varies from 5 to 9, $\therefore e^x$ varies from e^5 to e^9 .

Also, e^{iy} denotes angle variation.

So, image is given by region in figure.

So, the region is bounded by two circles of radii e^5 and e^9 .



1.4 If $f(z) = u + iv$ is an analytic function of $z = x + iy$ and $u - v = \frac{e^y - \cos x + \sin x}{\cosh y - \cos x}$, find $f(z)$ subject to the condition, $f\left(\frac{\pi}{2}\right) = \frac{3-i}{2}$.

(2011 : 12 Marks)

Solution:

Let

$$f(z) = u + iv$$

\Rightarrow

$$if(z) = iu - v$$

\Rightarrow

$$(1+i)f(z) = u - v + i(u + v) \\ = u + iv$$

\therefore

$$U = u - v = \frac{e^y - \cos x + \sin x}{\cosh y - \cos x}$$

$$= \frac{\cosh y + \sinh y - \cos x + \sin x}{\cosh y - \cos x}$$

$$= 1 + \frac{\sinh y + \sin x}{\cosh y - \cos x}$$

Let

$$\frac{\partial U}{\partial x} = \phi_1(x, y) \text{ and } \frac{\partial U}{\partial y} = \phi_2(x, y)$$

\therefore

$$\phi_1(x, y) = \frac{\partial U}{\partial x} = \frac{\cos x (\cosh y - \cos x) - \sin x (\sinh y + \sin x)}{(\cosh y - \cos x)^2}$$

\therefore

$$\phi_1(z, 0) = \frac{\cos z (1 - \cos z) - \sin^2 z}{(1 - \cos z)^2} = \frac{\cos z - 1}{(1 - \cos z)^2}$$

$$= \frac{-1}{1 - \cos z} = \frac{-1}{2} \operatorname{cosec}^2 \frac{z}{2}$$

$$\phi_2(x, y) = \frac{\partial U}{\partial y} = \frac{\cosh y (\cosh y - \cos x) - \sinh y (\sinh y + \sin x)}{(\cosh y - \cos x)^2}$$

\therefore

$$\phi_2(z, 0) = \frac{1 - \cos z}{(1 - \cos z)^2} = \frac{1}{1 - \cos z}$$

$$= \frac{1}{2} \operatorname{cosec}^2 \frac{z}{2}$$

∴ By Milne's Method

$$\begin{aligned}(1+i)f(z) &= \int [\phi_1(z, 0) - i\phi_2(z, 0)] dz + C \\&= \int \left(-\frac{1}{2} \operatorname{cosec}^2 \frac{z}{2} - i \frac{1}{2} \operatorname{cosec}^2 \frac{z}{2} \right) dz + C \\&= -\frac{1}{2}(1+i) \int \operatorname{cosec}^2 \frac{z}{2} dz + C \\&= (1+i) \cot \frac{z}{2} + C \\f(z) &= \cot \frac{z}{2} + \frac{C}{1+i}\end{aligned}$$

$$\text{At } z = \frac{\pi}{2},$$

$$f(z) = \frac{3-i}{2}$$

∴

$$d = f\left(\frac{\pi}{2}\right) - \cot \frac{\pi}{4} = \frac{1-i}{2}$$

∴

$$f(z) = \cot \frac{z}{2} + \frac{1-i}{2}$$

1.5 Show that the function defined by

$$f(z) = \begin{cases} \frac{x^3 y^5 (x+iy)}{x^6 + y^{10}}, & z \neq 0 \\ 0, & z = 0 \end{cases}$$

is not analytic at the origin though it satisfies Cauchy-Riemann equations at the origin.

(2012 : 12 Marks)

Solution:

It is given that

$$f(z) = \begin{cases} \frac{x^3 y^5 (x+iy)}{x^6 + y^{10}}, & z \neq 0 \\ 0, & z = 0 \end{cases}$$

We have,

$$\begin{aligned}\frac{f(z) - f(0)}{z} &= \frac{\frac{x^3 y^5 (x+iy)}{x^6 + y^{10}} - 0}{x+iy} \\&= \frac{x^3 y^5}{x^6 + y^{10}}\end{aligned}$$

If $z = 0$ along $y = x$, then

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z} = \lim_{x \rightarrow 0} \frac{x^8}{x^6 + x^{10}} = \lim_{x \rightarrow 0} \frac{x^2}{1+x^4} = 0 \quad \dots(i)$$

If $z = 0$ along $y^{10} = x^6$, then

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z} = \lim_{x \rightarrow 0} \frac{x^3 \cdot x^3}{x^6 + x^6} = \lim_{x \rightarrow 0} \frac{x^6}{2x^6} = \frac{1}{2} \quad \dots(ii)$$

Hence, $f(z)$ is not analytic as the limits in (i) and (ii) are not unique.

Again

$$\begin{aligned} f(z) &= \frac{x^3 y^5 (x + iy)}{x^6 + y^{10}} \\ &= \frac{x^4 y^5}{x^6 + y^{10}} + i \frac{x^3 y^6}{x^6 + y^{10}} = u + iv \end{aligned}$$

\therefore

$$\begin{aligned} \frac{\partial u}{\partial x} &= \lim_{x \rightarrow 0} \frac{u(x, 0) - u(0, 0)}{x} \\ &= \lim_{x \rightarrow 0} \frac{0 - 0}{x} = 0 \end{aligned}$$

$$\frac{\partial u}{\partial y} = \lim_{y \rightarrow 0} \frac{u(0, y) - u(0, 0)}{y} = \lim_{y \rightarrow 0} \frac{0 - 0}{y} = 0$$

$$\frac{\partial v}{\partial x} = \lim_{x \rightarrow 0} \frac{v(x, 0) - v(0, 0)}{x} = \lim_{x \rightarrow 0} \frac{0 - 0}{x} = 0$$

$$\frac{\partial v}{\partial y} = \lim_{y \rightarrow 0} \frac{v(0, y) - v(0, 0)}{y} = \lim_{y \rightarrow 0} \frac{0 - 0}{y} = 0$$

Since,

$$\frac{\partial y}{\partial x} = \frac{\partial v}{\partial y}$$

and

$$\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

\therefore Cauchy-Riemann equations are satisfied at the origin.

- 1.6 Use Cauchy integral formula to evaluate $\int_C \frac{e^{3z}}{(z+1)^4} dz$, where c is the circle $|z| = 2$.

(2012 : 15 Marks)

Solution:

From Cauchy Integral Formula, we have

$$f^{(n)}(a) = \frac{n!}{2\pi i} \int_C \frac{f(z) dz}{(z-a)^{n+1}}$$

Put $a = -1$, $n = 3$

$$f^{(3)}(-1) = \frac{3!}{2\pi i} \int_C \frac{f(z) dz}{(z+1)^4} \quad \dots(i)$$

Take $f(z) = e^{3z}$, then

$$f^{(n)}(z) = 3^n e^{3z}$$

\therefore

$$f^{(3)}(-1) = 3^3 e^{-3} = \frac{27}{e^3}$$

\therefore from (i),

$$\frac{27}{e^3} = \frac{3!}{2\pi i} \int_C \frac{e^{3z} dz}{(z+1)^4}$$

\Rightarrow

$$\int_C \frac{e^{3z} dz}{(z+1)^4} = \frac{9\pi i}{e^3}$$

- 1.7 Prove that if $be^{a+1} < 1$ where a and b are positive and real, then the function $z^n e^{-a} - be^z$ has n zeros in the unit circle.

Solution:

(2013 : 10 Marks)

Approach : We use Rouché's theorem which says that if $f(z)$ and $g(z)$ are analytic functions in a simply connected domain. And $|f(z) + g(z)| < |f(z)|$ on C a rectifiable curve in the domain then $f(z)$ and $g(z)$ has same number of zeros in the domain.

Let

and

\therefore

and

and since

\therefore

$$g(z) = z^n e^{-a} - be^z$$

$$f(z) = -z^n e^{-a}$$

$$f(z) + g(z) = -be^z$$

$$|f(z) + g(z)| = |-be^z| = be \text{ on } |z| = 1$$

$$|f(z)| = |-z^n e^{-a}| = e^{-a} |z^n| = e^{-a} \text{ on } |z| = 1$$

$$be^{a+1} < 1 \Rightarrow be < e^{-a} \text{ on } |z| = 1$$

$$|f(z) + g(z)| < |f(z)|$$

Also $f(z)$ and $g(z)$ are analytic on and inside the unit circle. Hence conditions of Rouché's theorem are satisfied and so $f(z)$ and $g(z)$ has same number of zeros inside unit circle, i.e.,

$$g(z) = z^n e^{-a} - be^z$$

has n zeros inside unit circle.

- 1.8 Prove that the function $f(z) = u + iv$, where

$$f(z) = \frac{x^3(1+i) - y^3(1-i)}{x^2 + y^2}, z \neq 0; f(0) = 0$$

satisfies Cauchy-Riemann equations at the origin, but the derivative of f at $z = 0$ does not exist.

(2014 : 10 Marks)

Solution:

Here,

$$u = \frac{x^3 y^3}{x^2 + y^2}$$

$$v = \frac{x^3 + y^3}{x^2 + y^2}$$

where $z \neq 0$

Here we see the both u and v are rational and finite for all values of $z \neq 0$. So u and v are continuous at all those points for which $z \neq 0$.

Hence $f(z)$ is continuous.

Where $z \neq 0$

At the origin $u = 0, v = 0$,

[since $f(0) = 0$]

Hence u and v are both continuous at the origin consequently $f(z)$ is continuous at the origin.

$\therefore f(z)$ is continuous every-where at the origin.

$$\frac{\partial u}{\partial x} = \lim_{x \rightarrow 0} \frac{u(x,0) - u(0,0)}{x} = \lim_{x \rightarrow 0} \left(\frac{x}{x} \right) = 1$$

$$\frac{\partial u}{\partial x} = \lim_{y \rightarrow 0} \frac{u(0,y) - u(0,0)}{y} = \lim_{y \rightarrow 0} \left(\frac{-y}{y} \right) = -1$$

$$\frac{\partial u}{\partial x} = \lim_{x \rightarrow 0} \frac{u(x,0) - v(0,0)}{x} = \lim_{x \rightarrow 0} \left(\frac{x}{x} \right) = 1$$

$$\frac{\partial u}{\partial y} = \lim_{y \rightarrow 0} \frac{v(0,y) - v(0,0)}{y} = \lim_{y \rightarrow 0} \left(\frac{y}{y} \right) = 1$$

\therefore We see that $\frac{\partial U}{\partial x} = \frac{\partial V}{\partial y}$ and $\frac{\partial U}{\partial y} = -\frac{\partial V}{\partial x}$

Hence Cauchy-Riemann conditions are satisfied at $z=0$.

Again,

$$\begin{aligned} f'(0) &= \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z} \\ &= \lim_{x \rightarrow 0} \left[\frac{x^3 - y^3 + i(x^3 + y^3)}{x^2 + y^2} \cdot \frac{1}{x + iy} \right] \end{aligned}$$

Now, let $z \rightarrow 0$ along $y=x$, then

$$\begin{aligned} f'(0) &= \lim_{x \rightarrow 0} \frac{x^3 - x^3 + i(x^3 + x^3)}{x^2 + x^2} \cdot \frac{1}{x + ix} \\ &= \lim_{x \rightarrow 0} \frac{2i}{2(1+i)} = \frac{1}{2}(1-i) \end{aligned}$$

Again let $z \rightarrow 0$ along $y=0$, then

$$f'(0) = \lim_{x \rightarrow 0} \frac{x^3(1+i)}{x^3} = 1+i$$

\therefore we see that $f'(0)$ is not unique.

i.e. the values of $f'(0)$ are not the same as $z \rightarrow 0$ along different curves.

\therefore

$f'(z) =$ does not exist at the origin.

- 1.9 Show that the function $v(x, y) = \ln(x^2 + y^2) + x + y$ is harmonic. Find its conjugate harmonic function $u(x, y)$. Also, find the corresponding analytic function $f(z) = u + iv$, in terms of z .

(2015 : 10 Marks)

Solution:

Given,

$$v(x, y) = \ln(x^2 + y^2) + x + y$$

$$\frac{\partial v}{\partial x} = \frac{2x}{x^2 + y^2} + 1, \quad \frac{\partial^2 v}{\partial x^2} = \frac{2(y^2 - x^2)}{(x^2 + y^2)^2}$$

$$\frac{\partial v}{\partial y} = \frac{2y}{x^2 + y^2} + 1, \quad \frac{\partial^2 v}{\partial y^2} = \frac{2(x^2 - y^2)}{(x^2 + y^2)^2}$$

Now,

$$\frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = \frac{2(y^2 - x^2)}{(x^2 + y^2)^2} + \frac{2(x^2 - y^2)}{(x^2 + y^2)^2} = 0$$

$\therefore v$ is a harmonic function.

Given, u is harmonic conjugate of v .

\therefore By Cauchy-Riemann equations,

$$u_x = v_y \text{ and } u_y = -v_x$$

$$\frac{\partial u}{\partial x} = \frac{2y}{x^2 + y^2} + 1$$

\Rightarrow

$$u = 2\tan^{-1}\left(\frac{x}{y}\right) + x + f_1(y) \quad \dots(i)$$

Similarly,

$$\frac{\partial u}{\partial y} = \frac{-2x}{x^2 + y^2} - 1$$

\Rightarrow

$$u = -2\tan^{-1}\left(\frac{y}{x}\right) - y + f_2(x) \quad \dots(ii)$$

From (i) and (ii)

$$u = 2 \tan^{-1} \left(\frac{x}{y} \right) + x - y + c, \text{ where } c \text{ is a constant.}$$

$$v_x = \frac{2x}{x^2 + y^2} + 1 = \psi_2$$

$$v_y = \frac{2y}{x^2 + y^2} + 1 = \psi_1$$

∴ By Milne Thomson's method

$$f(z) = \int (\psi_1(z, 0) + i(\psi_2(z, 0))) dz$$

$$= \int \left[1 + i \left(\frac{2z}{z^2} + 1 \right) \right] dz$$

$$f(z) = z + i(2 \log z + z) + k, \text{ where } k \text{ is a constant.}$$

- 1.10 Is $v(x, y) = x^3 - 3xy^2 + 2y$ a harmonic function? Prove your claim. If yes, find its conjugate harmonic function $u(x, y)$ and hence obtain the analytic function whose real and imaginary parts are u and v respectively.

(2016 : 10 Marks)

Solution:

Let f be the function

$$f = u + iv \quad (\text{given})$$

Given :

$$v = x^3 - 3xy^2 + 2y$$

∴

$$v_x = 3x^2 - 3y^2, v_{xx} = 6x \quad \dots(i)$$

$$v_y = -6xy + 2, v_{yy} = -6x \quad \dots(ii)$$

Now,

$$v_{xx} + v_{yy} = +6x - 6x = 0$$

⇒ v is a harmonic function.

u is harmonic conjugate of v .

By Cauchy-Riemann equations,

$$u_x = v_y \Rightarrow u_x = -6xy + 2 \quad (\text{from (ii)})$$

⇒

$$\frac{\partial u}{\partial x} = -6xy + 2$$

⇒

$$\int dy = \int (-6xy + 2) dx$$

⇒

$$u = -3x^2y + 2x + g(y) \quad \dots(iii)$$

where g is an arbitrary function of y

Also,

$$u_y = -v_x \Rightarrow u_y = -(3x^2 - 3y^2) \quad (\text{from (i)})$$

⇒

$$\frac{\partial u}{\partial y} = -3x^2 + 3y^2$$

⇒

$$\int du = \int (-3x^2 + 3y^2) dy$$

⇒

$$u = -3x^2y + y^3 + n(x), \text{ where } n(x) \text{ is an arbitrary function of } x. \dots(iv)$$

From (iii) & (iv)

$$u = -3x^2y + 2x + y^3 + c, \text{ where } c \text{ is a constant}$$

$$f = (-3x^2y + 2x + y^3 + c) + i(x^3 - 3xy^2 + 2y)$$

So,

- 1.11 Prove that every power series represents an analytic function inside its circle of convergence.

(2016 : 20 Marks)

Solution:

Let the series be $f(z) = \sum_{n=0}^{\infty} a_n z^n$ and R be its radius of convergence.

Then,

$$\frac{1}{R} = \overline{\lim} |a_n|^{1/n}$$

Let

$$\phi(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}$$

We know that radius of convergence of $\phi(z)$ and $f(z)$ would be the same. Let z be any arbitrary point inside circle of convergence. Then, we have $|z| < R$. So, there exists a positive number r such that $|z| < r < R$. Let's say $|z| = \alpha$, then it can be said that $\alpha < r$. We can choose a positive number $|h| = \beta$ such that

$$\alpha + \beta < r$$

Since, the power series $f(z)$ is convergent in the area $|z| < R$, therefore $a_n r^n$ is bounded for $0 < r < R$, so there exists a positive number M such that $|a_n r^n| < M$.

$$\begin{aligned} \text{Now, } \left| \frac{f(z+n) - f(z)}{n} - \phi(z) \right| &= \left| \sum_{n=0}^{\infty} a_n \left\{ \frac{(z+n)^n - z^n}{n} - nz^{n-1} \right\} \right| \\ &= \left| \sum_{n=0}^{\infty} \left[a_n \left\{ \frac{n(n-1)}{2} z^{n-2} h + \dots + h^{n-1} \right\} \right] \right| \\ &\leq \sum_{n=0}^{\infty} |a_n| \left\{ \frac{n(n-1)}{2} |\alpha|^{n-2} |\beta| + \dots + |\beta|^{n-1} \right\} \\ &\leq \sum_{n=0}^{\infty} \frac{M}{r^n} \left\{ \frac{n(n-1)}{2} \alpha^{n-2} \beta + \dots + \beta^{n-1} \right\} \\ &\leq \sum_{n=0}^{\infty} \frac{M}{r^n} \cdot \frac{1}{\beta} \{(\alpha + \beta)^n - \alpha^n - n\alpha^{n-1}\beta\} \\ &= \frac{M}{\beta} \sum_{n=0}^{\infty} \left\{ \left(\frac{\alpha + \beta}{r} \right)^n - \left(\frac{\alpha}{r} \right)^n - \frac{n\beta}{\alpha} \left(\frac{\alpha}{r} \right)^n \right\} \end{aligned} \quad \dots(i)$$

Now,

$$\begin{aligned} \sum_{n=0}^{\infty} \left(\frac{\alpha + \beta}{r} \right)^n &= 1 + \frac{\alpha + \beta}{r} + \left(\frac{\alpha + \beta}{r} \right)^2 + \dots \\ &= \left(1 - \frac{\alpha + \beta}{r} \right)^{-1} = \frac{r}{r - \alpha - \beta} \end{aligned}$$

and

$$\sum_{n=0}^{\infty} \left(\frac{\alpha}{r} \right)^n = 1 + \left(\frac{\alpha}{r} \right) + \left(\frac{\alpha}{r} \right)^2 + \dots = \left(1 - \frac{\alpha}{r} \right)^{-1} = \frac{r}{r - \alpha}$$

Again, putting $S = \sum_{n=0}^{\infty} n \left(\frac{\alpha}{r} \right)^n$, we have

$$S = \frac{\alpha}{r} + 2 \left(\frac{\alpha}{r} \right)^2 + 3 \left(\frac{\alpha}{r} \right)^3 + \dots \quad \dots(ii)$$

$$\Rightarrow S \left(\frac{\alpha}{r} \right) = \left(\frac{\alpha}{r} \right)^2 + 2 \left(\frac{\alpha}{r} \right)^3 + \dots \quad \dots(iii)$$

Subtracting (ii), (iii), we get

$$\begin{aligned} S\left(1-\frac{\alpha}{r}\right) &= \frac{\alpha}{r} + \left(\frac{\alpha}{r}\right)^2 + \left(\frac{\alpha}{r}\right)^3 + \dots \\ \Rightarrow \frac{S(r-\alpha)}{r} &= \frac{\alpha}{r} \left(1-\frac{\alpha}{r}\right)^{-1} = \frac{\alpha}{r-\alpha} \\ \Rightarrow S &= \frac{r\alpha}{(r-\alpha)^2} \end{aligned}$$

Substituting these values in (i), we get

$$\begin{aligned} \Rightarrow \left| \frac{f(z+n)-f(z)}{n} - \phi(z) \right| &\leq \frac{M}{\beta} \left[\frac{r}{r-\alpha-\beta} - \frac{r}{r-\alpha} - \frac{r\beta}{(r-\alpha)^2} \right] \\ &= \frac{M}{\beta} \left[\frac{r(r-\alpha)^2 - r(r-\alpha)(r-\alpha-\beta) - r\beta(r-\alpha-\beta)}{(r-\alpha)^2(r-\alpha-\beta)} \right] \\ &= \frac{M}{\beta} \cdot \frac{r\beta^2}{(r-\alpha-\beta)(r-\alpha)^2} \\ &= \frac{Mr\beta}{(r-\alpha-\beta)(r-\alpha)^2} \rightarrow 0 \text{ as } \beta \rightarrow 0 \end{aligned}$$

$$\therefore \lim_{n \rightarrow 0} \frac{f(z+n)-f(z)}{n} = \phi(z), \text{ which implies } f'(z) = \phi(z).$$

Thus, $f(z)$ is a single valued differentiable function. Consequently, it is analytic within the circle $|z| = R$. So, it is analytic inside its circle of convergence.

1.12 Let $f = u + iv$ be an analytic function on the unit disc $D = \{z \in \mathbb{C} : |z| < 1\}$. Show that $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 = \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2}$ at all points of D .

(2017 : 15 Marks)

Solution:

Every analytic function satisfies Cauchy-Riemann equations :

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \quad \dots(i)$$

Also, because u and v are the real and imaginary parts of an analytic function, so derivatives of u and v , of all orders, exist and are continuous functions of x and y .

So that we have

$$\frac{\partial^2 v}{\partial x \partial y} = \frac{\partial^2 v}{\partial y \partial x} \quad \dots(ii)$$

Differentiating (i)

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 v}{\partial x \partial y}; \frac{\partial^2 u}{\partial y^2} = -\frac{\partial^2 v}{\partial y \partial x}$$

Adding these two,

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad (\text{by virtue of (ii)})$$

$$\text{Similarly, } \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = 0$$

Hence, both u and v satisfy Laplace equation.

- 1.13 Prove that the function : $u(x, y) = (x - 1)^3 - 3xy^2 + 3y^2$ is harmonic and find its harmonic conjugate and corresponding analytic function $f(z)$ in terms of z .

(2018 : 10 Marks)

Solution:

Given :

$$u(x, y) = (x - 1)^3 - 3xy^2 + 3y^2$$

$$\frac{\partial y}{\partial x} = 3(x - 1)^2 - 3y^2, \quad \frac{\partial u}{\partial y} = -6xy + 6y$$

$$\frac{\partial^2 u}{\partial x^2} = 6(x - 1), \quad \frac{\partial^2 u}{\partial y^2} = -6x + 6$$

$$\Rightarrow \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 6(x - 1) - 6x + 6 = 0$$

So, $u(x, y)$ is harmonic.

Let $v(x, y)$ be its harmonic conjugate.

∴ By Cauchy-Euler equations,

$$u_x = v_y \text{ and } v_y = -v_x$$

$$\text{i.e.,} \quad 3(x - 1)^2 - 3y^2 = \frac{\partial v}{\partial y} \Rightarrow v = 3(x - 1)^2y - y^3 + f(x) \quad \dots(i)$$

$$\Rightarrow v = 3x^2y - 6xy + f(y) \quad \dots(ii)$$

From (i) and (ii), we get

So, harmonic conjugate,

By Milne-Thomson's method,

$$\begin{aligned} f(z) &= \int (u_x(z, 0) - u_y(z, 0)) dz \\ &= \int 3(z - 1)^2 dz \\ &= (z - 1)^3 + c \end{aligned}$$

(where c is a constant)

- 1.14 Suppose $f(z)$ is analytic function on a domain D in \mathbb{C} and satisfies the equation $\operatorname{Im} f(z) = (\operatorname{Re} f(z))^2$, $z \in D$. Show that $f(z)$ is constant in D .

(2019 : 10 Marks)

Solution:

Given $f(z)$ is an analytic function as domain D in \mathbb{C} .

Also

$$\operatorname{Im} f(z) = (\operatorname{Re} f(z))^2, z \in D \quad \dots(1)$$

If

$$f(z) = 4 + iv \quad \dots(1)$$

$$\operatorname{Im} f(z) = v \text{ and } \operatorname{Re} f(z) = 4 \quad \dots(2)$$

∴

$$\left. \begin{aligned} \frac{\partial y}{\partial x} &= \frac{\partial v}{\partial y} \\ \frac{\partial y}{\partial y} &= \frac{\partial v}{\partial x} \end{aligned} \right\} \quad \dots(3)$$

From (1) and (2),

$$v = u^2$$

∴

$$\left. \begin{aligned} \frac{\partial v}{\partial x} &= 24 \frac{\partial y}{\partial x} \\ \frac{\partial v}{\partial y} &= 24 \frac{\partial y}{\partial y} \end{aligned} \right\} \quad \dots(4)$$

From (3) and (4)

$$\frac{\partial v}{\partial x} = 0 \text{ and } \frac{\partial v}{\partial y} = 0$$

∴

$$\frac{\partial f}{\partial z} = \frac{\partial y}{\partial x} + \frac{\partial v}{\partial x} \Rightarrow \frac{\partial f}{\partial z} = 0$$

By integrating ⇒
Hence, the result.

$$f(z) = C(\text{Constant})$$

2. Taylor's Series

- 2.1 If the function $f(z)$ is analytic and one valued in $|z - a| < R$, prove that for $0 < r < R$, $f'(a) = \frac{1}{\pi r} \int_0^{2\pi} P(\theta) e^{-i\theta} d\theta$, where $P(\theta)$ is the real part of $f(a + re^{i\theta})$.

(2011 : 15 Marks)

Solution:

The function $f(z)$ is given to be analytic in $|z - a| < R$ and $r < R$, therefore $f(z)$ is also analytic inside the circle C defined by $|z - a| = r$.

$$\therefore f'(a) = \frac{1}{2\pi i} \int_C \frac{f(z)}{(z-a)^2} dz \quad \dots(i)$$

Expanding $f(z)$ in a Taylor's series about $z = a$, we have

$$f(z) = \sum_{n=0}^{\infty} a_n (z-a)^n$$

$$\Rightarrow f(z) = f(a + re^{i\theta}) = \sum_{n=0}^{\infty} a_n r^n e^{in\theta} \quad (\because z - a = re^{i\theta})$$

$$\Rightarrow \overline{f(z)} = \sum_{n=0}^{\infty} \bar{a}_n r^n e^{-in\theta}$$

$$\begin{aligned} \therefore \frac{1}{2\pi i} \int_C \frac{\overline{f(z)}}{(z-a)^2} dz &= \frac{1}{2\pi i} \int_0^{2\pi} \frac{\sum \bar{a}_n r^n e^{-in\theta}}{r^2 e^{i2\theta}} \cdot rie^{i\theta} d\theta \\ &= \frac{1}{2\pi} \sum \bar{a}_n r^{n-1} \int_0^{2\pi} e^{-i(n+1)\theta} d\theta = 0 \end{aligned} \quad \dots(ii)$$

From (i) and (ii), we have

$$\begin{aligned} f'(a) &= \frac{1}{2\pi i} \int_C \frac{f(z) + \overline{f(z)}}{(z-a)^2} dz \\ &= \frac{1}{2\pi i} \int_C \frac{2 \text{ real part of } f(z)}{(z-a)^2} dz \\ &= \frac{1}{\pi i} \int_0^{2\pi} \frac{\text{real part of } f(a + re^{i\theta}) ire^{i\theta}}{r^2 e^{i2\theta}} d\theta \quad [\because z = a + re^{i\theta}] \\ &= \frac{1}{\pi r} \int_0^{2\pi} P(\theta) e^{-i\theta} d\theta, \text{ where } P(\theta) \text{ is the real part of } f(a + re^{i\theta}). \end{aligned}$$

2.2 Describe the maximal ideals in the ring of Gaussian integers $\mathbb{Z}[i] = \{a + bi, b \in \mathbb{Z}\}$.

(2012 : 20 Marks)

Solution:

Let us define

$$\delta(a+ib) = |a+ib|^2 = a^2 + b^2 \text{ for all } a+ib \in \mathbb{Z}[i]$$

$$\delta(u) = \delta(a+ib) = a^2 + b^2 > 0 \text{ for all } u \neq 0 \text{ in } \mathbb{Z}[i]$$

Again for any $u, v \in \mathbb{Z}[i]$

$$\delta(uv) = |uv|^2 \geq |u|^2 = \delta(u)$$

Let $u, v \in \mathbb{Z}[i], v \neq 0$

Then,

$$u = a+ib, v = c+id, a, b, c, d \in \mathbb{Z} \text{ and } (c, d) \neq (0, 0)$$

∴

$$\begin{aligned} \frac{u}{v} &= \frac{a+ib}{c+id} = \frac{(a+ib)}{(c+id)} \times \frac{c+id}{c+id} \\ &= \frac{(a+ib)(c+id)}{c^2+d^2} = \alpha+i\beta \text{ (say)} \end{aligned}$$

where α and β are rational numbers. Then we can find integers m and n such that $|m - \alpha| \leq \frac{1}{2}$ and $|n - \beta| \leq \frac{1}{2}$.

∴

Now

$$\begin{aligned} u &= (\alpha+i\beta)v = (m+in)v + [(\alpha-m)+i(\beta-n)]v \\ [(\alpha-m)+i(\beta-n)]v &= [(\alpha+i\beta)-(m+in)]v \\ &= (\alpha+i\beta)v - (m+in)v \\ &= u - (m+in)v \in \mathbb{Z}[i] \end{aligned} \quad \dots(i)$$

as $\mathbb{Z}[i]$ is a ring and $u, v, m+in \in \mathbb{Z}[i]$

Let

Then

$$\begin{aligned} r &= [(\alpha-m)+i(\beta-n)]v \\ \delta(r) &= |r|^2 = [(\alpha-m)^2 + (\beta-n)^2] |v|^2 \\ &\leq \left(\frac{1}{4} + \frac{1}{4} \right) |v|^2 \end{aligned} \quad \dots(ii)$$

Taking $q = m+in$, we have,

$$u = vq + r \quad \text{(using (i) and (ii))}$$

where either $r = 0$ or $\delta(r) < \delta(v)$

Hence, $\mathbb{Z}[i]$ is a Euclidean domain.

As every Euclidean domain is a principal ideal domain (PID).

∴ $\mathbb{Z}[i]$ is PID.

But in a PID, every non-zero not unit element a is prime element iff $\langle a \rangle$ is a maximal ideal.

∴ Maximal ideals in $\mathbb{Z}[i]$ are the ideals generated by prime elements.

2.3 For a function $f: \mathbb{C} \rightarrow \mathbb{C}$ and $n \geq 1$, let f_n denote the n^{th} derivative of f and $f(0) = f$. Let f be an entire function such that for some $n \geq 1$, $f^n\left(\frac{1}{k}\right) = 0$ for all $K = 1, 2, 3, \dots$. Show that f is a polynomial.

(2017 : 15 Marks)

Solution:

Since $f(z)$ is entire, ∴ $f(z)$ is analytic. Hence, $f(z)$ can be expressed as Taylor's series around $z=0$ as

$$f(z) = \sum_{m=1}^{\infty} a_m \frac{1}{z^m} + \sum_{m=0}^{\infty} a_m z^m$$

$$f(z) = \sum_{m=1}^{\infty} (-1)^n (m)(m+1)\dots(m+(n-1))a_{-m} \frac{1}{z^{m+n}} + \sum_{m=n}^{\infty} (m)(m-1)\dots(m-(n-1))a_m z^{m-n}$$

Now,

$$f^n\left(\frac{1}{K}\right) = \sum_{m=1}^{\infty} (-1)^m (m)(m+1)\dots(m+(n-1))a_{-m} \cdot K^{m+n} + \sum_{m=n}^{\infty} m(m-1)\dots(m-n+1)a_m \left(\frac{1}{K}\right)^{m-n}$$

As $f^n\left(\frac{1}{K}\right) \rightarrow 0$ for all $K = 1, 2, 3, \dots$

Let us take $K \rightarrow \infty$

We get $a_{-n} \rightarrow 0$ and $a_n = 0$

Similarly, if we take $n+1$, $a_{n+1} = 0$

\therefore

$$a_n = a_{n+1} = a_{n+2} = \dots = 0$$

$$\therefore f(z) = a_0 + a_1 z + a_2 z^2 + \dots + a_{n-1} z^{n-1}$$

which is a polynomial function.

- 2.4 Show that an isolated singular point z_0 of a function $f(z)$ is a pole of order m if and only if $f(z)$ can be written in the form $f(z) = \frac{\phi(z)}{(z - z_0)^m}$ where $\phi(z)$ is analytic and non-zero at z_0 .

Moreover

$$\text{Res}_{z=z_0} f(z) = \frac{\phi^{(m-1)}(z_0)}{(m-1)!} \text{ if } m \geq 1$$

(2019 : 15 Marks)

Solution:

Since $f(z)$ has a pole of order m , then by definition, for $0 < |z - z_0| < R$.

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n + \frac{b_2}{z - z_0} + \frac{b_2}{(z - z_0)^2} + \frac{b_m}{(z - z_0)^m}, b_m \neq 0$$

$$\Rightarrow f(z) = \frac{1}{(z - z_0)^m} \left[\sum_{n=0}^{\infty} a_n (z - z_0)^{m+n} + b_1 (z - z_0)^{m-1} + b_2 (z - z_0)^{m-2} + b_2 (z - z_0)^{m-2} + b_m \right]$$

$$\Rightarrow f(z) = \frac{\phi(z)}{(z - z_0)^m}$$

Clearly, $\phi(z_0) = b_m \neq 0$ and is analytic at z_0 , as it has Taylor series expansion about z_0 .

Conversely, suppose $f(z)$ can be written in the form :

$$f(z) = \frac{\phi(z)}{(z - z_0)^m}, \text{ then}$$

$$\phi(z) = \phi(z_0) + \phi'(z_0)(z - z_0) + \frac{\phi''(z_0)}{2!}(z - z_0)^2 + \dots +$$

$$\frac{\phi^{(m-1)}(z_0)}{(m-1)!}(z - z_0)^{m-1} + \dots$$

Since,

$$\phi(z_0) \neq 0$$

$f(z)$ has a pole of order m .

With residue,

$$b_1 = \frac{\phi^{(m-1)}(z_0)}{(m-1)!}$$

In case of simple pole, i.e., $m=1$,

$\text{Res } z = z_0, f(z_0) = \phi(z_0)$. Hence, proved.

3. Singularity

- 3.1 Determine all entire functions $f(z)$ such that 0 is a removal singularity of $f\left(\frac{1}{z}\right)$.

(2017 : 10 Marks)

Solution:

For any complex valued function, $f(z)$

$$f(z) = \sum_{n=0}^{\infty} a_n z^n + \sum_{n=1}^{\infty} b_n z^{-n}$$

Since, $f(z)$ is entire, so all b_n are zero.

$$\begin{aligned} f(z) &= \sum_{n=0}^{\infty} a_n z^n \\ f\left(\frac{1}{z}\right) &= \sum_{n=0}^{\infty} a_n \left(\frac{1}{z}\right)^n \\ &= a_0 + \frac{a_1}{z} + \frac{a_2}{z^2} + \dots \end{aligned}$$

$f\left(\frac{1}{z}\right)$ is a removable singularity.

4. Laurant's Series

- 4.1 Find the Laurent series of the function $f(z) = \exp\left[\frac{\lambda}{2}\left(z - \frac{1}{z}\right)\right]$ as $\sum_{n=-\infty}^{\infty} c_n z^n$ for $0 < |z| < \infty$

where

$$c_n = \frac{1}{\pi} \int_0^\pi \cos(n\phi - \lambda \sin\phi) d\phi ; n = 0, \pm 1, \pm 2, \dots$$

with λ a given complex number and taking the unit circle c given by $z = e^{i\phi} (-\pi \leq \phi \leq \pi)$ as contour in this region.

(2010 : 15 Marks)

Solution:

O is the only point of singularity here. So, we can expand $\exp\left(\frac{\lambda}{2}\left(z - \frac{1}{z}\right)\right)$ as Laurent series valid for every value of $z \neq 0$, i.e., for $|z| > 0$.

Thus, we have

$$|z| \neq 0$$

$$\exp\left(\frac{\lambda}{2}\left(z - \frac{1}{z}\right)\right) = \sum_{n=-\infty}^{\infty} c_n z^n$$

when

$$c_n = \frac{1}{2\pi i} \int_c \frac{f(z)}{z^{n+1}} dz$$

where c is any circle with its centre at the origin. Taking c as the circle with radius unity, we have

$$c_n = \frac{1}{2\pi i} \int_c \frac{e^{\lambda/2(z-1)}}{z^{n+1}} dz$$

Putting $z = e^{i\phi}$, we obtain

$$c_n = \frac{1}{2\pi i} \int_0^{2\pi} \frac{e^{\lambda/2(e^{i\phi}-e^{-i\phi})}}{e^{i(n+1)\phi}} \cdot ie^{i\phi} d\phi$$

$$\Rightarrow c_n = \frac{1}{2\pi} \int_0^{2\pi} e^{\lambda \sin \phi} \cdot e^{-in\phi} d\phi$$

$$\Rightarrow c_n = \frac{1}{2\pi} \int_0^{2\pi} \cos(\lambda \sin \phi - n\phi) d\phi + \frac{i}{2\pi} \int_0^{2\pi} \sin(\lambda \sin \phi - n\phi) d\phi \quad \dots(1)$$

Taking $\phi = 2\pi - \theta$, we have

$$\begin{aligned} & \int_0^{2\pi} \sin(\lambda \sin \phi - n\phi) d\phi - \left(- \int_{2\pi}^0 \sin[-\lambda \sin \theta - 2n\pi + n\theta] d\theta \right) \\ &= \int_0^{2\pi} \sin(\lambda \sin \phi - n\phi) d\phi \end{aligned}$$

So that $\int_0^{2\pi} \sin(\lambda \sin \phi - n\phi) d\phi = 0 \quad \dots(2)$

$$\therefore c_n = \frac{1}{2\pi} \int_0^{2\pi} \cos(\lambda \sin \phi - n\phi) d\phi + 0 \quad (\text{Putting value from (2) in (1)})$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \cos(n\phi - \lambda \sin \phi) d\phi$$

$$= \frac{2}{2\pi} \int_0^\pi \cos(n\phi - \lambda \sin \phi) d\phi$$

$$\Rightarrow c_n = \frac{1}{\pi} \int_0^\pi \cos(n\phi - \lambda \sin \phi) d\phi \quad \dots(3)$$

So, given expression can be expressed as $\sum_{-\infty}^{\infty} c_n z^n$ where c_n is given by above equation (3).

4.2 Find the Laurent series for the function

$$f(z) = \frac{1}{1-z^2} \text{ with centre } z = 1.$$

(2011 : 15 Marks)

Solution:

We have

$$f(z) = \frac{1}{1-z^2} = \frac{1}{(1-z)(1+z)}$$

$$= \frac{1}{z} \left(\frac{1}{1-z} + \frac{1}{1+z} \right) \quad \dots(i)$$

First we will find Laurent expansion for $\phi(z) = \frac{1}{1+z}$ about $z = 1$.

We write,

$$\phi(z) = \sum_{n=0}^{\infty} a_n (z-1)^n \quad \dots(ii)$$

where

$$a_n = \frac{\phi^n(1)}{n!}$$

But

$$\begin{aligned} \phi^n(z) &= (-1)(-2) \dots (-n)(1+2)^{-(n+1)} \\ &= (-1)^n n! (1+z)^{-(n+1)} \end{aligned}$$

$$\therefore \frac{\phi^n(1)}{n!} = \frac{(-1)^n}{2^{n+1}}$$

$$\therefore a_n = \frac{\phi^n(1)}{n!} = \frac{(-1)^n}{2^{n+1}}$$

From (ii),

$$\begin{aligned} \phi(z) &= \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} (z-1)^n \\ &= \sum_{n=0}^{\infty} \frac{(1-z)^n}{2^{n+1}} \end{aligned}$$

\therefore From (i),

$$f(z) = \frac{1}{2} \left(\frac{1}{1-z} + \sum_{n=0}^{\infty} \frac{(1-z)^n}{2^{n+1}} \right)$$

This is the required expansion.

4.3 Expand the function $f(z) = \frac{1}{(z+1)(z+3)}$ in Laurent series valid for :

- (i) $1 < |z| < 3$; (ii) $|z| > 3$; (iii) $0 < |z+1| < 2$; (iv) $|z| < 1$

(2012 : 15 Marks)

Solution:

Given :

$$\begin{aligned} f(z) &= \frac{1}{(z+1)(z+3)} \\ &= \frac{1}{2(z+1)} - \frac{1}{2(z+3)} \quad (\text{using partial fractions}) \end{aligned}$$

(i)

\Rightarrow

$$\frac{1}{|z|} < 1 \text{ and } \frac{|z|}{3} < 1$$

\therefore

$$\begin{aligned} f(z) &= \frac{1}{2(z+1)} - \frac{1}{2(z+3)} \\ &= \frac{1}{2z \left(1 + \frac{1}{z} \right)} - \frac{1}{6 \left(1 + \frac{z}{3} \right)} \\ &= \frac{1}{2z} \left(1 + \frac{1}{z} \right)^{-1} - \frac{1}{6} \left(1 + \frac{z}{3} \right)^{-1} \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2z} \left(1 - \frac{1}{z} + \frac{1}{z^2} - \frac{1}{z^3} \dots \right) - \frac{1}{6} \left(1 - \frac{z}{3} + \frac{z^2}{9} - \frac{z^3}{27} \dots \right) \\
 &= \dots + \frac{1}{2z^3} - \frac{1}{2z^2} + \frac{1}{2z} + \frac{1}{6} - \frac{z}{18} + \frac{z^2}{54} - \frac{z^3}{162} \dots \\
 \text{(ii)} \quad |z| > 3
 \end{aligned}$$

$$\Rightarrow \frac{3}{|z|} < 1$$

$$\begin{aligned}
 \therefore f(z) &= \frac{1}{2z} \left(1 + \frac{1}{z} \right)^{-1} - \frac{1}{2z} \left(1 + \frac{3}{z} \right)^{-1} \\
 &= \frac{1}{2z} \left(1 - \frac{1}{z} + \frac{1}{z^2} - \frac{1}{z^3} + \dots \right) - \frac{1}{2z} \left(1 - \frac{3}{z} + \frac{9}{z^2} - \frac{27}{z^3} \dots \right) \\
 &= \frac{1}{z^2} - \frac{4}{z^3} + \frac{13}{z^4} - \frac{40}{z^5} + \dots
 \end{aligned}$$

$$\text{(iii)} \quad 0 < |z+1| < 2$$

$$\text{Put } z+1 = 4$$

Then

$$0 < |z+1| < 2 \Leftrightarrow 0 < |4| < 2$$

$$\begin{aligned}
 \therefore f(z) &= \frac{1}{(z+1)(z+3)} = \frac{1}{4(4+2)} \\
 &= \frac{1}{24} \left(1 + \frac{4}{2} \right)^{-1} \\
 &= \frac{1}{24} \left(1 - \frac{4}{2} + \frac{4^2}{4} - \frac{4^3}{8} \dots \right) \\
 &= \frac{1}{24} - \frac{1}{4} + \frac{4}{8} - \frac{4^2}{16} + \dots \\
 &= \frac{1}{24} - \frac{1}{4} + \frac{z+1}{8} - \frac{(z+1)^2}{16} + \dots
 \end{aligned}$$

$$\text{(iv)} \quad |z| < 1$$

Then

$$\begin{aligned}
 f(z) &= \frac{1}{2(z+1)} - \frac{1}{6 \left(1 + \frac{z}{3} \right)} \\
 &= \frac{1}{2} \left(1 - z + z^2 - z^3 \dots \right) - \frac{1}{6} \left(1 - \frac{z}{3} + \frac{z^2}{9} - \frac{z^3}{27} \dots \right) \\
 &= \frac{1}{3} - \frac{4}{9}z + \frac{13}{17}z^2 \dots
 \end{aligned}$$

4.4 Expand in Laurent series the function $f(z) = \frac{1}{z^2(z-1)}$ about $z=0$ and $z=1$.

(2014 : 10 Marks)

Solution:

Given that

$$f(z) = \frac{1}{z^2(z-1)}$$

About $z = 0$, the Laurent Series is given by

$$\begin{aligned}\frac{1}{z^2(z-1)} &= \frac{1}{-z^2(1-z)} = -\frac{1}{z^2}(1-z)^{-1} \\ &= -\frac{1}{z^2}(1+z+z^2+z+\dots) \\ &= -\left(\frac{1}{z^2} + \frac{1}{z} + 1+z+z^2+\dots\right)\end{aligned}$$

Let $z-1 = u \Rightarrow z = u+1$ and

$$\begin{aligned}\frac{1}{z^2(z-1)} &= \frac{1}{(u+1)^2 u} = \frac{1}{u}(1+u)^{-2} \\ &= \frac{1}{u} \left[1 - (2)(u) + \frac{(-2)(-3)}{2!}(u)^2 + \frac{(-2)(-3)(-4)}{3!}(u)^3 + \dots \right] \\ &= \frac{1}{u} [1 - 2u + 3u^2 - 4u^3 + \dots]\end{aligned}$$

- 4.5 Find all possible Taylor's and Laurent's series expansions of the function $f(z) = \frac{2z-3}{z^2-3z+2}$ about the point $z = 0$.

(2015 : 20 Marks)

Solution:

Given :

$$f(z) = \frac{2z-3}{(z^2-3z+2)} = \frac{2z-3}{(z-2)(z-1)}$$

Taylor Series :

$$\begin{aligned}f(z) &= \frac{1}{z-1} + \frac{1}{z-2} \\ &= \frac{1}{0-1} + (-z) \left(\frac{-1}{z-1} \right)_{z=0}^z + \frac{z^2}{2!} \times \frac{2}{(z-1)^3} \Big|_{z=0} + \frac{z^3}{3!} \times \frac{-6}{(z-1)^4} \Big|_{z=1} + \\ &\quad \dots + \frac{1}{0-2} + z \times \frac{-1}{(z-2)^2} \Big|_{z=0} + \frac{2z^2}{2!z} \times \frac{1}{(z-2)^3} \Big|_{z=0} \\ &\quad + \frac{z^3}{3!} \times \frac{-6}{(z-2)^4} \Big|_{z=0} \\ &= (-1 - z - z^2 - z^3 - \dots) + \left(\frac{-1}{2} - \frac{z}{4} - \frac{z^2}{8} - \frac{z^3}{16} - \dots \right)\end{aligned}$$

Laurent Series :

$$\begin{aligned}f(z) &= \frac{1}{z-1} + \frac{1}{z-2} \\ &= \frac{-1}{1-z} - \frac{1}{z-2} \\ &= -\left(1-z\right)^{-1} - \frac{1}{2} \left(1-\frac{2}{z}\right)^{-1}\end{aligned}$$

$$= -(1+z+z^2+z^3+\dots) - \frac{1}{2} \left(1 + \frac{z}{2} + \left(\frac{z}{2}\right)^2 + \left(\frac{z}{3}\right)^3 + \dots \right)$$

If $|z| < 1$ and $\left|\frac{z}{2}\right| < 1 \Rightarrow |z| < 2$

$$\therefore f(z) = -(1+z+z^2+\dots) - \frac{1}{2} \left(1 + \frac{z}{2} + \frac{z^2}{4} + \dots \right) \text{ around point } O.$$

- 4.6 Find the Laurent's series which represent function $\frac{1}{(1+z^2)(z+2)}$ which (i) $|z| < 1$ (ii) $1 < |z| < 2$
 (iii) $|z| > 2$

(2018 : 15 Marks)

Solution:

Let

$$f(z) = \frac{1}{(1+z^2)(z+2)} = \frac{1}{5} \left(\frac{1}{z+2} - \frac{z-2}{z^2+1} \right)$$

- (i) $|z| < 1$

$$f(z) = \frac{1}{5} \left[\frac{1}{2} \left(1 + \frac{z}{2} \right)^{-1} - (z-2)(1+z^2)^{-1} \right]$$

$$\text{as } \left| \frac{z}{2} \right| < 1 \Rightarrow |z| < 2 \text{ and } |z|^2 < 1 \Rightarrow |z| < 1$$

Both conditions are satisfied in this region.

So,

$$f(z) = \frac{1}{10} \left[1 - \frac{z}{2} + \frac{z^2}{2^2} - \dots \right] - \frac{z-2}{5} [1 - z^2 + z^4 - z^6 + \dots]$$

- (ii) $1 < |z| < 2$, we have

$$f(z) = \frac{1}{5} \left[\frac{1}{2} \left(1 + \frac{z}{2} \right)^{-1} - \frac{z-2}{z^2} \left(1 + \frac{1}{z^2} \right)^{-1} \right]$$

$$\text{Here, } \left| \frac{z}{2} \right| < 1 \Rightarrow |z| < 2 \text{ and } \left| \frac{1}{z^2} \right| < 1 \text{ or } |z| > 1$$

Both conditions are satisfied in this region.

∴

$$f(z) = \frac{1}{10} \left(1 - \frac{z}{2} + \frac{z^2}{2^2} + \frac{z^3}{2^3} + \dots \right) - \frac{z-2}{5z^2} \left(1 - \frac{1}{z^2} + \frac{1}{z^4} + \dots \right)$$

- (iii) $|z| > 2$

$$f(z) = \frac{1}{5} \left[\frac{1}{2} \left(1 + \frac{2}{z} \right)^{-1} - \frac{1}{z^2} (z-2) \left(1 + \frac{1}{z^2} \right)^{-1} \right]$$

$$\Rightarrow f(z) = \frac{1}{5z} \left(1 - \frac{2}{z} + \frac{2^2}{z^2} - \dots \right) - \frac{z-2}{5z^2} \left(1 - \frac{1}{z^2} + \frac{1}{z^4} - \dots \right) \quad (\text{for } |z| > 2)$$

- 4.7 Obtain the first three terms of the Laurent series expansion of the function $f(z) = \frac{1}{(e^z - 1)}$ about the point $z = 0$ valid in the region $0 < |z| < 2\pi$.

(2019 : 10 Marks)

$$\begin{aligned}
 &= \frac{1}{8}[2 + (1 + \cos 40) - 2\cos 20] \\
 &= \frac{1}{8}[\cos 40 - 2\cos 20 + 3] \\
 I &= \int_0^\pi \sin^4 \theta d\theta = \frac{1}{2} \int_0^{2\pi} \sin^4 \theta d\theta \\
 &= \frac{1}{2} \int_0^{2\pi} \frac{1}{8} [\cos 4\theta - 2\cos 2\theta + 3] d\theta \\
 &= \frac{1}{16} \int_0^{2\pi} [\cos 4\theta - 2\cos 2\theta + 3] d\theta \\
 &= \text{Real part of } \frac{1}{16} \int_0^{2\pi} (e^{i4\theta} - 2e^{i2\theta} + 3) d\theta
 \end{aligned}$$

Let $z = e^{i\theta}$,

$$\begin{aligned}
 dz &= ie^{i\theta} d\theta \Rightarrow d\theta = \frac{dz}{iz} \\
 &= \text{R.P. of } \frac{1}{16} \oint_C \frac{(z^4 - 2z^2 + 3) dz}{iz}
 \end{aligned}$$

where C is the unit circle.

$$\oint_C \frac{z^4 - 2z^2 + 3}{z} dz = 2\pi i \text{ Sum of Residue of } f(z)$$

$$\text{where } f(z) = \frac{z^4 - 2z^2 + 3}{z}$$

Now $f(z)$ has one pole at $z = 0$ inside unit circle.

$$\begin{aligned}
 \text{Residue at } z = 0 &= \lim_{z \rightarrow 0} zf(z) \\
 &= 3
 \end{aligned}$$

$$\oint f(z) = 6\pi i$$

$$\frac{1}{16i} \oint_C f(z) = \frac{3}{8}\pi$$

$$\therefore \int_0^\pi \sin^4 \theta d\theta = \text{R.P. } \frac{3}{8}\pi = \frac{3\pi}{8}$$

5.2 Evaluate the integral $\int_0^\pi \frac{d\theta}{\left(1 + \frac{1}{2}\cos \theta\right)^2}$ using residues.

(2014 : 20 Marks)

Solution:

Let

$$I = \int_0^\pi \frac{d\theta}{\left(1 + \frac{1}{2}\cos \theta\right)^2}$$

$$\begin{aligned}
 &= \frac{1}{2} \int_0^{2\pi} \frac{4d\theta}{(2+\cos\theta)^2} \\
 &= \int_0^{2\pi} \frac{2d\theta}{(2+\cos\theta)^2}
 \end{aligned}$$

Let the contour 'C' be the unit circle $|z| = 1$ with centre at the origin.

Let $z = e^{i\theta}$ then

$$\cos\theta = \frac{1}{2}\left(z + \frac{1}{z}\right)$$

\Rightarrow

$$dz = ie^{i\theta} d\theta$$

\Rightarrow

$$d\theta = \frac{dz}{iz}$$

\therefore

$$I = \int_0^{2\pi} \frac{2d\theta}{(2+\cos\theta)^2} = \int_0^{2\pi} \frac{2dz}{iz \left(2 + \frac{z^2+1}{2z}\right)^2}$$

$$= \frac{1}{i} \int_0^{2\pi} \frac{8z}{(z^2+4z+1)^2} dz$$

$$= \frac{8}{i} \int_C \frac{z}{(z^2+4z+1)^2} dz$$

$$= \frac{8}{i} \int_C f(z) dz$$

where

$$f(z) = \frac{z}{(z^2+4z+1)^2} \quad \dots(i)$$

Now the poles of $f(z)$ are given by

$$(z^2+4z+1)^2 = 0 \Rightarrow z = \frac{-4 \pm \sqrt{16-4}}{2} = \frac{-4 \pm \sqrt{12}}{2} = -2 \pm \sqrt{3} \text{ (twice)}$$

$\therefore f(z)$ has poles of order at $z = -2 \pm \sqrt{3}$ (twice)

Let $\alpha = -2 + \sqrt{3}$, $\beta = -2 - \sqrt{3}$

Clearly $|\beta| > 1$

Since $|\alpha\beta| = 1 \Rightarrow |\alpha| < 1$

Hence, the θdy pole inside C is $z = \alpha$ of order 2.

($\because |\beta| > 1$)

$$\begin{aligned}
 \therefore \int_C f(z) dz &= \int \frac{z dz}{(z^2+4z+1)^2} \\
 &= 2\pi i \text{ (residue at } z = \alpha)
 \end{aligned}$$

Now the residue at $z = \alpha$ is

$$\begin{aligned}
 \lim_{z \rightarrow \alpha} \frac{d}{dz} (z - \alpha)^2 \frac{z}{(z^2+4z+1)^2} &= \lim_{z \rightarrow \alpha} \frac{d}{dz} \frac{z}{(z - \beta)^2} \\
 &= \lim_{z \rightarrow \alpha} \frac{-(\beta + z)}{(z - \beta)^3} \\
 &= \frac{-(\alpha + \beta)}{(\alpha - \beta)^3}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{-(-4)}{(2\sqrt{3})^3} \\
 &= \frac{(4)}{8(3\sqrt{3})} = \frac{1}{6\sqrt{3}} \\
 \therefore \int_C f(z) dz &= 2\pi i \left(\frac{1}{6\sqrt{3}} \right) = \frac{2\pi i}{6\sqrt{3}} = \frac{\sqrt{3}i}{3\sqrt{3}} \\
 \therefore \text{from (i)} \quad \int_0^{2\pi} \frac{2d\theta}{(2+\cos\theta)^2} &= \frac{8}{i} \left(\frac{\pi i}{3\sqrt{3}} \right) = \frac{8\pi}{3\sqrt{3}} \\
 \therefore I &= \int_0^\pi \frac{d\theta}{\left(1 + \frac{\cos\theta}{2}\right)^2} = \frac{8\pi}{3\sqrt{3}}
 \end{aligned}$$


5.3 State Cauchy's Residue theorem. Using it, evaluate the integral $\int_C \frac{e^z + 1}{z(z+1)(z-i)^2} dz$. {C : |z| = 2} (2015 : 15 Marks)

Solution:

Cauchy's Residue theorem states that in a given region, integral of a function along the closed curve is equal to $2\pi i$ times sum of its residues.

Given,

$$\begin{aligned}
 I &= \int_C \frac{e^z + 1}{z(z+1)(z-i)^2} dz, C: |z| = 2 \\
 &= \int_C f(z) dz
 \end{aligned}$$



In the given region, $|z| = 2$, $f(z)$ is analytic everywhere except at $z = 0, -1, i$.

$z = 0$ is a pole of order 1.

$z = -1$ is a pole of order 1.

$z = i$ is a pole of order 2.

Now, Residue at $z = 0$:

$$\begin{aligned}
 \lim_{z \rightarrow 0} \frac{z(e^z + 1)}{z(z+1)(z-i)^2} &= \frac{1+1}{1(-i)^2} = \frac{2}{(-1)} \\
 &= -2
 \end{aligned}$$

Residue at $z = -1$:

$$\begin{aligned}
 \lim_{z \rightarrow -1} \frac{(z+1)(e^z + 1)}{z(z+1)(z-i)^2} &= \frac{1+e^{-1}}{(-1)(-1-i)^2} = \frac{-(1+e^{-1})}{1-1+2i} \\
 &= \frac{(1+e^{-1})}{2}i
 \end{aligned}$$

Residue at $z = i$:

$$\begin{aligned}
 \lim_{z \rightarrow i} \frac{d}{dz} (z-i)^2 \frac{e^z + 1}{z(z+1)(z-i)^2} &= \lim_{z \rightarrow i} \frac{d}{dz} \cdot \frac{e^z + 1}{z(z+1)} \\
 &= \lim_{z \rightarrow i} \left\{ \frac{-(e^z + 1)}{z^2(z+1)} - \frac{(e^z + 1)}{z(z+1)^2} + \frac{e^z}{z(z+1)} \right\}
 \end{aligned}$$

$$= \frac{+(1+e^i)}{(1+i)(1+i)} - \frac{(1+e^i)}{i(1+i)^2} + \frac{e^i}{i(i+1)}$$

$$= \frac{1+e^i}{1+i} - \frac{1+e^i}{2} + \frac{e^i}{i(i+1)}$$

$$= \frac{1-i}{2} + \frac{1+e^i}{2} = \frac{2-i+e^i}{2}$$

$$\int_C f(z) dz = 2\pi i (\text{sum of residues})$$

$$= 2\pi i \left(-2 + \frac{(1+e^{-1})i}{2} + \frac{2-i+e^i}{2} \right)$$

$$= 2\pi i \left(-2 + \frac{i+ie^{-1}+2-i+e^i}{2} \right)$$

∴

$$\int_C f(z) dz = \pi i (e^i + ie^{-1} + 2 - 4)$$

$$= \pi i \left(e^i + \frac{i}{e} - 2 \right)$$

∴

$$\int_C f(z) dz = \pi i \left(e^i + \frac{i}{e} - 2 \right)$$

5.4 Show by applying residue theorem that

$$\int_0^\infty \frac{dx}{(x^2 + a^2)^2} = \frac{\pi}{4a^3}, \quad a > 0$$

(2018 : 15 Marks)

Solution:

$$\text{Consider } \int_{-R}^R \frac{dz}{(z^2 + a^2)^2} + \int_z \frac{dz}{(z^2 + a^2)^2}$$

Now,

$$\int_{-c}^a \frac{dz}{(z^2 + a^2)^2} = \int_c^a \frac{dz}{(z+ai)^2(z-ai)^2}$$

In the above region, $z = ai$ is a pole of order 2.

$$\therefore \text{Residue at } z = ai : \lim_{z \rightarrow ai} \frac{d}{dz} \frac{(z-ai)^2}{(z+ai)^2(z-ai)^2}$$

$$= \lim_{z \rightarrow ai} \frac{-2}{(z+ai)^3} = \frac{2}{8 \times a^3 \times i} = \frac{-i}{4a^3}$$


By Cauchy's residue theorem,

$$\int_C \frac{dz}{(z^2 + a^2)^2} = 2\pi i (\text{sum of residues})$$

$$= 2\pi i \times \frac{-i}{4a^3} = \frac{\pi}{2a^3} \quad \dots (i)$$

Now,

$$\int_L \frac{dz}{(z^2 + a^2)^2} = \int_{-R}^R \frac{dz}{(z^2 + a^2)^2} + \int_{\tau} \frac{dz}{(z^2 + a^2)^2}$$



as $R \rightarrow \infty$,

$$\int_{\gamma} \frac{dz}{(z^2 + a^2)^2} = 0$$

$$\therefore \int_C \frac{dz}{(z^2 + a^2)^2} = \int_{-\infty}^{\infty} \frac{dz}{(z^2 + a^2)^2} = 2 \int_0^{\infty} \frac{dz}{(z^2 + a^2)^2}$$

∴ From (i)

$$2 \int_0^{\infty} \frac{dz}{(z^2 + a^2)^2} = \frac{\pi}{2a^3}$$

$$\Rightarrow \int_0^{\infty} \frac{dz}{(z^2 + a^2)^2} = \frac{\pi}{4a^3}$$

$$\text{or } \int_0^{\infty} \frac{dx}{(x^2 + a^2)^2} = \frac{\pi}{4a^3}, a > 0$$

6. Contour Integration

6.1 If α, β, γ are real numbers such that $\alpha^2 > \beta^2 + \gamma^2$, show that :

$$\int_0^{2\pi} \frac{d\theta}{\alpha + \beta \cos \theta + \gamma \sin \theta} = \frac{2\pi}{\sqrt{\alpha^2 - \beta^2 - \gamma^2}}$$

(2009 : 30 Marks)

Solution:

Proof : Let

then,


$$z = e^{i\theta}$$

$$dz = ie^{i\theta} d\theta = iz d\theta$$

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i} = \frac{z - z^{-1}}{2i}$$

$$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2} = \frac{z + z^{-1}}{2}$$

Put values of $d\theta$, $\cos \theta$ and $\sin \theta$ in the L.H.S. and the curve becomes a circle : $|z| = 1$



i.e.,

$$\begin{aligned} \int_0^{2\pi} \frac{d\theta}{\alpha + \beta \cos \theta + \gamma \sin \theta} &= \oint_C \frac{dz}{iz \left\{ \alpha + \beta \left(\frac{z+z^{-1}}{2} \right) + \gamma \left(\frac{z-z^{-1}}{2i} \right) \right\}} \\ &= \oint_C \frac{dz}{iz [2i\alpha + \beta i(z+z^{-1}) + \gamma(z-z^{-1})]} \\ &= \oint_C \frac{2z dz}{z[2i\alpha z + \beta i(z^2+1) + 8(z^2-1)]} \\ &= \oint_C \frac{2dz}{z^2(\gamma + \beta i) + 2i\alpha z + \beta i - \gamma} \\ &= \oint_C \frac{2dz}{(\gamma + \beta i) \left\{ z^2 + \frac{2i\alpha z}{\gamma + \beta i} + \frac{\beta i - \gamma}{\gamma + \beta i} \right\}} = \oint_C f(z) dz \end{aligned}$$

where C is the circle of unit radius with centre at the origin.

Now, poles of $f(z)$ are given by

$$z^2 + \frac{2i\alpha z}{\gamma + \beta_i} + \frac{\beta_i - \gamma}{\gamma + \beta_i} = 0$$

$$z = \frac{\frac{-2i\alpha}{\gamma + \beta_i} \pm \sqrt{\left(\frac{2i\alpha}{\gamma + \beta_i}\right)^2 - \frac{4(\beta_i - \gamma)}{(\gamma + \beta_i)}}}{2}$$

$$z = \frac{\frac{-2i\alpha}{2(\gamma + \beta_i)} \pm \frac{2}{2} \sqrt{\frac{i^2\alpha^2 - (\beta_i - \gamma)(\gamma + \beta_i)}{(\gamma + \beta_i)^2}}}{2}$$

$$z = \frac{\frac{-i\alpha \pm \sqrt{-\alpha^2 - \{(\beta_i)^2 - \gamma^2\}}}{(\gamma + \beta_i)}}{2}$$

$$z = \frac{\frac{-i\alpha \pm \sqrt{-\alpha^2 + \gamma^2 + \beta^2}}{\gamma + \beta_i}}{2}$$

$$z = \frac{\frac{-i\alpha \pm \sqrt{-1}\sqrt{\alpha^2 - \beta^2 - \gamma^2}}{\gamma + \beta_i}}{2}$$

$$z = \frac{\frac{-i\alpha \pm i\sqrt{\alpha^2 - \beta^2 - \gamma^2}}{\gamma + \beta_i}}{2} \quad (\text{as given that } \alpha^2 > \beta^2 + \gamma^2)$$

Now only

$$z = \frac{\frac{-i\alpha + i\sqrt{\alpha^2 - \beta^2 - \gamma^2}}{\gamma + \beta_i}}{2} \text{ lies inside the circle : } |z| = 1$$

$\Rightarrow z_0$ (say)

Now, residue at $z = z_0$

$$= \lim_{z \rightarrow z_0} \frac{(z - z_0)2}{(\gamma + \beta_i) \left(z^2 + \frac{2i\alpha z}{\gamma + \beta_i} + \frac{\beta_i - \gamma}{\gamma + \beta_i} \right)}$$

$$= \lim_{z \rightarrow z_0} \frac{2(z - z_0)}{(\gamma + \beta_i)(z - z_0)(z - z_1)} \left(\text{where } z_1 = \frac{-i\alpha - i\sqrt{\alpha^2 - \beta^2 - \gamma^2}}{\gamma + \beta_i} \right)$$

$$= \frac{2}{(\gamma + \beta_i)(z_0 - z_1)}$$

where,

$$z_0 - z_1 = \left(\frac{\frac{-i\alpha + i\sqrt{\alpha^2 - \beta^2 - \gamma^2}}{\gamma + \beta_i}}{2} \right) - \left(\frac{\frac{-i\alpha - i\sqrt{\alpha^2 - \beta^2 - \gamma^2}}{\gamma + \beta_i}}{2} \right)$$

$$= \frac{\frac{2i\sqrt{\alpha^2 - \beta^2 - \gamma^2}}{\gamma + \beta_i}}{2}$$

$$= \frac{1}{i\sqrt{\alpha^2 - \beta^2 - \gamma^2}}$$

Now,

$$\int_0^{2\pi} \frac{d\theta}{\alpha + \beta \cos \theta + \gamma \sin \theta} = \oint_C f(z) dz$$


$$\begin{aligned}
 &= 2\pi i(\text{Residue at } z = z_0) \\
 &= 2\pi i \times \frac{1}{i\sqrt{\alpha^2 - \beta^2 - \gamma^2}} \quad (\text{By Cauchy's Residue Theorem}) \\
 &= \frac{2\pi}{\sqrt{\alpha^2 - \beta^2 - \gamma^2}}. \text{ Proved.}
 \end{aligned}$$

6.2 Evaluate by Contour integration $\int_0^1 \frac{dx}{(x^2 - x^3)^{1/3}}$.

(2011 : 15 Marks)

Solution:

$$\begin{aligned}
 \int_0^1 \frac{dx}{(x^2 - x^3)^{1/3}} &= \int_{\infty}^1 \frac{-\frac{dx}{x^2}}{\left(\frac{1}{x^2} - \frac{1}{x^3}\right)^{1/3}} \\
 &= \int_1^{\infty} \frac{dx}{x(x-1)^{1/3}} = \int_0^{\infty} \frac{x^{-1/3} dx}{x+1} = \int_0^{\infty} \frac{x^{2/3-1} dx}{x+1}
 \end{aligned}$$



Let

$$\int_C f(z) dz = \int_C \frac{z^{a-1}}{1-z} dz$$

where C is the contour consisting of a large semi-circle Γ , $|z| = R$ in the upper half plane indented at $z = 0$, $z = 1$.

γ_1 and γ_2 are the semi-circles in the upper half plane with radii P_1 and P_2 and centres $z = 0$, $z = 1$ respectively. By Cauchy's residue theorem, we have

$$\int_C f(z) dz = \int_{\Gamma} f(z) dz + \int_{-R}^{-P_1} f(x) dx + \int_{\gamma_1} f(z) dz + \int_{P_1}^{1-P_2} f(x) dx + \int_{\gamma_2} f(z) dz + \int_{1+P_2}^R f(x) dx = 0 \quad \dots(i)$$

(∴ $f(z)$ has no pole inside C)

Now,

$$\begin{aligned}
 \left| \int_{\Gamma} f(z) dz \right| &\leq \int_0^{\pi} \left| \frac{R^{a-1} e^{i(a-1)\theta}}{1-R e^{i\theta}} \cdot i R e^{i\theta} \right| d\theta \leq \int_0^{\pi} \frac{R^a}{R-1} d\theta \\
 &= \frac{R^a \pi}{R-1} \rightarrow 0 \text{ as } R \rightarrow \infty \text{ as } a < 1.
 \end{aligned}$$

Since,

$$\lim_{z \rightarrow 0} z^a f(z) = \lim_{z \rightarrow 0} \frac{z^a}{1-z} = 0, a > 0$$

$$\lim_{P_1 \rightarrow 0} \int_{\gamma_1} f(z) dz = -i(\pi - 0)0 = 0$$

$$\lim_{z \rightarrow 1} (z-1)f(z) = \lim_{z \rightarrow 1} (z-1) \frac{z^{a-1}}{1-z} = -1$$

$$\therefore \lim_{\rho_2 \rightarrow 0} \int_{V_2} f(z) dz = i\pi$$

Hence, as $\rho_1 \rightarrow 0$, $\rho_2 \rightarrow 0$, $R \rightarrow \infty$, we have from (i)

$$\int_{-\infty}^0 f(x) dx + \int_0^1 f(x) dx + i\pi + \int_1^\infty f(x) dx = 0$$

$$\Rightarrow \int_{-\infty}^0 f(x) dx + \int_0^\infty f(x) dx = -i\pi$$

$$\Rightarrow \int_0^\infty f(-x) dx + \int_0^\infty f(x) dx = -i\pi \quad (\text{Putting } -x \text{ for } x \text{ in the first integral})$$

$$\Rightarrow \int_0^\infty \frac{(-x)^{a-1}}{1+x} dx + \int_0^\infty \frac{x^{a-1}}{1-x} dx = -i\pi$$

$$\Rightarrow \int_0^\infty \frac{(-1)^{a-1} x^{a-1}}{1+x} dx + \int_0^\infty \frac{x^{a-1}}{1-x} dx = -i\pi$$

$$\Rightarrow \int_0^\infty \frac{(e^{-i\pi})^{a-1} x^{a-1}}{1+x} dx + \int_0^\infty \frac{x^{a-1}}{1-x} dx = -i\pi$$

$$\Rightarrow \int_0^\infty \frac{e^{-i\pi} e^{i\alpha\pi} x^{a-1}}{1+x} dx + \int_0^\infty \frac{x^{a-1}}{1-x} dx = -i\pi$$

$$\Rightarrow - \int_0^\infty \frac{e^{i\alpha\pi} x^{a-1}}{1+x} dx + \int_0^\infty \frac{x^{a-1}}{1-x} dx = -i\pi$$

Equating imaginary parts, we have

$$- \int \frac{\sin a\pi x^{a-1}}{1+x} dx = -\pi$$

$$\Rightarrow \int \frac{x^{a-1}}{1+x} dx = \frac{\pi}{\sin a\pi}$$

Putting $a = \frac{2}{3}$, we have

$$\int \frac{x^{-1/3}}{1+x} dx = \frac{\pi}{\sin \frac{2\pi}{3}} = \frac{\pi}{\sin \frac{2\pi}{3}} = \frac{\pi}{\sqrt{3}/2} = \frac{2\pi}{\sqrt{3}}$$

6.3 Evaluate by contour Integration :

$$I = \int_0^{2\pi} \frac{d\theta}{1-2a\cos\theta+a^2}, a^2 < 1$$

(2012 : 15 Marks)

Solution:

Given :

$$I = \int_0^{2\pi} \frac{d\theta}{1-2a\cos\theta+a^2}, a^2 < 1$$

Put $z = e^{i\theta} \Rightarrow$

$$dz = ie^{i\theta} d\theta = iz d\theta$$

Then

$$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2} = \frac{1}{2} \left(z + \frac{1}{z} \right)$$

Let C be a unit circle.

Then,

$$\begin{aligned} I &= \int_C \frac{1}{1+a^2 - \left(z + \frac{1}{z} \right) a} \cdot \frac{dz}{iz} \\ &= -\frac{1}{ia} \int_C \frac{dz}{c(z-a)\left(z-\frac{1}{a} \right)} = \int_C f(z) dz \end{aligned}$$

Now, $z = a$ and $z = \frac{1}{a}$ are simple poles of $f(z)$.

Since, $0 < a < 1$, therefore the simple pole $z = a$ lies inside C .

Residue at $z = a$ is

$$\lim_{z \rightarrow a} (z-a)f(z) = \lim_{z \rightarrow a} \frac{-(z-a)}{ia(z-a)\left(z-\frac{1}{a} \right)} = \frac{1}{i(1-a^2)}$$

Hence by Cauchy's residue theorem, we have

$$\begin{aligned} \int_0^{2\pi} \frac{d\theta}{1+a^2-2a\cos\theta} &= 2\pi i (\text{sum of the residues inside } C) \\ &= 2\pi i \cdot \frac{1}{i(1-a^2)} = \frac{2\pi}{1-a^2} \end{aligned}$$

6.4 Let $r : [0, 1] \rightarrow C$ be the curve

$$r(t) = e^{2\pi it}, 0 \leq t \leq 1$$

Find giving justifications, the value of the contour integral $\int_r \frac{dz}{4z^2-1}$.

(2016 : 15 Marks)

Solution:

Given : $r : [0, 1] \rightarrow C$

and

$$r(t) = e^{2\pi it} (0 \leq t \leq 1)$$

r represents the curve as shown in figure on complex plane.

Now, let

$$I = \int_r \frac{dz}{4z^2-1}$$


\Rightarrow

$$I = \int_r \frac{dz}{(2z+1)(2z-1)} = \int_r f(x) dx = \int_r \frac{dz}{4\left(z+\frac{1}{2} \right)\left(z-\frac{1}{2} \right)}$$

$$f(z) = \frac{1}{4x^2-1}$$

Now, in the region bounded by r , $z = -\frac{1}{2}$, $z = \frac{1}{2}$ are poles of order 1.

\therefore Residue at $z = \frac{1}{2}$, R :



$$R_1 = \lim_{z \rightarrow \frac{1}{2}} \frac{\left(z - \frac{1}{2}\right) \times 1}{4 \left(z + \frac{1}{2}\right) \left(z - \frac{1}{2}\right)} = \frac{1}{4 \left(\frac{1}{2} + \frac{1}{2}\right)} = \frac{1}{4}$$

Residue at $z = -\frac{1}{2}$: R_2 :

$$R_2 = \lim_{z \rightarrow -\frac{1}{2}} \frac{\left(z + \frac{1}{2}\right) \times 1}{4 \left(z - \frac{1}{2}\right) \left(z + \frac{1}{2}\right)} = -\frac{1}{4}$$

By Cauchy's residue theorem,

$$\begin{aligned} \int_C \frac{dz}{4z^2 - 1} &= 2\pi i (\text{sum of residues}) \\ &= 2\pi i (R_1 + R_2) = 2\pi i \left(\frac{1}{4} - \frac{1}{4}\right) \end{aligned}$$

$$\Rightarrow \int_C \frac{dz}{4z^2 - 1} = 0$$

6.5 Using contour integral method, prove that $\int_0^\infty \frac{x \sin mx}{a^2 + x^2} dx = \frac{\pi}{2} e^{-ma}$.

(2017 : 15 Marks)


Solution:

Let

$$I = \int_C \frac{e^{imz}}{a^2 + z^2} dz = \int_C f(z) dz$$

where C is the contour consisting of a large semi-circle, T of radius R containing all the poles of the integrand in the upper half plane and the part of real axis from $-R$ to R .

By Cauchy-Residue Theorem,



$$\begin{aligned} \int_C f(z) dz &= \int_{-R}^R \frac{e^{imx}}{a^2 + x^2} dx + \int_T \frac{e^{imz}}{a^2 + z^2} dz \\ &= 2\pi i (\text{sum of residues}) \end{aligned}$$

Since,

$$\lim_{z \rightarrow \infty} \frac{1}{(z^2 + a^2)} = 0$$

Therefore,

$$\lim_{z \rightarrow \infty} \int_C f(z) dz = 0, \text{ by Jordan Lemma}$$

$$\therefore \lim_{R \rightarrow \infty} \int_{-R}^R \frac{e^{-imx}}{a^2 + x^2} dx = 2\pi i (\text{sum of residues})$$

or

$$\int_{-\infty}^{\infty} \frac{e^{imx}}{a^2 + x^2} dx = 2\pi i (\text{sum of residues})$$

$z = \pm ai$ are simple poles of $f(z)$.

The pole, $z = ai$ lies inside, C .

Residue at $z = ai$ is

$$\begin{aligned}\lim_{z \rightarrow ai} (z - ai)f(z) &= \lim_{z \rightarrow ai} \frac{(z - ai)e^{imz}}{a^2 + z^2} \\ &= \lim_{z \rightarrow ai} \frac{e^{imz}}{z + ai} = \frac{e^{-ma}}{2ia}\end{aligned}$$

From (i)

$$\int_{-\infty}^{\infty} \frac{e^{imx}}{x^2 + a^2} dx = 2\pi i \left(\frac{e^{-ma}}{2ia} \right) = \frac{\pi}{a} e^{-ma}$$

Equating real parts on both sides,

$$\int_{-\infty}^{\infty} \frac{\cos mx}{x^2 + a^2} dx = \frac{\pi}{a} e^{-ma}$$

or

$$\int_0^{\infty} \frac{\cos mx}{x^2 + a^2} dx = \frac{\pi}{2a} e^{-ma}$$

Differentiating both sides w.r.t. m

$$\int_0^{\infty} \frac{x \sin mx}{x^2 + a^2} dx = \frac{\pi}{2} e^{-ma}$$

■ ■ ■