

MATHEMATICS

Paper—I

Time Allowed : Three Hours

Maximum Marks : 200

QUESTION PAPER SPECIFIC INSTRUCTIONS

Please read each of the following instructions carefully before attempting questions.

There are **EIGHT** questions in all, out of which **FIVE** are to be attempted.

Question no. 1 and 5 are compulsory. Out of the remaining **SIX** questions, **THREE** are to be attempted selecting at least **ONE** question from each of the two Sections A and B.

Attempts of questions shall be counted in chronological order. Unless struck off, attempt of a question shall be counted even if attempted partly. Any page or portion of the page left blank in the answer book must be clearly struck off.

All questions carry equal marks. The number of marks carried by a question / part is indicated against it.

Answers must be written in **ENGLISH** only.

Unless other-wise mentioned, symbols and notations have their usual standard meanings.

Assume suitable data, if necessary and indicate the same clearly.

SECTION—A

Q. 1(a) Find the dimension and a basis of the solution space W of the system

$$x + 2y + 2z - s + 3t = 0, \quad x + 2y + 3z + s + t = 0, \quad 3x + 6y + 8z + s + 5t = 0$$

8

Q. 1(b) Find the characteristic equation of the matrix $A = \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix}$ and hence find the matrix

$$\text{represented by } A^8 - 5A^7 + 7A^6 - 3A^5 + A^4 - 5A^3 + 8A^2 - 2A + I.$$

8

Q. 1(c) Evaluate the integral $\int_0^\infty \int_0^x x e^{-x^2/y} dy dx$ by changing the order of integration.

8

Q. 1(d) Find the surface generated by the straight line which intersects the lines $y = z = a$ and $x + 3z = a = y + z$ and is parallel to the plane $x + y = 0$.

8

Q. 1(e) Find C of the Mean value theorem, if $f(x) = x(x - 1)(x - 2)$, $a = 0$, $b = \frac{1}{2}$ and C has usual meaning.

8

Q. 2(a) Let V be the vector space of 2×2 matrices over \mathbb{R} and let $M = \begin{bmatrix} 1 & -1 \\ -2 & 2 \end{bmatrix}$.

Let $F : V \rightarrow V$ be the linear map defined by $F(A) = MA$. Find a basis and the dimension of

(i) the kernel of F

(ii) the image U of F . 10

Q. 2(b) Locate the stationary points of the function $x^4 + y^4 - 2x^2 + 4xy - 2y^2$ and determine their nature. 10

Q. 2(c) Find an orthogonal transformation of co-ordinates which diagonalizes the quadratic form

$$q(x, y) = 2x^2 - 4xy + 5y^2. \quad 10$$

Q. 2(d) Discuss the consistency and the solutions of the equations

$$x + ay + az = 1, \quad ax + y + 2az = -4, \quad ax - ay + 4z = 2$$

for different values of a . 10

Q. 3(a) Prove that if $a_0, a_1, a_2, \dots, a_n$ are the real numbers such that

$$\frac{a_0}{n+1} + \frac{a_1}{n} + \frac{a_2}{n-1} + \dots + \frac{a_{n-1}}{2} + a_n = 0$$

then there exists at least one real number x between 0 and 1 such that

$$a_0 x^n + a_1 x^{n-1} + a_2 x^{n-2} + \dots + a_{n-1} x + a_n = 0. \quad 10$$

Q. 3(b) Reduce the following equation to its canonical form and determine the nature of the conic

$$4x^2 + 4xy + y^2 - 12x - 6y + 5 = 0. \quad 10$$

Q. 3(c) Let F be a subfield of complex numbers and T a function from $F^3 \rightarrow F^3$ defined by

$$T(x_1, x_2, x_3) = (x_1 + x_2 + 3x_3, 2x_1 - x_2, -3x_1 + x_2 - x_3).$$

What are the conditions on (a, b, c) such that (a, b, c) be in the null space of T ? Find the nullity of T . 10

Q. 3(d) Find the equations to the tangent planes to the surface

$$7x^2 - 3y^2 - z^2 + 21 = 0, \text{ which pass through the line } 7x - 6y + 9 = 0, z = 3. \quad 10$$

Q. 4(a) Evaluate

$$\int_0^{\pi/2} \frac{x \sin x \cos x \, dx}{\sin^4 x + \cos^4 x}. \quad 10$$

Q. 4(b) Let $H = \begin{bmatrix} 1 & i & 2+i \\ -i & 2 & 1-i \\ 2-i & 1+i & 2 \end{bmatrix}$ be a Hermitian matrix. Find a non-singular matrix P such

that $P^t H \bar{P}$ is diagonal and also find its signature. 10

Q. 4(c) Find the magnitude and the equations of the line of shortest distance between the lines

$$\frac{x-8}{3} = \frac{y+9}{-16} = \frac{z-10}{7}$$

$$\frac{x-15}{3} = \frac{y-29}{8} = \frac{z-5}{-5}$$

10

Q. 4(d) Find all the asymptotes of the curve

$$x^4 - y^4 + 3x^2y + 3xy^2 + xy = 0.$$

10

SECTION—B

Q. 5(a) Solve :

$$\frac{dy}{dx} + x \sin 2y = x^3 \cos^2 y.$$

8

Q. 5(b) A particle is performing a simple harmonic motion of period T about centre O and it passes through a point P, where $OP = b$ with velocity v in the direction of OP. Find the time which elapses before it returns to P. 8

Q. 5(c) \vec{F} being a vector, prove that

$$\text{curl curl } \vec{F} = \text{grad div } \vec{F} - \nabla^2 \vec{F}$$

$$\text{where } \nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}.$$

8

Q. 5(d) A triangular lamina ABC of density ρ floats in a liquid of density σ with its plane vertical, the angle B being in the surface of the liquid, and the angle A not immersed. Find ρ/σ in terms of the lengths of the sides of the triangle. 8

Q. 5(e) A heavy uniform rod rests with one end against a smooth vertical wall and with a point in its length resting on a smooth peg. Find the position of equilibrium and discuss the nature of equilibrium. 8

Q. 6(a) Solve the differential equation

$$\frac{d^2y}{dx^2} - 4x \frac{dy}{dx} + (4x^2 - 1)y = -3e^{x^2} \sin 2x$$

by changing the dependent variable.

13

Q. 6(b) Evaluate $\int_S \vec{F} \cdot d\vec{s}$, where $\vec{F} = 4x\vec{i} - 2y^2\vec{j} + z^2\vec{k}$ and s is the surface bounding the region

$$x^2 + y^2 = 4, z = 0 \text{ and } z = 3. \quad 13$$

Q. 6(c) Two bodies of weights w_1 and w_2 are placed on an inclined plane and are connected by a light string which coincides with a line of greatest slope of the plane; if the coefficient of friction between the bodies and the plane are respectively μ_1 and μ_2 , find the inclination of the plane to the horizontal when both bodies are on the point of motion, it being assumed that smoother body is below the other. 14

Q. 7(a) Solve

$$(D^3 + 1)y = e^{\frac{x}{2}} \sin\left(\frac{\sqrt{3}}{2}x\right)$$

where $D = \frac{d}{dx}$. 13

Q. 7(b) A body floating in water has volumes v_1 , v_2 and v_3 above the surface, when the densities of the surrounding air are respectively ρ_1 , ρ_2 , ρ_3 . Find the value of :

$$\frac{\rho_2 - \rho_3}{v_1} + \frac{\rho_3 - \rho_1}{v_2} + \frac{\rho_1 - \rho_2}{v_3}. \quad 13$$

Q. 7(c) A particle is projected vertically upwards with a velocity u , in a resisting medium which produces a retardation kv^2 when the velocity is v . Find the height when the particle comes to rest above the point of projection. 14

Q. 8(a) Apply the method of variation of parameters to solve

$$\frac{d^2y}{dx^2} - y = 2(1 + e^x)^{-1}. \quad 13$$

Q. 8(b) Verify the Divergence theorem for the vector function

$$\vec{F} = (x^2 - yz)\vec{i} + (y^2 - xz)\vec{j} + (z^2 - xy)\vec{k}$$

taken over the rectangular parallelopiped

$$0 \leq x \leq a, \quad 0 \leq y \leq b, \quad 0 \leq z \leq c. \quad 14$$

Q. 8(c) A particle is projected with a velocity v along a smooth horizontal plane in a medium whose resistance per unit mass is double the cube of the velocity. Find the distance it will describe in time t . 13