

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Mains Test Series - 2019

Test number - 12

Section-A

Ques: 1. a} Find whether the following statements are true or false. Justify your answer.

- (i) $\mathbb{Z} \times \mathbb{Z}$ is a cyclic group.
- (ii) $\{a + b\sqrt{2} \in \mathbb{R} \mid a, b \text{ are rational numbers}\}$ is a cyclic group under usual addition of real numbers.
- (iii) $G = \{(1,1), (1,-1), (-1,1), (-1,-1)\}$ is a group under the operation $(a,b)(c,d) = (ac, bd)$ but not a cyclic group.
- (iv) The number of elements of the subgroup $\langle a^{10} \rangle$ in the cyclic group $\langle a \rangle$ of order 30 is 10.
- (v) The symmetric group S_n contains a cyclic group of order 'n'.

Solution :-

(i) False:

Suppose, it is cyclic, with generator (a, b) where $a, b \in \mathbb{Z}$. Then every element of $\mathbb{Z} \times \mathbb{Z}$ would be of the form (na, nb) for some $n \in \mathbb{Z}$, which is not possible.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

ii) False:

Suppose it were cyclic with generator $(m+n\sqrt{2})$ where $m, n \in \mathbb{Q}$, then every element of the set $\{a+b\sqrt{2} \in \mathbb{R} \mid a, b \in \mathbb{Q}\}$ would be of the form $(pm+pn\sqrt{2})$ for some $p \in \mathbb{Q}$ which is not possible.

iii) True:

Consider, composition table for G

*	(1,1)	(1,-1)	(-1,1)	(-1,-1)
(1,1)	(1,1)	(1,-1)	(-1,1)	(-1,-1)
(1,-1)	(1,-1)	(1,1)	(-1,-1)	(-1,1)
(-1,1)	(-1,1)	(-1,-1)	(1,1)	(1,-1)
(-1,-1)	(-1,-1)	(-1,1)	(1,-1)	(1,1)

clearly, we observe that G is a group.

Also, since all elements have order 2 [i.e. no

element has order $4 = O(G)$]

Therefore, G is not a cyclic group.

iv) False:

$$\text{Since, } O(\langle a^{10} \rangle) = \frac{O(a)}{(O(a), 10)} = \frac{30}{(30, 10)} = 3.$$

i.e; number of elements of the subgroup $\langle a^{10} \rangle$ in the cyclic group $\langle a \rangle$ of order 30 is 3.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

(M) True:

A symmetric group S_n contains a cycle of length 'n'. Also order of a cycle of length 'n' is 'n'. Hence, cyclic group of this cycle as a generator has order 'n'.

Ques: 1(b) Let R be commutative ring. If $a, b \in R$ are nilpotent then show that so are $a+b$ and ab for any $\lambda \in R$. Give an example of a non-commutative ring in which a, b are nilpotent but $a+b$ is not.

Solution:

If a and b are nilpotent in a commutative ring R ,

then $a^m = b^n = 0$ for some $m, n \in \mathbb{N}$

② Now,

$$\begin{aligned}
 (a+b)^{m+n} &= a^{m+n} + {}^m C_1 a^{m+n-1} b^1 + {}^m C_2 a^{m+n-2} b^2 \\
 &\quad + \dots + {}^m C_n a^m b^n + \dots \\
 &\quad + b^{m+n} \\
 &= a^m \cdot a^n + {}^m C_1 a^m \cdot a^{n-1} \cdot b^1 + {}^m C_2 a^m b^2 a^{n-2} + \\
 &\quad \dots + {}^m C_n a^m b^n + \dots + b^n \cdot b^m \\
 &\quad [\because R \text{ is commutative}] \\
 &= 0. \quad [\because a^m = b^n = 0]
 \end{aligned}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Hence, $a+b$ are nilpotent.

$$\begin{aligned}
 \textcircled{b} \quad (ax)^m &= (ax)(ax)(ax)\dots(ax) \quad (\text{m-times}) \\
 &= a^m x^m \quad [\because R \text{ is commutative}] \\
 &= 0 \quad [\because a^m = 0]
 \end{aligned}$$

Hence, ax is nilpotent, where $a \in R$ is nilpotent and $x \in R$ (arbitrary).

\textcircled{c} Consider a Ring $R = M_2(\mathbb{C})$

$$\text{let } A = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \text{ and } B = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\text{Clearly, } A^2 = B^2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = 0$$

$\Rightarrow A$ and B are nilpotent elements of R .

$$\text{Now; } A+B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

which is not a nilpotent element of R as there does not exist any positive integer ' n ' such that $(A+B)^n = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$.

Also, $M_2(\mathbb{C})$ is a non-commutative ring.

since, $A \cdot B \neq B \cdot A$.

Hence the result

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques-1(c): Test the convergence of the series:-

$$1 + \frac{x^2}{2} + \frac{x^4}{4} + \frac{x^6}{6} + \dots, x > 0$$

Solution:-

$$\text{Given series} - 1 + \frac{x^2}{2} + \frac{x^4}{4} + \frac{x^6}{6} + \dots ; x > 0$$

Leaving the first term

$$u_n = \frac{x^{2n}}{2^n}$$

$$\therefore u_{n+1} = \frac{x^{2n+2}}{2^{n+2}}$$

$$\frac{u_n}{u_{n+1}} = \frac{2n+2}{2n} \cdot \frac{1}{x^2} = \left(1 + \frac{1}{n}\right) \cdot \frac{1}{x^2}$$

$$\lim_{n \rightarrow \infty} \frac{u_n}{u_{n+1}} = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right) \cdot \frac{1}{x^2} = \frac{1}{x^2}$$

\therefore By Ratio test, the series is convergent if

$$\frac{1}{x^2} > 1 ; \text{i.e } x^2 < 1 \text{ i.e if } x < 1$$

The series is divergent if $\frac{1}{x^2} < 1$ i.e $x^2 > 1$

$$\text{i.e. } x > 1.$$

When $x^2 = 1$ i.e $x=1$, the Ratio test fails

However, when $x=1$, the series becomes

$$1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \dots$$

Leaving first term

$$u_n = \frac{1}{2^n}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Taking $v_n = \frac{1}{n}$

$$\Rightarrow \frac{u_n}{v_n} = \frac{\frac{1}{2n}}{\frac{1}{n}} = \frac{1}{2}.$$

\Rightarrow Let $\lim_{n \rightarrow \infty} \frac{u_n}{v_n} = \frac{1}{2}$; which is finite and non-zero.

\therefore By comparison Test, the series

$\sum u_n$ and $\sum v_n$ converges or diverges together.

But, the series $\sum v_n = \sum \frac{1}{n}$ diverges (by p-test)

$$\text{as } p = 1$$

$\therefore \sum u_n$ diverges.

Hence, $\sum u_n$ converges when $x < 1$ and diverges when $x \geq 1$.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 1) d) If $f(z) = u + iv$ is analytic function and $u - v = e^x \cdot (\cos y - \sin y)$, find $f(z)$ in terms of z .

Solution:- we have;

$$u - v = e^x \cdot (\cos y - \sin y),$$

$$\text{and } f(z) = u + iv$$

$$\text{Then, } \frac{\partial u}{\partial x} - \frac{\partial v}{\partial x} = e^x (\cos y - \sin y) \quad \text{--- (1)}$$

$$\text{and } \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} = e^x (-\sin y - \cos y)$$

$$\Rightarrow \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} = -e^x (\sin y + \cos y)$$

$$\text{or } -\frac{\partial v}{\partial x} - \frac{\partial u}{\partial x} = -e^x (\sin y + \cos y)$$

[By Cauchy-Riemann equations]

$$\text{or } \frac{\partial v}{\partial x} + \frac{\partial u}{\partial x} = e^x (\sin y + \cos y) \quad \text{--- (2)}$$

from (1) and (2); we get

$$\frac{\partial u}{\partial x} = e^x \cos y = \phi_1(x, y).$$

$$\frac{\partial v}{\partial x} = e^x \sin y = \phi_2(x, y)$$

$$\text{Now, } f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \phi_1(z, 0) + i\phi_2(z, 0)$$

$$\therefore f(z) = \int [\phi_1(z, 0) + i\phi_2(z, 0)] dz + C$$

$$f(z) = \int [e^z \cdot \cos 0 + i e^z \cdot \sin 0] dz + C$$

$$f(z) = \int e^z dz + C \Rightarrow f(z) = e^z + C$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Alternative Solution:-

We have as above

$$\frac{\partial V}{\partial x} = e^x \sin y \quad \dots \quad (3)$$

$$\text{and } \frac{\partial U}{\partial x} = e^x \cos y \quad \dots \quad (4)$$

Integrating (3), we get

$$V = e^x \sin y + f(y)$$

Differentiating it w.r.t 'y'

$$\frac{\partial V}{\partial y} = e^x \cos y + f'(y)$$

Also (4) gives. $e^x \cos y = \frac{\partial U}{\partial x} = \frac{\partial V}{\partial y}$

Then (5) and (6) give

$$f'(y) = 0 \text{ and } f(y) = C$$

$$\therefore V = e^x \sin y + C$$

$$\text{and } U = V + e^x (\cos y - \sin y) = e^x \cos y + C$$

$$\therefore f(z) = e^x \cos y + C + i(e^x \sin y + C)$$

$$= e^x (\cos y + i \sin y) + C + iC$$

$$= e^x \cdot e^{iy} + C$$

$$f(z) = e^{x+iy} + C_1$$

$f(z) = e^z + C_1$

which is required solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 1.e} A firm has two bottling plants, one located at Coimbatore and other at Chennai. Each plant produces three drinks, Coca-Cola, Fanta, and Thumps-up named A, B and C respectively. The number of bottles produced per day are, as follows:-

	Plant at	
	Coimbatore (E)	Chennai (F)
Coca-Cola (A)	15,000	15,000
Fanta (B)	30,000	40,000
Thumps-Up (C)	20,000	50,000

A market survey indicates that, during the month of April, there will be a demand of 200,000 bottles of Coca-Cola, 400,000 bottles of Fanta, and 440,000 bottles of Thumps-up. The operating cost per day for plants at Coimbatore, and Chennai is 600 and 400 monetary units respectively. For how many days each plant be run in April so as to, minimize the production cost, while still meeting the market demand?

Solution:

Let, the plants at Coimbatore and Chennai be run for x_1 and x_2 days.

The objective function is given by

$$\text{Minimize } Z = 600x_1 + 400x_2.$$

S.C.

$$15000x_1 + 15000x_2 \geq 200,000$$

$$30000x_1 + 10000x_2 \geq 400,000$$

$$20000x_1 + 50,000x_2 \geq 440,000$$


INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Subject to constraints are re-written as—

$$15x_1 + 15x_2 \geq 200$$

$$30x_1 + 10x_2 \geq 400$$

$$20x_1 + 50x_2 \geq 440$$



Hence, get three optimal point

$$(x_1, x_2) = (0, 40) \Rightarrow Z_{\min} = 600 \times 0 + 400 \times 40 = 16000$$

$$(x_1, x_2) = (10, 6) \Rightarrow Z_{\min} = 600 \times 10 + 400 \times 6 = 6000 + 2400 \\ = 8400$$

$$(x_1, x_2) = (24, 0) \Rightarrow Z_{\min} = 600 \times 24 + 400 \times 0 \\ = 14400$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

1. Optimal feasible Solution.

$$Z_{\min} \text{ at } (10, 6) = 600 \times 10 + 400 \times 6 \\ = 6000 + 2400$$

$Z_{\min} \text{ at } (10, 6) = 8400$

Which is required solution

INSTITUTE OF MATHEMATICAL SCIENCES

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:- 2(a) (i) Show that any infinite cyclic group is isomorphic to $\langle \mathbb{Z}, + \rangle$ the group of integers.

Solution:- Let, $G = \langle a \rangle$ be any infinite cyclic group

Define $f: G \rightarrow \mathbb{Z}$, such that

$$f(a^i) = i \quad i \in \mathbb{Z}$$

Since, $G = \langle a \rangle$ is of infinite order

$a^i \in G$ for all $i \in \mathbb{Z}$

and $a^i = a^j$ for no $i \neq j$.

Thus; $a^i = a^j \Rightarrow i = j$

$\Rightarrow f(a^i) = f(a^j)$ or that f is well defined.

Again; $f(a^i) = f(a^j) \Rightarrow i = j$

$\Rightarrow a^i = a^j \Rightarrow f$ is 1-1

$$f(a^i \cdot a^j) = f(a^{i+j}) = i+j$$

$$f(a^i \cdot a^j) = f(a^i) + f(a^j)$$

Show that f is a homomorphism.

f is obviously onto and hence the isomorphism is established.

Cor. : Every subgroup of an infinite cyclic group is an infinite cyclic group which is isomorphism to the group itself.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 2. a.(ii) Any finite cyclic group of order n is isomorphic to \mathbb{Z}_n , the group of integers addition modulo n .

Solution:- Let, $G = \langle a \rangle$ be a cyclic group, such that

$$o(G) = o(a) = n$$

$$\text{then } G = \{e, a, a^2, \dots, a^{n-1}\}$$

$$\mathbb{Z}_n = \{0, 1, 2, \dots, n-1\}$$

Define $f: G \rightarrow \mathbb{Z}_n$; such that

f is clearly well defined 1-1 onto mapping:

Again;

$$\begin{aligned} f(a^i \cdot a^j) &= f(a^{i+j}) = i+j \\ &= f(a^i) + f(a^j) \end{aligned}$$

Thus, f is a homomorphism and hence an isomorphism.

Remark:- Any two cyclic groups of same order (finite or infinite) are isomorphism.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 2(a).iii) Prove that if an ideal 'U' of a ring 'R' contains a unit of R, then $U=R$.

Solution:-

Consider a Ring R and an ideal U of R which contains a unit of R.

By definition of an ideal,
we know that

$$U \subseteq R \quad \text{--- (1)}$$

Further, Let unit of R be 1 (without loss of generality).
we have, $1 \in U$

also if $r \in R$

$$1 \cdot r \in U \quad [\because U \text{ is an ideal of } R]$$

$$\Rightarrow r \in U.$$

$$\therefore R \subseteq U. \quad \text{--- (2)}$$

from (1) and (2)

$$U = R$$

Hence proved

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 2 (b) i) Determine the values of a, b, c for which the function:

$$f(x) = \begin{cases} \frac{\sin(a+1)x + \sin x}{x} & ; \text{ for } x < 0 \\ c & ; \text{ for } x = 0 \\ \frac{(x+bx^2)^{1/2} - x^{1/2}}{bx^{3/2}} & ; \text{ for } x > 0 \end{cases}$$

is continuous at $x=0$

Solution:-

given function; $f(x) = \begin{cases} \frac{\sin(a+1)x + \sin x}{x} & ; \text{ for } x < 0 \\ c & ; \text{ for } x = 0 \\ \frac{(x+bx^2)^{1/2} - x^{1/2}}{bx^{3/2}} & ; \text{ for } x > 0 \end{cases}$

$$\text{Here; } f(0+0) = \lim_{h \rightarrow 0} f(0+h) = \lim_{h \rightarrow 0} \frac{(h+bh^2)^{1/2} - h^{1/2}}{bh^{3/2}}$$

$$= \lim_{h \rightarrow 0} \frac{h^{1/2}[(1+bh)^{1/2} - 1]}{bh^{3/2} \cdot bh}$$

$$= \lim_{h \rightarrow 0} \frac{\left[1 + \frac{1}{2}bh + \dots\right] - 1}{bh} = \frac{1}{2}.$$

which is independent of 'b' and so 'b' may have any real value except '0'.

$$\text{Again; } f(0-0) = \lim_{h \rightarrow 0} f(0-h) = \lim_{h \rightarrow 0} \frac{\sin(a+1)(-h) + \sin(-h)}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{\sin(a+1)h + \sin h}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2 \sin(\frac{a}{2} + 1) \cdot h \cdot \cos(ah/2)}{h}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$= \lim_{h \rightarrow 0} \frac{\sin \frac{f(a+2)h}{2} h}{\frac{f(a+2)h}{2} h} \cdot \underbrace{(a+2)}_1 \underbrace{\cos \left(\frac{ah}{2}\right)}_1$$

$$= a+2$$

∴ For continuity at $x=0$,
 we have $f(0+0) = f(0-0) = f(0)$

$$\frac{1}{2} = a+2 = c$$

$$\therefore c = \frac{1}{2} \text{ and } a = -\frac{3}{2} \text{ and } b \neq 0$$

Which is required result

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:-2(b)-(iii) Show that the function f defined by
 $f(x) = \frac{1}{x^2}$, $x \neq 0$, is uniformly continuous at $[a, \infty)$
where $a > 0$, but not uniformly continuous at $(0, \infty)$.

Solution:- given function: $f(x) = \frac{1}{x^2}$; $x \neq 0$

Let, $\epsilon > 0$ be given.

Let x_1, x_2 be any two points of $[a, \infty)$ so that

$$x_1 \geq a \quad \text{and} \quad x_2 \geq a$$

$$|x_1| \geq a \quad \text{and} \quad |x_2| \geq a$$

$$\frac{1}{|x_1|} \leq \frac{1}{a} \quad \text{and} \quad \frac{1}{|x_2|} \leq \frac{1}{a}$$

$$\begin{aligned} \text{Now;} \quad |f(x_1) - f(x_2)| &= \left| \frac{1}{x_1^2} - \frac{1}{x_2^2} \right| \\ &= \left| \frac{\frac{1}{x_1} + \frac{1}{x_2}}{x_1 x_2} \right| \left| \frac{1}{x_1} - \frac{1}{x_2} \right| \\ &\leq \left(\frac{1}{|x_1|} + \frac{1}{|x_2|} \right) \left| \frac{x_2 - x_1}{x_1 x_2} \right| \\ &\leq \left(\frac{1}{|x_1|} + \frac{1}{|x_2|} \right) \left(\frac{|x_2 - x_1|}{|x_1| |x_2|} \right) \\ &\leq \left(\frac{1}{a} + \frac{1}{a} \right) \frac{|x_1 - x_2|}{a \cdot a} \\ &= \frac{2}{a^3} |x_1 - x_2| < \epsilon \end{aligned}$$

Whenever; $|x_1 - x_2| < \frac{a^3}{2} \cdot \epsilon$

Choose; $\delta = \frac{a^3}{2} \cdot \epsilon > 0$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

then; $|f(x_1) - f(x_2)| < \varepsilon$; whenever $|x_1 - x_2| < \delta$

$\Rightarrow f$ is uniformly continuous on $[a, \infty)$.

Now, we shall show that f is not uniformly continuous on $(0, \infty)$.

Let $\varepsilon > 0$ be given.

We shall show that for each $\delta > 0$ $\exists x_1, x_2 \in (0, \infty)$ such that

$$|x_1 - x_2| < \delta \Rightarrow |f(x_1) - f(x_2)| \geq \varepsilon$$

The sequence $\langle a_n \rangle$ defined by

$$a_n = \frac{1}{\sqrt{n+2\varepsilon}} - \frac{1}{\sqrt{n}} ; \text{ converges to } 0$$

Since; $a_n \rightarrow 0$ and $n \rightarrow \infty$

\therefore Given $\delta > 0$, \exists a positive integer m , such that

$$|a_n - 0| < \delta \quad \forall n \geq m$$

$$\Rightarrow \left| \frac{1}{\sqrt{n+2\varepsilon}} - \frac{1}{\sqrt{n}} \right| < \delta \quad \forall n \geq m$$

Let, us take $x_1 = \frac{1}{\sqrt{m+2\varepsilon}}$ and $x_2 = \frac{1}{\sqrt{m}}$

then $|x_1 - x_2| < \delta$

whereas, $|f(x_1) - f(x_2)| = |(m+2\varepsilon) - m|$

$$= |m+2\varepsilon - m| = 2\varepsilon > \varepsilon$$

Hence, f is not uniformly continuous on $(0, \infty)$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:- 2(c) } Evaluate $\int_0^{2\pi} \frac{d\theta}{(a+b\cos\theta)^2}$, $a > b > 0$

Solution:-

$$\text{Let ; } I = \int_0^{2\pi} \frac{d\theta}{(a+b\cos\theta)^2} ; a > b > 0$$

$$\begin{aligned} I &= \frac{1}{i} \int_C \frac{dz}{z^2 [a + \frac{1}{2}b(z + \frac{1}{z})]^2} \\ &= \frac{4}{T} \int_C \frac{z dz}{(bz^2 + 2az + b)^2} = \frac{4}{ib^2} \int_C \frac{z dz}{(z^2 + 2\frac{a}{b}z + 1)^2} \\ &= \frac{4}{ib^2} \int_C f(z) dz \quad \text{--- say.} \end{aligned}$$

where, C is the unit circle.

$f(z)$ has poles of order two given by.

$$z^2 + 2\frac{a}{b}z + 1 = 0$$

$$\text{or. } z = \frac{1}{2} \left[-\frac{2a}{b} \pm \sqrt{\frac{4a^2}{b^2} - 4} \right]$$

$$z = \frac{1}{2} \left[-\frac{2a}{b} \pm \frac{1}{b} \sqrt{4a^2 - 4b^2} \right]$$

$$z = \frac{-a \pm \sqrt{a^2 - b^2}}{b}$$

$$\text{Taking } \alpha = \frac{-a + \sqrt{a^2 - b^2}}{b} \text{ and } \beta = \frac{-a - \sqrt{a^2 - b^2}}{b}$$

Thus, $z = \alpha, \beta$ are double poles of $f(z)$.

Since; $a > b > 0$, $|\beta| > 1$

Also; we have $|\alpha\beta| = 1$; therefore $|\alpha| < 1$.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Thus; $z = \alpha$ is the pole of order two inside C .

$$\begin{aligned} \text{We have; } f(z) &= \frac{4}{ib^2} \cdot \frac{z}{[z^2 + 2(\alpha/b)z + 1]^2} \\ &= \frac{4z}{ib^2} \left[\frac{1}{(z-\alpha)^2(z-\beta)^2} \right] = \frac{1}{(z-\alpha)^2} \phi(z) \end{aligned}$$

$$\begin{aligned} \text{Now; } \phi'(z) &= [\phi'(z)]_{z=\alpha} \\ &= \frac{4}{ib^2} \left[\frac{(z-\beta)^2 - 2z(z-\beta)}{(z-\beta)^4} \right]_{z=\alpha} \\ &= \frac{4}{ib^2} \left[\frac{(\alpha-\beta)^2 - 2\alpha(\alpha-\beta)}{(\alpha-\beta)^4} \right] \\ &= \frac{4}{ib^2} \left[\frac{\alpha^2 + \beta^2 - 2\alpha\beta + 2\alpha\beta - 2\alpha^2}{(\alpha-\beta)^4} \right] \\ &= \frac{4}{ib^2} \frac{(\alpha+\beta)}{(\alpha-\beta)^3} = -\frac{ai}{(\alpha^2-b^2)^{3/2}} \end{aligned}$$

∴ Residue at the double pole $z = \alpha$ is

$$\frac{1}{1!} \phi'(\alpha) = -\frac{ai}{(\alpha^2-b^2)^{3/2}}$$

Hence; $\int_0^{2\pi} \frac{d\theta}{(a+b\cos\theta)^2} = 2\pi i \cdot \text{sum of residues at poles inside } C$.

$$\therefore \int_0^{2\pi} \frac{d\theta}{(a+b\cos\theta)^2} = \frac{2\pi a}{(\alpha^2-b^2)^{3/2}}$$

Which is required solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Q.3(a)(i) Show that the set of all polynomials with even coefficients is a prime ideal in $\mathbb{Z}[x]$?

Solution:-

First, we will show that A is an ideal of $\mathbb{Z}[x]$, when A is the set of all polynomials with even integers.

$$\text{Let } f(x) = (2kx^2 + 2jx + 2l) \text{ where } k, j, l \in \mathbb{Z}$$

$$\text{Let } g(x) = (ax^2 + bx + c) \text{ where } a, b, c \in \mathbb{Z}$$

since, all the coefficients of $f(x)$ are even,
 $f(x)g(x)$ must be in A. Now.

$$f(x)g(x) = (2kx^2 + 2jx + 2l)(ax^2 + bx + c)$$

$$f(x)g(x) = 2kax^4 + 2kbx^3 + 2kcx^3 + 2jax^3 + 2jbx^2 \\ + 2jxc + 2alx^2 + 2blx + 2cl$$

$$f(x)g(x) = 2(ka)x^4 + 2(kb+ja)x^3 + 2(kc+jb+la)x^2 \\ + 2(jc+bl)x + 2lc.$$

since, all the coefficients of the products are even, the result is in A, so

A is an Ideal.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Next, we will show that A is a prime ideal. To do this, we need to show that if two polynomials are not in A , then their product cannot be in A .

Let, there be two polynomials, both with odd coefficients of the form $(2k+1)$.

When finding the product of these two polynomials, one of the first product is

$$\begin{aligned} & ((2k+1)x^2) * ((2h+1)x^2) \\ &= (4kh + 2k + 2h + 1)x^4 \\ &= (4(kh + 2(k+h)+1))x^4. \end{aligned}$$

Since, the coefficient ends in $+1$, it is an odd number, so the polynomial cannot be in A .

Hence, 'A' is a prime ideal.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 3(a)(ii)) Let $\mathbb{Z}[i]$ denote the set of all complex numbers of the form $a+bi$ where a and b are integers. Determine all the prime elements of $\mathbb{Z}[i]$.

Solution :-

$$\text{Given } \mathbb{Z}[i] = \{a+bi \mid a, b \in \mathbb{Z}\}.$$

Let us suppose that $a+bi$ is prime.

$$\text{then } (a+bi) \mid (a+bi)(a-bi) = N(a+bi)$$

$$= a^2 + b^2$$

Thus, every prime divides a natural number a^2+b^2 .

[Now, writing this number as a product of primes in \mathbb{N} and keeping in mind that $a+bi$ is a prime in $\mathbb{Z}[i]$, we find that $a+bi$ must divide one of the prime factors of a^2+b^2 .

Therefore, we have the following primes in $\mathbb{Z}[i]$:

- (i) ~~$1+i$~~ , the prime dividing 2;
- (ii) $a+bi$ and $a-bi$, where $p=a^2+b^2=1 \pmod{4}$;
- (iii) rational primes $q \equiv 3 \pmod{4}$.

In particular, $\mathbb{Z}[i]$ has infinitely many primes.

Hence the result

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques. 3(b)) Show that the differential co-efficient of

$$\sum_{n=1}^{\infty} \frac{1}{n^3 + n^4 x^2} \text{ is } -2x \sum_{n=1}^{\infty} \frac{1}{n^2(1+nx^2)^2} \text{ for all real } x.$$

Solve:-

$$\text{Here; } f_n(x) = \frac{1}{n^3 + n^4 x^2} = \frac{1}{n^3(1+nx^2)}$$

$$\Rightarrow f_n'(x) = \frac{1}{n^3} \left[\frac{-2nx}{(1+nx^2)^2} \right] = \frac{-2x}{n^2(1+nx^2)^2}$$

$$\Rightarrow \sum_{n=1}^{\infty} f_n'(x) = -2x \sum_{n=1}^{\infty} \frac{1}{n^2(1+nx^2)^2}$$

Now, $f_n'(x)$ is maximum when $\frac{d}{dx} f_n'(x) = 0$

$$\text{i.e. when } -\frac{2}{n^2} \cdot \frac{(1+nx^2)^2 \cdot 1 - x \cdot 2(1+nx^2) \cdot 2nx}{(1+nx^2)^4} = 0$$

or when ; $1 - 3nx^2 = 0$

$$\Rightarrow x = \frac{1}{\sqrt{3n}}$$

$$\therefore \text{Maximum value of } |f_n'(x)| = \frac{2 \cdot \frac{1}{\sqrt{3n}}}{n^2(1+\frac{1}{3})^2} = \frac{3\sqrt{3}}{8n^{5/2}}$$

$$\Rightarrow |f_n'(x)| \leq \frac{3\sqrt{3}}{8n^{5/2}} = \frac{1}{n^{5/2}} \forall x$$

Since; $\sum_{n=1}^{\infty} \frac{1}{n^{5/2}}$ is convergent

Therefore, by Weierstrass's M-Test, the series

$\sum_{n=1}^{\infty} f_n'$ is uniformly convergent for all real x .

(13)

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

and Hence,

$\sum_{n=1}^{\infty} f_n$ can be differentiated term by term.

$$\therefore \left(\sum_{n=1}^{\infty} f_n \right)' = \sum_{n=1}^{\infty} f'_n$$

$$\Rightarrow \left[\sum_{n=1}^{\infty} \frac{1}{n^3 + n^4 x^2} \right]' = -2x \sum_{n=1}^{\infty} \frac{1}{n^2 (1+nx^2)^2}$$

which is required result

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:- 3(c)) Solve the following LPP by simplex method.

$$\text{Minimize } Z = 4x_1 + x_2$$

Subject to constraints

$$3x_1 + x_2 = 3$$

$$4x_1 + 3x_2 \geq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1, x_2 \geq 0$$

Solution :-

$$\text{Given: Minimize } Z = 4x_1 + x_2$$

for simplex, we need to convert Z into
Maximize form.

$$Z_{\max} = -Z_{\min}$$

$$\therefore Z_{\max} = -4x_1 - x_2$$

Now, we write the given LPP in the standard
form -

$$\text{Max } Z' = -4x_1 - x_2 + OS_1 + OS_2 - MA_1 - MA_2$$

Subject to constraints

$$3x_1 + x_2 + OS_1 + OS_2 + A_1 + A_2 = 3$$

$$4x_1 + 3x_2 - S_1 + OS_2 + OA_1 + A_2 = 6$$

$$x_1 + 2x_2 + OS_1 + S_2 + OA_1 + OA_2 = 4$$

$$x_1, x_2, S_1, S_2, A_1, A_2 \geq 0$$

Where, S_1 - surplus variable

S_2 - slack variable

A_1, A_2 - Artificial variable.

as; S_1 is surplus variable, which should not be a basic variable, since its value $S_1 = -6$ when all variables are zero is negative, which is not feasible.

(14)

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$\therefore S_1$ must be prevented from appearing in initial solution. This is done by taking $S_1 = 0$. By setting other non-basic variables

$$x_1 = x_2 = 0$$

We obtain the IBFS as

$$x_1 = x_2 = 0, S_1 = 0, A_1 = 3, A_2 = 6 \text{ and } S_2 = 4$$

Thus the initial simplex table is :-

		C_j	-4	-1	0	0	-M	-M		
C_B	Basis		x_1	x_2	S_1	S_2	A_1	A_2	b	θ
-M	A_1		3	1	0	0	1	0	3	$3/3 = 1 \rightarrow$
-M	A_2		4	3	-1	0	0	1	6	$6/4 = 3/2$
0	S_2		1	2	0	1	0	0	4	$4/1 = 4.$
$Z_j = \sum C_B a_{ij}$			-7M	-4M	M	0	-M	-M	-9M	
$C_j = C_j - Z_j$			7M-4	4M+1-M	0	0	0	0		



From above table ; $x_1 \rightarrow$ incoming variable
 $A_1 \rightarrow$ Outgoing variable.

Here (3) is the key element and convert it into unity and all other elements in its column to zero. Then the new simplex table is :-

		C_j	-4	-1	0	0	-M	-M		
C_B	Basis		x_1	x_2	S_1	S_2	A_1	A_2	b	θ
-4	x_1		1	$\frac{1}{3}$	0	0	$\frac{1}{3}$	0	1	$\frac{1}{3} \geq 3$
-M	A_2		0	$(\frac{5}{3})$	-1	0		1	2	$2/\frac{5}{3} = \frac{6}{5} \rightarrow$
0	S_2		0	$\frac{5}{3}$	0	1		0	3	$3/\frac{5}{3} = \frac{9}{5}$
$Z_j = \sum C_B a_{ij}$			-4	$-\frac{4}{3} - \frac{5}{3}M + M$	0		-M	-4-2M		
$C_j = C_j - Z_j$			0	$(\frac{5}{3}M - \frac{1}{3})$	-M	0	0			



Outgone

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

From above table ; $x_2 \rightarrow$ Incoming variable
 $A_2 \rightarrow$ Outgoing variable.

Here $(5/3)$ is the key element and make it unity and all the other elements in its column equal to zero.

Then the revised simplex table is :

		C_j	-4	-1	0	0	
C_B	Basis		x_1	x_2	S_1	S_2	b
-4	x_1		1	0	$1/5$	0	$3/5$
-1	x_2		0	1	$-3/5$	0	$6/5$
0	S_2		0	0	1	1	1
$Z_j = C_B \cdot a_{ij}$			-4	-1	$-1/5$	0	$-18/5$
$C_j - Z_j$			0	0	$4/5$	0	

From ; the above table all C_j 's ≤ 0 except S_1 column, which should not be incoming variable. as $S_1 = 0$.

∴ The solution is an optimal BFS to the problem and is given by.

$$x_1 = 3/5, x_2 = 6/5 \text{ and } S_2 = 1$$

$$\text{Max. } Z' = -18/5$$

$$\therefore \text{Minimize } Z = 4x_1 + x_2$$

$$= 4 \times \frac{3}{5} + \frac{6}{5} = \frac{18}{5},$$

$$\therefore \text{Minimize } Z = 18/5$$

which is required solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:- 4(a) show by an example that in a finite commutative ring, every maximal ideal need not be prime.

Solution:- consider,

the Ring $R = \{0, 2, 4, 6\}$ under addition and multiplication modulo 8.

$$\text{Let, } M = \{0, 4\},$$

then M is easily seen to be an ideal of R .

$$\text{Again } 2 \otimes 6 = 4 \in M$$

$$\text{but } 2, 6 \notin M$$

we find M is not a prime ideal.

we show M is maximal.

Let, $M \subseteq N \subseteq R$, where N is an ideal of R

since $\langle M, + \rangle$ will be a subgroup of $\langle N, + \rangle$
 by Lagrange's theorem $O(M) | O(N)$.

$$\text{Similarly } O(N) | O(R) = 4.$$

$$\text{i.e. } 2 | O(N), O(N) | 4$$

$$\text{i.e. } O(N) = 2 \text{ or } 4$$

if $O(N) = 2$, then $M = N$ as $M \subseteq N$

if $O(N) = 4$, then $N = R$ as $N \subseteq R$

Hence, M is a maximal ideal of R .

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 4(b)) Examine the convergence of the integrals:

$$(i) \int_1^\infty \frac{\tan^{-1} x}{x^2} dx$$

$$(ii) \int_0^1 \frac{dx}{(1+x)^2 (1-x)^3}$$

Solution:-

$$(i) I = \int_1^\infty \frac{\tan^{-1} x}{x^2} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{\tan^{-1} x}{x^2} dx$$

$$\text{put } x = \tan \theta ; \quad dx = \sec^2 \theta d\theta$$

$$\Rightarrow \int \frac{\tan^{-1} x}{x^2} dx = \int \frac{\theta}{\tan^2 \theta} \cdot \sec^2 \theta d\theta.$$

$$\Rightarrow \int \theta \cdot \frac{\cos^2 \theta}{\sin^2 \theta} \cdot \frac{1}{\cos^2 \theta} d\theta = \int \theta \csc^2 \theta d\theta$$

$$\Rightarrow \theta (-\cot \theta) - \int 1 (-\cot \theta) d\theta$$

$$\Rightarrow -\theta \cot \theta + \log \sin \theta.$$

$$= -\frac{\tan^{-1} x}{x} + \log \frac{x}{\sqrt{1+x^2}}$$

$$\therefore \int_1^\infty \frac{\tan^{-1} x}{x^2} dx = \lim_{t \rightarrow \infty} \left[-\frac{\tan^{-1} x}{x} + \log \frac{x}{\sqrt{1+x^2}} \right]_1^t$$

$$= \lim_{t \rightarrow \infty} \left[-\frac{\tan^{-1} t}{t} + \log \frac{t}{\sqrt{1+t^2}} + \tan^{-1} 1 - \log \frac{1}{\sqrt{2}} \right]$$

$$= 0 + \lim_{t \rightarrow \infty} \log \frac{1}{\sqrt{1+t^2}} + \frac{\pi}{4} + \frac{1}{2} \log 2$$

$$= \log 1 + \frac{\pi}{4} + \frac{1}{2} \log 2 = \frac{\pi}{4} + \frac{1}{2} \log 2.$$

which is finite.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$\Rightarrow \int_1^{\infty} \frac{\tan^{-1} x}{x^2} dx$ is convergent and its value is

$$\int_1^{\infty} \frac{\tan^{-1} x}{x^2} dx = \frac{\pi}{4} + \frac{1}{2} \log 2$$

$$(ii) \text{ Here; } f(x) = \frac{1}{(1+x)^2(1-x)^3}$$

x_1 is the only point of infinite discontinuity of f on $[0, 1]$.

$$\text{Take; } g(x) = \frac{1}{(1-x)^3}$$

$$\text{Then; } \lim_{x \rightarrow 1^-} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 1^+} \frac{1}{(1+x)^2} = \frac{1}{(1+1)^2} = \frac{1}{4}.$$

which is non-zero and finite.

\therefore By comparison test,

$\int_0^1 f(x) dx$ and $\int_0^1 g(x) dx$ converges or diverges together.

$$\text{But } \int_0^1 g(x) dx = \int_0^1 \frac{dx}{(1-x)^3} \quad [\text{form } \int_a^b \frac{dx}{(b-a)^n} \text{ with } \because n=3 > 1] \\ \& b=1]$$

$\therefore g(x)$ diverges.

$\therefore \int_0^1 f(x) dx = \int_0^1 \frac{dx}{(1+x)^2(1-x)^3}$ is divergent.

Which is required result.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:- 4(c) Expand $f(z) = \frac{1}{(z+1)(z+3)}$ in a Laurent's series

valid for the regions :

(i) $|z| < 1$

(ii) $1 < |z| < 3$

(iii) $|z| > 3$

(iv) $0 < |z+1| < 2$

Solution:- we have; $f(z) = \frac{1}{(z+1)(z+3)}$

Resolving into partial fraction, we get

$$f(z) = \frac{1}{2(z+1)} - \frac{1}{2(z+3)}$$

(i) $|z| < 1$

we have ; $f(z) = \frac{1}{2}(z+1)^{-1} - \frac{1}{6}(1+\frac{z}{3})^{-1}$

$$f(z) = \frac{1}{2}[1-z+z^2-z^3+\dots] - \frac{1}{6}\left[1-\frac{z}{3}+\frac{z^2}{9}-\frac{z^3}{27}+\dots\right]$$

$$f(z) = \left(\frac{1}{2}-\frac{1}{6}\right) - \left(\frac{1}{2}-\frac{1}{18}\right)z + \left(\frac{1}{2}-\frac{1}{54}\right)z^2 + \dots$$

$$\boxed{f(z) = \frac{1}{3} - \frac{4}{9}z + \frac{13}{27}z^2 - \dots}$$

(ii) $1 < |z| < 3$

Then, we have ; $\frac{1}{|z|} < 1$ and $\frac{|z|}{3} < 1$.

$$\text{Now, } \frac{1}{2(z+1)} = \frac{1}{2z(1+\frac{1}{z})} = \frac{1}{2z} \left(1 + \frac{1}{z}\right)^{-1}$$

$$= \frac{1}{2z} \left[1 - \frac{1}{z} + \frac{1}{z^2} - \frac{1}{z^3} + \dots\right]$$

$$= \frac{1}{2z} - \frac{1}{2z^2} + \frac{1}{2z^3} - \frac{1}{2z^4} + \dots$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\begin{aligned} \text{and } \frac{1}{2(z+3)} &= \frac{1}{6(1+\frac{z}{3})} = \frac{1}{6} \left[1 + \frac{z}{3} \right]^{-1} \\ &= \frac{1}{6} \left[1 - \frac{z}{3} + \frac{z^2}{9} - \frac{z^3}{27} + \dots \right] \\ &= \frac{1}{6} - \frac{1}{18}z + \frac{1}{54}z^2 - \frac{1}{162}z^3 + \dots \end{aligned}$$

Thus; the Laurent's Series valid for the region
 $|z| < |z| < 3$ is

$$\therefore f(z) = \dots + \frac{1}{2z^3} - \frac{1}{2z^2} + \frac{1}{2z} + \frac{1}{6} - \frac{1}{18}z + \frac{1}{54}z^2 - \dots$$

(iii) $|z| > 3$

Then; $\frac{3}{|z|} < 1$

$$\therefore f(z) = \frac{1}{2z} \left(1 + \frac{1}{z} \right)^{-1} - \frac{1}{2z} \left(1 + \frac{3}{z} \right)^{-1}$$

$$f(z) = \frac{1}{2z} \left[1 - \frac{1}{z} + \frac{1}{z^2} - \frac{1}{z^3} + \dots \right] - \frac{1}{2z} \left[1 - \frac{3}{z} + \frac{9}{z^2} - \frac{27}{z^3} + \dots \right]$$

$$f(z) = \left[\frac{1}{2z} - \frac{1}{2z^2} + \frac{1}{2z^3} - \frac{1}{2z^4} + \dots \right] + \left[\frac{1}{2z} + \frac{3}{6z^2} - \frac{9}{2z^3} + \frac{27}{2z^4} + \dots \right]$$

$$f(z) = \frac{1}{z^2} - \frac{4}{z^3} + \frac{13}{z^4} - \frac{40}{z^5} + \dots$$

(iv) $0 < |z+1| < 2$

Let $z+1 = u$; then $0 < |u| < 2$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\therefore f(z) = \frac{1}{(z+1)(z+3)} = \frac{1}{u(u+2)} = \frac{1}{24} \left(1 + \frac{1}{u+2} \right)^{-1}$$

$$f(z) = \frac{1}{24} \left[1 - \frac{1}{2} + \frac{u^2}{4} - \frac{u^3}{8} + \dots \right]$$

$$f(z) = \frac{1}{2u} - \frac{1}{4} + \frac{u}{8} - \frac{u^2}{16} + \dots$$

$$f(z) = \frac{1}{2(z+1)} - \frac{1}{4} + \frac{z+1}{8} - \frac{(z+1)^2}{16} + \dots$$

which is required result

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 4(d)) A department head has four subordinates, and four tasks to be performed. The subordinates differ in efficiency, and the tasks in their intrinsic difficulty. His estimate of the times each man would take to perform each task is given in the effectiveness matrix below. How should the tasks be allocated, one to a man, so as to minimize the total men hours?

		Subordinates			
		I	II	III	IV
Task	A	8	26	17	11
	B	13	28	4	26
	C	38	19	18	15
	D	19	26	24	10

Solution:- Given ; Effective matrix

	I	II	III	IV
A	8	26	17	11
B	13	28	4	26
C	38	19	18	15
D	19	26	24	10

Step-1) Subtract the least element of the row
and from all the elements of that row;

The reduced Table

	I	II	III	IV
A	0	18	9	3
B	9	24	0	22
C	23	4	3	0
D	9	16	14	0

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Now, check whether all the rows and each column have at least one zero; which is not here; hence, subtract the least element of each column from all other elements of that column.

The reduced table is

	I	II	III	IV
A	0	14	9	3
B	9	20	0	22
C	23	0	3	0
D	9	12	14	0

Now, cover all the zeros using minimum number of Horizontal and vertical lines.

Here; no. of lines = no. of zeros = 4.

Hence, optimality achieved.

For optimal solution, the

Table to choose only one zero from each row with least no. of zeros, & discard others in respective row and column.

	I	II	III	IV
A	0	14	9	3
B	9	20	0	22
C	23	0	3	X
D	9	12	14	0

$\therefore A \rightarrow I$
 $B \rightarrow II$
 $C \rightarrow II$
 $D \rightarrow IV$.

$$\therefore \text{Minimized Men hours} = 8 + 4 + 19 + 10 \\ = 31 \text{ hours.}$$

$$Z_{\min} = 31 \text{ hours}$$

which is required solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Section-B

Ques 5(a)) find the complete integral of

$$(x+y)(p+q)^2 + (x-y)(p-q)^2 = 1.$$

Solution:-

Let x and y be two new variables such that

$$x^2 = x+y \quad \text{and} \quad y^2 = x-y \quad \dots \quad (1)$$

∴ Given equation is $(x+y)(p+q)^2 + (x-y)(p-q)^2 = 1 \quad \dots \quad (2)$

$$\text{Now; } p = \frac{\partial z}{\partial x} = \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial x} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial x}$$

$$p = \frac{1}{2x} \frac{\partial z}{\partial x} + \frac{1}{2y} \cdot \frac{\partial z}{\partial y} \quad \dots \quad (3)$$

$$\therefore \text{from (1), } \frac{\partial x}{\partial x} = \frac{1}{2x} \text{ and } \frac{\partial y}{\partial x} = \frac{1}{2y}$$

$$\text{and } q = \frac{\partial z}{\partial y} = \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial y} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial y}$$

$$q = \frac{1}{2x} \frac{\partial z}{\partial x} - \frac{1}{2y} \frac{\partial z}{\partial y} \quad \dots \quad (4)$$

$$[\because \text{from (1); } \frac{\partial x}{\partial y} = \frac{1}{2x}; \frac{\partial y}{\partial y} = \frac{-1}{2y}]$$

From (3) & (4) \Rightarrow

$$p+q = \frac{1}{x} \frac{\partial z}{\partial x} \quad \text{and} \quad p-q = \frac{1}{y} \frac{\partial z}{\partial y} \quad \dots \quad (5)$$

From, using (5) and (1), (2) reduces to

$$x^2 \cdot \frac{1}{x^2} \left(\frac{\partial z}{\partial x} \right)^2 + y^2 \cdot \frac{1}{y^2} \left(\frac{\partial z}{\partial y} \right)^2 = 1$$

$$P^2 + Q^2 = 1$$

$$\dots \quad (6)$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Where; $P = \frac{\partial z}{\partial x}$ and $Q = \frac{\partial z}{\partial y}$

(4) is of the form $f(P, Q) = 0$

\therefore Solution of (4) is

$$z = ax + by + c \quad \text{--- (7)}$$

where; $a^2 + b^2 = 1$

$$\text{or } b = \sqrt{1-a^2}$$

Putting a for P and b for Q in (6)

\therefore from (7), the required complete integral is

$$z = ax + y\sqrt{1-a^2} + c$$

$$z = a\sqrt{x+y} + \sqrt{x-y}\sqrt{1-a^2} + c \quad \text{--- by (1)}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:- 5(b) Solve $(D^2 + DD' - 6D'^2)z = x^2 \sin(x+y)$

Solve:- Given equation

$$(D^2 + DD' - 6D'^2)z = x^2 \sin(x+y)$$

$$\Rightarrow (D+3D')(D-2D')z = x^2 \sin(x+y)$$

Its auxillary equation is

$$(m+3)(m-2) = 0$$

$$m = -3, 2$$

$$\therefore C.F = \phi_1(y-3x) + \phi_2(y+2x)$$

where, ϕ_1 and ϕ_2 being arbitrary function.

$$\begin{aligned}
 P.I &= \frac{1}{(D+3D')(D-2D')} x^2 \sin(x+y) \\
 &= \frac{1}{(D+3D')} \left\{ \frac{1}{(D-2D')} x^2 \cdot \sin(x+y) \right\} \\
 &= \frac{1}{(D+3D')} \int x^2 \sin(x+c-2x) dx ; \quad \text{where } y = c-2x \\
 &= \frac{1}{(D+3D')} \int x^2 \sin(c-x) dx \\
 &= \frac{1}{(D+3D')} \left[x^2 \cos(c-x) - \int 2x \cos(c-x) dx \right] \\
 &= \frac{1}{(D+3D')} \left[x^2 \cos(c-x) - \left\{ -2x \sin(c-x) + \int 2 \sin(c-x) dx \right\} \right] \\
 &= \frac{1}{(D+3D')} \left[x^2 \cos(c-x) + 2x \sin(c-x) - 2 \cos(c-x) \right] \\
 &= \frac{1}{(D+3D')} \left[(x^2-2) \cos(c-x) + 2x \sin(c-x) \right] \\
 &= \frac{1}{(D+3D')} \left[(x^2-2) \cos(x+y) + 2x \sin(x+y) \right] \left[\because y = 2x-c \right. \\
 &\quad \left. \therefore c = y+2x \right]
 \end{aligned}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$= \int [x^2 - 2] [\cos(x+c'+3x) + 2x \sin(x+c'+3x)] dx$$

$$\text{where; } c' = y - 3x$$

$$= \int [x^2 - 2] \cos(4x+c') + 2x \sin(4x+c') dx$$

$$= (x^2 - 2) \frac{\sin(4x+c')}{4} - \int 2x \frac{\sin(4x+c')}{4} dx + 2 \int x \sin(4x+c') dx.$$

Integrating by parts 1st integral and keeping the second integral unchanged.

$$= \frac{1}{4}(x^2 - 2) \sin(4x+c') + \frac{3}{2} \int x \sin(4x+c') dx$$

$$= \frac{(x^2 - 2) \sin(4x+c')}{4} + \frac{3}{2} \left[-x \frac{\cos(4x+c')}{4} + \int \frac{\cos(4x+c')}{4} dx \right]$$

$$= \frac{x^2 - 2}{4} \sin(4x+c') + \left(-\frac{3}{8} x \cos(4x+c') + \frac{3}{24} \sin(4x+c') \right)$$

$$\Rightarrow \frac{1}{4}(x^2 - 2) \sin(4x+c') - \frac{3}{8} x \cos(4x+c') + \frac{3}{24} \sin(4x+c')$$

$$\Rightarrow \frac{1}{4}(x^2 - 2) \sin(x+y) - \frac{3}{8} x \cos(x+y) + \frac{3}{24} \sin(x+y).$$

$$\therefore P.I. = \left(\frac{x^2}{4} - \frac{13}{32} \right) \sin(x+y) - \frac{3}{8} x \cos(x+y)$$

∴ The required solution is $z = C.F + P.I.$

$$z = \Phi_1(y-3x) + \Phi_2(y+2x) + \left(\frac{x^2}{4} - \frac{13}{32} \right) \sin(x+y) - \frac{3}{8} x \cos(x+y)$$

Required Solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:-5(c) } The quadratic equation $x^4 - 4x^2 + 4 = 0$ has double root at $x = \sqrt{2}$. Starting with $x_0 = 1.5$, compute three successive iterations to the roots by Newton-Raphson method. Does the result converges quadratically or linearly?

Solution:-

The Newton-Raphson method for determining a multiple root of multiplicity m is given by

$$x_{k+1} = x_k - \frac{m f_k}{f'_k}$$

Since, the quadratic equation $x^4 - 4x^2 + 4 = 0$ has double root at $x = \sqrt{2}$; i.e $m=2$, we have

$$x_{n+1} = x_n - \frac{2 f(x_n)}{f'(x_n)} \Rightarrow f(x) = x^4 - 4x^2 + 4 \\ f'(x) = 4x^3 - 8x$$

and given, $x_0 = 1.5$.

Then,

$$x_1 = x_0 - \frac{2 f(x_0)}{f'(x_0)} = 1.5 - 2 \frac{f(1.5)}{f'(1.5)}$$

$$x_1 = 1.5 - 2 \times \frac{0.0625}{1.5} = 1.5 - 0.08333$$

$$x_1 = 1.41667$$

for 2nd Iteration

$$x_2 = x_1 - \frac{2 f(x_1)}{f'(x_1)} = 1.41667 - 2 \frac{f(1.41667)}{f'(1.41667)}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$x_2 = 1.41667 - 2 \cdot \left[\frac{0.0001}{0.03864} \right]$$

$$x_2 = 1.41667 - 0.005071 = 1.4116$$

for third Iteration

$$x_3 = x_2 - \frac{2f(x_2)}{f'(x_2)} = 1.4116 - \frac{2f(1.4116)}{f'(1.4116)}$$

$$x_3 = 1.4116 - 2 \left[\frac{0.00004}{-0.0417} \right]$$

$$x_3 = 1.4116 + 0.001918 = 1.41352$$

$$\therefore x_1 = 1.41667 ; x_2 = 1.4116 ; x_3 = 1.41352$$

$$\& [x_3 = 1.41352 \approx 1.414 = \sqrt{2}]$$

Thus, the method $x_{k+1} = x_k - \frac{mf_k}{f'_k}$ has

quadratic rate of convergence.

[\because In the error term

$$\epsilon_{k+1} = \left(1 - \frac{\alpha}{m}\right) \epsilon_k + \frac{\alpha}{m^2(m+1)f^m(\xi)} \epsilon_k^2 + O(\epsilon_k^3)$$

The coefficient of ϵ_k vanishes

$$\text{i.e. } \alpha = m = 2$$

]

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 5(d) Convert the following:

(i) $(41.6875)_{10}$ to binary number

Solve:-

2	41	
2	20	1
2	10	0
2	5	0
2	2	1
1	1	0

$$\begin{aligned}
 0.6875 \times 2 &= 1.3750 \rightarrow 1 \\
 0.3750 \times 2 &= 0.7500 \rightarrow 0 \\
 0.7500 \times 2 &= 1.5000 \rightarrow 1 \\
 0.5000 \times 2 &= 1.0000 \rightarrow 1 \\
 0.0000 &
 \end{aligned}$$

$$\therefore (41.6875)_{10} \leftrightarrow (101001.1011)_2$$

(ii) $(101101)_2$ to decimal number.

$$\begin{aligned}
 \text{Solve:- } 1 \times 2^5 + 0 \times 2^4 + 1 \times 2^3 + 1 \times 2^2 + 0 \times 2^1 + 1 \times 2^0 \\
 = 32 + 0 + 8 + 4 + 0 + 1 \\
 = 45.
 \end{aligned}$$

$$\therefore (101101)_2 \leftrightarrow (45)_{10}$$

(iii) $(AF63)_{16}$ to decimal number.

$$\begin{aligned}
 \text{Solve:- } A \times 16^3 + F \times 16^2 + 6 \times 16^1 + 3 \times 16^0 \\
 = 10 \times 16^3 + 15 \times 256 + 6 \times 16 + 3 \\
 = 40960 + 3840 + 96 + 3 = 44899.
 \end{aligned}$$

$$\therefore (AF63)_{16} \leftrightarrow (44899)_{10}$$

(iv) $(101111011111)_2$ to Hexadecimal number

$$\begin{array}{ccc}
 \text{Solve:- } (1011 & 1101 & 1111)_2 & \leftrightarrow & (BDF)_{16} \\
 \text{B} & \text{D} & \text{F}
 \end{array}$$

$$\therefore (101111011111)_2 \leftrightarrow (BDF)_{16}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:- 5 (e) > The x -component of velocity is

$$u = x^3 + z^4 + 6 \text{ and } y \text{ component is } v = y^3 + z^4.$$

Find the simplest z -component of velocity that satisfy continuity.

Solution:-

Given; x -component of velocity is

$$u = x^3 + z^4 + 6$$

y -component of velocity is

$$v = y^3 + z^4.$$

Hence; for simplest z -component of velocity that satisfy equation of continuity.

i.e.
$$\boxed{\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = 0} \quad \text{--- (1)}$$

$$\frac{\partial u}{\partial x} = 3x^2 \quad \text{--- (A)}$$

$$\frac{\partial v}{\partial y} = 3y^2 \quad \text{--- (B)}$$

Put (A) & (B) in (1), we get.

$$\frac{\partial w}{\partial z} = -(3x^2 + 3y^2)$$

$$\Rightarrow w = -(3x^2 + 3y^2) z + f(x, y).$$

∴ Simplest z -component of a velocity

$$\boxed{w = -(3x^2 + 3y^2) z}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 6(a) Find the integral surface of

$x^2 p + y^2 q + z^2 = 0$, $p = \frac{\partial z}{\partial x}$ and $q = \frac{\partial z}{\partial y}$ which passes through the hyperbola $xy = x+y$, $z=1$.

Solution:-

$$\text{Given, } x^2 p + y^2 q + z^2 = 0$$

$$x^2 p + y^2 q = -z^2 \quad \dots \textcircled{1}$$

$$\text{Given curve; } xy = x+y ; z=1 \quad \dots \textcircled{2}$$

Here Lagrange's auxiliary equations are

$$\frac{dx}{x^2} = \frac{dy}{y^2} = \frac{dz}{-z^2} \quad \dots \textcircled{3}$$

from first and third fractions:

$$\frac{dx}{x^2} + \frac{dz}{z^2} = 0$$

$$\Rightarrow -\frac{1}{x} - \frac{1}{z} = -c_1 \Rightarrow \frac{1}{x} + \frac{1}{z} = c_1 \quad \dots \textcircled{4}$$

from 2nd and 3rd fractions:

$$\frac{dy}{y^2} + \frac{dz}{z^2} = 0$$

$$\Rightarrow \frac{1}{y} + \frac{1}{z} = c_2 \quad \dots \textcircled{5}$$

$$\text{Adding } \textcircled{4} \text{ and } \textcircled{5}, \quad \frac{1}{x} + \frac{1}{y} + \frac{2}{z} = c_1 + c_2$$

$$\Rightarrow \frac{x+y}{xy} + \frac{2}{z} = c_1 + c_2 \Rightarrow \frac{xy}{xy} + \frac{2}{z} = c_1 + c_2 \quad (\text{from } \textcircled{2})$$

$$1 + \frac{2}{z} = c_1 + c_2 \Rightarrow 1 + 2 = c_1 + c_2 \quad [\text{from } \textcircled{2}]$$

$$\therefore \frac{1}{x} + \frac{1}{y} + \frac{2}{z} = 3 \Rightarrow \boxed{yz + 2xy + xz = 3xyz} \quad \begin{matrix} \text{required} \\ \text{solution.} \end{matrix}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 6(b)} Reduce the equation

$x^2(y-1)r + (-x)(y^2-1)s + y(y-1)t + xy p - q = 0$
 to canonical form and hence solve it.

Solution:- Given

$$x^2(y-1)r - x(y^2-1)s + y(y-1)t + xy p - q = 0 \quad \text{--- (1)}$$

Comparing (1) with

$$Rr + Ss + Tt + f(x, y, z, p, q) = 0, \text{ we get}$$

$$R = x^2(y-1); S = -x(y^2-1); T = y(y-1)$$

$\therefore \lambda$ -quadratic $R\lambda^2 + S\lambda + T = 0$ reduces to

$$x^2(y-1)\lambda^2 - x(y^2-1)\lambda + y(y-1) = 0$$

$$\therefore \lambda_1 = y/x, \lambda_2 = 1/x \text{ (real & distinct)}$$

$$\text{so; } (\frac{dy}{dx}) + \lambda_1 = 0 \text{ and } (\frac{dy}{dx}) + \lambda_2 = 0$$

$$\Rightarrow \frac{dy}{dx} + \frac{y}{x} = 0$$

$$\frac{dy}{dx} + \frac{1}{x} = 0$$

$$\text{Integrating these, } xy = C_1$$

$$xe^y = C_2$$

So, for canonical form, we take

$$u = xy \text{ and } v = xe^y \quad \text{--- (2)}$$

$$\therefore p = \frac{\partial z}{\partial x} = \frac{\partial z}{\partial u} \cdot \frac{\partial u}{\partial x} + \frac{\partial z}{\partial v} \cdot \frac{\partial v}{\partial x}$$

$$p = \frac{\partial z}{\partial x} = y \frac{\partial z}{\partial u} + e^y \cdot \frac{\partial z}{\partial v} \quad (\text{by 2}) \quad \text{--- (3)}$$

$$q = \frac{\partial z}{\partial y} = \frac{\partial z}{\partial u} \cdot \frac{\partial u}{\partial y} + \frac{\partial z}{\partial v} \cdot \frac{\partial v}{\partial y}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$q = \frac{\partial z}{\partial y} = x \frac{\partial z}{\partial u} + x e^y \cdot \frac{\partial z}{\partial v} \quad \text{--- (4)}$$

$$r = \frac{\partial^2 z}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial x} \right) = \frac{\partial}{\partial x} \left(y \frac{\partial z}{\partial u} + e^y \frac{\partial z}{\partial v} \right) \quad (\text{by (3)})$$

$$r = y \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial u} \right) + e^y \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial v} \right)$$

$$\begin{aligned} r = y & \left[\frac{\partial}{\partial u} \left(\frac{\partial z}{\partial u} \right) \cdot \frac{\partial u}{\partial x} + \frac{\partial}{\partial v} \left(\frac{\partial z}{\partial u} \right) \frac{\partial v}{\partial x} \right] + e^y \left[\frac{\partial}{\partial u} \left(\frac{\partial z}{\partial v} \right) \cdot \frac{\partial u}{\partial x} \right. \\ & \left. + \frac{\partial}{\partial v} \left(\frac{\partial z}{\partial v} \right) \cdot \frac{\partial v}{\partial x} \right] \end{aligned}$$

$$r = y^2 \frac{\partial^2 z}{\partial u^2} + 2y e^x \frac{\partial^2 z}{\partial u \partial v} + e^{2y} \frac{\partial^2 z}{\partial v^2}$$

$$s = \frac{\partial^2 z}{\partial x \partial u} = \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial u} \right) = \frac{\partial}{\partial x} \left(x \cdot \frac{\partial z}{\partial u} + x e^y \frac{\partial z}{\partial v} \right)$$

$$s = \frac{\partial z}{\partial u} + x \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial u} \right) + e^y \frac{\partial z}{\partial v} + x e^y \frac{\partial}{\partial x} \left(\frac{\partial z}{\partial v} \right)$$

$$s = \frac{\partial z}{\partial u} + e^y \frac{\partial z}{\partial v} + x \left[\frac{\partial}{\partial u} \left(\frac{\partial z}{\partial u} \right) \frac{\partial u}{\partial x} + \frac{\partial}{\partial v} \cdot \left(\frac{\partial z}{\partial u} \right) \frac{\partial v}{\partial x} \right]$$

$$+ x e^y \left[\frac{\partial}{\partial u} \left[\frac{\partial z}{\partial v} \right] \frac{\partial u}{\partial x} + \frac{\partial}{\partial v} \left(\frac{\partial z}{\partial v} \right) \frac{\partial v}{\partial x} \right]$$

$$s = \frac{\partial z}{\partial u} + e^y \frac{\partial z}{\partial v} + x y \frac{\partial^2 z}{\partial u^2} + (y x e^y + e^y x) \frac{\partial^2 z}{\partial u \partial v} + x e^{2y} \frac{\partial^2 z}{\partial v^2}$$

$$\text{and } t = \frac{\partial^2 z}{\partial y^2} = \frac{\partial}{\partial y} \left(\frac{\partial z}{\partial y} \right) = \frac{\partial}{\partial y} \left[x \frac{\partial z}{\partial u} + x e^y \frac{\partial z}{\partial v} \right]$$

$$t = x \frac{\partial}{\partial y} \left(\frac{\partial z}{\partial u} \right) + x e^y \frac{\partial z}{\partial v} + x \cdot e^y \frac{\partial}{\partial y} \left(\frac{\partial z}{\partial v} \right)$$

$$\begin{aligned} t = x & \left[\frac{\partial}{\partial u} \left(\frac{\partial z}{\partial u} \right) \frac{\partial u}{\partial y} + \frac{\partial z}{\partial v} \left(\frac{\partial z}{\partial u} \right) \frac{\partial v}{\partial y} \right] + x e^y \frac{\partial z}{\partial v} + \\ & x e^y \left[\frac{\partial}{\partial u} \left(\frac{\partial z}{\partial v} \right) \left(\frac{\partial u}{\partial y} \right) + \frac{\partial}{\partial v} \left(\frac{\partial z}{\partial v} \right) \frac{\partial v}{\partial y} \right] \end{aligned}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$t = x^2 \frac{\partial^2 z}{\partial u^2} + 2x^2 e^y \frac{\partial^2 z}{\partial u \partial v} + x^2 e^{2y} \frac{\partial^2 z}{\partial y^2} + x e^y \frac{\partial z}{\partial v}$$

Substituting the values of P, Q, R, S, t in ①
 and simplifying, we get

$$\frac{\partial^2 z}{\partial u \partial v} = 0 ; \text{ which is canonical form of } ①$$

Integrating it w.r.t 'u' -

$$\frac{\partial z}{\partial v} = \phi(v)$$

Integrating it w.r.t 'v' -

$$z = \phi_1(v) + \phi_2(u)$$

$$z = \phi_1(xe^y) + \phi_2(xy)$$

- by ②

required solution

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 6(c) > find the characteristics of the equation $pq = z$ and determine the integral surface which passes through straight line $x=1, z=y$.

Solution:-

If the initial data curve is given in parametric form as -

$$x_0(s) = 1 ; \quad y_0(s) = s ; \quad z_0(s) = s$$

then ordinarily the solution is sought in parametric form as -

$$x = x(t, s) ; \quad y = y(t, s) , \quad z = z(t, s)$$

Thus, using the given data, the differential equation becomes

$$p_0(s) q_0(s) - s = 0 = F \quad \text{--- (1)}$$

and the strip condition gives

$$1 = p_0(0) + q_0(1) \quad \text{or} \quad q_0 = 1 \quad \text{--- (2)}$$

Therefore;

$$q_0 = 1 , \quad p_0 = s \quad (\text{unique initial strip}) \quad \text{--- (3)}$$

Now, the characteristic equations for the given PDE are,

$$\left. \begin{aligned} \frac{dx}{dt} &= q , \quad \frac{dy}{dt} = p , \quad \frac{dz}{dt} = 2pq \\ \frac{dp}{dt} &= p , \quad \frac{dq}{dt} = q . \end{aligned} \right\} - \text{--- (4)}$$

On integration, we get

$$\left. \begin{aligned} p &= C_1 \exp(t) , \quad q = C_2 \exp(t) , \quad x = C_2 \exp(t) + C_3 \\ y &= C_1 \exp(t) + C_4 ; \quad z = 2C_1 C_2 \exp(2t) + C_5 \end{aligned} \right\} - \text{--- (5)}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Now, taking into account the initial conditions

$$x_0 = 1, y_0 = s, z_0 = s, p_0 = s, q_0 = 1 \quad \text{--- (6)}$$

We can determine the constants of integration
and obtain

$$(\text{Since } c_2 = 1, c_3 = 0)$$

$$p = s \exp(t), q = \exp(t), x = \exp(t) \quad \}$$

$$y = s \exp(t), z = s \exp(2t) \quad \text{--- (7)}$$

Consequently, the required integral surface is obtained from Eq.(7) as

$$\boxed{z = xy}$$

which is required solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 6(d)} A uniform string of line density ' ρ ' is stretched to tension ρc^2 and executes a small transverse vibration in a plane through the undisturbed line of string. The ends $x=0$, L of the string are fixed. The string is at rest, with the point $x=b$ drawn aside through a small distance ' ϵ ' and released at time $t=0$. Find the expression for the displacement $y(x, t)$.

Solution:-

The transverse vibration of the string is described by -

$$\text{P.D.E: } y_{xx} = \frac{1}{c^2} y_{tt} \quad \dots \quad (1)$$

The boundary and initial conditions are

$$\text{BCs: } y(0, t) = y(L, t) = 0$$

$$\text{IC: } y_t(x, 0) = 0$$

Using the variables separable method, let

$$y(x, t) = X(x) T(t)$$

then, we have from eq (1)


$$\frac{X''}{X} = \frac{1}{c^2} \cdot \frac{T''}{T} = \pm \lambda^2$$

The equation of the string is given by

$$y(x, 0) = \begin{cases} \frac{\epsilon x}{b}, & 0 \leq x \leq b \\ \frac{\epsilon(x-L)}{(b-L)}, & b \leq x \leq L \end{cases}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

The solution to the given problem, discussed now for various values of λ .



Case I) Taking the constant $\lambda=0$, we have

$$X'' = T'' = 0$$

whose general solution is

$$X = Ax + B, \quad T = Ct + D$$

Therefore;

$$y(x,t) = (Ax+B)(Ct+D)$$

Using, the BCs, at $x=0, L$,

we observe that $A=B=0$, implies a trivial solution.

Case II) Taking the constant as $+\lambda^2$, we have

$$X'' - \lambda^2 X = 0 = T'' - c^2 \lambda^2 T$$

Thus, the general solution is

$$y(x,t) = (A \cosh \lambda x + B \sinh \lambda x)(C \cosh ct + D \sinh ct)$$

Now, the BCs!

$$y(0,t) = 0 \text{ gives } A=0$$

and; $y(L,t) = 0 \text{ gives } B \sinh \lambda L = 0$

which is possible only if $B=0$.

Thus, we are again getting only a trivial solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Case II) If the constant is $-\lambda^2$, then we have

$$x'' + \lambda^2 x = 0 = T'' + C^2 \lambda^2 T = 0$$

In this case, the general solution is

$$y(x,t) = (A \cos \lambda x + B \sin \lambda x)(P \cos \omega t + Q \sin \omega t)$$

Using the BCs:-

$$y(0,t) = 0 \quad \text{gives } A = 0$$

$$y(L,t) = 0 \quad \text{gives } B \sin \lambda L = 0$$

for a non-trivial solution, $B \neq 0 \Rightarrow \lambda L = n\pi$.

$$\text{Therefore, } \lambda = \frac{n\pi}{L}, \quad n = 1, 2, 3, \dots$$

Also using the IC: $y_t(x,0) = 0$

we can notice that $Q = 0$.

Hence, the acceptable non-trivial solution is

$$y(x,t) = BP \sin \frac{n\pi x}{L} \cdot \cos \frac{cn\pi t}{L}; \quad n=1,2,3,\dots$$

Using the principle of superposition, we have

$$y(x,t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{L} \cdot \cos \frac{cn\pi t}{L}$$

which gives;

$$y(x,0) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{L}$$

This is half-range sine series, where

$$b_n = \frac{2}{L} \int_0^L y(x,0) \sin \frac{n\pi x}{L} dx$$

$$= \frac{2}{L} \int_0^b \frac{E}{b} x \sin \frac{n\pi x}{L} dx + \frac{2}{L} \int_b^L \frac{E}{b-L} (x-L) \sin \frac{n\pi x}{L} dx$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\begin{aligned}
 &= \frac{2\varepsilon}{Lb} \left[-\frac{\cos(n\pi x/L)}{n\pi/L} \cdot x \right]_0^b - \frac{2\varepsilon}{Lb} \left[-\frac{\sin(n\pi x/L)x}{n^2\pi^2/L^2} \right]_0^L \\
 &\quad + \frac{2\varepsilon}{L(b-L)} \left[-\frac{\cos(n\pi x/L)}{n\pi/L} (x-L) \right]_0^L - \frac{2\varepsilon}{L(b-L)} \left[-\frac{\sin(n\pi x/L)}{n^2\pi^2/L^2} \right]_0^L
 \end{aligned}$$

$$b_n = \frac{2\varepsilon L^2}{n^2\pi^2 b(L-b)} \cdot \sin \frac{n\pi b}{L}$$

Hence, the subsequent motion of the string is given by

$$y(x,t) = \sum_{n=1}^{\infty} \frac{2\varepsilon L^2}{n^2\pi^2 b(L-b)} \cdot \sin \frac{n\pi b}{L} \cdot \sin \frac{n\pi x}{L} \cdot \cos \frac{cn\pi t}{L}$$

Which is required solution

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 7(a)} find Lagrange's interpolation polynomial fitting the points $y(1) = -3$, $y(3) = 0$, $y(4) = 30$, $y(6) = 132$. Find $y(5)$.

Solution:-

From given dat.

$$\begin{array}{ll} y(1) = -3 & y(3) = 0 \\ y(4) = 30 & y(6) = 132 \end{array}$$

i.e; $x_0 = 1$, $x_1 = 3$, $x_2 = 4$, $x_3 = 6$ & $x = 5$

$$f(x_0) = -3 ; f(x_1) = 0 ; f(x_2) = 30 ; f(x_3) = 132$$

By Lagrange's interpolation formula:-

$$\begin{aligned} f(x) &= \frac{(x-x_1)(x-x_2)(x-x_3)}{(x_0-x_1)(x_0-x_2)(x_0-x_3)} \cdot f(x_0) + \\ &\quad \frac{(x-x_0)(x-x_2)(x-x_3)}{(x_1-x_0)(x_1-x_2)(x_1-x_3)} f(x_1) + \frac{(x-x_0)(x-x_1)(x-x_3)}{(x_2-x_0)(x_2-x_1)(x_2-x_3)} f(x_2) \\ &\quad + \frac{(x-x_0)(x-x_1)(x-x_2)}{(x_3-x_0)(x_3-x_1)(x_3-x_2)} \cdot f(x_3) \end{aligned}$$

$$\begin{aligned} f(5) &= \frac{(5-3)(5-4)(5-6)}{(1-3)(1-4)(1-6)} \cdot (-3) + \frac{(5-1)(5-4)(5-6)}{(3-1)(3-4)(3-6)} (0) \\ &\quad + \frac{(5-1)(5-3)(5-6)}{(4-1)(4-3)(4-6)} \cdot (30) + \frac{(5-1)(5-3)(5-4)}{(6-1)(6-3)(6-4)} \cdot (132) \end{aligned}$$

$$\begin{aligned} f(5) &= \frac{2 \times 1 \times 1}{2 \times 3 \times 5} \times -3 + \frac{2 \times 1 \times 1}{2 \times 1 \times 3} \times 0 \\ &\quad + \frac{4 \times 2 \times 1}{3 \times 1 \times 2} \times 30 + \frac{4 \times 2 \times 1}{5 \times 3 \times 2} \times 132 \end{aligned}$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$f(5) = -\frac{1}{5}x_1 + \left(-\frac{2}{3}\right)x_0 + 4x_{10} + \frac{4}{15}x_{13} 2$$

$$f(5) = -\frac{1}{5} + 0 + 40 + \frac{528}{15}$$

$$f(5) = -\frac{3}{15} + \frac{600}{15} + \frac{528}{15}$$

$$f(5) = \frac{600 + 528 - 3}{15} = \frac{1125}{15}$$

$$\boxed{y(5) = 75}$$

which is required result.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 7(b)} A reservoir discharging water through sluices at a depth 'h' below the water surface has a surface area 'A' for various values of 'h' as given below:

$h(\text{ft.})$	10	11	12	13	14
$A(\text{sq.ft})$	950	1070	1200	1350	1530

If t denotes time in minutes, the rate of fall of the surface is given by $\frac{dh}{dt} = -48\sqrt{h}/A$.

Estimate the time taken for the water level to fall from 14 to 10 ft. above the sluices.

Solution:-

Given; The rate of fall of the surface is given by

$$\frac{dh}{dt} = -48\sqrt{h}/A$$

$$\Rightarrow dt = -\frac{1}{48} \frac{A}{\sqrt{h}} dh$$

$$\Rightarrow t = -\frac{1}{48} \int_{14}^{10} \frac{A}{\sqrt{h}} dh.$$

$$t = \frac{1}{48} \int_{10}^{14} \frac{A}{\sqrt{h}} dh.$$

Given table

height(h)(ft)	10	11	12	13	14
Area(A)- Sq.ft	950	1070	1200	1350	1530

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

which improves into given table

h	10	11	12	13	14
\sqrt{h}	3.1623	3.3166	3.464	3.6056	3.7417
A	950	1070	1200	1350	1530
$\frac{A}{48\sqrt{h}}$	6.2587 (y ₀)	6.7212 (y ₁)	7.2169 (y ₂)	7.8005 (y ₃)	8.519 (y ₄)

Using Simpson's $\frac{1}{3}$ rd rule

$$t = \frac{h}{3} [(y_0 + y_4) + 4(y_1 + y_3) + 2y_2]$$

$$\begin{aligned} t &= \frac{1}{3} [6.2587 + 8.519) + 4(6.7212 + 7.8005) \\ &\quad + 2(7.2169)] \end{aligned}$$

$$t = \frac{1}{3} [87.319]$$

$$t = 29.1 \approx 29 \text{ minutes.}$$

\therefore Time required = 29 minutes.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 7(c) Use the classical fourth-order Runge-Kutta method with $h=0.2$ to calculate a solution at $x=0.4$ for initial value problem $\frac{dy}{dx} = 4-x^2+4$
 $y(0)=0$, on the interval $[0, 0.4]$.

Solution:-

given; $y(0)=0$, $h=0.2$

$$\frac{dy}{dx} = f(x, y) = 4-x^2+4$$

on interval $[0, 0.4]$.

$$y[0.4] = ? \quad x=0$$

as $h=0.2$, we need two iterations.

$$\therefore y(0.2) = y(0) + K$$

$$K = \frac{1}{6} [K_1 + K_4 + 2(K_2 + K_3)]$$

$$\text{Where } K_1 = h f(x_0, y_0) = 0.2 f(0, 0) = 0.2 \times 4 \\ = 0.8$$

$$K_2 = h f\left(x_0 + \frac{1}{2}h, y_0 + \frac{1}{2}K_1\right) = 0.2 f(0.1, 0.4) \\ = 0.8782$$

$$K_3 = h f\left(x_0 + \frac{1}{2}h, y_0 + \frac{1}{2}K_2\right) = 0.2 f(0.1, 0.439) \\ = 0.8858$$

$$K_4 = h f(x_0 + h, y_0 + K_3) = 0.2 f(0.2, 0.8858) \\ = 0.96916$$

$$\therefore K = \frac{1}{6} [0.96916 + 0.8 + 2(0.8782 + 0.8858)]$$

$$K = 0.8827933 \approx 0.8828$$

$$y(0.2) = y(0) + 4K = 0 + 0.8828 = 0.8828.$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Now, for $u(0.4) = ?$

$$u(0.2) = 0.8828 \quad , \quad x = 0.2 \quad , \quad h = 0.2$$

$$K_1 = h f(0.2, 0.8828) = 0.2 (4 - 0.04 + 0.8828) \\ = 0.96856$$

$$K_2 = h f(0.3, 1.36708) = 0.2 (5.36708 - 0.09) \\ = 1.055416$$

$$K_3 = h f(0.3, 1.410508) = 0.2 (5.410508 - 0.09) \\ = 1.0641016$$

$$K_4 = h f(0.4, 1.9469) = 0.2 (5.9469 - 0.16) \\ = 1.15738$$

$$K = \frac{1}{6} [K_1 + K_4 + 2(K_2 + K_3)]$$

$$K = \frac{1}{6} [0.96856 + 1.15738 + 2(1.055416 \\ + 1.0641016)]$$

$$K = \frac{1}{6} (6.3649352)$$

$$K = 1.060823 \approx 1.0608$$

$$\therefore u(0.4) = u(0.2) + K$$

$$u(0.4) = 0.8828 + 1.0608$$

$$u(0.4) = 1.9436$$

Which is required solution.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna


Ques: 7(d)(i) Draw the circuit diagram for $\bar{F} = A\bar{B}C + \bar{C}B$ using NAND to NAND logic long.

Solution:

$$\bar{F} = A\bar{B}C + \bar{C}B$$

$$\bar{F} = \overline{\overline{A}\overline{B}C} + \overline{\overline{C}B}$$

$$\boxed{\bar{F} = \overline{\overline{A}\overline{B}C \cdot \overline{B}\overline{C}}}$$



(iii) Design a logic circuit having three inputs A, B, C such that output is 1 when $A=0$ or whenever $B=C=1$. Also obtain logic circuit using only NAND gates?

Solution

A	B	C	output = y	m_i (output).
0	0	0	1	$m_0 = \bar{A}\bar{B}\bar{C}$
0	0	1	1	$m_1 = \bar{A}\bar{B}C$
0	1	0	1	$m_2 = \bar{A}B\bar{C}$
0	1	1	1	$m_3 = \bar{A}BC$
1	0	0	0	$m_4 = A\bar{B}\bar{C}$
1	0	1	0	$m_5 = A\bar{B}C$
1	1	0	0	$m_6 = AB\bar{C}$
1	1	1	1	$m_7 = ABC$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$y = 1 \quad (\text{output})$$

$$y = m_0 + m_1 + m_2 + m_3 + m_7$$

$$y = \bar{A}\bar{B}\bar{C} + \bar{A}\bar{B}C + \bar{A}B\bar{C} + \bar{A}BC + ABC$$


$$y = \bar{A}[\bar{B}\bar{C} + \bar{B}C + B\bar{C} + BC] + ABC$$

$$y = \bar{A}[\bar{B}(\bar{C}+C) + B(\bar{C}+C)] + ABC$$

$$[\because \bar{C}+C=1]$$

$$y = \bar{A}[\bar{B}+B] + ABC \quad [\because \bar{B}+B=1]$$

$y = \bar{A} + ABC$



Ques: 7(d) ii) In a Boolean Algebra B, for any a and b prove that $ab' + a'b = 0$ if and only if $a=b$.

Solution: Given, $ab' + a'b = 0$

$$\Leftrightarrow ab' = (a'b)' \quad [-(a'b) = (a'b)']$$

$$ab' = a'' + b'$$

$$ab' = a + b'$$

$$ab' - (b')' = a$$

$$b'(a-1) = a$$

$$-(b')' = a$$

$$(b')' = a \Rightarrow \boxed{b=a}$$

Hence proved.


Ques: 8(a) > (i) Find the M.I. of a right solid cone of mass M , height h and radius of whose base is a , about its axis.

Solution:- Let, 'O' be the vertex of the right solid cone of mass M , height h and radius of whose base is a . If α is the semi-vertical angle and ρ the density of the cone, then

$$M = \frac{1}{3} \pi \rho h^3 \tan^2 \alpha \quad \text{--- (1)}$$

Consider an elementary disc

PQ of thickness δx , parallel to the base AB and at a distance 'x' from the vertex 'O'.



\therefore Mass of the disc,

$$\delta m = \rho \pi x^2 \tan^2 \alpha \delta x$$

M.I. of this elementary disc about axis OD.

$$= \frac{1}{2} \delta m CP^2 = \frac{1}{2} (\rho \pi x^2 \tan^2 \alpha \delta x) \cdot x^2 \tan^2 \alpha$$

$$= \frac{1}{2} \rho \pi x^4 \tan^4 \alpha \delta x$$

\therefore M.I. of the cone about the axis OD.

$$= \int_0^h \frac{1}{2} \rho \pi x^4 \tan^4 \alpha dx = \frac{\rho \pi}{10} h^5 \tan^4 \alpha$$

$$= \frac{3}{10} M h^2 \tan^2 \alpha \quad (\text{from (1)}).$$

$$\therefore \text{M.I. of cone about the axis OD} = \frac{3}{10} M a^2 \quad \left[\because \tan \alpha = \frac{a}{h} \right]$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques:- 8(a) ii) write Hamilton's equations for a particle of mass 'm' moving in a plane under a force which is some function of distance from the origin?

Solution:-

Let, $P(r, \theta)$ be the co-ordinates of a particle of mass 'm' at time 't' referred to the pole at O and OX as initial line. The components of velocity at P along OP and perpendicular to it are \dot{r} and $r\dot{\theta}$ respectively.

$$\therefore (\text{Vel})^2 \text{ of mass } m \text{ at } P \Rightarrow V^2 = \dot{r}^2 + (r\dot{\theta})^2$$

$$\therefore K.E, T = \frac{1}{2} m V^2 = \frac{1}{2} m (\dot{r}^2 + (r\dot{\theta})^2)$$

Since, the force at P is some function of \dot{r} , therefore the potential V is function of r alone and will be independent of θ . i.e $V = V(r)$,

$$\text{Thus, } L = T - V = \frac{1}{2} m (\dot{r}^2 + (r\dot{\theta})^2) - V(r)$$

Here ' r ' and ' θ ' are the generalised co-ordinates,

$$\begin{aligned} \therefore P_r &= \frac{\partial L}{\partial \dot{r}} = m\dot{r} \\ P_\theta &= \frac{\partial L}{\partial \dot{\theta}} = mr^2\dot{\theta} \end{aligned} \quad] \quad \text{--- (1)}$$

Since, L does not contain ' t ' explicitly

$$\therefore H = T + V = \frac{1}{2} m (\dot{r}^2 + (r\dot{\theta})^2) + V(r)$$

$$H = T + V = \frac{1}{2} m [(P_r/m)^2 + r^2(P_\theta/mr^2)^2] + V(r) \rightarrow \text{from (1)}$$

(33)

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\therefore H = \frac{1}{2m} \left(p_x^2 + p_\theta^2 / r^2 \right) + V(r)$$

Hence, four Hamilton's equations are

$$\dot{p}_x = -\frac{\partial H}{\partial x} = \frac{p_x^2}{mr^3} - \frac{dV}{dr} \quad \text{--- } H_1$$

$$\dot{x} = -\frac{\partial H}{\partial p_x} = -\frac{p_x}{m} \quad \text{--- } H_2$$

$$\dot{p}_\theta = -\frac{\partial H}{\partial \theta} = 0 \quad \text{--- } H_3$$

$$\dot{\theta} = \frac{\partial H}{\partial p_\theta} = \frac{p_\theta}{mr^2} \quad \text{--- } H_4.$$

Which is required results!

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

Ques: 8(b)} A particle of mass 'm' moves in a conservative forces field. find -
 (i) the Lagrangian function and
 (ii) the equation of motion in cylindrical co-ordinates (ρ, ϕ, z) .

Solution:-

Let, P be the position of the particle of mass 'm' whose cylindrical coordinates referred to axes OX, OY, OZ are (ρ, ϕ, z) .


\therefore If (x, y, z) are its cartesian co-ordinates, then

$$x = OA = \rho \cos \phi$$

$$y = OB = \rho \sin \phi, \quad z = z$$

If i, j, k are the unit vectors along OX, OY, OZ respectively, then

$$\overrightarrow{OP} = r = \rho \cos \phi \hat{i} + \rho \sin \phi \hat{j} + z \hat{k}$$



(34)

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

If \hat{r}_1 and $\hat{\phi}_1$ are the unit vectors in the directions of ρ and ϕ increasing respectively, then

$$\hat{r}_1 = \frac{\partial \mathbf{r}}{\partial \rho} / \left| \frac{\partial \mathbf{r}}{\partial \rho} \right| = \frac{\cos \phi \hat{i} + \sin \phi \hat{j}}{\sqrt{\cos^2 \phi + \sin^2 \phi}}$$

$$\boxed{\hat{r}_1 = \cos \phi \hat{i} + \sin \phi \hat{j}}$$

$$\hat{\phi}_1 = \frac{\partial \mathbf{r}}{\partial \phi} / \left| \frac{\partial \mathbf{r}}{\partial \phi} \right| = \frac{-\rho \sin \phi \hat{i} + \rho \cos \phi \hat{j}}{\sqrt{(\rho^2 \sin^2 \phi + \rho^2 \cos^2 \phi)}}$$

$$\hat{\phi}_1 = \frac{-\rho \sin \phi \hat{i} + \rho \cos \phi \hat{j}}{\rho \sqrt{\sin^2 \phi + \cos^2 \phi}} = -\sin \phi \hat{i} + \cos \phi \hat{j}$$

Now; $\mathbf{v} = \dot{\mathbf{r}} = (\dot{\rho} \cos \phi - \rho \sin \phi \dot{\phi}) \hat{i} +$
 $(\dot{\rho} \sin \phi + \rho \cos \phi \dot{\phi}) \hat{j} + \dot{z} \hat{k}$

$$\mathbf{v} = \dot{\mathbf{r}} = \dot{\rho} (\cos \phi \hat{i} + \sin \phi \hat{j}) + \rho \dot{\phi} (-\sin \phi \hat{i} + \cos \phi \hat{j}) + \dot{z} \hat{k}$$

$$\mathbf{v} = \dot{\mathbf{r}} = (\dot{\rho}) \hat{r}_1 + \rho \dot{\phi} (\hat{\phi}_1) + \dot{z} \hat{k}$$

$$\therefore v^2 = \dot{\rho}^2 + (\rho \dot{\phi})^2 + \dot{z}^2$$

Total K.E, $\Rightarrow T = \frac{1}{2} m v^2 = \frac{1}{2} m (\dot{\rho}^2 + \rho^2 \dot{\phi}^2 + \dot{z}^2)$

Let; $V = V(\rho, \phi, z)$ be the potential function.

\therefore (i) Lagrangian function; $L = T - V$

i.e. $\boxed{L = \frac{1}{2} m (\dot{\rho}^2 + \rho^2 \dot{\phi}^2 + \dot{z}^2) - V(\rho, \phi, z)}$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

(ii) Lagrange's ρ equation is ,

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\rho}} \right) - \frac{\partial L}{\partial \rho} = 0$$

$$\text{or } \frac{d}{dt} (m\dot{\rho}) - \left(m\rho \dot{\phi}^2 - \frac{\partial V}{\partial \rho} \right) = 0$$

$$\text{i.e. } m\ddot{\rho} + (-m\rho \dot{\phi}^2) = -\frac{\partial V}{\partial \rho}$$

Lagrange's ϕ equation is $\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\phi}} \right) - \frac{\partial L}{\partial \phi} = 0$

$$\text{or } \frac{d}{dt} (m\rho^2 \dot{\phi}) - \left(-\frac{\partial V}{\partial \phi} \right) = 0$$

$$\text{or } \frac{d}{dt} (m\rho^2 \dot{\phi}) = -\frac{\partial V}{\partial \phi}$$

and Lagrange's z equation is

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{z}} \right) - \frac{\partial L}{\partial z} = 0$$

$$\text{or } \frac{d}{dt} (m\dot{z}) - \left(-\frac{\partial V}{\partial z} \right) = 0$$

$m\ddot{z} = -\frac{\partial V}{\partial z}$

which is required result.

Ques:- 8(c) } Prove that liquid motion is possible when velocity at (x, y, z) is given by

$$u = \frac{3x^2 - r^2}{r^5}, v = \frac{3xy}{r^5}, w = \frac{3xz}{r^5}$$

where, $r^2 = x^2 + y^2 + z^2$ are the stream lines are the intersection of the surfaces, $(x^2 + y^2 + z^2)^3 = C(y^2 + z^2)^2$ by the planes passing through Ox. Is this motion irrotational.

Solution :-

Step-I } To prove that the liquid motion is possible for this we have to show that the equation of continuity namely -

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = 0 \quad \dots \quad (1)$$

is satisfied.

$$r^2 = x^2 + y^2 + z^2$$

$$\Rightarrow \frac{\partial r}{\partial x} = \frac{x}{r}, \quad \frac{\partial r}{\partial y} = \frac{y}{r}, \quad \frac{\partial r}{\partial z} = \frac{z}{r}$$

$$\Rightarrow \frac{\partial u}{\partial x} = \frac{(6x - 2x)r^5 - 5r^3x(3x^2 - r^2)}{r^{10}}$$

$$\frac{\partial v}{\partial y} = \frac{3x}{r^{10}}(r^5 - 5r^3y^2)$$

$$\frac{\partial w}{\partial z} = \frac{3x}{r^{10}}(r^5 - 5r^3z^2)$$

$$\text{This } \Rightarrow \frac{\partial u}{\partial x} = \frac{3x}{r^7}(3r^2 - 5x^2)$$

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

$$\frac{\partial v}{\partial y} = \frac{3x}{r^7} (x^2 - 5y^2)$$

$$\frac{\partial w}{\partial z} = \frac{3x}{r^7} (x^2 - 5z^2)$$

$$\Rightarrow \boxed{\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = 0}$$

Hence, the result

Step-II) To determine stream lines

Stream lines are the solutions of $\frac{dx}{u} = \frac{dy}{v} = \frac{dz}{w}$

Putting the values;

$$\begin{aligned} \frac{dx}{3x^2 - r^2} &= \frac{dy}{3xy} = \frac{dz}{3xz} \\ &= \frac{x dx + y dy + z dz}{x(3x^2 - r^2)} = \frac{y dy + z dz}{3x(y^2 + z^2)}. \end{aligned}$$

$$\Rightarrow \frac{dx}{3x^2 - r^2} = \frac{dy}{3xy} = \frac{dz}{3xz} = \frac{x dx + y dy + z dz}{x \cdot 2r^2} = \frac{y dy + z dz}{3x(y^2 + z^2)}$$

$$\Rightarrow \frac{dy}{3xy} = \frac{dz}{3xz}$$

$$\Rightarrow \frac{dy}{y} = \frac{dz}{z} \Rightarrow \log y = \log z + \log C$$

$$\Rightarrow \boxed{y = zC} \quad — \textcircled{2}$$

and $\frac{x dx + y dy + z dz}{x \cdot 2(x^2 + y^2 + z^2)} = \frac{y dz + z dy}{3x(y^2 + z^2)} \quad — \textcircled{3}$

Integrating $\textcircled{3}$, we get.

INSTITUTE FOR IAS/IFoS/CSIR/GATE EXAMINATIONS
MATHEMATICS by K. Venkanna

(36)

$$\Rightarrow \frac{1}{2} \log(x^2 + y^2 + z^2) = \frac{1}{3} \log(y^2 + z^2) + \frac{1}{6} \log b$$

$$\Rightarrow (x^2 + y^2 + z^2)^{\frac{3}{2}} = b(y^2 + z^2)^2$$

which is required result.

INSTITUTE OF MATHEMATICAL SCIENCES