

Q.1(c) Evaluate. $I = \int_0^1 \sqrt[3]{x \cdot \log \frac{1}{x}} dx$.

Sol.

Sol.

let us substitute,

$$\log\left(\frac{1}{x}\right) = t$$

$$\Rightarrow \frac{1}{1/x} \left(\frac{-1}{x^2} \right) dx = dt.$$

$$\Rightarrow \frac{-1}{x} \cdot dx = dt \Rightarrow dx = -e^{-t} dt$$

$$I = \int_0^1 x^{1/3} \cdot [\log(\frac{1}{x})]^{1/3} dx.$$

$$= \int_{\infty}^0 -e^{-t/3} \cdot t^{1/3} \cdot e^{-t} dt.$$

$$I = \int_0^{\infty} e^{-4t/3} \cdot t^{1/3} \cdot dt$$

$$\text{let, } \frac{4t}{3} = p \Rightarrow \frac{4}{3} dt = dp$$

$$= \int_0^{\infty} e^{-p} \left(\frac{3p}{4} \right)^{1/3} \cdot \frac{3}{4} dp.$$

$$= \left(\frac{3}{4} \right)^{4/3} \int_0^{\infty} e^{-p} \cdot p^{1/3} dp = \left(\frac{3}{4} \right)^{4/3} \int_0^{\infty} e^{-p} \cdot p^{\frac{4}{3}-1} dp.$$

$$\text{we know, } \int_0^{\infty} t^{x-1} \cdot e^{-t} dt = \Gamma x$$

$$\text{so, } I = \left(\frac{3}{4} \right)^{4/3} \Gamma \frac{4}{3}.$$

3-a) Find the maximum and minimum values of $x^2+y^2+z^2$ subject to the condition $\frac{x^2}{4} + \frac{y^2}{25} + \frac{z^2}{25} = 1$, & $x+y-z=0$.

* Let's consider a function

$$f = x^2+y^2+z^2 + \lambda \left(\frac{x^2}{4} + \frac{y^2}{25} + \frac{z^2}{25} - 1 \right) + \mu (x+y-z)$$

20 Marks

∴ The stationary points are found out by $df=0$

$$\Rightarrow f_x dx + f_y dy + f_z dz = 0$$

$$\Rightarrow \underline{f_x = 0}, \underline{f_y = 0}, \underline{f_z = 0}$$

$$\begin{aligned} 2x + \frac{2x\lambda}{4} + \mu &= 0 \\ 2y + \frac{2y\lambda}{25} + \mu &= 0 \\ 2z + \frac{2z\lambda}{25} - \mu &= 0 \end{aligned} \quad \left| \quad \begin{aligned} 2x^2 + \frac{2x^2\lambda}{4} + \mu x &= 0 \\ 2y^2 + \frac{2y^2\lambda}{25} + \mu y &= 0 \\ 2z^2 + \frac{2z^2\lambda}{25} - \mu z &= 0 \end{aligned} \right.$$

adding we have

$$2(x^2+y^2+z^2) + 2\lambda \left(\frac{x^2}{4} + \frac{y^2}{25} + \frac{z^2}{25} \right) + \mu(x+y-z) = 0$$

$$\Rightarrow 2x^2 + 2\lambda \times 1 + \mu \times 0 = 0$$

$$\Rightarrow \boxed{\lambda = -x^2}$$

$$\Rightarrow \boxed{x^2+y^2+z^2 = x}$$

$$\begin{aligned} 2x \left(1 + \frac{\lambda}{4} \right) &= -\mu \\ 2y \left(1 + \frac{\lambda}{25} \right) &= -\mu \\ 2z \left(1 + \frac{\lambda}{25} \right) &= \mu \end{aligned}$$

$$2(x-y) + 2\lambda \left(\frac{x}{4} - \frac{y}{25} \right) = 0$$

$$\Rightarrow \lambda = \frac{x-y}{\frac{y}{25} - \frac{x}{4}}$$

$$2x = \frac{-\mu}{1 + \frac{\lambda}{4}}, \quad 2y = \frac{-\mu}{1 + \frac{\lambda}{25}}, \quad 2z = \frac{\mu}{1 + \frac{\lambda}{25}}$$

$$\Rightarrow x+y-z=0 \Rightarrow \frac{-\mu}{2 \left(1 + \frac{\lambda}{4} \right)} - \frac{\mu}{1 + \frac{\lambda}{25}} - \frac{\mu}{1 + \frac{\lambda}{25}} = 0 \Rightarrow \mu$$

$$\frac{4}{1+\lambda} + \frac{5}{5+\lambda} + \frac{25}{25+\lambda} = 0 \Rightarrow 4(25+\lambda)(5+\lambda) + 5(4+\lambda)(25+\lambda) + 25(4+\lambda)(5+\lambda) = 0$$

$$\Rightarrow 4(25 + 30\lambda + \lambda^2) + 5(100 + 89\lambda + \lambda^2) + 25(20 + \lambda^2 + 9\lambda) = 0$$

$$\Rightarrow 34\lambda^2 + 490\lambda + 1500 = 0$$

$$\Rightarrow \lambda = \frac{-490 \pm \sqrt{490^2 - 4 \times 34 \times 1500}}{2 \times 34} = \frac{-490 \pm \sqrt{36100}}{68}$$

$$= \frac{-490 \pm 190}{68} = -10 \text{ or } -\frac{75}{17}$$

Now putting this value on the

$$\frac{4}{1+\frac{\lambda}{4}} = -2x \quad \frac{4}{1+\frac{\lambda}{5}} = -2y, \quad \frac{4}{1+\frac{\lambda}{25}} = +2z$$

for $\lambda = -10$
 ~~$4 = -2x$~~

$$x = \frac{-4}{2(1+\frac{\lambda}{4})} = \frac{-4}{2(1-\frac{10}{4})} = \frac{-4 \times 2}{-8} = \frac{4}{3}$$

$$y = \frac{-1}{2} \frac{4}{1-\frac{10}{5}} = -\frac{1}{2} \times \frac{4 \times 5}{-5} = \frac{4}{2}$$

$$z = +\frac{1}{2} \frac{4}{1-\frac{10}{25}} = \frac{4}{2} \times \frac{25 \times 5}{15 \times 3} = \frac{54}{6}$$

$$\frac{x^2}{4} + \frac{y^2}{5} + \frac{z^2}{25} = 1$$

$$\Rightarrow \frac{u^2}{9 \times 4} + \frac{u^2}{4 \times 5} + \frac{25u^2}{36 \times 25} = 1$$

$$\Rightarrow u^2 = \frac{180}{19} \Rightarrow u = \pm \sqrt{\frac{180}{19}}$$

$$\begin{aligned} &1.05263 \times 2.3684 \\ &= 2.489 \\ &< 10 \end{aligned}$$

The 4 stationary points are $\left(\pm \frac{1}{3} \sqrt{\frac{180}{19}}, \pm \frac{1}{2} \sqrt{\frac{180}{19}}, \pm \frac{5}{6} \sqrt{\frac{180}{19}} \right)$

Similarly for $\lambda = -\frac{35}{17}$

$$x = \frac{-4}{2\left(1 + \frac{\lambda}{4}\right)} = \frac{-4}{2\left(1 - \frac{35}{68}\right)} = \frac{-4}{2\left(\frac{33}{68}\right)} = \frac{324}{11}$$

$$y = \frac{-4}{2\left(1 + \frac{\lambda}{5}\right)} = \frac{-4}{2\left(1 - \frac{35}{5 \times 17}\right)} = \frac{-4 \times 25 \times 17}{2 \times 102}$$

$$\boxed{y = \frac{-417}{4}}$$

$$z = \frac{4}{2\left(1 + \frac{\lambda}{25}\right)} = \frac{4}{2\left(1 - \frac{35}{17 \times 25}\right)} = \frac{4 \times 17}{2 \times 14}$$

$$\therefore \frac{x^2}{4} + \frac{y^2}{5} + \frac{z^2}{25} = 1$$

$$\Rightarrow \frac{324^2}{11^2 \times 4} + \frac{17^2 \times 4^2}{4^2 \times 5} + \frac{17^2 \times 4^2}{25^2 \times 25} = 1$$

$$\Rightarrow \mu = 0.5203$$

$$\therefore x = \frac{32}{11} \times 0.52, \quad y = \frac{-17}{4} \times 0.5203, \quad z = \frac{17}{25} \times 0.52$$

$$\therefore x^2 + y^2 + z^2 = 1.5136^2 + 2.211235^2 + 0.3157^2$$

$$\text{on } \lambda = -\frac{35}{17} = \underline{\underline{7.28}}$$

$$\text{if } \underline{\underline{\lambda = -10}}, \quad x^2 + y^2 + z^2 = \underline{\underline{10}}$$

So the max^m value of the $x^2 + y^2 + z^2 = 10$ if min^m = $\underline{\underline{\frac{35}{17}}}$

3-5) Let $f(x,y) = \begin{cases} \frac{2x^4y - 5x^2y^2 + y^5}{(x^2+y^2)^2}, & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0) \end{cases}$

Find a $\delta > 0$ such that $|f(x,y) - f(0,0)| < 0.01$ whenever $\sqrt{x^2+y^2} < \delta$. 15 Marks

Ans

$$f(x,y) = \frac{2x^4y - 5x^2y^2 + y^5}{(x^2+y^2)^2}$$

$$|f(x,y) - f(0,0)| = \left| \frac{2x^4y - 5x^2y^2 + y^5}{(x^2+y^2)^2} \right|$$

$$= \left| \frac{y(2x^4 + y^4) - 5x^2y^2}{(x^2+y^2)^2} \right| \leq \left| \frac{y(2x^4 + y^4)}{(x^2+y^2)^2} \right|$$

$$= \left| \frac{y(\sqrt{2}x^2 + y^2)^2 - 2\sqrt{2}x^2y^2 - 5x^2y^2}{(x^2+y^2)^2} \right|$$

$$= \left| \frac{y(\sqrt{2}x^2 + y^2)^2 - 2\sqrt{2}x^2y^2 - 5x^2y^2}{(x^2+y^2)^2} \right|$$

$$\leq \frac{y(\sqrt{2}x^2 + y^2)^2}{(x^2+y^2)^2} \leq \frac{y(x^2+y^2)^2}{(x^2+y^2)^2} \leq y$$

if $y < 0$

$$\leq \left| \frac{y(2x^4 + y^4)}{(x^2+y^2)^2} \right| = \left| \frac{y(\sqrt{2}x^2 + y^2)^2 - 2\sqrt{2}x^2y^2 - 5x^2y^2}{(x^2+y^2)^2} \right|$$

if $y > 0$

$$\leq \left| \frac{y(\sqrt{2}x^2 + y^2)^2 - 2\sqrt{2}x^2y^2 - 5x^2y^2}{(x^2+y^2)^2} \right|$$

$$|f(x,y)| = \frac{|2x^4y + y^5 - 5x^2y^2|}{(x^2+y^2)^2}$$

Since the power is $\left(\frac{5}{2}\right)$

CONCEPT

$$= (x^2+y^2)^{5/2} = \frac{y^5 \left(1 + \frac{2x^4}{y^4} - \frac{5x^2}{y^3}\right)}{(x^2+y^2)^2}$$

$$\leq \frac{y^5 \left(1 + \frac{2x^4}{y^4}\right)}{(x^2+y^2)^2} \leq \frac{y^5 \left(1 + \frac{5}{2} \frac{x^2}{y^2} + \frac{25}{4} \frac{x^4}{y^4}\right)}{(x^2+y^2)^2}$$

$$\leq \frac{y^5 \left(1 + \frac{x^2}{y^2}\right)^{5/2}}{(x^2+y^2)^2} \Rightarrow \sqrt{(x^2+y^2)} < \epsilon = 0.01 = \underline{\underline{\delta}}$$

3(c)

20/6

Ans.

Find the surface area of the plane $x + 2y + 2z = 12$ cut off by $x = 0$, $y = 0$ and $x^2 + y^2 = 16$.

Plane $x + 2y + 2z = 12$

$$\text{or } \frac{x}{12} + \frac{y}{6} + \frac{z}{6} = 1$$

cuts the coordinates at a distance of 12, 6 and 6 from origin.

Cylinder: $x^2 + y^2 = 16$

Planes: $x = 0$, $y = 0$


$$\text{Surface Area} = \iint_A \sqrt{1 + z_x^2 + z_y^2} \, dx \, dy$$

$$= \iint_A \sqrt{1 + \left(-\frac{1}{2}\right)^2 + (-1)^2} \, dx \, dy$$

$$= \frac{3}{2} \iint_A dx \, dy$$

$$= \frac{3}{2} \cdot \left[\frac{1}{4} \pi (4)^2 \right]$$

$$= 6\pi$$



$$\therefore z = -\frac{x}{2} - y + 6$$

$$z_x = -\frac{1}{2}, z_y = -1$$

(A: Projection of surface on xy-plane)

$$x^2 + y^2 \leq 16$$


$$x \geq 0, y \geq 0$$

4f) Evaluate $\iint_R f(x,y) \, dx \, dy$ over the rectangle $R = [0,1; 0,1]$

where

$$f(x,y) = \begin{cases} x+y, & x^2 < y < 2x^2 \\ 0, & \text{elsewhere.} \end{cases}$$

Ans



Area of the region:

$$\begin{aligned} \therefore \iint_R f(x,y) \, dxdy &= \int_{x=0}^{2x^2} \int_{y=x^2}^{2x^2} (x+y) \, dy \, dx = \int_0^1 \left[yx + \frac{y^2}{2} \right]_{y=x^2}^{2x^2} dx \\ &= \int_0^1 \left[x(x^2) + \frac{1}{2}(4x^4 - x^4) \right] dx \\ &= \frac{x^4}{4} + \frac{3}{2} \frac{x^5}{5} \bigg|_0^1 = \frac{1}{4} + \frac{3}{10} = \frac{5+6}{20} = \frac{11}{20} \text{ (Ans)} \end{aligned}$$

2017:

1 Feb - 2016 :

1-(a) ~~prove that all bijective function from a non-empty set X onto itself is a~~

1-(b) Show that $\frac{x}{1+x} < \log(1+x) < x \quad \forall x > 0$.

Ans Let's consider $f(x) = \frac{x}{1+x} - \log(1+x)$

$$\Rightarrow f(x) = \frac{(1+x) - x}{(1+x)^2} - \frac{1}{1+x} = \frac{1}{(1+x)^2} - \frac{1}{(1+x)} = \frac{1-x-x}{(1+x)^2}$$

$\Rightarrow f(x)$ is decreasing function

$$\Rightarrow \underline{f(x) < f(0)} \quad \text{for } \underline{x > 0}$$

$$\Rightarrow \underline{\frac{x}{1+x} - \log(1+x) < 0} \Rightarrow \underline{\frac{x}{1+x} < \log(1+x)} \quad \text{--- (1)}$$

Let $g(x) = x - \log(1+x)$

$$\Rightarrow g'(x) = 1 - \frac{1}{1+x} > 0 \quad \forall \underline{x > 0}$$

$\Rightarrow g(x)$ is increasing function.

$$\Rightarrow g(x) > g(0), \text{ for } x > 0$$

$$\Rightarrow \underline{x - \log(1+x) > 0} \Rightarrow \underline{\log(1+x) < x} \quad \text{--- (2)}$$

from 1 & 2 \Rightarrow $\frac{x}{1+x} < \log(1+x) < x$ Ans.

(c) Examine if the function, $f(x,y) = \frac{xy}{x^2+y^2}$, $(x,y) \neq (0,0)$ and $f(0,0) = 0$ is continuous at $(0,0)$. Find $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ at points other than origin.

$$f(x,y) = \begin{cases} \frac{xy}{x^2+y^2}, & (x,y) \neq (0,0) \\ 0 & (x,y) = (0,0) \end{cases}$$

now Let's take $x = r \cos \theta$, $y = r \sin \theta$.

$$\therefore f(r,\theta) = \frac{r^2 \sin \theta \cos \theta}{r^2} = \frac{1}{2} \sin 2\theta$$

$$\therefore |f(x,y) - f(0,0)| = \left| \frac{\sin 2\theta}{2} \right| \leq \left(\frac{1}{2} \right)$$

50 (A) 50% Chand Bali
30 100% Moon Sand

(20) $\rightarrow 10$ 50:50

Nilgiri P. 0
Special
4 - N.P.
3 \rightarrow WLS

But if we take $\epsilon = \frac{1}{4}$ then $|f(r,\theta) - f(0,0)| > \epsilon$

Hence the function is not continuous.

$$\text{now } \frac{\partial f}{\partial x} = \lim_{h \rightarrow 0} \frac{f(h,0) - f(0,0)}{h}$$

$$= \lim_{h \rightarrow 0} (0) = \underline{\underline{0}}$$

$$\frac{\partial f}{\partial y} = \lim_{k \rightarrow 0} \frac{f(0,k) - f(0,0)}{k} = 0.$$

So the partial derivatives exist & is 0 for points (x,y)

2-(a) After changing the order of integration of

$$\int_0^{\infty} \int_0^{\infty} e^{-xy} \sin(nu) \, du \, dy, \text{ show that } \int_0^{\infty} \frac{\sin nu}{n} \, du = \frac{\pi}{2}.$$

Ans ~~1~~ $I = \int_{y=0}^{\infty} \int_{u=0}^{\infty} e^{-xy} \sin(nu) \, du \, dy,$

$$= \int_0^{\infty} \sin(nu) \left(\frac{1}{x} \right) e^{-xy} \Big|_{y=0}^{\infty} \, du.$$

$$= - \int_0^{\infty} \frac{\sin nu}{n} e^{-xy} \Big|_{y=0}^{\infty} \, du$$

$$= - \int_0^{\infty} \frac{\sin nu}{n} (0 - 1) \, du.$$

$$= \int_0^{\infty} \frac{\sin nu}{n} \, du. \quad \text{--- (1)}$$

Now First integrating w.r.t x :

$$\int_0^{\infty} \left\{ -\frac{1}{y} e^{-xy} \sin(nu) \Big|_{x=0}^{\infty} + \frac{1}{y} e^{-xy} n \cos(nu) \Big|_{x=0}^{\infty} \right\} dy.$$

$$= \int_0^{\infty} \left\{ \frac{n}{y} \left(0 - \frac{1}{y} e^{-xy} \cos(nu) \Big|_{x=0}^{\infty} - \int_0^{\infty} \frac{e^{-xy}}{y} n \sin(nu) \right) \right\} dy$$

$$= \int_0^{\infty} \left\{ \frac{n}{y} \left(0 + \frac{1}{y} - \frac{n}{y} I' \right) \right\} dy.$$

$$\Rightarrow \frac{n}{y^2} - \frac{n^2}{y^2} I' = 0 \Rightarrow \left(1 + \frac{n^2}{y^2} \right) I = \frac{n}{y^2} \Rightarrow I = \frac{n}{n^2 + y^2}$$

$$= \int_0^{\infty} \frac{y}{n^2 + y^2} dy = \frac{1}{n} \tan^{-1}\left(\frac{y}{n}\right) \Big|_0^{\infty} = \frac{\pi}{2} - 0 = \frac{\pi}{2}$$

$$\therefore I = \int_0^{\infty} \frac{\sin nx}{n} dn = \frac{\pi}{2} \quad (*)$$

Q-4) Using MVT, find a point on the curve $y = \sqrt{x-2}$, defined on $[2, 3]$, where the tangent is parallel to the chord joining the end points of the curve.

Ans

$$y = \sqrt{x-2} \Rightarrow y^2 = x-2 \Rightarrow x = y^2 + 2$$

$y = \sqrt{x-2}$ is continuous and differentiable in $[2, 3]$.

Hence by Lagrange's MVT:

$$f'(c) = \frac{f(3) - f(2)}{3-2} \quad \text{where } c \in (2, 3)$$

$$= \frac{1}{1} \Rightarrow \underline{f'(c) = 1}$$

$$f'(x) = \frac{1}{2\sqrt{x-2}} \Rightarrow f'(c) = 1 \Rightarrow \frac{1}{2\sqrt{c-2}} = 1 \Rightarrow \sqrt{c-2} = \frac{1}{2} \Rightarrow c-2 = \frac{1}{4}$$

$$\Rightarrow c = 2 + \frac{1}{4} = \underline{\underline{\frac{9}{4}}} \in (2, 3)$$

Hence at $x = \frac{9}{4}$, the tangent is parallel to the chord joining the end points of the curve. (*)

Q-5) Using Lagrange's Multiplier, find the point on the plane $2x + 3y + 4z = 5$, which is closest to the point $(1, 0, 0)$.

Ans Let $P(x, y, z)$ be any point on the plane $2x + 3y + 4z = 5$.
Such that $f(x, y, z) = (x-1)^2 + y^2 + z^2$ is the minimum.

Now let's consider the function

$$F(x, y, z) = (x-1)^2 + y^2 + z^2 + \lambda(2x + 3y + 4z - 5)$$

$$\therefore dF = 0 \Rightarrow F_x dx + F_y dy + F_z dz = 0$$

$$\Rightarrow F_x = 0 \Rightarrow 2(x-1) + 2\lambda = 0 \Rightarrow \boxed{x = 1-\lambda}$$

$$F_y = 0 \Rightarrow 2y + 3\lambda = 0 \Rightarrow \boxed{y = -\frac{3\lambda}{2}}$$

$$F_z = 0 \Rightarrow 2z + 4\lambda = 0 \Rightarrow \boxed{z = -2\lambda}$$

$$\therefore 2(1-\lambda) + 3\left(-\frac{3\lambda}{2}\right) + 4(-2\lambda) = 5$$

$$\Rightarrow 2 - 2\lambda - \frac{9\lambda}{2} - 8\lambda = 5$$

$$\Rightarrow -10\lambda - \frac{9\lambda}{2} = 3 \Rightarrow \boxed{\lambda = -\frac{6}{29}}$$

$$x = 1 + \frac{6}{29} = \frac{35}{29}$$

$$y = -\frac{3}{2} \times \frac{6}{29} = -\frac{9}{29}$$

$$z = -2 \times \frac{6}{29} = -\frac{12}{29}$$

which is the reqd stationary points

$$d^2F = F_{xx}(dx)^2 + F_{yy}(dy)^2 + F_{zz}(dz)^2 + F_{xy}dx dy + F_{xz}dx dz + F_{yz}dy dz + F_{yx}dy dx + F_{zy}dz dy + F_{zx}dz dx + F_{xy}dy dx$$

70

Hence the function $F(x, y, z)$ is maximum at $\left(\frac{35}{29}, -\frac{9}{29}, -\frac{12}{29}\right)$

which is the required point. (Ans)

3-c) obtain the area between the curve $r = 3(\sec\theta + \csc\theta)$ and its asymptote $x = 3$. **CHECK**

$$x = r \cos\theta = 3(1 + \cos^2\theta)$$

$$y = r \sin\theta = 3(\tan\theta + \sin\theta \csc\theta)$$

$$\Rightarrow \frac{x}{y} = \frac{\csc\theta}{\sin\theta} = \cot\theta$$

$$\Rightarrow \boxed{y = x \tan\theta}$$

$$\frac{x}{3} = 1 + \cos 2\theta$$

$$\frac{y}{3} = \frac{\sin 2\theta}{\cos 2\theta} (1 + \cos 2\theta)$$

$$r = \frac{3(1 + \cos 2\theta)}{\cos 2\theta}$$

$$\theta = 0 \Rightarrow r = 3 \times 2 = 6, \quad \theta = \pi \Rightarrow r = \frac{3 \times 2}{-1} = -6$$


$$\theta = \frac{\pi}{2} \Rightarrow r = \infty$$

$$\theta = \frac{3\pi}{2} \Rightarrow r = \infty$$

$$\theta = \frac{\pi}{4} \Rightarrow r = \frac{3(1 + \frac{1}{\sqrt{2}})}{\frac{1}{\sqrt{2}}} = \frac{9}{2} \times \sqrt{2} = \frac{9\sqrt{2}}{2}$$

$$x = r \cos \theta = 3$$

$$\theta = \frac{\pi}{3} \Rightarrow r = \frac{3(1 + \frac{1}{2})}{\frac{1}{2}} = \frac{15}{1} \times \frac{2}{2} = 7.5$$



$$A = 4 \times \int_{\theta=0}^{\pi/2} \int_{r=3\sec\theta}^{3(\sec\theta+\cos\theta)} r dr d\theta = \int_0^{\pi/2} \frac{r^2}{2} \Big|_{3\sec\theta}^{3(\sec\theta+\cos\theta)} d\theta$$

$$= \frac{1}{2} \times 9 \int_0^{\pi/2} \left\{ (\sec\theta + \cos\theta)^2 - \sec^2\theta \right\} d\theta$$

$$= \frac{9}{2} \int_0^{\pi/2} (\sec^2\theta + \cos^2\theta + 2 - \sec^2\theta) d\theta$$

$$= \frac{9}{2} \int_0^{\pi/2} (2 + \cos^2\theta) d\theta = \frac{9}{2} \times \left(\frac{\pi}{2} + \frac{1}{2} \times \frac{\pi}{2} \right) = \frac{3\pi}{4} \times \frac{9}{2} = \frac{27\pi}{8}$$

asymptote $\Rightarrow \theta = \frac{\pi}{2}$

$$\text{So total area} = 4 \times \frac{27\pi}{8} = \frac{27\pi}{2} \text{ (sq. units)}$$

4-b) Show that the integral $\int_0^\infty e^{-x} x^{\alpha-1} dx$, $\alpha > 0$ exists, by separately taking the cases for $\alpha > 1$, and $0 < \alpha < 1$.

~~For $\alpha > 1$, $\int_0^\infty e^{-x} x^{\alpha-1} dx$ is an improper integral~~
~~Let $f(x) = e^{-x} x^{\alpha-1}$~~

$$\text{Let } I = \int_0^{\infty} e^{-x} x^{\alpha+1} dx = \int_0^a e^{-x} x^{\alpha+1} dx + \int_a^{\infty} e^{-x} x^{\alpha+1} dx.$$

(I₁) (I₂)

for $\alpha > 1$: I₁ is a proper integral.

I₂ is an improper integral.

$$\therefore \int_a^{\infty} e^{-x} x^{\alpha+1} dx \quad \text{let } f(x) = x^{\alpha+1} e^{-x}.$$

Let's take $g(x) = \frac{1}{x^2}$

$$\therefore \lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \infty} \frac{x^{\alpha+1} e^{-x}}{\frac{1}{x^2}} = \lim_{x \rightarrow \infty} x^{\alpha+1} e^{-x} = \lim_{x \rightarrow \infty} \frac{x^{\alpha+1}}{e^x} \left(\frac{\infty}{\infty} \right)$$

$$= \frac{(\alpha+1)x^{\alpha}}{e^x} \left(\frac{\infty}{\infty} \right) \dots = \frac{(\alpha+1)!}{e^x} = 0$$

Hence since $\int_a^{\infty} \frac{1}{x^2} dx$ exists $\Rightarrow \int_a^{\infty} \frac{dx}{e^x} x^{\alpha+1}$ is convergent & exists

for $\alpha > 1$.

for $0 < \alpha < 1$:

$$I = \int_0^1 e^{-x} x^{\alpha+1} dx + \int_1^{\infty} e^{-x} x^{\alpha+1} dx$$

(I₁) (I₂)

I₁ is an improper integral: I₂ is also an improper integral
 & point of non-convergence $\Rightarrow x=0$

$$I_1 = \int_0^1 e^{-x} x^{\alpha+1} dx \quad \text{let } f(x) = \frac{e^{-x}}{x^{1-\alpha}}$$

let's do the comparison with $g(x) = \frac{1}{x^{1-\frac{1}{2}}}$

where $\int_0^1 \frac{1}{x^{1-\frac{1}{2}}} dx$ is convergent

$$\therefore \lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 0} \frac{e^{-x}}{x^{1-\alpha}} \times x^{\alpha+1} = \lim_{x \rightarrow 0} \frac{e^{-x}}{x^{1-\alpha-\alpha-1}} = 0$$

So the integral is convergent.

For I_2 : $\int_0^{\infty} e^{-x} x^{\alpha+1} dx$, $0 < \alpha < 1$

Let's take $g(x) = \frac{1}{x^2}$

$$\therefore \lim_{x \rightarrow \infty} \frac{e^{-x}}{x^{1-\alpha}} x^2 = \frac{e^{-x}}{x^{1-\alpha-2}} = x^{(1+\alpha)} e^{-x} = \frac{x^{1+\alpha}}{e^x} \left(\frac{\infty}{\infty} \right)$$

$$= \frac{(1+\alpha) x^{\alpha}}{e^x} = 0 \text{ as } x \rightarrow \infty$$

Hence since $\int_1^{\infty} \frac{dx}{x^2}$ is convergent, by comparison we have $\int_0^{\infty} e^{-x} x^{\alpha+1} dx$ is also convergent.

Hence this integral $\int_0^{\infty} e^{-x} x^{\alpha+1} dx$ exists for $\alpha > 0$.

4-c) Prove that $\Gamma(2n) = \frac{2^{2n-1}}{\sqrt{\pi}} \Gamma(n) \Gamma(n+\frac{1}{2})$

$\Gamma(n) = (n-1)!$

$\Gamma(n+1) = n \Gamma(n) = n!$

$$\begin{aligned} \Gamma(n+\frac{1}{2}) &= \Gamma\left(1 + (n-\frac{1}{2})\right) = (n-\frac{1}{2}) \Gamma(n-\frac{1}{2}) = (n-\frac{1}{2})(n-\frac{3}{2}) \Gamma(n-\frac{3}{2}) \\ &= \dots = (n-\frac{1}{2})(n-\frac{3}{2})(n-\frac{5}{2}) \dots \left(n-\frac{2n-1}{2}\right) \Gamma(\frac{1}{2}). \end{aligned}$$

$$\therefore \frac{2^{2n-1}}{\sqrt{\pi}} \Gamma(n) \Gamma(n+\frac{1}{2})$$

$\Gamma(2n) = (2n-1)!$

$$= \frac{2^{2n-1}}{\sqrt{\pi}} \times (n-1)! \times \frac{\sqrt{\pi}}{2^n} \frac{1 \times 3 \times 5 \times \dots \times (2n-1)}{2^n}$$

$$= 2^{2n-1-n} (n-1)! (1 \times 3 \times 5 \times \dots \times (2n-1))$$

$$= 2^{n-1} (n-1)! \cdot 1 \times 3 \times 5 \times \dots \times (2n-1)$$

$$= (2n-1)! \cdot n-1$$

We know that $B(m, n) = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)}$, $m > 0, n > 0$

So let's take $m=n$.

$$\therefore B(n, n) = \frac{(\Gamma(n))^2}{\Gamma(2n)} = 2 \int_0^{\pi/2} \sin^{2n-1} \theta \cos^{2n-1} \theta d\theta = \int_0^1 x^{n-1} (1-x)^{n-1} dx$$

$$\Rightarrow \frac{1}{2} \frac{\Gamma(n) \Gamma(n)}{\Gamma(2n)} = \int_0^{\pi/2} \sin^{2n-1} \theta \cos^{2n-1} \theta d\theta$$

$$\frac{(\Gamma(n))^2}{\Gamma(2n)} = \int_0^1 x^{n-1} (1-x)^{n-1} dx \quad \text{let } x = \sin^2 \theta$$

$$\Rightarrow dx = \sin 2\theta d\theta$$

$$\Rightarrow \frac{\Gamma(n) \Gamma(n)}{\Gamma(2n)} = \int_0^{\pi/2} 2 \sin^{2n-2} \theta \cos^{2n-2} \theta \sin \theta \cos \theta d\theta$$

$$= 2 \int_0^{\pi/2} (\sin \theta \cos \theta)^{2n-1} d\theta$$

$$= \frac{2}{2^{2n-1}} \int_0^{\pi/2} (\sin 2\theta)^{2n-1} d\theta$$

$$= \frac{2}{2^{2n-1}} \int_0^{\pi/2} (\sin 2\theta)^{2n-1} d\theta$$

$$\text{let } 2\theta = \alpha$$

$$\Rightarrow 2d\theta = d\alpha$$

$$\Rightarrow B(n, n) = \frac{1}{2^{2n-2}} \int_0^{\pi} (\sin \alpha)^{2n-1} \frac{d\alpha}{2}$$

$$= \frac{1}{2^{2n-1}} \int_0^{\pi} \sin^{2n-1} \alpha d\alpha = \frac{2}{2^{2n-1}} \int_0^{\pi/2} \sin^{2n-1} \alpha d\alpha$$

$$\because \int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx$$

$$\text{if } f(2a-x) = f(x)$$

$$= \frac{1}{2^{2n-2}} \int_0^{\pi/2} \sin^{2n-1} \alpha \cos^0 \alpha d\alpha = \frac{1}{2^{2n-2}} \frac{\Gamma(n) \cdot \Gamma(\frac{1}{2})}{2 \Gamma(n+\frac{1}{2})}$$

$$2n-1=0 \Rightarrow n=1$$

$$\Rightarrow \frac{\Gamma(n) \Gamma(n)}{\Gamma(2n)} = \frac{1}{2 \cdot 2^{2n-2}} \frac{\sqrt{\pi} \Gamma(n)}{\Gamma(n+\frac{1}{2})}$$

$$\Rightarrow \Gamma(2n) = \Gamma(n) \Gamma(n+\frac{1}{2}) \frac{2^{2n-1}}{\sqrt{\pi}}$$

Proved