

## 6

## Vector Analysis

## 1. Scalar and Vector Fields

- 1.1 Prove that the vectors  $\vec{a} = 3i + j - 2k$ ,  $\vec{b} = -i + 3j + 4k$ ,  $\vec{c} = 4i - 2j - 6k$  can form the sides of a triangle. Find the lengths of the medians of the triangle.

(2016 : 10 Marks)

**Solution:**

Here, we find that


$$\begin{aligned}\vec{b} + \vec{c} &= (-i + 3j + 4k) + (4i - 2j - 6k) \\ &= 3i + j - 2k \\ &= \vec{a}\end{aligned}$$

i.e.,

$$\vec{b} + \vec{c} = \vec{a}$$

And also we notice that these three vectors are not collinear (components are not proportional). Hence, these form the sides of a triangle. Let  $AX$ ,  $BY$  and  $CZ$  be medians.

By triangle law of vector addition :



$$\begin{aligned}\overrightarrow{AX} &= \overrightarrow{AB} + \overrightarrow{BX} = \vec{c} - \frac{\vec{a}}{2} = \frac{1}{2}(5i - 5j - 14k) \\ |\overrightarrow{AX}| &= \sqrt{\frac{1}{4}(25+25+196)} = \sqrt{\frac{246}{4}} = \frac{\sqrt{246}}{2} \\ \overrightarrow{BY} &= \overrightarrow{BA} + \overrightarrow{AY} = -\left(\vec{c} + \frac{\vec{b}}{2}\right) = -\frac{1}{2}(7i - j - 8k) \\ |\overrightarrow{BY}| &= \frac{1}{2}\sqrt{49+1+64} = \frac{\sqrt{114}}{2} \\ \overrightarrow{CZ} &= \overrightarrow{CA} + \overrightarrow{AZ} = \vec{b} + \frac{\vec{c}}{2} = \frac{1}{2}(2i + 4j + 2k) \\ |\overrightarrow{CZ}| &= |i + 2j + k| = \sqrt{1+4+1} = \sqrt{6}\end{aligned}$$

## 2. Differentiation of a Vector Field of a Scalar Variable

- 2.1 For two vectors  $\vec{a}$  and  $\vec{b}$  given respectively by

$$\vec{a} = 5t^2\hat{i} + t\hat{j} - t^3\hat{k}$$

$$\vec{b} = \sin t\hat{i} - \cos t\hat{j}$$

and

Determine : (i)  $\frac{d}{dt}(\vec{a} \cdot \vec{b})$  and  $\frac{d}{dt}(\vec{a} \times \vec{b})$

(2009 : 10 Marks)

Solution :

$$\begin{aligned}
 \vec{a} &= 5t^2\hat{i} + t\hat{j} - t^3\hat{k} \\
 \vec{b} &= \sin 5t\hat{i} - \cos t\hat{j} \\
 \vec{a} \cdot \vec{b} &= 5t^2 \sin 5t - t \cos t \quad (\because \hat{i} \cdot \hat{i} = 1 \text{ etc.}, \hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0) \\
 \therefore \frac{d}{dt}(\vec{a} \cdot \vec{b}) &= \frac{d}{dt}(5t^2 \sin 5t - t \cos t) \\
 &= 5(2t \sin 5t + t^2 \cdot 5 \cos 5t) - (1 \cdot \cos t - t \sin t) \\
 &= 10t \sin 5t + 25t^2 \cos 5t - \cos t + t \sin t \\
 \vec{a} \times \vec{b} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 5t^2 & t & -t^3 \\ \sin 5t & -\cos t & 0 \end{vmatrix} \\
 &= \hat{i}(0 - t^3 \cos t) + \hat{j}(-t^3 \sin 5t - 0) + \hat{k}(-5t^2 \cos t - t \sin 5t) \\
 &= -t^3 \cos t \hat{i} - t^3 \sin 5t \hat{j} - (5t^2 \cos t + t \sin 5t) \hat{k} \\
 \therefore \frac{d}{dt}(\vec{a} \times \vec{b}) &= (-3t^2 \cos t + t^3 \sin t) \hat{i} - (3t^2 \sin 5t + 5t^3 \cos t) \hat{j} - \\
 &\quad (10t \cos t - 5t^2 \sin t + t \cos 5t + 1 \cdot \sin 5t) \hat{k}
 \end{aligned}$$

2.2 If

$$\begin{aligned}
 \vec{A} &= x^2yz\hat{i} - 2xz^3\hat{j} + xz^2\hat{k} \\
 \vec{B} &= 2z\hat{i} + y\hat{j} - x^2\hat{k}
 \end{aligned}$$

find the value of  $\frac{\partial^2}{\partial x \partial y}(\vec{A} \times \vec{B})$  at  $(1, 0, -2)$ .

(2012 : 12 Marks)

Solution:

Given :

$$\begin{aligned}
 \vec{A} &= x^2yz\hat{i} - 2xz^3\hat{j} + xz^2\hat{k} \\
 \vec{B} &= 2z\hat{i} + y\hat{j} - x^2\hat{k} \\
 \therefore \vec{A} \times \vec{B} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ x^2yz & -2xz^3 & xz^2 \\ 2z & y & -x^2 \end{vmatrix} \\
 &= \hat{i}(2x^3z^3 - xyz^2) + \hat{j}(2xz^3 + x^4yz) + \hat{k}(x^2y^2z + 4xz^4) \\
 \frac{\partial}{\partial y}(\vec{A} \times \vec{B}) &= \hat{i}(-xz^2) + \hat{j}(x^4z) + \hat{k}(2x^2yz) \\
 \Rightarrow \frac{\partial^2}{\partial x \partial y}(\vec{A} \times \vec{B}) &= \hat{i}(-z^2) + \hat{j}(4x^3z) + \hat{k}(4xyz)
 \end{aligned}$$

∴ At  $(1, 0, -2)$

$$\frac{\partial^2}{\partial x \partial y}(\vec{A} \times \vec{B}) = -4\hat{i} - 8\hat{j}$$

- 2.3 The position vector of a moving point at time  $t$  is,  $\vec{r} = (\sin t)i + (\cos 2t)j + (t^2 + 2t)k$ . Find the components of acceleration  $\vec{a}$  in the directions parallel to the velocity vector  $\vec{v}$  and perpendicular to the plane of  $\vec{r}$  and  $\vec{v}$  at time  $t = 0$ .

(2017 : 10 Marks)

Solution:

$$\vec{r} = (\sin t)i + (\cos 2t)j + (t^2 + 2t)k \quad \dots(i)$$

$$\vec{v} = \frac{d\vec{r}}{dt} = (\cos t)i - 2 \sin 2t j + (2t + 2)k \quad \dots(ii)$$

$$\vec{a} = \frac{d^2\vec{r}}{dt^2} = (-\sin t)i - 4 \cos 2t j + 2k \quad \dots(iii)$$

At  $t = 0$ ,

$$\vec{r} = j, \vec{v} = i + 2k, \vec{a} = -4j + 2k$$

Component of  $\vec{a}$  in direction of  $\vec{v}$ 

$$\vec{a}_v = \frac{\vec{a} \cdot \vec{v}}{|\vec{v}|} \cdot \frac{\vec{v}}{|\vec{v}|} = \frac{+4}{(1+4)} \vec{v} = \frac{4}{5}(i+2k)$$

Component of  $\vec{a}$  in the direction of vector perpendicular to the plane of  $\vec{r}$  and  $\vec{v}$ .Let  $\vec{p}$  be the vector perpendicular to  $\vec{r}$  and  $\vec{v}$ 

$$\vec{p} = \vec{r} \times \vec{v} = 2\hat{i} - \hat{k}, |\vec{p}|^2 = 5$$

$$\vec{A}_p = \frac{\vec{p} \cdot \vec{a}}{|\vec{p}|} \cdot \frac{\vec{p}}{|\vec{p}|} = \frac{-2}{5}(2\hat{i} - \hat{k})$$

### 3. Gradient, Divergence and Curl in Cartesian and Cylindrical Coordinate and Directional Derivative

- 3.1 Find the directional derivative of :

(i)  $4xz^3 - 3x^2y^2z^2$  at  $(2, -1, 2)$  along  $z$ -axis(ii)  $x^2yz + 4xz^2$  at  $(1, -2, 1)$  in the direction of  $2\hat{i} - \hat{j} - 2\hat{k}$ 

(2009 : 6 + 6 = 12 Marks)

Solution:

Approach : The directional derivative in any direction is the dot product of the gradient with the unit direction.

$$(i) f(x, y, z) = 4xz^3 - 3x^2y^2z^2$$

$$\nabla f = \frac{\partial f}{\partial x}\hat{i} + \frac{\partial f}{\partial y}\hat{j} + \frac{\partial f}{\partial z}\hat{k}$$

$$= (4z^3 - 6xy^2z^2)\hat{i} - 6x^2yz^2\hat{j} + (12xz^2 - 6x^2y^2z)\hat{k}$$

Directional derivative along  $z$ -axis

$$= \nabla f \cdot \hat{k} = 12xz^2 - 6x^2y^2z$$

Directional derivative along  $z$ -axis at  $(2, -1, 2)$ 

$$= (\nabla f \cdot \hat{k})|_{(2,-1,2)} = 12 \cdot 2 \cdot 2^2 - 6 \cdot 2^2 \cdot (-1)^2 \cdot 2 = 48$$

(ii)

$$f(x, y, z) = x^2yz + 4xz^2$$

$$\nabla f = (2xyz + 4z^2)\hat{i} + (x^2z)\hat{j} + (x^2y + 8xz)\hat{k}$$

$$\nabla f|_{(1,-2,1)} = \hat{j} + 6\hat{k}$$

∴ Directional derivative along  $(2\hat{i} - \hat{j} - 2\hat{k})$

$$\begin{aligned} &= (\hat{j} + 6\hat{k}) \cdot \left( \frac{2\hat{i} - \hat{j} - 2\hat{k}}{3} \right) \\ &= \left( \frac{-1 - 12}{3} \right) = \frac{-13}{3} \end{aligned}$$

**3.2** Examine whether the vectors  $\nabla u$ ,  $\nabla v$  and  $\nabla w$  are coplanar, where  $u$ ,  $v$  and  $w$  are the scalar functions defined by :

$$u = x + y + z$$

$$v = x^2 + y^2 + z^2$$

and

$$w = yz + zx + xy$$

(2011 : 15 Marks)

**Solution:**

$$\begin{aligned} \nabla u &= \left( \frac{\partial}{\partial x} \vec{i} + \frac{\partial}{\partial y} \vec{j} + \frac{\partial}{\partial z} \vec{k} \right) (x + y + z) \\ &= \vec{i} + \vec{j} + \vec{k} \end{aligned}$$

$$\nabla v = 2x\vec{i} + 2y\vec{j} + 2z\vec{k}$$

Similarly,

$$\nabla w = (y+z)\vec{i} + (x+z)\vec{j} + (x+y)\vec{k}$$

Now,  $\nabla u$ ,  $\nabla v$ ,  $\nabla w$  would be co-planar if their scalar triple product is zero.

$$\begin{aligned} \nabla v \times \nabla w &= \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2x & 2y & 2z \\ y+z & x+z & x+y \end{vmatrix} \\ &= \vec{i}(2xy + 2y^2 - 2xz - 2z^2) + \vec{j}(2yz + 2z^2 - 2x^2 - 2xy) + \\ &\quad \vec{k}(2x^2 + 2xz - 2y^2 - 2yz) \\ \therefore \nabla u(\nabla v \times \nabla w) &= (\vec{i} + \vec{j} + \vec{k}) \cdot [(2xy + 2y^2 - 2xz - 2z^2)\vec{i} \\ &\quad + (2yz + 2z^2 - 2x^2 - 2xy)\vec{j} + (2x^2 + 2xz - 2y^2 - 2yz)\vec{k}] \\ &= 2xy + 2y^2 - 2x^2 - 2z^2 + 2yz + 2z^2 - 2x^2 - 2xy + 2x^2 + 2xz - 2y^2 - 2yz \\ &= 0 \end{aligned}$$

∴ The vector  $\nabla u$ ,  $\nabla v$ ,  $\nabla w$  are co-planar.

**3.3** If  $\vec{r}$  be the position vector of a point, find the value(s) of  $n$  for which the vector  $r^n \vec{r}$  is (i) irrotational, (ii) solenoidal.

(2011 : 15 Marks)

**Solution:**

A vector  $\vec{V}$  is said to be solenoidal if divergence of  $\vec{V} = 0$ .

i.e.,

$$\nabla \cdot \vec{V} = 0$$

Also,

$$\operatorname{div}(\phi \vec{V}) = (\operatorname{grad} \phi) \cdot \vec{V} + \phi \operatorname{div} \vec{V}$$

 $\therefore r^{\gamma} \vec{r}$  will be solenoidal if

$$\operatorname{div}(r^{\gamma} \vec{r}) = 0$$

$$\Rightarrow (\operatorname{grad} r^{\gamma}) \cdot \vec{r} + r^{\gamma} \operatorname{div}(\vec{r}) = 0$$

$$\Rightarrow (nr^{\gamma-1} \operatorname{grad} r) \cdot \vec{r} + r^{\gamma} \cdot 3 = 0$$

$$\begin{aligned}\therefore \operatorname{div} \vec{r} &= \left( \hat{i} \frac{\partial}{\partial x} + \hat{j} \frac{\partial}{\partial y} + \hat{k} \frac{\partial}{\partial z} \right) (xi + yj + zk) \\ &= 1 + 1 + 1 = 3\end{aligned}$$

and

$$\operatorname{grad} f(4) = f'(4) \operatorname{grad} u$$

$$\Rightarrow \left( nr^{\gamma-1} \cdot \frac{\vec{r}}{r} \right) \cdot \vec{r} + 3r^{\gamma} = 0 \quad \left[ \because \operatorname{grad} r = \frac{\vec{r}}{r} \right]$$

$$\Rightarrow nr^{\gamma-2} (\vec{r} \cdot \vec{r}) + 3r^{\gamma} = 0$$

$$\Rightarrow nr^{\gamma-2} \cdot r^2 + 3r^{\gamma} = 0$$

$$\Rightarrow r^{\gamma}(n+3) = 0 \Rightarrow n = -3$$

A vector  $\vec{V}$  is said to be irrotational if

$$\nabla \times \vec{V} = 0$$

Also,

$$\nabla \times (\phi \vec{V}) = (\operatorname{grad} \phi) \times \vec{V} + \phi (\nabla \times \vec{V})$$

 $\therefore r^{\gamma} \vec{r}$  will be irrotational if

$$\nabla \times (r^{\gamma} \vec{r}) = 0$$

$$\Rightarrow (\operatorname{grad} r^{\gamma}) \times \vec{r} + r^{\gamma} (\nabla \times \vec{r}) = 0$$

$$\Rightarrow \left( nr^{\gamma-1} \frac{\vec{r}}{r} \right) \times \vec{r} + r^{\gamma} \cdot 0 = 0 \quad (\because \nabla \times \vec{r} = 0)$$

Hence,  $r^{\gamma} \vec{r}$  is irrotational for all the real values of  $n$ .

## 3.4 A vector field is given by

$$\vec{F} = (x^2 + xy^2)i + (y^2 + x^2y)j$$

Verify that the field  $\vec{F}$  is irrotational or not. Find the scalar potential.

(2015 : 12 Marks)

Solution:

A vector field  $\vec{F}$  is said to be irrotational if  $\operatorname{curl} \vec{F} = 0$ , i.e.,

$$\nabla \times \vec{F} = 0$$

$$\vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^2 + xy^2 & y^2 + x^2y & 0 \end{vmatrix}$$

$$= \hat{i}(0-0) - \hat{j}(0-0) + \hat{k}(2xy - 2xy)$$

$$= \vec{0}$$

$\Rightarrow \vec{F}$  is irrotational.

Now, it can be written as grad of a scalar field, i.e., to find  $\phi$  so that

$$\nabla\phi = \vec{F}$$

i.e.,

$$\hat{i}\frac{\partial\phi}{\partial x} + \hat{j}\frac{\partial\phi}{\partial y} = (x^2 + xy^2)\hat{i} + (y^2 + x^2y)\hat{k}$$

$$\therefore \frac{\partial\phi}{\partial x} = x^2 + xy^2; \quad \frac{\partial\phi}{\partial y} = y^2 + x^2y \quad \dots (*)$$

$$\Rightarrow \phi = \frac{x^3}{3} + \frac{x^2y^2}{2} + f(y)$$

Differentiating w.r.t.  $y$  and comparing with  $(*)$

$$\frac{\partial\phi}{\partial y} = x^2y + f'(y)$$

$$\Rightarrow f'(y) = y^2$$

$$f(y) = \frac{y^3}{3} + C$$

$$\therefore \phi(x, y) = \frac{x^3}{3} + \frac{y^3}{3} + \frac{x^2y^2}{2} + C$$

### 3.5 For what values of the constants $a$ , $b$ and $c$ the vector

$$\vec{V} = (x+y+az)\hat{i} + (bx+2y-z)\hat{j} + (-x+cy+2z)\hat{k}$$

is irrational. Find the divergence in cylindrical coordinates of this vector with these values.

(2017 : 10 Marks)

**Solution:**

$$\text{Irrational} \Rightarrow \text{Curl } \vec{V} = 0$$

$$\text{Curl } \vec{V} = \nabla \times \vec{V} \begin{bmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ f & g & h \end{bmatrix} \text{ (if } V = fi + gj + hk\text{)}$$

$$= \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x+y+az & bx+2y-z & -x+cy+2z \end{vmatrix}$$

$$= i(c-(-1)) - j(-1-a) + k(b-1)$$

$$= (c+1)i + (a+1)j + (b-1)k$$

$$= 0$$

$$\therefore a = -1, b = 1, c = -1$$

$$\vec{V} = (x+y-z)\hat{i} + (x+2y-z)\hat{j} + (-x-y+2z)\hat{k}$$

We find  $\text{div } \vec{V}$  and express it in cylindrical coordinates.

$$\vec{V} = (x + y - z)i + (x + 2y - z)j + (-x - y + 2z)k$$

$$\operatorname{div} \vec{V} = \nabla \cdot \vec{V}$$

$$= \frac{\partial V_x}{\partial x} + \frac{\partial V_y}{\partial y} + \frac{\partial V_z}{\partial z}$$

$$= 1 + 2 + 2 = 5 \text{ (constant)}$$

Hence, divergence in cylindrical co-ordinates = 5.

- 3.6 Let  $\vec{v} = v_1\hat{i} + v_2\hat{j} + v_3\hat{k}$ . Show that  $\operatorname{curl}(\operatorname{curl} \vec{v}) = \operatorname{grad}(\operatorname{div} \vec{v}) - \nabla^2 \vec{v}$ .

(2018 : 12 Marks)

Solution:

$$\operatorname{Curl}(\operatorname{Curl} \vec{v}) = \nabla \times (\nabla \times \vec{v})$$

Now, we know that

$$\vec{A} \times (\vec{B} \times \vec{C}) = (\vec{A} \cdot \vec{C})\vec{B} - (\vec{A} \cdot \vec{B})\vec{C}$$

Given :

$$v_1\hat{i} + v_2\hat{j} + v_3\hat{k} = \vec{v}$$

$\therefore$

$$\nabla \times \vec{v} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_1 & v_2 & v_3 \end{vmatrix}$$

$$= \hat{i}\left(\frac{\partial v_3}{\partial y} - \frac{\partial v_2}{\partial z}\right) - \hat{j}\left(\frac{\partial v_3}{\partial x} - \frac{\partial v_1}{\partial z}\right) + \hat{k}\left(\frac{\partial v_2}{\partial x} - \frac{\partial v_1}{\partial y}\right)$$

$\therefore$

$$\nabla \times (\nabla \times \vec{v}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{\partial v_3}{\partial y} - \frac{\partial v_2}{\partial z} & \frac{\partial v_1}{\partial z} - \frac{\partial v_3}{\partial x} & \frac{\partial v_2}{\partial x} - \frac{\partial v_1}{\partial y} \end{vmatrix}$$

$$= \hat{i}\left(\frac{\partial^2 v_2}{\partial y \partial x} - \frac{\partial^2 v_1}{\partial y^2} - \frac{\partial^2 v_1}{\partial z^2} + \frac{\partial v_3}{\partial z \partial x}\right) - \hat{j}\left(\frac{\partial^2 v_2}{\partial x^2} - \frac{\partial^2 v_1}{\partial x \partial y} - \frac{\partial^2 v_3}{\partial y \partial z} + \frac{\partial^2 v_2}{\partial z^2}\right) + \hat{k}\left(\frac{\partial^2 v_1}{\partial x \partial z} - \frac{\partial^2 v_3}{\partial x^2} - \frac{\partial^2 v_3}{\partial y^2} + \frac{\partial^2 v_2}{\partial y \partial z}\right)$$

$$= \hat{i}\left(\frac{\partial^2 v_2}{\partial y \partial x} + \frac{\partial^2 v_3}{\partial z \partial x}\right) + \hat{j}\left(\frac{\partial^2 y}{\partial x \partial y} + \frac{\partial^2 v_3}{\partial y \partial z}\right) + \hat{k}\left(\frac{\partial^2 v_1}{\partial x \partial z} + \frac{\partial^2 v_2}{\partial y \partial z}\right)$$

$$= -\hat{i}\left(\frac{\partial^2 v_1}{\partial y^2} + \frac{\partial^2 v_1}{\partial z^2}\right) - \hat{j}\left(\frac{\partial^2 v_2}{\partial x^2} + \frac{\partial^2 v_2}{\partial z^2}\right) - \hat{k}\left(\frac{\partial^2 v_3}{\partial x^2} + \frac{\partial^2 v_3}{\partial y^2}\right)$$

$$= \hat{i}\left(\frac{\partial^2 v_1}{\partial x^2} + \frac{\partial^2 v_2}{\partial x \partial y} + \frac{\partial^2 v_3}{\partial x \partial z}\right) + \hat{j}\left(\frac{\partial^2 v_1}{\partial y \partial x} + \frac{\partial^2 v_2}{\partial y^2} + \frac{\partial^2 v_3}{\partial y \partial z}\right) + \hat{k}\left(\frac{\partial^2 v_2}{\partial x^2} + \frac{\partial^2 v_2}{\partial y^2} + \frac{\partial^2 v_2}{\partial z^2}\right) - \hat{i}\left(\frac{\partial^2 y}{\partial x^2} + \frac{\partial^2 v_1}{\partial y^2} + \frac{\partial^2 v_1}{\partial z^2}\right) -$$

$$\hat{j}\left(\frac{\partial^2 v_2}{\partial x^2} + \frac{\partial^2 v_2}{\partial y^2} + \frac{\partial^2 v_2}{\partial z^2}\right) - \hat{k}\left(\frac{\partial^2 v_3}{\partial x^2} + \frac{\partial^2 v_3}{\partial y^2} + \frac{\partial^2 v_3}{\partial z^2}\right)$$

$$= \nabla(\nabla \cdot \vec{v}) - \nabla^2 \vec{v} = \operatorname{grad}(\operatorname{div} \vec{v}) - \nabla^2 \vec{v}$$

- 3.7 Find the directional derivative of the function  $xy^2 + yz^2 + zx^2$  along the tangent to the curve  $x = t$ ,  $y = t^2$ ,  $z = t^3$  at the point  $(1, 1, 1)$ .

(2019 : 10 Marks)

**Solution:**

Let

$$\phi(x, y, z) = xy^2 + yz^2 + zx^2$$

Then,

$$\text{grad } \phi = \left( \frac{\partial \phi}{\partial x} \right) i + \left( \frac{\partial \phi}{\partial y} \right) j + \left( \frac{\partial \phi}{\partial z} \right) k$$

$$\text{grad } \phi = (y^2 + 2zx)i + (z^2 + 2xy)j + (x^2 + 2yz)k$$

$$\text{grad } \phi_{\text{at}(1,1,1)} = 3i + 3j + 3k$$

Also, for the curve  $x = t$ ,  $y = t^2$  and  $z = t^3$ 

We have,

$$\frac{dx}{dt} = 1, \frac{dy}{dt} = 2t, \frac{dz}{dt} = 3t^2$$

At point  $(1, 1, 1)$  on the curve  $x = t$ ,  $y = t^2$ ,  $z = t^3$ 

we have,

$$t = 1$$

Now, vector along the tangent to the above curve at the point  $(x, y, z)$ 

$$\begin{aligned} &= \left( \frac{dx}{dt} \right) \hat{i} + \left( \frac{dy}{dt} \right) \hat{j} + \left( \frac{dz}{dt} \right) \hat{k} \\ &= \hat{i} + 2\hat{j} + 3t^2\hat{k} \end{aligned}$$

Putting  $t = 1$ , a vector along the tangent to the curve at the point  $(1, 1, 1)$ , we have

$$= \hat{i} + 2\hat{j} + 3\hat{k}$$

If  $\hat{a}$  be the unit vector in the direction of this tangent then

$$\begin{aligned} \hat{a} &= \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{\|\hat{i} + 2\hat{j} + 3\hat{k}\|} = \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{1+4+9}} \\ \hat{a} &= \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{14}} \end{aligned}$$

∴ The required directional derivative

$$\begin{aligned} &= \hat{a} \text{ grad } \phi \text{ at } (1, 1, 1) \\ &= \frac{\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{14}} (3\hat{i} + 3\hat{j} + 3\hat{k}) \\ &= \frac{3+6+9}{\sqrt{14}} = \frac{18}{\sqrt{14}} \\ \therefore \hat{a} \text{ grad } \phi &= \frac{18}{\sqrt{14}}. \end{aligned}$$

#### 4. Vector Identity and Vector Equations

- 4.1 Show that  $\text{div}(\text{grad } r^n) = n(n+1)r^{n-2}$

(2009 : 12 Marks)

**Solution:**

$$\nabla \cdot (\nabla r^n) = n(n+1)r^{n-2}$$

$$r^2 = x^2 + y^2 + z^2$$

Differentiating partially with respect to  $x$  on both sides

$$2r \frac{\partial r}{\partial x} = 2x \Rightarrow \frac{\partial r}{\partial x} = \frac{x}{r}$$

Similarly,

$$\frac{\partial r}{\partial y} = \frac{y}{r}; \quad \frac{\partial r}{\partial z} = \frac{z}{r}$$

$$\begin{aligned}\text{grad } r^n &= \sum \frac{\partial}{\partial x} i(r^n) = \sum \frac{\partial r^n}{\partial x} \hat{i} \\ &= \sum n r^{n-1} \frac{\partial r}{\partial x} \hat{i} = n r^{n-1} \sum \frac{x}{r} \hat{i} \\ &= n r^{n-2} \vec{r}\end{aligned}$$

where,

$$\vec{r} = x \hat{i} + y \hat{j} + z \hat{k}$$

$$\begin{aligned}\text{div}(\text{grad } r^n) &= \sum \frac{\partial}{\partial x} i \cdot n r^{n-2} \sum x \hat{i} = \sum \frac{\partial}{\partial x} (n r^{n-2} x) \\ &= \sum \left( n r^{n-2} + n(n-2) r^{n-3} \cdot x \frac{\partial r}{\partial x} \right) \\ &= 3 n r^{n-2} + \sum n(n-2) r^{n-3} \frac{x^2}{r} \\ &= 3 n r^{n-2} + n(n-2) r^{n-4} \sum x^2 \\ &= n(n-2) r^{n-2} + 3 n r^{n-2} = n(n+1) r^{n-2}\end{aligned}$$

#### 4.2 Find the directional derivative of

$$f(x, y) = x^2 y^3 + xy$$

at the point (2, 1) in the direction of a unit vector which makes an angle of  $\frac{\pi}{3}$  with the x-axis.

(2010 : 12 Marks)

**Solution:**

Given :

$$f(x, y) = x^2 y^3 + xy$$

∴

$$\begin{aligned}\nabla f &= \frac{\partial f}{\partial x} \hat{i} + \frac{\partial f}{\partial y} \hat{j} = \frac{\partial(x^2 y^3 + xy)}{\partial x} \hat{i} + \frac{\partial(x^2 y^3 + xy)}{\partial y} \hat{j} \\ &= (2xy^3 + y)\hat{i} + (3x^2 y^2 + x)\hat{j}\end{aligned}$$

Now,

$$\begin{aligned}(\nabla f)_{(2,1)} &= (2 \times 2 \times 1^3 + 1)\hat{i} + (3 \times 2^2 \times 1^2 + 2)\hat{j} \\ &= 5\hat{i} + 14\hat{j}\end{aligned}$$

$(\nabla f)_{(2,1)}$  is the direction of unit vector at angle  $\frac{\pi}{3}$  with x-axis is

$$(\nabla f)_{(2,1)} \cdot \left( \cos \frac{\pi}{3} \hat{i} + \sin \frac{\pi}{3} \hat{j} \right) = (5\hat{i} + 14\hat{j}) \cdot \left( \frac{1}{2} \hat{i} + \frac{\sqrt{3}}{2} \hat{j} \right) = \frac{5}{2} + 7\sqrt{3}$$

#### 4.3 Show that the vector field defined by the vector function

$$\vec{v} = xyz(y\hat{i} + x\hat{j} + z\hat{k})$$

is conservative.

(2010 : 12 Marks)

**Solution:**

Given :

$$\vec{v} = xyz(\hat{yzi} + \hat{xzj} + \hat{xyk})$$

$$= xy^2z^2\hat{i} + x^2yz^2\hat{j} + x^2y^2z\hat{k}$$

$$\nabla \times \vec{v} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ xy^2z^2 & x^2yz^2 & x^2y^2z \end{vmatrix}$$

$$= \hat{i}\left(\frac{\partial}{\partial y}x^2y^2z - \frac{\partial}{\partial z}x^2yz^2\right) - \hat{j}\left(\frac{\partial}{\partial x}x^2y^2z - \frac{\partial}{\partial z}xy^2z^2\right) + \hat{k}\left(\frac{\partial}{\partial x}x^2yz^2 - \frac{\partial}{\partial y}xy^2z^2\right)$$

$$= \hat{i}(2x^2y^2 - 2x^2y^2) - \hat{j}(2x^2y^2z - 2xy^2z) + \hat{k}(2xyz^2 - 2xyz^2) = 0$$

as  $\nabla \times \vec{v} = 0$ .  $\therefore \vec{v}$  is conservative.

**4.4 Prove that :**

$$\operatorname{div}(f\vec{v}) = f(\operatorname{div} \vec{v}) + (\operatorname{grad} f) \cdot \vec{v}$$

where  $f$  is a scalar function.

(2011 : 20 Marks)

**Solution:**

$$\text{LHS} = \operatorname{div}(f\vec{v})$$

$$\text{RHS} = f \operatorname{div} \vec{v} + (\operatorname{grad} f) \cdot \vec{v}$$

Take LHS :

$$\begin{aligned} \operatorname{div}(f\vec{v}) &= \sum i \frac{\partial}{\partial x}(f\vec{v}) \\ &= \left( \sum i \frac{\partial f}{\partial x} \right) \cdot \vec{v} + f \sum i \frac{\partial}{\partial x}(\vec{v}) \\ &= \operatorname{grad} f \cdot \vec{v} + f(\operatorname{div} \vec{v}) \\ &= f(\operatorname{div} \vec{v}) + (\operatorname{grad} f) \cdot \vec{v} \\ &= \text{RHS} \end{aligned}$$

So,

LHS = RHS. Hence Proved.

**4.5 If  $u$  and  $v$  are two scalar fields and  $\vec{f}$  is a vector field, such that**

$$u\vec{f} = \operatorname{grad} v,$$

**find the value of  $\vec{f} \cdot \operatorname{curl} \vec{f}$ .**

(2011 : 10 Marks)

**Solution:**

Given :

$$u\vec{f} = \operatorname{grad} v$$

$\Rightarrow$

$$\vec{f} = \frac{1}{u} \cdot \operatorname{grad} v$$

$\therefore$

$$\vec{f} \cdot \operatorname{curl} \vec{f} = \left( \frac{1}{u} \operatorname{grad} v \right) \cdot \operatorname{curl} \left( \frac{1}{u} \operatorname{grad} v \right)$$

$$\begin{aligned}
 &= \left( \frac{1}{u} \operatorname{grad} v \right) \cdot \left[ \left( \operatorname{grad} \frac{1}{u} \right) \times (\operatorname{grad} v) + \frac{1}{u} (\operatorname{curl} \operatorname{grad} v) \right] \\
 &\quad [ \because \operatorname{curl} (\phi \vec{A}) = (\operatorname{grad} \phi) \times \vec{A} + \phi \operatorname{Curl} \vec{A} ] \\
 &= \left( \frac{1}{u} \operatorname{grad} v \right) \cdot \left[ \left( \operatorname{grad} \frac{1}{u} \right) \times (\operatorname{grad} v) + 0 \right] \quad (\because \operatorname{Curl} \operatorname{grad} \phi = 0) \\
 &= \frac{1}{u} \left[ \operatorname{grad} v \cdot \operatorname{grad} \frac{1}{u} - \operatorname{grad} u \cdot \operatorname{grad} v \right] = 0
 \end{aligned}$$

[ $\because$  If a vector repeats in a scalar triple product, then its value is zero].

- 4.6 Calculate  $\nabla^2(r^n)$  and find its expression in terms of  $r$  and  $n$ ,  $r$  being the distance of any point  $(x, y, z)$  from the origin,  $n$  being a constant and  $\nabla^2$  being the Laplace operator.

(2013 : 10 Marks)

Solution:

$$\begin{aligned}
 \nabla^2(r^n) &= \nabla \cdot \nabla(r^n) \\
 \nabla(r^n) &= \sum \frac{\partial}{\partial x} (r^n) \hat{i} = nr^{n-1} \sum \frac{\partial r}{\partial x} \hat{i} = nr^{n-1} \sum \frac{\partial r}{\partial x} \hat{i} \\
 \text{Now } r^2 &= x^2 + y^2 + z^2 \Rightarrow 2r \frac{\partial r}{\partial x} = 2x \Rightarrow \frac{\partial r}{\partial x} = \frac{x}{r}
 \end{aligned}$$

$$\therefore \nabla(r^n) = nr^{n-1} \sum \frac{x}{r} \hat{i} = nr^{n-2} \vec{r}$$

$$\text{where } \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\begin{aligned}
 \nabla \cdot \nabla(r^n) &= \left( \sum \frac{\partial}{\partial x} \hat{i} \right) \cdot (nr^{n-2} \sum x \hat{i}) \\
 &= \sum \frac{\partial}{\partial x} (nr^{n-2} x) \\
 &= \sum \left( nr^{n-2} + n(n-2)r^{n-3} \frac{\partial r}{\partial x} \cdot x \right) \\
 &= \left( 3nr^{n-2} + \sum n(n-2)r^{n-3} \frac{x^2}{r} \right) \\
 &= 3nr^{n-2} + n(n-2)r^{n-4} \sum x^2 \\
 &= 3nr^{n-2} + n(n-2)r^{n-2} \\
 &= n(n+1)r^{n-2}
 \end{aligned}$$

- 4.7 Find  $f(r)$  such that  $\nabla f = \frac{\vec{r}}{r^5}$  and  $f(1) = 0$ .

(2016 : 10 Marks)

Solution:

We know that

$$\nabla f = f'(r) \nabla r = f'(r) \frac{\vec{r}}{r} \quad \left( \because \nabla r = \frac{\vec{r}}{r} \right)$$

$$\nabla f = \frac{\vec{r}}{r^5}$$

We have,

$$\therefore f'(r) \frac{\vec{r}}{r} = \frac{\vec{r}}{r^5}$$

$$\Rightarrow \vec{r} \left[ \frac{f'(r)}{r} - \frac{1}{r^5} \right] = 0$$

Since,  $\vec{r} \neq 0$ ,  $\therefore$

$$f(r) = \frac{1}{r^4}$$

Integrating, we get

$$f(r) = \frac{-1}{3r^3} + C$$

$$f(1) = 0 \Rightarrow 0 = \frac{-1}{3 \cdot 1} + c \Rightarrow c = \frac{1}{3}$$

$\therefore$

$$f(r) = \frac{1}{3} \left( 1 - \frac{1}{r^3} \right)$$

$$\begin{aligned} \nabla &= i \frac{\partial}{\partial x} + j \frac{\partial}{\partial y} + k \frac{\partial}{\partial z} \\ \vec{r} &= xi + yj + zk \\ r &= \sqrt{x^2 + y^2 + z^2} \end{aligned}$$

## 5. Application to Geometry

- 5.1 A curve in space is defined by the vector equation  $\vec{r} = t^2\hat{i} + 2t\hat{j} - t^3\hat{k}$ . Determine the angle between the tangents to this curve at the  $t = +1$  and  $t = -1$ .

(2013 : 10 Marks)

**Solution:**

$$\vec{r} = t^2\hat{i} + 2t\hat{j} - t^3\hat{k}$$

$$\frac{d\vec{r}}{dt} = 2t\hat{i} + 2\hat{j} - 3t^2\hat{k}$$

$$\frac{ds}{dt} = \left| \frac{d\vec{r}}{dt} \right| = \sqrt{4t^2 + 4 + 9t^4}$$

$$\therefore \hat{t} = \frac{d\vec{r}}{ds} = \frac{\frac{d\vec{r}}{dt}}{\frac{ds}{dt}} = \frac{2\hat{i} + 2\hat{j} - 3t^2\hat{k}}{\sqrt{4t^2 + 4 + 9t^4}}$$

where  $\hat{t}$  is the direction of the tangent

$$\hat{t}\Big|_{t=1} = \frac{2\hat{i} + 2\hat{j} - 3\hat{k}}{\sqrt{17}}$$

$$\hat{t}\Big|_{t=-1} = \frac{-2\hat{i} + 2\hat{j} - 3\hat{k}}{\sqrt{17}}$$

Let  $\theta$  be the angle

Then

$$\cos \theta = \hat{t}_1 \cdot \hat{t}_2 = \frac{1}{17} (2\hat{i} + 2\hat{j} - 3\hat{k}) \cdot (-2\hat{i} + 2\hat{j} - 3\hat{k}) = \frac{-4 + 4 + 9}{17} = \frac{9}{17}$$

$\therefore$

$$\theta = \cos^{-1} \frac{9}{17}$$

- 5.2 Find the angle between the surfaces  $x^2 + y^2 + z^2 - 9 = 0$  and  $z = x^2 + y^2 - 3$  at  $(2, -1, 2)$ .  
(2015 : 10 Marks)

**Solution:**

Angle between two surfaces at a point is the angle between the normals to the surfaces at that point.

Let

$$f_1 = x^2 + y^2 + z^2$$

and

$$f_2 = x^2 + y^2 - z$$

Then

$$\text{grad}(f_1) = 2xi + 2yj + 2zk$$

$$\text{grad}(f_2) = 2xi + 2yj - k$$

Let

$$n_1 = \text{grad } f_1 \text{ at point } (2, -1, 2) = 4i - 2j + 4k$$

$$n_2 = \text{grad } f_2 \text{ at point } (2, -1, 2) = 4i - 2j - k$$

The vectors  $n_1$  and  $n_2$  are along normals to the two surfaces at the point  $(2, -1, 2)$ . If  $\theta$  is the angle between these two vectors then,

$$\begin{aligned} \cos \theta &= \frac{n_1 \cdot n_2}{|n_1| |n_2|} \\ &= \frac{16+4-4}{\sqrt{16+4+16} \sqrt{16+4+1}} = \frac{16}{6\sqrt{21}} \\ \therefore \theta &= \cos^{-1} \frac{8}{3\sqrt{21}} \end{aligned}$$

- 5.3 Find the values of  $\lambda$  and  $\mu$  so that the surfaces  $\lambda x^2 - \mu yz = (\lambda + 2)x$  and  $4x^2y + z^3 = 4$  may intersect orthogonally at  $(1, -1, 2)$ .

(2015 : 12 Marks)

**Solution:**

Let

$$f_1 = \lambda x^2 - (\lambda + 2)x - \mu yz$$

$$f_2 = 4x^2y + z^3$$

$$\text{grad } f_1 = \nabla f_1 = (2\lambda x - \lambda - 2)i - \mu zj - \mu yk$$

$$\nabla f_2 = 8xyi + 4x^2j + 3z^2k$$

At point  $(1, -1, 2)$

$$n_1 = (\lambda - 2)i - 2\mu j + \mu k$$

$$n_2 = -8i + 4j + 12k$$

$$n_1 \cdot n_2 = 0$$

(intersect orthogonally)

$$-8(\lambda - 2) - 8\mu + 12\mu = 0$$

$$-8\lambda + 4\mu = -16 \Rightarrow 2\lambda - \mu = 4$$

... (i)

Point  $(1, -1, 2)$  lies on surface  $\lambda x^2 - \mu yz = (\lambda + 2)x$

$$\therefore \lambda + 2\mu = (\lambda + 2) \Rightarrow \mu = 1$$

$$\therefore 2\lambda - 1 = 4 \Rightarrow \lambda = \frac{5}{2}$$

- 5.4 Find the curvature vector and its magnitude at any point  $\vec{r} = (\theta)$  of the curve  $\vec{r} = (a \cos \theta, a \sin \theta, a\theta)$ . Show that the locus of the feet of the perpendicular from the origin to the tangent is a curve that completely lies on the hyperboloid  $x^2 + y^2 - z^2 = a^2$ .

(2017 : 16 Marks)

**Solution:**

$$\vec{r} = a \cos \theta i + a \sin \theta j + a\theta k$$

$$\frac{d\vec{r}}{d\theta} = -a \sin \theta i + a \cos \theta j + ak$$

$$\frac{ds}{d\theta} = \left| \frac{d\vec{r}}{d\theta} \right| = \sqrt{2a}$$

$$\vec{\tau} = \frac{d\vec{r}}{ds} = \frac{\frac{d\vec{r}}{d\theta}}{\frac{ds}{d\theta}}$$

$$= \frac{1}{\sqrt{2}}(-\sin \theta i + \cos \theta j + k)$$

$$\frac{d\vec{\tau}}{d\theta} = \frac{1}{\sqrt{2}}(-\cos \theta i - \sin \theta j)$$

∴ Curvature vector,

$$\vec{k} = \frac{d\vec{\tau}}{ds} = \frac{\cancel{d\vec{\tau}/d\theta}}{\cancel{ds/d\theta}}$$


$$= \frac{1}{\sqrt{2a}} \cdot \frac{1}{\sqrt{2}}(-\cos \theta i - \sin \theta j)$$

$$= \frac{-\cos \theta i}{2a} - \frac{\sin \theta j}{2a}$$

Magnitude,

$$|k| = \frac{1}{2a}$$

Locus of feet of perpendicular from origin to tangent



$$\overrightarrow{AC} = \vec{r} - (\vec{r} \cdot \vec{\tau})\vec{\tau}$$

C = Foot of perpendicular

$$\vec{r} \cdot \vec{\tau} = (a \cos \theta i + a \sin \theta j + a \theta k) \cdot (-\sin \theta i + \cos \theta j + k) \frac{1}{\sqrt{2}}$$

$$= \frac{a\theta}{\sqrt{2}}$$

$$\vec{r} - (\vec{r} \cdot \vec{\tau})\vec{\tau} = (a \cos \theta i + a \sin \theta j + a \theta k) - \frac{a\theta}{\sqrt{2}} \cdot \frac{1}{\sqrt{2}}(-\sin \theta i + \cos \theta j + k)$$

$$= \left( a \cos \theta + \frac{a \theta \sin \theta}{2} \right) i + \left( a \sin \theta - \frac{a \theta \cos \theta}{2} \right) j + \frac{a \theta}{2} k$$

Let foot of perpendicular be C(x, y, z)

∴

$$x = a \cos \theta + \frac{a \theta \sin \theta}{2}$$

$$y = a \sin \theta - \frac{a \theta \cos \theta}{2}$$

$$z = \frac{a \theta}{2}$$

$$\therefore x^2 + y^2 - z^2 = a^2 + \frac{a^2 \theta^2}{4} - \frac{a^2 \theta^2}{4} = a^2$$

$x^2 + y^2 - z^2 = a^2$  is the required locus.

- 5.5 Find the angle between tangent at general point of curve whose equations are  $x = 3t$ ,  $y = 3t^2$ ,  $z = 3t^3$  and the line  $z - x = y = 0$ .

(2018 : 10 Marks)

**Solution:**The radius vector of given curve,  $\vec{r}$  can be written as

$$\vec{r} = 3\hat{i} + 3t^2\hat{j} + 3t^3\hat{k}$$

$$\frac{d\vec{r}}{dt} = 3\hat{i} + 6t\hat{j} + 9t^2\hat{k}$$

$$\frac{ds}{dt} = \left| \frac{d\vec{r}}{dt} \right| = \sqrt{9 + 36t^2 + 81t^4} = 3\sqrt{1 + 4t^2 + 9t^4}$$

Tangent vector,

$$\vec{T} = \frac{d\vec{r}}{ds} = \frac{d\vec{r}/dt}{ds/dt} = \frac{1}{3\sqrt{1+4t^2+9t^4}} \times (3\hat{i} + 6t\hat{j})$$

 $\Rightarrow$ 

$$\vec{T} = \frac{\hat{i} + 2t\hat{j} + 3t^2\hat{k}}{\sqrt{1+4t^2+9t^4}}$$

The given line can be written as

$$\frac{x}{1} = \frac{y}{0} = \frac{z}{1}$$

 $\therefore$  angle between line and tangent vector,  $\theta$ 

$$\cos \theta = \frac{1 \times 1 + 2t \times 0 + 3t^2 \times 1}{\sqrt{2} \cdot \sqrt{1+4t^2+9t^4}} = \frac{1+3t^2}{\sqrt{2} \cdot \sqrt{1+4t^2+9t^4}}$$


$$\theta = \cos^{-1} \left( \frac{1+3t^2}{\sqrt{2} \cdot \sqrt{1+4t^2+9t^4}} \right)$$

- 5.6 Find the circulation of  $\vec{F}$  round the curve  $C$ , where  $\vec{F} = (2x+y^2)\hat{i} + (3y-4x)\hat{j}$  and  $C$  is the curve  $y = x^2$  from  $(0, 0)$  to  $(1, 1)$  and the curve  $y^2 = x$  from  $(1, 1)$  to  $(0, 0)$ .

(2019 : 15 Marks)

**Solution:**

Here the closed curve ' $C$ ' consists of arcs  $OAP$  and  $PBO$ . Let  $C_1$  denote the arc  $OAP$  and  $C_2$  denote arc  $PBO$ . Along  $C_1$ , we have  $y = x^2$  so that  $dy = 2x dx$  and  $x$  varies from  $0 \rightarrow 1$  along  $C_2$ , we have  $x = y^2$  so that  $dx = 2y dy$  and  $y$  varies from  $1$  to  $0$  also.



$$\vec{F} \cdot d\vec{r} = [(2x+y^2)\hat{i} + (3y-4x)\hat{j}][(dx\hat{i} + dy\hat{j})]$$

$$\vec{F} \cdot d\vec{r} = (2x+y^2)dx + (3y-4x)dy$$

Now circulation of  $\vec{F}$  round  $C$ .

∴

$$\begin{aligned}\oint_C \vec{F} \cdot d\vec{r} &= \oint_{C_1} \vec{F} \cdot d\vec{r} + \oint_{C_2} \vec{F} \cdot d\vec{r} \\ &= \oint_{C_1} (2x+y^2)dx + (3y-4x)dy + \oint_{C_2} (2x+y^2)dx + (3y-4x)dy \\ &= \int_{x=0}^1 (2x+x^4)dx + (3x^2-4x)2xdx + \\ &\quad \int_{y=1}^0 (2y^2+y^2)2ydy + (3y-4y^2)dy \\ &= \int_0^1 (2x-8x^2+6x^3+x^4)dx + \int_0^1 (3y-4y^2+6y^3)dy \\ &= \left[ x^2 - \frac{8}{3}x^3 + \frac{3}{2}x^4 + \frac{x^5}{5} \right]_0^1 + \left[ \frac{3}{2}y^2 - \frac{4y^3}{3} + \frac{8}{2}y^4 \right]_1^0 \\ &= \left[ 1 - \frac{8}{3} + \frac{3}{2} + \frac{1}{5} \right] - \left[ \frac{3}{2} - \frac{4}{3} + \frac{3}{2} \right] \\ &= \left[ \frac{30-80+45+6}{30} \right] - \left[ \frac{9-8+9}{6} \right] \\ &= \frac{1}{30} - \frac{5}{3} = \frac{1-50}{30} = \frac{-49}{30}\end{aligned}$$

∴

$$\oint_C \vec{F} \cdot d\vec{r} = \frac{-49}{30}$$

## 6. Curves in Space

### 6.1 Find $k/z$ for the curve

$$\vec{r}(t) = a \cos t \hat{i} + a \sin t \hat{j} + bt \hat{k}$$

(2010 : 12 Marks)

**Solution:**

Given :

$$\vec{r}(t) = a \cos t \hat{i} + a \sin t \hat{j} + bt \hat{k}$$

∴

$$\frac{d\vec{r}(t)}{dt} = -a \sin t \hat{i} + a \cos t \hat{j} + b \hat{k}$$

$$\frac{d^2\vec{r}(t)}{dt^2} = -a \cos t \hat{i} - a \sin t \hat{j}$$

$$\frac{d^3\vec{r}(t)}{dt^3} = a \sin t \hat{i} - a \cos t \hat{j}$$

$$\frac{d\vec{r}(t)}{dt} \times \frac{d^2\vec{r}(t)}{dt^2} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -a \sin t & a \cos t & b \\ -a \cos t & -a \sin t & 0 \end{vmatrix}$$

$$\begin{aligned}
 &= i(0 + ab\sin t) - j(0 + abc\cos t) + k(a^2 \sin^2 t + a^2 \cos^2 t) \\
 &= ab\sin t i - abc\cos j + a^2 k
 \end{aligned}$$

$$k = \frac{\left| \frac{dr}{dt} \times \frac{d^2r}{dt^2} \right|}{\left| \frac{dr}{dt} \right|^3} = \frac{|ab\sin t i - abc\cos j + a^2 k|}{|-as\sin t i + ac\cos j + b k|^3}$$

$$\begin{aligned}
 &= \frac{(a^2 b^2 \sin^2 t - a^2 b^2 \cos^2 t + a^4)^{1/2}}{(a^2 \sin^2 t + a^2 \cos^2 t + b^2)^{3/2}} = \frac{(a^2 b^2 + a^4)^{1/2}}{(a^2 + b^2)^{3/2}} \\
 &= \frac{a(a^2 + b^2)^{1/2}}{(a^2 + b^2)^{3/2}} = \frac{a}{a^2 + b^2}
 \end{aligned}$$

$$z = \frac{\left[ \frac{dr}{dt}, \frac{d^2r}{dt^2}, \frac{d^3r}{dt^3} \right]}{\left| \frac{dr}{dt} \times \frac{d^2r}{dt^2} \right|^2} = \frac{\left( \frac{dr}{dt} \times \frac{d^2r}{dt^2} \right) \cdot \frac{d^3r}{dt^3}}{\left| \frac{dr}{dt} \times \frac{d^2r}{dt^2} \right|^2}$$

$$\begin{aligned}
 &= \frac{(ab\sin t i - abc\cos j + a^2 k) \cdot (as\sin t i - ac\cos j)}{|ab\sin t i - abc\cos j + a^2 k|^2} \\
 &= \frac{a^2 b \sin^2 t + a^2 b \cos^2 t}{(a^2 b^2 + a^4)} = \frac{a^2 b}{a^2 (a^2 + b^2)} = \frac{b}{a^2 + b^2}
 \end{aligned}$$

$$\frac{k}{z} = \frac{\frac{a}{a^2 + b^2}}{\frac{b}{a^2 + b^2}} = \frac{a}{b}$$

- 6.2 Derive the Frenet-Serret formulae. Define the curvature and torsion for a space curve. Compute them for the space curve

$$x = t, y = t^2, z = \frac{2}{3}t^3$$

Show that the curvature and torsion are equal for this curve.

(2012 : 20 Marks)

**Solution:**

The following three relations are known as Serret-Frenet formulae :

$$t' = kn \quad \dots(i)$$

$$n' = \tau b - kt \quad \dots(ii)$$

$$b' = -\tau n \quad \dots(iii)$$

where,  $k$  is the magnitude of curvature,  $\tau$  is the magnitude of torsion,  $t, n, b$  are the unit tangent vector, the unit principal normal vector and the unit bi-normal vector respectively and '1' denotes the differentiation w.r.t. the arc length  $S$ .

**Derivation :** We know

$$t^2 = 1$$

Differentiating it w.r.t. the arc length  $S$ ,

$$\mathbf{t} \cdot \mathbf{t}' = 0 \Rightarrow \mathbf{t}' \text{ is perpendicular to } \mathbf{t}.$$

The equation of the oscillating plane at a point  $P(r)$  of the curve is

$$[\mathbf{R} - \mathbf{r}, \mathbf{t}, \mathbf{t}'] = 0$$

This equation shows that  $\mathbf{t}'$  lies in the oscillating plane and hence  $\mathbf{t}'$  is perpendicular to the binormal  $\mathbf{b}$  (since oscillating plane is perpendicular to  $\mathbf{b}$ ).

$\therefore \mathbf{t}'$  is parallel to  $\mathbf{b} \times \mathbf{t}$ .

$\Rightarrow \mathbf{t}'$  is parallel to  $\mathbf{n}$ .  $(\because \mathbf{n} = \mathbf{b} \times \mathbf{t})$

$$\Rightarrow \mathbf{t}' = \pm kn$$

But we choose the direction  $\mathbf{n}$  so that the curvature  $k$  is always positive, i.e., we take  $\mathbf{t}' = kn$  which proves (i).

Again, we know that  $b^2 = 1$ .

Differentiating w.r.t. the arc length  $S$ ,

$$\mathbf{b} \cdot \mathbf{b}' = 0 \Rightarrow \mathbf{b}' \text{ is perpendicular to } \mathbf{b}.$$

Also,

$$\mathbf{b} \cdot \mathbf{t} = 0$$

... (iv)

Differentiating w.r.t.  $S$ ,

$$\mathbf{b}' \cdot \mathbf{t} + \mathbf{b} \cdot \mathbf{t}' = 0 \Rightarrow \mathbf{b}' \cdot \mathbf{t} + \mathbf{b} \cdot kn = 0$$

i.e.,

$$\mathbf{b}' \cdot \mathbf{t} = 0$$

$(\because \mathbf{b} \cdot \mathbf{n} = 0)$

$\Rightarrow$

$$\mathbf{b}' \perp \mathbf{t}$$

... (v)

From (iv) and (v),  $\mathbf{b}'$  is perpendicular to both  $\mathbf{t}$  and  $\mathbf{b}$ .

$\Rightarrow \mathbf{b}'$  is parallel to  $\mathbf{b} \times \mathbf{t}$ .

$\Rightarrow \mathbf{b}'$  is parallel to  $\mathbf{n}$ .

We may write,

$$\mathbf{b}' = \tau \mathbf{n}$$

By convention, we have

$$\mathbf{b}' = -\tau \mathbf{n}, \text{ which proves (iii).}$$

We know,

$$\mathbf{n} = \mathbf{b} \times \mathbf{t}$$

Differentiating w.r.t.  $S$ ,

$$\mathbf{n}' = \mathbf{b}' \times \mathbf{t} + \mathbf{b} \times \mathbf{t}'$$

$$= -\tau \mathbf{n} \times \mathbf{t} + \mathbf{b} \times kn$$

(from (i) and (iii))

$$= \tau \mathbf{b} - kt, \text{ which proves (ii).}$$

**Curvature :** The arc rate at which the tangent changes direction as the point  $P(r)$  moves along the curve is called the curvature vector of the curve and its magnitude is denoted by  $k$ .

**Torsion :** The arc rate at which the bi-normal changes direction as the point  $P(r)$  moves along the curve is called the torsion vector of the curve and its magnitude is denoted by  $\tau$ .

Given :

$$x = t, y = t^2, z = \frac{2}{3}t^3$$

Let

$$\vec{r} = \left( t, t^2, \frac{2}{3}t^3 \right)$$

$$\vec{r}' = (1, 2t, 2t^2), \vec{r}'' = (0, 2, 4t)$$

$$\vec{r} \times \vec{r}'' = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 2t & 2t^2 \\ 0 & 2 & 4t \end{vmatrix}$$

$$= \vec{i}(8t^2 - 4t^2) + \vec{j}(0 - 4t) + \vec{k}(2)$$

$$= (4t_1^2 - 4t_1 + 2)$$

$$|\vec{r} \times \vec{r}''| = \sqrt{16t^4 + 4t^2 + 4} = 2(2t^2 + 1)$$

$$[\vec{r}, \vec{r}', \vec{r}''] = \vec{r} \times \vec{r} \times \vec{r}''$$

$$= (4t_1^2 - 4t_1 2)(0, 0, 4) \\ = 8$$

$$\therefore \text{Torsion, } K = \frac{|\vec{r} \times \ddot{\vec{r}}|}{|\vec{r}|} = \frac{2(2t^2 + 1)}{(1 + 4t^2 + 4t^4)^{3/2}}$$

$$= \frac{2}{(2t^2 + 1)^3}$$

$$\text{Curvature, } \tau = \frac{[\vec{r}, \vec{r}, \vec{r}]}{|\dot{\vec{r}} \times \ddot{\vec{r}}|^2}$$

$$= \frac{8}{4(2t^2 + 1)^2} = \frac{2}{(2t^2 + 1)^2}$$

- 6.3 Show that the curve  $\vec{x}(t) = t\hat{i} + \left(\frac{1+t}{t}\right)\hat{j} + \frac{(1-t^2)}{t}\hat{k}$  lies in a plane.

(2013 : 10 Marks)

**Solution:**

For the curve to lie in a plane the binormal  $B$  must point in a constant direction as one moves along the curve.

i.e.,  $\frac{dB}{ds} = 0$

$$\Rightarrow -\tau N = 0 \Rightarrow \tau = 0$$

i.e., torsion must be zero.

$$\tau = \frac{[\dot{\vec{r}} \ddot{\vec{r}} \ddot{\vec{r}}]}{(\dot{\vec{r}} \times \ddot{\vec{r}}) \cdot (\dot{\vec{r}} \times \ddot{\vec{r}})}$$

Now,  $\dot{\vec{r}} = \frac{d\vec{x}}{dt} = \hat{i} + (-t^{-2})\hat{j} - (t^{-2} + 1)\hat{k}$

$$\ddot{\vec{r}} = \frac{d^2\vec{x}}{dt^2} = 2t^{-3}\hat{j} + 2t^{-3}\hat{k}$$

$$\ddot{\vec{r}} = \frac{d^3\vec{x}}{dt^3} = -6t^{-4}\hat{j} - 6t^{-4}\hat{k}$$

$$\ddot{\vec{r}} \times \ddot{\vec{r}} = \begin{vmatrix} i & j & k \\ 0 & 2t^{-3} & 2t^{-3} \\ 0 & -6t^{-4} & -6t^{-4} \end{vmatrix} = 0$$

$$[\dot{\vec{r}} \ddot{\vec{r}} \ddot{\vec{r}}] = 0$$

$$\therefore \tau = 0$$

So, the curve lies in a plane.

- 6.4 Find the curvature vector at any point of the curve  $\vec{r}(t) = t \cos t \hat{i} + t \sin t \hat{j}$ ,  $0 \leq t \leq 2\pi$ . Give its magnitude also.

(2014 : 10 Marks)

**Solution:**

The position vector  $\vec{r}$  of any point on the given curve is  $\vec{r}$ .

$$\vec{r} = t \cos \hat{i} + t \sin \hat{j}$$

i.e.,

$$\vec{r} = (t \cos t, t \sin t)$$

∴

$$\frac{d\vec{r}}{dt} = (\cos t - t \sin t + \sin t + t \cos t)$$

Now,

$$\frac{ds}{dt} = \left| \frac{d\vec{r}}{dt} \right|$$

$$= \sqrt{\cos^2 t + t^2 \sin^2 t - 2t \cos t \sin t + \sin^2 t + t^2 \cos^2 t + 2t \cos t \sin t}$$

$$= \sqrt{1+t^2}$$

Hence,  $T$

$$= \frac{d\vec{r}}{ds} = \frac{d\vec{r}/dt}{ds/dt} = \frac{1}{\sqrt{1+t^2}} (\cos t - t \sin t, \sin t + t \cos t)$$

Differentiating this w.r.t.  $S$ , we get

$$\begin{aligned} \frac{dT}{dS} &= kN = \frac{dT/dt}{dS/dt} \\ &= \frac{1}{\sqrt{1+t^2}} \cdot \frac{dT}{dt} \\ &= \frac{1}{\sqrt{1+t^2}} \left[ (-\sin t - \sin t - t \cos t) \sqrt{1+t^2} - \frac{1}{2\sqrt{1+t^2}} (2t)(\cos t - t \sin t), \right. \\ &\quad \left. (\cos t + \cos t - t \sin t) \sqrt{1+t^2} - \frac{1}{2\sqrt{1+t^2}} (2t)(\sin t + t \cos t) \cdot \frac{1}{(1+t^2)} \right] \\ &= \frac{1}{\sqrt{1+t^2}} \cdot \frac{1}{(1+t^2)} \left[ \frac{(-2\sin t - t \cos t)(1+t^2) - t(\cos t + \sin t)}{\sqrt{1+t^2}}, \right. \\ &\quad \left. \frac{(2\cos t - t \sin t)(1+t^2) - t(t \sin t + t \cos t)}{\sqrt{1+t^2}} \right] \end{aligned}$$

$$kN = \frac{1}{(1+t^2)^2} (-2(1+t^2)(\sin t + \cos t), (2+1+t^2)(\cos t - \sin t))$$

$$kN = \frac{(2+t^2)}{(1+t^2)^2} (\sin t + \cos t) \hat{i} + \left( \frac{2+t}{(1+t)^2} \cdot \cos t - \sin t \right) \hat{j}$$

which is the required curvature vector.

∴

$$k = \|kN\|$$

$$\begin{aligned} &= \sqrt{\left[ \frac{2+t^2}{(1+t^2)^2} \right]^2 ((\sin t + \cos t)^2 + (\cos t - \sin t)^2)} \\ &= \frac{2+t^2}{(1+t^2)^2} \sqrt{2\sin^2 t + 2\cos^2 t} \\ &= \frac{\sqrt{2}(2+t^2)}{(1+t^2)^2} \end{aligned}$$

## 6.5 Find the curvature and torsion of the curve :

$$\vec{r} = a(u - \sin u) \hat{i} + a(1 - \cos u) \hat{j} + bu \hat{k}$$

(2018 : 12 Marks)

Solution:

Given,

$$\vec{r} = a(u - \sin u)\hat{i} + a(1 - \cos u)\hat{j} + bu\hat{k}$$

$$\frac{d\vec{r}}{du} = a(1 - \cos u)\hat{i} + a\sin u\hat{j} + b\hat{k}$$

$$\frac{d^2\vec{r}}{du^2} = a\sin u\hat{i} + a\cos u\hat{j}$$

$$\frac{d^3\vec{r}}{du^3} = a\cos u\hat{i} - a\sin u\hat{j}$$

$$\frac{d\vec{r}}{du} \times \frac{d^2\vec{r}}{du^2} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a(1 - \cos u) & a\sin u & b \\ a\sin u & a\cos u & 0 \end{vmatrix}$$

$$= \hat{i}(-ab\cos u) - \hat{j}(-ab\sin u) + \hat{k}(a^2(\cos u - \cos^2 u) - a^2\sin^2 u)$$

$$= (-ab\cos u)\hat{i} + ab\sin u\hat{j} + \hat{k}(a^2(\cos u - 1))$$

$$\text{Curvature, } K = \frac{\left| \frac{d\vec{r}}{du} \times \frac{d^2\vec{r}}{du^2} \right|}{\left| \frac{d\vec{r}}{du} \right|^3}$$

$$= \frac{\sqrt{a^2b^2\cos^2 u + a^2b^2\sin^2 u + a^2(\cos u - 1)^2}}{\left( \sqrt{a^2(1 - \cos u)^2 + a^2\sin^2 u + b^2} \right)^3}$$

$$= \frac{\sqrt{a^2b^2 + a^4(\cos u - 1)^2}}{\left( \sqrt{2a^2 - 2a^2\cos u + b^2} \right)^3} = \frac{a\sqrt{b^2 + a^2(\cos u - 1)^2}}{\left( \sqrt{2a^2(1 - \cos u) + b^2} \right)^3}$$

$$\text{Torsion, } \tau = \frac{\left[ \frac{d\vec{r}}{du} \frac{d^2\vec{r}}{du^2} \frac{d^3\vec{r}}{du^3} \right]}{\left| \frac{d\vec{r}}{du} \times \frac{d^2\vec{r}}{du^2} \right|^2}$$

$$= \frac{-a^2b\cos^2 u - a^2b\sin^2 u}{a^2b^2 + a^4(\cos u - 1)^2} = \frac{-a^2b}{a^2b^2 + a^4(\cos u - 1)^2}$$

$$K = \frac{a\sqrt{b^2 + a^2(\cos u - 1)^2}}{(2a^2(1 - \cos u) + b^2)^{3/2}}$$

$$\tau = \frac{-a^2b}{a^2b^2 + a^4(\cos u - 1)^2}$$

Thus,

6.6 Find the radius of curvature and radius of torsion of the helix  $x = a \cos u$ ,  $y = a \sin u$ ,  $z = au \tan \alpha$ .  
(2019 : 15 Marks)

Solution:

Given :

$$x = a \cos u, y = a \sin u, z = au \tan \alpha$$

$$r = x\hat{i} + y\hat{j} + z\hat{k}$$

$$r = a \cos u \hat{i} + a \sin u \hat{j} + a \tan \alpha \hat{k}$$

$$k = \frac{\left| \frac{dr}{du} \times \frac{d^2r}{du^2} \right|}{\left| \frac{dr}{du} \right|^3} \quad \dots(2)$$

$$F = \frac{\left| \frac{dr}{du} \times \frac{d^2r}{du^2} \times \frac{d^3r}{du^3} \right|}{\left| \frac{dr}{du} \times \frac{d^2r}{du^2} \right|^2} \quad \dots(3)$$

From (1), differentiate w.r.t.  $u$

$$\frac{dr}{du} = -a \sin u \hat{i} + a \cos u \hat{j} + a \tan \alpha \hat{k}$$

$$\frac{d^2r}{du^2} = -a \cos u \hat{i} - a \sin u \hat{j} + 0 \hat{k}$$

$$\frac{d^3r}{du^3} = a \sin u \hat{i} - a \cos u \hat{j}$$

$$\begin{aligned} \left| \frac{dr}{dt} \right| &= \sqrt{(-a \sin u)^2 + (a \cos u)^2 + (a \tan \alpha)^2} \\ &= \sqrt{a^2(\sin^2 u + \cos^2 u + \tan^2 \alpha)} \\ &= a \sqrt{1 + \tan^2 \alpha} = a \sqrt{\sec^2 \alpha} \end{aligned}$$

$$\left| \frac{dr}{dt} \right| = a \sec \alpha$$

$$\frac{dr}{du} \times \frac{d^2r}{du^2} = \begin{vmatrix} i & j & k \\ -a \sin u & a \cos u & a \tan \alpha \\ -a \cos u & -a \sin u & 0 \end{vmatrix}$$

$$= a^2 \begin{vmatrix} i & j & k \\ -\sin u & \cos u & \tan \alpha \\ -\cos u & -\sin u & 0 \end{vmatrix}$$

$$= a^2 [i(-\sin u \tan \alpha) - \cos u \tan \alpha j + k]$$

$$\frac{dr}{du} \times \frac{d^2r}{du^2} = \sqrt{a^2(\sin^2 u \tan^2 \alpha + \cos^2 u \tan^2 \alpha) + a^4}$$

$$= a^2 \sqrt{\tan^2 \alpha (\sin^2 u + \cos^2 u)} + 1$$

$$= a^2 \sqrt{\tan^2 \alpha + 1} = a^2 \sec \alpha$$

$$\text{Curvature, } K = \frac{a^2 \sec \alpha}{(a \sec \alpha)^3}$$

$$K = \frac{1}{a} \cdot \frac{1}{\sec^2 \alpha} = \frac{1}{a} \cos^2 \alpha$$

$$\text{Radius of curvature} = \frac{1}{K} \cdot a \sec^2 \alpha$$

$$\left[ \frac{dr}{du} \times \frac{d^2r}{du^2} \times \frac{d^3r}{du^3} \right] = \begin{vmatrix} a \sin u & -a \cos u & 0 \\ -a \sin u & a \cos u & a \tan \alpha \\ a \cos u & a \sin u & 0 \end{vmatrix}$$

$$= a^3 \begin{vmatrix} \sin u & -\cos u & 0 \\ -\sin u & \cos u & \tan \alpha \\ \cos u & \sin u & 0 \end{vmatrix} \quad R_2 \rightarrow R_2 + R_1$$

$$= a^3 \begin{vmatrix} \sin u & -\cos u & 0 \\ 0 & 0 & \tan \alpha \\ \cos u & \sin u & 0 \end{vmatrix}$$

$$\begin{aligned} &= a^3 [\sin u(-\sin u \tan \alpha) + \cos u(-\cos u \tan \alpha) + 0] \\ &= a^3 [-(\sin^2 u \tan \alpha + \cos^2 u \tan \alpha)] \\ &= -a^3 [\tan \alpha (\sin^2 u + \cos^2 u)] \\ &= -a^3 \tan \alpha \quad [\because \sin^2 u + \cos^2 u = 1] \end{aligned}$$

$$T = \frac{|-a^3 \tan \alpha|}{(a^2 \sec \alpha)^2} = \frac{a^3 \tan \alpha}{a^4 \sec^2 \alpha}$$

$$T = \frac{1}{a} \cdot \frac{1}{2} \sin^2 \alpha = \frac{1}{2a} \sin 2\alpha$$

$$\text{Radius of tension} = \frac{1}{T} = \frac{a \sec^2 \alpha}{\tan \alpha}$$

$$\text{Radius tension} = \frac{2(a)}{\sin 2\alpha} \quad (\text{Required Solution})$$

## 7. Gauss-Divergence Theorem, Stokes Theorem and Green's Identity

- 7.1 Find the work done in moving the particle one round the ellipse  $\frac{x^2}{25} + \frac{y^2}{16} = 1, z = 0$  under the field of force given by

$$\vec{F} = (2x - y + z)\hat{i} + (x + y - z)\hat{j} + (3x - 2y + 4z)\hat{k}$$

(2009 : 20 Marks)

Solution:

Let  $C: \frac{x^2}{25} + \frac{y^2}{16} = 1, z = 0$  denote the ellipse.

$$\text{Work done} = \oint_C \vec{F} \cdot d\vec{r}$$

By Stoke's theorem

$$\oint_C \vec{F} \cdot d\vec{r} = \iint_S (\nabla \times \vec{F}) \cdot \hat{n} dS$$

where  $S$  is the surface of the ellipse

$$\begin{aligned}\nabla \times \vec{F} &= \left| \begin{array}{ccc} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2x-y+z & x+y-z & 3x-2y+4z \end{array} \right| \\ &= -\hat{i} - 2\hat{j} + 2\hat{k}\end{aligned}$$

For  $S$  normal is along positive  $z$ -axis

$$\therefore \hat{n} = \hat{k}$$

$$(\nabla \times \vec{F}) \cdot \hat{n} = 2$$

$$\begin{aligned}\therefore \iint_S (\nabla \times \vec{F}) \cdot \hat{n} dS &= \iint_S 2 dS = 2 \times \text{Area of ellipse} \\ &= 2\pi ab = 2\pi \cdot 5 \cdot 4 \\ &= 40\pi\end{aligned}$$

**7.2** Using divergent theorem evaluate  $\iint_S \vec{A} \cdot d\vec{S}$  where  $\vec{A} = x^3\hat{i} + y^3\hat{j} + z^3\hat{k}$  and  $S$  is the surface of the sphere  $x^2 + y^2 + z^2 = a^2$ .

(2009 : 20 Marks)

**Solution:**

By gauss divergence theorem

$$\begin{aligned}\iint_S \vec{A} \cdot d\vec{S} &= \iiint_V (\nabla \cdot \vec{A}) dV \\ \nabla \cdot \vec{A} &= \sum \frac{\partial}{\partial x} \hat{i} \cdot \vec{A} = 3x^2 + 3y^2 + 3z^2 \\ &= 3(x^2 + y^2 + z^2)\end{aligned}$$

$S$  is surface of sphere  $x^2 + y^2 + z^2 = a^2$ .

$\therefore V$  is the volume compressing the whole sphere.

$$\begin{aligned}\iint_S \vec{A} \cdot d\vec{S} &= \iiint_V (\nabla \cdot \vec{A}) dV \\ &= \iiint_V 3(x^2 + y^2 + z^2) dV\end{aligned}$$

Converting to spherical polar coordinates.

$$\begin{aligned}x^2 + y^2 + z^2 &= r^2 \\ dV &= r^2 \sin \theta dr d\theta d\phi\end{aligned}$$

$$\begin{aligned}I &= \int_{\phi=0}^{2\pi} \int_{\theta=0}^{2\pi} \int_{r=0}^a 3r^2 \cdot r^2 \sin \theta dr d\theta d\phi \\ &= \int_0^{2\pi} \left( \int_0^{2\pi} \left( \int_0^a (3r^4 dr) \sin \theta d\theta \right) d\phi \right)\end{aligned}$$

$$\begin{aligned}
 &= \int_0^{\pi} \left[ \frac{3a^5}{5} \int_0^{\pi} \sin \theta d\theta \right] d\phi \\
 &= \frac{3a^5}{5} [-\cos \theta]_0^{\pi} \pi = \frac{6\pi a^5}{5}
 \end{aligned}$$

- 7.3 Find the value of  $\iint_S (\nabla \times \vec{F}) \cdot d\vec{S}$  taken over the upper portion of the surface  $x^2 + y^2 - 2ax + az = 0$  and the bounding curve lies on the plane  $z = 0$  when

$$\vec{F} = (y^2 + z^2 - x^2)\hat{i} + (z^2 + x^2 - y^2)\hat{j} + (x^2 + y^2 - z^2)\hat{k}$$

(2009 : 20 Marks)

**Solution:**The given surface  $S$ ,

$$\begin{aligned}
 x^2 + y^2 - 2ax + az &= 0 \\
 \Rightarrow (x-a)^2 + y^2 &= a^2 - az
 \end{aligned}$$

is of an inverted paraboloid as there is maximum value of  $z_{\max} = a$ . Assumption  $a = +ve$ . Had 'a' been -ve, it would not be inverted paraboloid.

Consider the part of the paraboloid enclosed by  $S_1$  and  $S_2$  i.e., the circle of intersection in the XY-plane.

$$S_2(x-a)^2 + y^2 = a^2, z=0$$

These surfaces enclose a volume.


Using gauss divergence theorem

$$\begin{aligned}
 \iint_{S=S_1+S_2} (\nabla \times \vec{F}) \cdot d\vec{S} &= \iiint_V \nabla \cdot (\nabla \times \vec{F}) dV \\
 &= 0 \text{ as } \nabla \cdot (\nabla \times \vec{F}) = 0
 \end{aligned}$$

$$\therefore \iint_{S_1} (\nabla \times \vec{F}) \cdot dS = - \iint_{S_2} (\nabla \times \vec{F}) \cdot dS$$

Now for  $S_2$ ,

$\hat{n} = -\hat{k}$  (outward normal is towards downward direction).



$$\begin{aligned}
 \nabla \times \vec{F} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y^2 + z^2 - x^2 & z^2 + x^2 - y^2 & x^2 + y^2 - z^2 \end{vmatrix} \\
 &= (2y - 2z)\hat{i} + 2(z - x)\hat{j} + 2(x - y)\hat{k}
 \end{aligned}$$

$$\therefore (\nabla \times \vec{F}) \cdot \hat{n} = 2(y - x)$$

$$\therefore I = - \iint_{S_2} 2(y - x) dS = \iint_{S_2} 2(x - y) dS$$

Converting to polar coordinates.

$$x = a + r \cos \theta$$

$$y = r \sin \theta$$

$$dS = r dr d\theta$$

$$I = 2 \int_{\theta=0}^{2\pi} \int_{r=0}^a (a + r \cos \theta - r \sin \theta) r dr d\theta$$

$$\begin{aligned}
 &= 2 \int_{\theta=0}^{2\pi} \left[ \frac{ar^2}{2} + \frac{r^3}{3}(\cos\theta - \sin\theta) \right]_0^{\theta} d\theta \\
 &= 2 \int_{\theta=0}^{2\pi} \frac{a^3}{2} + \frac{a^3}{3}(\cos\theta - \sin\theta) d\theta \\
 &= 2 \cdot \frac{a^3}{2} \cdot 2\pi \text{ (as integral of } \sin\theta \text{ and } \cos\theta \text{ over } 0 \text{ to } 2\pi \text{ is zero).} \\
 &= 2\pi a^3
 \end{aligned}$$


## 7.4 Verify Green's Theorem for

$$e^{-x} \sin y dx + e^{-x} \cos y dy$$

the path of integration being the boundary of the square whose vertices are  $(0, 0)$ ,  $\left(\frac{\pi}{2}, 0\right)$ ,  $\left(\frac{\pi}{2}, \frac{\pi}{2}\right)$  and  $\left(0, \frac{\pi}{2}\right)$ .

(2010 : 20 Marks)

Solution:



By Green's Theorem,

$$\int (M dx + N dy) = \iint \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx dy$$

Given equation is  $e^{-x} \sin y dx + e^{-x} \cos y dy$ 

Here,

$$M = e^{-x} \sin y$$

$$N = e^{-x} \cos y$$

$$\therefore \int (e^{-x} \sin y dx + e^{-x} \cos y dy) = \iint \left( \frac{\partial e^{-x} \cos y}{\partial x} - \frac{\partial e^{-x} \sin y}{\partial y} \right) dx dy \quad (\text{By Green's Theorem})$$

$$\begin{aligned}
 \text{R.H.S.} : & \iint \left( \frac{\partial e^{-x} \cos y}{\partial x} - \frac{\partial e^{-x} \sin y}{\partial y} \right) dx dy \\
 &= \iint (-e^{-x} \cos y - e^{-x} \cos y) dx dy \\
 &= -2 \int_{y=0}^{\pi/2} \int_{x=0}^{\pi/2} e^{-x} \cos y dx dy \\
 &= -2 \left[ \frac{e^{-x}}{-1} \right]_0^{\pi/2} [\sin y]_0^{\pi/2} = \frac{-2}{+1} (e^{-\pi/2} - 1)(1 - 0) = 2(e^{-\pi/2} - 1)
 \end{aligned}$$

$$\text{L.H.S. : } \int e^{-x} \sin y dx + e^{-x} \cos y dy$$

In figure, from  $A$  to  $B$ :

$$\int (e^{-x} \sin y dx + e^{-x} \cos y dy) = \int 0 dx + 0 = 0$$

as

$$\sin y = 0$$

and

$$dy = 0$$

from  $B$  to  $C$ :

$$\begin{aligned} \int (e^{-x} \sin y dx + e^{-x} \cos y dy) &= 0 + \int_{y=0}^{\pi/2} e^{-\pi/2} \cos y dy \\ &= e^{-\pi/2} [\sin y]_0^{\pi/2} = e^{-\pi/2} \end{aligned} \quad \left( dx = 0; x = \frac{\pi}{2} \right)$$

from  $C$  to  $D$ :

$$\begin{aligned} \int (e^{-x} \sin y dx + e^{-x} \cos y dy) &= \int_{x=\frac{\pi}{2}}^0 e^{-x} dx \\ &= -[e^{-x}]_{\pi/2}^0 = -[1 - e^{-\pi/2}] = e^{-\pi/2} - 1 \end{aligned} \quad \left( dy = 0, y = \frac{\pi}{2} \right)$$

from  $D$  to  $A$ :

$$\begin{aligned} \int (e^{-x} \sin y dx + e^{-x} \cos y dy) &= 0 + \int e^{-x} \cos y dy \\ &= \int_{y=\frac{\pi}{2}}^0 \cos y dy = [\sin y]_{\pi/2}^0 = 0 - 1 = -1 \end{aligned} \quad (dx = 0, x = 0)$$

$\therefore$  Line integral along  $A \rightarrow B \rightarrow C \rightarrow D \rightarrow A$  is  $0 + e^{-\pi/2} + e^{-\pi/2} - 1 - 1$

$$\begin{aligned} &= 2e^{-\pi/2} - 2 = 2(e^{-\pi/2} - 1) \\ &= \text{R.H.S.} \end{aligned}$$

$$\therefore \text{L.H.S.} = \text{R.H.S.}$$

Green's theorem is verified.

7.5 If  $\vec{u} = 4y\hat{i} + x\hat{j} + 2z\hat{k}$ , calculate the double integral  $\iint (\nabla \times \vec{u}) \cdot d\vec{s}$  over the hemisphere given by

$$x^2 + y^2 + z^2 = a^2, z \geq 0.$$

(2011 : 15 Marks)

Solution:

Let  $C$  be the boundary of the sphere  $x^2 + y^2 + z^2 = a^2, z \geq 0$ . Then  $C$  is a circle in the  $xy$ -plane of radius  $a$  and centre origin.

The equations of the curve  $C$  are

$$x^2 + y^2 = a^2, z = 0$$

Suppose  $x = a \cos t, y = a \sin t, z = 0; 0 \leq t < 2\pi$  are parametric equations of  $C$ .

Then using Stoke's theorem,

$$\begin{aligned} \iint_S \text{Curl } u \cdot d\vec{s} &= \oint_C \vec{u} \cdot d\vec{r} \\ &= \oint_C (4y\hat{i} + x\hat{j} + 2z\hat{k}) \cdot (dx\hat{i} + dy\hat{j} + dz\hat{k}) \\ &= \oint_C 4ydx + xdy + 2zdz \end{aligned}$$

$$\begin{aligned}
 &= \oint_C (4ydx + xdy), \text{ as on } C, z = 0 \text{ and } dz = 0. \\
 &= \int_{t=0}^{2\pi} 4a \sin t (-a \sin t) dt + \int_{t=0}^{2\pi} a \cos t (a \cos t) dt \\
 &= -2a^2 \int_0^{2\pi} 2 \sin^2 t dt + \frac{a^2}{2} \int_0^{2\pi} 2 \cos^2 t dt \\
 &= -2a^2 \int_0^{2\pi} (1 - \cos 2t) dt + \frac{a^2}{2} \int_0^{2\pi} (1 + \cos 2t) dt \\
 &= -2a^2 \left[ t + \frac{\sin 2t}{2} \right]_0^{2\pi} + \frac{a^2}{2} \left[ t - \frac{\sin 2t}{2} \right]_0^{2\pi} \\
 &= -2a^2 [2\pi + 0] + \frac{a^2}{2} [2\pi - 0] \\
 &= -4\pi a^2 + \pi a^2 = -3\pi a^2
 \end{aligned}$$

### 7.6 Verify gauss' Divergence Theorem for the vector

$$\vec{V} = x^2 \hat{i} + y^2 \hat{j} + z^2 \hat{k}$$

taken over the cube

$$0 \leq x, y, z \leq 1$$

(2011 : 15 Marks)

**Solution:**

We need to prove that

$$\iiint_V \nabla \cdot \vec{V} dV = \iint_S \vec{V} \cdot \vec{n} dS$$

Given :

$$\vec{V} = x^2 \hat{i} + y^2 \hat{j} + z^2 \hat{k} \text{ taken over the cube } 0 \leq x, y, z \leq 1.$$

$$\begin{aligned}
 \nabla \cdot \vec{V} &= \left( \hat{i} \frac{\partial}{\partial x} + \hat{j} \frac{\partial}{\partial y} + \hat{k} \frac{\partial}{\partial z} \right) \cdot (x^2 \hat{i} + y^2 \hat{j} + z^2 \hat{k}) \\
 &= 2x + 2y + 2z = 2(x + y + z)
 \end{aligned}$$

$$\therefore \iiint_V \nabla \cdot \vec{V} dV = \iiint_V 2(x + y + z) dV$$


$$= 2 \int_{x=0}^1 \int_{y=0}^1 \int_{z=0}^1 (x + y + z) dz dy dx$$

$$= 2 \int_{x=0}^1 \int_{y=0}^1 \left( xz + yz + \frac{z^2}{2} \right)_0^1 dy dx$$

$$= 2 \int_{x=0}^1 \left( xy + \frac{y^2}{2} + \frac{y}{2} \right)_0^1 dx$$

$$= 2 \int_{x=0}^1 \left( x + \frac{1}{2} + \frac{1}{2} \right) dx = 2 \int_{x=0}^1 (x + 1) dx$$

$$= 2 \left( \frac{x^2}{2} + x \right)_0^1 = 2 \left( \frac{1}{2} + 1 \right) = 2 \left( \frac{3}{2} \right) = 3$$



Now we will calculate  $\iint_S \vec{V} \cdot \hat{n} dS$  over the six faces of the cube.

Over the face DEFG,  $\hat{n} = \hat{i}$ ,  $x = 1$

$$\begin{aligned}\therefore \iint_{DEFG} \vec{V} \cdot \hat{n} dS &= \int_{z=0}^1 \int_{y=0}^1 (x^2 \hat{i} + y^2 \hat{j} + z^2 \hat{k}) \cdot \hat{i} dy dz \\ &= \int_{z=0}^1 \int_{y=0}^1 1 \cdot dy dz = \int_{z=0}^1 (y)_0^1 dz \\ &= \int_{z=0}^1 dz = (z)_0^1 = 1\end{aligned}$$

Over the face ABCO,  $\hat{n} = -\hat{i}$ ,  $x = 0$

$$\therefore \iint_{ABCO} \vec{V} \cdot \hat{n} dS = \int_{z=0}^1 \int_{y=0}^1 0 dy dz = 0$$

Over the face ABEF,  $\hat{n} = \hat{j}$ ,  $y = 1$

$$\begin{aligned}\therefore \iint_{ABEF} \vec{V} \cdot \hat{n} dS &= \int_{z=0}^1 \int_{x=0}^1 (x^2 \hat{i} + y^2 \hat{j} + z^2 \hat{k}) \cdot \hat{j} dx dz \\ &= \int_{z=0}^1 \int_{x=0}^1 dx dz = 1\end{aligned}$$

Over the face OGDC,  $\hat{n} = -\hat{j}$ ,  $y = 0$

$$\therefore \iint_{OGDC} \vec{V} \cdot \hat{n} dS = 0$$

Over the face BCDE,  $\hat{n} = \hat{k}$ ,  $z = 1$

$$\therefore \iint_{BCDE} \vec{V} \cdot \hat{n} dS = \int_{x=0}^1 \int_{y=0}^1 dx dy = 1$$

Over the face AFGO,  $\hat{n} = -\hat{k}$ ,  $z = 0$

$$\therefore \iint_{AFGO} \vec{V} \cdot \hat{n} dS = 0$$

Adding the six surface integrals, we get,

$$\iint_S \vec{F} \cdot \hat{n} dS = 1 + 0 + 1 + 0 + 1 + 0 = 3$$

Hence, the theorem is verified.

### 7.7 Verify Green's theorem in the plane for

$$\oint_C [(xy + y^2)dx + x^2dy]$$

where  $C$  is the closed curve of the region bounded by  $y = x$  and  $y = x^2$ .

(2012 : 20 Marks)


**Solution:**

By Green's theorem, if  $R$  is the plane region bounded by a simple closed curve  $C$ , then

$$\iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx dy = \oint_C (M dx + N dy)$$

Here,

$$M = xy + y^2, N = x^2$$



$$\text{We have, } \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx dy$$

$$\begin{aligned} &= \iint_R \left( \frac{\partial}{\partial x}(x^2) - \frac{\partial}{\partial y}(xy + y^2) \right) dx dy \\ &= \iint_R (\partial x - x - \partial y) dx dy = \iint_R (x - \partial y) dx dy \\ &= \int_{x=0}^1 \int_{y=x^2}^x (x - \partial y) dy dx = \int_{x=0}^1 [xy - y^2]_{y=x^2}^x dx \\ &= \int_0^1 (x^4 - x^3) dx = \frac{-1}{20} \end{aligned}$$

Again, along  $y = x^2$ ,  $dy = 2x dx$

$\therefore$  along  $y = x$ , the line integral becomes

$$\int_1^0 (x \cdot x + x^2) dx + x^2 dx = \int_1^0 3x^2 dx = -1$$

$$\therefore \text{The required line integral} = \frac{19}{20} - 1 = -\frac{1}{20}$$

Hence, the theorem is verified.

7.8 If  $\vec{F} = y\vec{i} + (x - 2xz)\vec{j} - xy\vec{k}$ , evaluate

$$\iint_S (\nabla \times \vec{F}) \cdot \vec{n} dS$$

where S is the surface of the sphere  $x^2 + y^2 + z^2 = a^2$  above the xy-plane.

(2012 : 20 Marks)

Solution:

$$\begin{aligned} \nabla \times \vec{F} &= \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y & x - 2xz & -xy \end{vmatrix} \\ &= \vec{i}(-x + 2z) + \vec{j}(0 + y) + \vec{k}(1 - 2z - 1) \\ &= x\vec{i} + y\vec{j} - 2z\vec{k} \end{aligned} \quad \dots(i)$$

A normal to the surface  $x^2 + y^2 + z^2 = a^2$  is

$$\begin{aligned} \nabla(x^2 + y^2 + z^2) &= \left( \vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z} \right)(x^2 + y^2 + z^2) \\ &= 2x\vec{i} + 2y\vec{j} + 2z\vec{k} \end{aligned}$$

$\therefore \vec{i}$  = a unit normal to the surface  $x^2 + y^2 + z^2 = a^2$  is given by

$$\vec{i} = \frac{2x\vec{i} + 2y\vec{j} + 2z\vec{k}}{\sqrt{4x^2 + 4y^2 + 4z^2}} = \frac{x\vec{i} + y\vec{j} + z\vec{k}}{a} \quad [\because x^2 + y^2 + z^2 = a^2]$$

The projection of S on the xy-plane is the region R bounded by the circle  $x^2 + y^2 = a^2$ ,  $z = 0$ .

Then

$$\begin{aligned} \iint_S (\nabla \times \vec{F}) \cdot \vec{i} dS &= \iint_R (\nabla \times \vec{F}) \cdot \vec{i} \frac{dxdy}{|\vec{i} \cdot \vec{k}|} \\ &= \iint_R (x\vec{i} + y\vec{j} - 2z\vec{k}) \cdot \left( \frac{x\vec{i} + y\vec{j} + z\vec{k}}{a} \right) \frac{dxdy}{z/a} \\ &= \int_{-a}^a \int_{-\sqrt{a^2-x^2}}^{\sqrt{a^2-x^2}} \frac{3(x^2 + y^2) - 2a^2}{\sqrt{a^2 - x^2 - y^2}} dy dx \end{aligned}$$

Changing to polar co-ordinates, by substituting,

$$x = r \cos \theta, y = r \sin \theta$$

and

$$dxdy = r dr d\theta$$

$$\begin{aligned} \iint_S (\nabla \times \vec{F}) \cdot \vec{i} dS &= \int_{0}^{2\pi} \int_{0}^a \frac{3r^2 - 2a^2}{\sqrt{a^2 - r^2}} \cdot r dr d\theta \\ &= \int_{0}^{2\pi} \int_{0}^a \frac{3(r^2 - a^2) + a^2}{\sqrt{a^2 - r^2}} \cdot r dr d\theta \end{aligned}$$

$$\begin{aligned}
 &= \int_{\theta=0}^{2\pi} \int_0^a \left( -3r\sqrt{a^2 - r^2} + \frac{dr}{\sqrt{a^2 - r^2}} \right) dr d\theta \\
 &= \int_{\theta=0}^{2\pi} \left[ (a^2 - r^2)^{3/2} - a^2 \sqrt{a^2 - r^2} \right]_0^a d\theta \\
 &= \int_{\theta=0}^{2\pi} (a^3 - a^3) d\theta = 0
 \end{aligned}$$

- 7.9 By using Divergence Theorem of Gauss, evaluate the surface integral  $\iint_S (a^2x^2 + b^2y^2 + c^2z^2)^{-1/2} dS$   
where  $S$  is the surface of the ellipsoid  $ax^2 + by^2 + cz^2 = 1$ ,  $a, b, c$  being all positive constants.  
(2013 : 15 Marks)

**Solution:**

We have to write the integral in the form  $\iint_S \vec{F} \cdot \hat{n} dS$

So we first find suitable  $\vec{F}$

Now

$\phi(x, y, z) = ax^2 + by^2 + cz^2 = 1$  is the given surface.

Normal to surface  $= \nabla \phi = 2ax\hat{i} + 2by\hat{j} + 2cz\hat{k}$

$$\therefore \hat{n} = \text{unit normal} = \frac{2ax\hat{i} + 2by\hat{j} + 2cz\hat{k}}{(a^2x^2 + b^2y^2 + c^2z^2)^{1/2}}$$

Taking

$$\vec{F} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\vec{F} \cdot \hat{n} = \frac{ax^2 + by^2 + cz^2}{(a^2x^2 + b^2y^2 + c^2z^2)^{1/2}} = \frac{1}{(a^2x^2 + b^2y^2 + c^2z^2)^{1/2}} \text{ on } S$$

By Gauss divergence theorem

$$\begin{aligned}
 \iiint_V \vec{F} \cdot \hat{n} dV &= \iiint_V (\nabla \cdot \vec{F}) dV = \iiint_V (\partial_x x + \partial_y y + \partial_z z) dV \\
 &= 3 \iiint_V dV = 3 \cdot \frac{4}{3} \pi \frac{1}{\sqrt{abc}} \\
 &= \frac{4\pi}{\sqrt{abc}}
 \end{aligned}$$

- 7.10 Use Stoke's theorem to evaluate the line integral  $\int_C (-y^2 dx + x^2 dy - z^2 dz)$  where  $C$  is the intersection of the cylinder  $x^2 + y^2 = 1$  and the plane  $x + y + z = 1$ .

(2013 : 15 Marks)

**Solution:**

By Stoke's theorem

$$\int_C \vec{F} \cdot d\vec{r} = \iint_S (\nabla \times \vec{F}) \cdot \hat{n} dS$$

$$\begin{aligned}
 \vec{F} \cdot d\vec{r} &= F_x dx + F_y dy + F_z dz \\
 &= -y^2 dx + x^2 dy - z^2 dz
 \end{aligned}$$

$$\vec{F} = -y^2\hat{i} + x^2\hat{j} - z^2\hat{k}$$



$$\nabla \times \vec{F} = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ -y^3 & x^3 & -z^3 \end{vmatrix} = 3(x^2 + y^2)\hat{k}$$

$\delta$  is part of the plane  $x + y + z = 1$ .

$$\therefore \text{Normal to } S = \frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}} \quad [(1, 1, 1) \text{ are direction ratio of normal}]$$

$$\begin{aligned} \therefore \int_C \vec{F} \cdot d\vec{r} &= \iint_S 3(x^2 + y^2)\hat{k} \cdot \frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}} dS \\ &= \iint_S \sqrt{3}(x^2 + y^2) dS \\ &= \iint_S \sqrt{3}(x^2 + y^2) dA \quad (\text{where } dA \text{ is projection on } XY\text{plane}) \\ &= \int_{\theta=0}^{2\pi} \int_{r=0}^1 \sqrt{3}r^2 r dr d\theta = \frac{\sqrt{3}}{4} \cdot 2\pi = \frac{\sqrt{3}}{2}\pi \end{aligned}$$

7.11 Evaluate the Stoke's theorem  $\int_{\Gamma} (ydx + zdy + xdz)$  where  $\Gamma$  is the curve given by  $x^2 + y^2 + z^2 - 2ax - 2ay = 0$ ,  $x + y = 2a$ , starting from  $(2a, 0, 0)$  and then going below the  $z$ -plane.

(2014 : 20 Marks)

Solution:

The centre of the sphere  $x^2 + y^2 + z^2 - 2ax - 2ay = 0$  is the point  $(a, a, 0)$ . Since the plane  $x + y = 2a$  passes through the point  $(a, a, 0)$ . Therefore the circle  $C$  is great circle of this sphere

$$\therefore \text{Radius of the circle } C = \text{Radius of the sphere} = \sqrt{(a^2 + a^2)} = a\sqrt{2}$$

$$\begin{aligned} \text{Now } \int_C (ydx + zdy + xdz) &= \int_C (yi + zj + xk) \cdot dr \\ &= \iint_S [\text{curl}(yi + zj + xk)] \cdot n ds \end{aligned}$$

where  $S$  is any surface of which circle  $C$  is boundary [Stoke's theorem]

$$\text{Now } \text{curl } (yi + zj + xk) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y & z & x \end{vmatrix} = -\hat{i} - \hat{j} - \hat{k} = -(\hat{i} + \hat{j} + \hat{k})$$

Let us take  $S$  as the surface of the plane  $x + y = 2a$  bounded by the circle  $C$ . Then a vector normal to  $S$  is  $\text{grad } (x + y) = \hat{i} + \hat{j}$ .

$$\therefore n = \text{Unit normal to } \delta = \frac{1}{2}(\hat{i} + \hat{j})$$

$$\begin{aligned} \therefore \int_C (ydx + zdy + xdz) &= \iint_C (\hat{i} + \hat{j} + \hat{k}) \cdot \left( \frac{1}{\sqrt{2}}\hat{i} + \frac{1}{\sqrt{2}}\hat{j} \right) ds \\ &= \frac{-2}{\sqrt{2}} \iint_S ds = \frac{-2}{\sqrt{2}} \text{ (area of the circle of radius } a\sqrt{2} \text{)} \\ &= -\sqrt{2}(2\pi a^2) \end{aligned}$$

7.12 Evaluate :  $\oint_C e^{-x}(\sin y \, dx + \cos y \, dy)$ , where  $C$  is the rectangle with vertices  $(0, 0)$ ,  $(\pi, 0)$ ,  $\left(\pi, \frac{\pi}{2}\right)$ ,  $\left(0, \frac{\pi}{2}\right)$ .

(2015 : 12 Marks)

**Solution:**

By Green's theorem in plane,

$$\oint_C (M \, dx + N \, dy) = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx \, dy$$

Here,

$$M = e^{-x} \sin y,$$

$$N = e^{-x} \cos y$$

$$\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} = -e^{-x} \cos y - e^{-x} \cos y = -2e^{-x} \cos y$$


$$\therefore \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx \, dy = \iint_R -2e^{-x} \cos y \, dx \, dy$$

$$= -2 \int_{x=0}^{\pi} e^{-x} dx \int_{y=0}^{\pi/2} \cos y \, dy$$

$$= -2[-e^{-x}]_0^\pi [\sin y]_0^{\pi/2}$$

$$= 2(e^{-\pi} - 1) \cdot \left( \sin \frac{\pi}{2} - \sin 0 \right)$$

$$= 2(e^{-\pi} - 1)$$



7.13 Prove that  $\oint_C f d\vec{r} = \iint_S d\vec{S} \times \nabla f$ .

(2016 : 15 Marks)

**Solution:**Let  $\vec{p}$  be any arbitrary constant vector and let  $\vec{F} = f\vec{p}$ .Using Stoke's theorem for  $\vec{F}$ .

$$\begin{aligned} \oint_C \vec{F} \cdot d\vec{r} &= \iint_S (\nabla \times \vec{F}) \cdot \hat{n} dS = \iint_S (\nabla \times f\vec{p}) \cdot \hat{n} dS \\ &= \iint_S [\nabla f \times \vec{p} + f(\nabla \times \vec{p})] \cdot d\vec{S} \\ &= \iint_S (\nabla f \times \vec{p}) \cdot d\vec{S} \quad (\because \text{Curl } \vec{p} = 0) \end{aligned}$$

$$\therefore \oint_C f\vec{p} \cdot d\vec{r} = \iint_S (\nabla f \times \vec{p}) \cdot d\vec{S}$$

$$\Rightarrow \vec{p} \cdot \left[ \oint_C f d\vec{r} - \iint_S d\vec{S} \times \nabla f \right] = 0 \quad [ \because (\nabla f \times \vec{p}) \cdot d\vec{S} = [\nabla f \vec{p} d\vec{S}] = [d\vec{S} \nabla f \vec{p}] = (d\vec{S} \times \nabla f) \cdot \vec{p} ]$$

Since,  $\vec{p}$  was arbitrary, hence

$$\oint_C f d\vec{r} = \iint_S d\vec{S} \times \nabla f$$

- 7.14 (i) Evaluate the integral:  $\iint_S \vec{F} \cdot \hat{n} dS$  where  $\vec{F} = 3xy^2\mathbf{i} + (yx^2 - y^3)\mathbf{j} + 3zx^2\mathbf{k}$  and  $S$  is a surface of the cylinder  $y^2 + z^2 \leq 4$ ,  $-3 \leq x \leq 3$ , using divergence theorem.

(2017 : 9 Marks)

- (ii) Using Green's theorem, evaluate  $\int_C \vec{F}(\vec{r}) \cdot d\vec{r}$  counterclockwise, where

$$\vec{F}(\vec{r}) = (x^2 + y^2)\mathbf{i} + (x^2 - y^2)\mathbf{j} \text{ and } d\vec{r} = dx\mathbf{i} + dy\mathbf{j}$$

and curve  $C$  is the boundary of the region,  $R = \{(x, y) : 1 \leq y \leq 2 - x^2\}$ .

(2017 : 8 Marks)

**Solution:**

- (i) Let  $V$  be the volume enclosed by surface,  $S$ . By divergence theorem,

$$\iint_S \vec{F} \cdot \hat{n} dS = \iiint_V \operatorname{div} \vec{F} \cdot dV \quad \dots(i)$$

$$\begin{aligned} \operatorname{div} \vec{F} &= \frac{\partial}{\partial x}(3xy^2) + \frac{\partial}{\partial y}(yx^2 - y^3) + \frac{\partial}{\partial z}(3zx^2) \\ &= 4x^2 \end{aligned}$$

$$\therefore \iint_S \vec{F} \cdot \hat{n} dS = \iiint_V 4x^2 dV, \quad V \text{ bounded by } y^2 + z^2 = 4, x = \pm 3$$

In cylindrical co-ordinates

$$y = r \cos \theta, z = r \sin \theta, x = x$$

$$r : 0 \text{ to } 2, \theta : 0 \text{ to } 2\pi, x : -3 \text{ to } 3$$

$$\begin{aligned} \iint_S \vec{F} \cdot \hat{n} dS &= 4 \int_{r=0}^2 \int_{\theta=0}^{2\pi} \int_{x=-3}^3 x^2 r dr d\theta dx \\ &= 4 \int_0^2 r dr \int_0^{2\pi} d\theta \int_{-3}^3 x^2 dx \\ &= 4 \left[ \frac{r^2}{2} \right]_0^2 [ \theta ]_0^{2\pi} \left[ \frac{x^3}{3} \right]_{-3}^3 \\ &= 4 \times \frac{4}{2} \times 2\pi \times 2 \times \frac{27}{3} = 288\pi \end{aligned}$$

(ii) We write


$$\begin{aligned} \int_C \vec{F}(\vec{r}) \cdot d\vec{r} &= \int_C (x^2 + y^2) dx + (x^2 - y^2) dy \\ &= \int_C M dx + N dy \\ M &= x^2 + y^2 \\ N &= x^2 - y^2 \end{aligned}$$

$C$ : Curve  $ABCA$

$S$ : Area shaded bounded by  $y \geq 1$  and  $y \leq 2 - x^2$

Using Green's Theorem

$$\begin{aligned} \int_C M dx + N dy &= \iint_S \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dS \\ &= 2 \iint_S (x - y) dxdy \end{aligned}$$



$$\begin{aligned}
 &= 2 \int_{x=-\sqrt{2-y}}^{\sqrt{2-y}} \int_{y=1}^2 (x-y) dx dy \\
 &= 2 \int_1^2 \left[ \frac{x^2}{2} - yx \right]_{x=-\sqrt{2-y}}^{\sqrt{2-y}} dy \\
 &= -4 \int_1^2 y \sqrt{2-y} dy \quad \dots(i) \\
 I &= -4 \int_1^2 y \sqrt{2-y} dy
 \end{aligned}$$

Put  $y = t + 1$ ,  $dy = dt$

$$I = -4 \int_0^1 (1+t) \sqrt{1-t} dt$$

Let  $t = \sin^2 \theta$ ,

$$dt = 2 \sin \theta \cos \theta d\theta, \theta : 0 \rightarrow \frac{\pi}{2}$$

$$\begin{aligned}
 &= -4 \int_0^{\pi/2} (1+\sin^2 \theta) \cos \theta \cdot 2 \sin \theta \cos \theta d\theta \\
 &= -8 \left[ \int_0^{\pi/2} \sin \theta \cos^2 \theta d\theta + \int_0^{\pi/2} \sin^3 \theta \cos^2 \theta d\theta \right] \\
 &= -8 \left[ \frac{\Gamma\left(\frac{1+1}{2}\right)\Gamma\left(\frac{2+1}{2}\right)}{2\Gamma\left(\frac{1+2+2}{2}\right)} + \frac{\Gamma\left(\frac{3+1}{2}\right)\Gamma\left(\frac{2+1}{2}\right)}{2\Gamma\left(\frac{3+2+2}{2}\right)} \right] \\
 &= -8 \left[ \frac{1}{3} + \frac{2}{15} \right] = -\frac{56}{15}
 \end{aligned}$$

$$\therefore \int_0^{\pi/2} \sin^p \theta \cos^q \theta d\theta = \frac{\Gamma\left(\frac{p+1}{2}\right)\Gamma\left(\frac{q+1}{2}\right)}{2\Gamma\left(\frac{p+q+2}{2}\right)}$$

7.15 If  $S$  is the surface of sphere  $x^2 + y^2 + z^2 = a^2$ , then evaluate

$$\iint (x+2)dydz + (y+2)dzdx + (z+2)dxdy$$

using Gauss's Divergence Theorem.

(2018 : 12 Marks)

**Solution:**

$$\text{Given: } \iint (x+z)dydz + (y+z)dzdx + (x+y)dxdy$$

$$= \iint \underbrace{((x+z)\hat{i} + (y+z)\hat{j} + (x+y)\hat{k}) \cdot \hat{n} dS}_{\vec{F}}$$

Here,

$$\hat{n} = \frac{2x\hat{i} + 2y\hat{j} + 2z\hat{k}}{\sqrt{4x^2 + 4y^2 + 4z^2}} = \frac{x\hat{i} + y\hat{j} + z\hat{k}}{a}$$

By Gauss's Divergence theorem,

$$\iint \vec{F} \cdot \hat{n} dS = \iiint \nabla \cdot \vec{F} dV$$

$$\nabla \cdot \vec{F} = 1 + 1 + 0 = 2$$

$$\iiint \nabla \cdot \vec{F} dV = \iiint 2 dV = 2 \iiint dV = 2 \times \frac{4}{3} \pi a^3$$

as volume of sphere is  $\frac{4\pi}{3} a^3$ .

$$\iiint \nabla \cdot \vec{F} dV = \frac{8\pi}{3} a^3$$

- 7.16 Evaluate the line integral  $\int_C -y^3 dx + x^3 dy + z^3 dz$  using Stokes' theorem. Here,  $C$  is the intersection of the cylinder  $x^2 + y^2 = 1$  and the plane  $x + y + z = 1$ . The orientation on  $C$  corresponds to counter-clockwise motion in  $xy$ -plane.

(2018 : 13 Marks)

**Solution:**


Given :

$$\int_C -y^3 dx + x^3 dy + z^3 dz = \int_C (-y^3 \hat{i} + x^3 \hat{j} + z^3 \hat{k}) \cdot (dx\hat{i} + dy\hat{j} + dz\hat{k})$$

$$= \int_C \vec{F} \cdot d\vec{r}$$

where,

$$\vec{F} = -y^3 \hat{i} + x^3 \hat{j} + z^3 \hat{k}$$



By Stoke's theorem,

$$\int_C \vec{F} \cdot d\vec{r} = \iint_S (\nabla \times \vec{F}) \cdot d\vec{S}$$

$$\nabla \times \vec{F} = 3(x^2 + y^2) \hat{k}$$

$$\hat{n} = \frac{\nabla(x + y + z - 1)}{\|\nabla(x + y + z - 1)\|} = \frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$$

$$\begin{aligned}\iint \nabla \times \vec{F} \cdot dS &= \iint 3(x^2 + y^2)\hat{k} \cdot \hat{n} dS \\ &= 3 \iint (x^2 + y^2)\hat{k} \cdot \frac{(\hat{i} + \hat{j} + \hat{k})}{\sqrt{3}} dS \\ &= 3 \iint (x^2 + y^2) dS\end{aligned}$$

Taking projection of  $\hat{n}$  in  $xy$ -plane, it forms a circle with equation  $x^2 + y^2 = 1$ , as plane of intersection is ellipse.

$$\begin{aligned}\iint \nabla \times \vec{F} \cdot dS &= \sqrt{3} \iint (x^2 + y^2) \frac{dxdy}{\hat{n} \cdot \hat{k}} \\ &= \sqrt{3} \iint (x^2 + y^2) \frac{dxdy}{\sqrt{3}} \\ &= 3 \iint (x^2 + y^2) dxdy\end{aligned}$$

In the region  $x^2 + y^2 \leq 1$ , let  $x = r \cos \theta$ ,  $y = r \sin \theta$ , the integral becomes

$$\begin{aligned}\iint \nabla \times \vec{F} \cdot dS &= 3 \iint r^2 \times r dr d\theta = 3 \int_{\theta=0}^{2\pi} \int_{r=0}^1 r^3 dr d\theta \\ &= 3 \times \frac{1}{4} \times 2\pi = \frac{3\pi}{2} \\ \therefore \int \vec{F} \cdot dr &= \iint (\nabla \times \vec{F}) dS = \frac{3\pi}{2}\end{aligned}$$

7.17 Let  $\vec{F} = xy^2\hat{i} + (y+x)\hat{j}$ . Integrate  $(\nabla \times \vec{F})\hat{k}$  over the region in first quadrant bounded by the curves  $y = x^2$  and  $y = x$  using Green's theorem.

(2018 : 13 Marks)


**Solution:**

Given :

$$\vec{F} = xy^2\hat{i} + (y+x)\hat{j}$$

$$\nabla \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ xy^2 & y+x & 0 \end{vmatrix} = \hat{i}(0) - \hat{j}(0) + \hat{k}(1-2xy) = (1-2xy)\hat{k}$$

$$\therefore (\nabla \times \vec{F}) \cdot \hat{k} = 1 - 2xy$$



Now,

$$\iint (\nabla \times \vec{F}) \cdot \hat{k} dx dy = \int P dx + Q dy$$

(By Green's theorem)

$$\Rightarrow \iint \left( \frac{\partial(y+x)}{\partial x} - \frac{\partial(xy^2)}{\partial y} \right) dx dy = \iint \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$$

Comparing LHS and RHS, we get

$$Q = y - x, P = xy^2$$

Therefore, we have to find  $\int (xy^2) dx + (y+x) dy$  over  $C_1$  and  $C_2$ .

Over  $C_1$ :

$$\begin{aligned} \int_{C_1} (P dx + Q dy) &= \int_{C_1} (xy^2) dx + (y+x) dy = \int_{C_1} (xy^2) dx + \int_{C_1} (y+x) dy \\ &= \int_{C_1} x^5 dx + \int_{C_1} (x^2 + x) \times 2nd \, x \quad (\text{as } y = x^2) \\ &= \int_{x=0}^1 x^5 dx + 2 \int_{x=0}^1 (x^3 + x^2) dx = \frac{1}{6} + 2 \times \frac{1}{4} + 2 \times \frac{1}{3} \\ &= \frac{1}{6} + \frac{1}{2} + \frac{2}{3} = \frac{8}{6} = \frac{4}{3} \end{aligned}$$

Over  $C_2$ :

$$\begin{aligned} \int_{C_2} P dx + Q dy &= \int_{C_2} (xy^2) dx + \int_{C_2} (y+x) dy \\ &= \int_{C_2} x \cdot x^2 dx + \int_{C_2} 2x dx \quad (\text{as } y = x) \\ &= \int_{x=1}^0 x^3 dx + \int_{x=1}^0 2x dx = \left[ \frac{x^4}{4} \right]_1^0 + [x^2]_1^0 \\ &= \frac{-1}{4} - 1 = \frac{-5}{4} \end{aligned}$$

∴

$$\begin{aligned} \int_C P dx + Q dy &= \int_{C_1} P dx + Q dy + \int_{C_2} P dx + Q dy \\ &= \frac{4}{3} - \frac{5}{4} = \frac{16 - 15}{12} = \frac{1}{12} \end{aligned}$$

- 7.18 (i) State Gauss divergence theorem. Verify this theorem for  $\vec{F} = 4x\hat{i} - 2y^2\hat{j} + z^2\hat{k}$ , taken over the region bounded by  $x^2 + y^2 = 4$ ,  $z = 0$  and  $z = 3$

(2019 : 15 Marks)

- (ii) Evaluate by Stokes' theorem  $\oint_C e^x dx + 2y dy - dz$ , where  $C$  is the curve  $x^2 + y^2 = 4$ ,  $z = 2$ .

(2019 : 5 Marks)

Solution:

- (i) **Gauss Divergence Theorem** states that if  $V$  is the volume bounded by a closed surface  $S$  and ' $A$ ' is a vector function of position with continuous derivatives, then

$$\iiint_V \Delta \cdot \vec{A} dV = \iint_S \vec{A} \cdot \hat{n} dS$$

where ' $n$ ' is the positive outward drawn normal to  $S$ .

$$\vec{F} = 4x\hat{i} + 2y^2\hat{j} + z^2\hat{k}$$

Now,

taken over the region bounded by  $x^2 + y^2 = 4$ ,  $z = 0$  and  $z = 3$ .

$\therefore$  By Gauss divergence theorem

$$\begin{aligned} \iiint_V \Delta \cdot \vec{F} dV &= \iint_S \vec{F} \cdot \hat{n} dS \\ \text{Volume integral} &= \iiint_V \Delta \cdot \vec{F} du \\ &= \iiint_V \left[ \frac{\partial}{\partial x}(ux) + \frac{\partial}{\partial y}(-2y^2) + \frac{\partial}{\partial z}(z^2) \right] dV \\ &= \iiint_V (4 - 4y - 2z) dV \\ &= \int_{x=-2}^2 \int_{y=-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{z=0}^3 (4 - 4y + 2z) dz dy dx \\ &= \int_{x=-2}^2 \int_{y=-\sqrt{4-x^2}}^{\sqrt{4-x^2}} [4z - 4yz + z^2]_0^3 dy dx \\ &\quad \left[ \because \text{Radius of circle : } x^2 + y^2 = 4, r = 2, y = \pm\sqrt{4-x^2} \right] \\ &= \int_{x=-2}^2 \int_{y=-\sqrt{4-x^2}}^{\sqrt{4-x^2}} [12 - 12y + 9] dy dx \end{aligned}$$


Using polar coordinates

$$\begin{aligned} x &= r \cos \theta; y = r \sin \theta \\ dr &= r(-\sin \theta) d\theta; dy = r \cos \theta d\theta \\ dx dy &= r d\theta dr \\ &= \int_{0=0}^{2\pi} \int_0^2 (21 - 12r \sin \theta) r d\theta dr \\ &= \int_0^{2\pi} \left[ \frac{21r^2}{2} - \frac{r^3}{3} \sin \theta \right]_0^2 d\theta \\ &= \int_0^{2\pi} \left[ 42 - \frac{8}{3} \sin \theta \right] d\theta = \left[ 42\theta + \frac{8}{3} \cos \theta \right]_0^{2\pi} \\ &= \left[ 84\pi + \left[ \frac{8}{3}(1) - \frac{8}{3}(1) \right] \right] = 84\pi \end{aligned}$$

$$\Rightarrow \iiint_V \Delta \cdot \vec{F} dV = 84\pi$$

The surface  $S$  of the cylinder consists of a base  $S_1(z=0)$  the top  $S_2(z=3)$  and the convex portion  $S_3(x^2 + y^2 = 4)$  then

$$\begin{aligned} \text{Surface Integral} &= \iint_S \vec{F} \cdot \hat{n} dS \\ &= \iint_{S_1} \vec{F} \cdot \hat{n} dS_1 + \iint_{S_2} \vec{F} \cdot \hat{n} dS_2 + \iint_{S_3} \vec{F} \cdot \hat{n} dS_3 \end{aligned}$$



On  $S_2 (z = 3)$ :  $\hat{n} = k$ ,

$$\vec{F} = 4xi - 2y^2j + 9k$$

$$\vec{F} \cdot \hat{n} = 9, \text{ so that}$$

$$\iint_{S_2} \vec{F} \cdot \hat{n} dS_2 = \iint_{S_2} 9 dS_2 = 9 \cdot \pi r^2 = 9\pi(2)^2 [\because r = 2]$$

$$\iint_{S_2} \vec{F} \cdot \hat{n} dS_2 = 36\pi$$

On  $S_3 (x^2 + y^2 = u)$

A perpendicular to  $x^2 + y^2 = u$  has the direction

$$\nabla(x^2 + y^2) = 2xi + 2yj$$

Then a unit normal is

$$\hat{n} = \frac{2xi + 2yj}{\sqrt{4x^2 + 4y^2}}$$

$$\hat{n} = \frac{2(xi + yj)}{2\sqrt{x^2 + y^2}} = \frac{xi + yj}{\alpha} \quad [\because x^2 + y^2 = 4]$$

$$\vec{F} \cdot \hat{n} = (4xi - 2y^2j + z^2k) \left( \frac{xi + yj}{2} \right)$$

$$\vec{F} \cdot \hat{n} = 2x^2 - y^3$$

$$\iint_{S_3} \vec{F} \cdot \hat{n} dS = \int_{\theta=0}^{2\pi} \int_{z=0}^3 [2(2\cos\theta)^2 - (2\sin\theta)^3] 2 dz d\theta$$

$$= \int_{\theta=0}^{2\pi} (48\cos^2\theta - 48\sin^3\theta) d\theta = \int_{\theta=0}^{2\pi} 48\cos^2\theta d\theta = 48\pi$$

$\therefore$  Then the surface integral

$$= 0 + 36\pi + 48\pi$$

$$= 84\pi$$

$$\therefore \iiint_V \vec{F} \cdot \hat{n} dS = \iiint_V \vec{F} dV = 84\pi. \text{ Hence verified.}$$

(ii)

$C$  is the curve

Using Stokes theorem

$$\vec{F} = e^x dx + 2y dy - dz$$

$$x^2 + y^2 = 4; z = 2$$

$$\oint_C \vec{F} dr = \iint_S (\nabla \times \vec{F}) \cdot \hat{n} dS$$

$$\nabla \times \vec{F} = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ e^x & 2y & -1 \end{vmatrix}$$

$$\nabla \times \vec{F} = \hat{i}(0) - \hat{j}(0) + \hat{k}(0)$$

$$\nabla \times F = 0$$

$$\oint_C e^x dx + 2y dy - dz = (\nabla \times \vec{F}) \cdot \hat{n} dS = 0 \text{ which is the required result.}$$



**18 BOOKS**


**COMPLETE MODULES of  
GENERAL STUDIES**

*for*  
**Civil Services Exam**

*Revised & Improved*

**nEXT IAS**  
BIG LEARNINGS MADE EASY

- 1. History of Medieval India
- 2. History of the World
- 3. History of Modern India
- 4. Physical & World Geography
- 5. Indian Economy
- 6. International Relations
- 7. Internal Security
- 8. Art & Culture
- 9. Indian Polity
- 10. Post-Independence History
- 11. History of Ancient India
- 12. Life Sciences
- 13. Indian Society
- 14. Science and Technology
- 15. Ethics, Integrity & Aptitude
- 16. Ecology & Environment
- 17. Indian Geography and Disaster Management
- 18. Governance in India



**Books available at retail stores as well as major online shopping portals.**

Written/reviewed  
by CSE selected  
candidates

Coloured printing  
with relevant  
infographics

Concise yet  
comprehensive

Relevant for  
Pre and Main  
Exam

CSE previous  
year questions  
included

Updated  
with recent  
current affairs



To buy online, visit: [www.madeeasypublications.org](http://www.madeeasypublications.org)