


Krishna's Series

DIFFERENTIAL CALCULUS



KRISHNA Prakashan Media (P) Ltd. Meerut

Copyrighted material

Scanned with CamScanner

Krishna's Series

DIFFERENTIAL CALCULUS

(Fully Solved)

(For Degree and Honours Students of All Indian Universities, various Competitive Examinations like P.C.S., I.A.S. and Engineering Students)

By

A.R. Vasishtha

*Retired Head, Deptt. of Mathematics
Meerut College, Meerut*

Dr. S.K. Sharma

*Retired Head, Deptt. of Mathematics
N.A.S. College, Meerut*

&

A.K. Vasishtha

*M.Sc., Ph.D.,
C.C.S. University, Meerut*



KRISHNA Prakashan Media (P) Ltd.

KRISHNA HOUSE, 11, Shivaji Road, Meerut - 250 001 (U.P.), INDIA

Krishna's Series

DIFFERENTIAL CALCULUS
[FULLY SOLVED]

First Edition : 1991

Twenty First Edition : 2009

This book or any part thereof may not be reproduced in any form without the written permission of the publishers and authors.

Book Code No. : 446-21

Price : Rs. 250.00 Only

Published by :

Satyendra Rastogi "Mitra"
for KRISHNA Prakashan Media (P) Ltd.
11, Shivaji Road, Meerut - 250 001 (U.P.) India.
Phones : 0121-2642946, 2644766 Fax : 0121-2645855
Website : www.krishnaprakashan.com
e-mail : info@krishnaprakashan.com

Printed at :

Raj Printers, Meerut

Copyrighted material

CONTENTS

<i>Chapters</i>	<i>Pages</i>
<u>UNIT - I</u>	
1. Differentiation.	1 — 18
1. Derivative of a function. 2. Some standard results. 3. Differential coefficient of a function of a function. 4. Hyperbolic functions. 5. Inverse hyperbolic functions. 6. Some more methods of differentiation (logarithmic differentiation, implicit functions, parametric equations, trigonometrical transformations, differentiation of a function w.r.t. a function).	
2. Successive Differentiation.	19 — 57
1. Definition. 2. Standard results. 3. Leibnitz's Theorem. 4. n th derivative for $x=0$.	
3. Expansions of Functions.	58 — 87
1. Taylor's series. 2. Maclaurin's series. 3. Expansions of functions (Maclaurin's theorem, Taylor's theorem) 4. Some important Expansions (Exponential series, Sine series, Cosine series, Binomial series, Logarithmic expansion).	
4. Partial Differentiation.	88 — 116
1. Functions of two or more variables. 2. Homogeneous functions. 3. Euler's theorem on homogeneous functions. 4. Total derivatives.	
5. Indeterminate Forms.	117 — 158
1. Definition. 2. The form $0/0$. 3. Algebraic methods. 4. Form ∞/∞ . 5. Form $0 \times \infty$. 6. Form $\infty - \infty$. 7. The forms $0^0, 1^\infty, \infty^0$.	
6. Tangents and Normals.	159 — 205
1. Tangent. 2. Tangents parallel and perpendicular to the x -axis. 3. Normal. 4. Angle of intersection of two curves. 5. Length of cartesian tangent, normal, subtangent and subnormal. 6. Polar co-ordinates. 7. Angle between radius vector and tangent. 8. Angle of intersection of two polar curves. 9. Polar sub-tangent and polar sub-normal 10. Length of perpendicular from pole to tangent 11. Pedal equation. 12. Differential Coefficient of arc length (Cartesian co-ordinates). Differential Coefficient of arc length (polar co-ordinates).	
7. Curvature.	206 — 254
1. Definition. 2. Radius of curvature of intrinsic curves. 3. Radius of curvature of cartesian curves. 4. Radius of curvature of parametric curves. 5. Radius of curvature of pedal curves. 6. Radius of curvature of polar curves. 7. Tangential polar formula for radius of curvature. 8. Miscellaneous formulae for radius of curvature when x and y are functions of arc length. 9. Radius of curvature at the origin (Newton's method, expansion method, radius of curvature at the pole).	

Chapters	Pages
10. Co-ordinates of centre of curvature. 11. Chord of curvature through the origin. 12. Chords of curvature parallel to co-ordinate axes.	255 — 263
8. Asymptotes.	255 — 303
1. Definition. 2. Determination of Asymptotes. 3. Asymptotes of general algebraic curves. 4. Non-existence of asymptotes. 5. Case of parallel asymptotes. 6. Asymptotes parallel to the co-ordinates axes. 7. Total number of asymptotes of a curve. 8. Complete working rule of finding the asymptotes of rational algebraic curves. 9. Asymptotes by expansion. 10. Alternative methods of finding asymptotes of algebraic curves. 11. Asymptotes by inspection. 12. Intersection of a curve and its asymptotes. 13. Asymptotes of polar curves. 14. Circular asymptotes.	255 — 303
9. Change of the Independent Variables.	304 — 332
1. Introduction. 2. To change the independent variable into the dependent variable. 3. To change independent variable x into another variable z , when $x = f(z)$. 4. Change of both the dependent and independent variables. 5. Transformation in case of two independent variables. 6. Transformation from cartesian to polar co-ordinates and vice versa.	304 — 332
Miscellaneous Solved Problems	334 — 336
UNIT - II	
1. Maxima and Minima	1 — 32
(Of functions of a single independent Variable)	
1. Definition. 2. Properties of Maxima and Minima. 3. Condition for maximum and minimum values. 4. Working rule of maxima and minima of $f(x)$. 5. Application of maxima and minima to geometrical and other problems.	1 — 32
2. Maxima and Minima.	33 — 53
(Of Functions of two independent variables)	
1. Definition. 2. Necessary conditions for the existence of a maximum or a minimum of $f(x, y)$ at $x = a, y = b$. 3. Stationary and Extreme points. 4. Sufficient conditions for maxima or minima. 5. Working rule for maxima and minima.	33 — 53
3. Maxima and Minima.	54 — 97
(Of functions of several variables)	
Maxima and minima of functions of several variables and Lagrange's method of undetermined multipliers.	54 — 97
4. Envelopes and Evolutes.	98 — 142
1. One parameter family of curves. 2. Envelope of a one parameter family of curves. 3. Method for finding envelope. 4. Envelope in	98 — 142

<u>Chapters</u>	<u>Pages</u>
case the equation or the family of curves is a quadratic in the parameter. 5. Geometrical significance of the envelope. 6. Evolute of a curve. 7. Length of arc of an evolute.	
5. Jacobians	143 — 175
1. Definition. 2. Case of functions of functions. 3. Jacobian of implicit functions. 4. Condition of a Jacobian to vanish.	
6. Singular Points.	176 — 217
1. Definition. 2. Concavity and Convexity. 3. Point of inflexion. 4. Test for point of inflexion. 5. Multiple points. 6. Classification of double points (Node, Cusp, Conjugate Point). 7. Species of Cusps. 8. Tangents at origin. 9. Change of origin. 10. Tangent at any point to a curve. 11. Position and character of double points. 12. Nature of a cusp at the origin. 13. Nature of a cusp at any point.	
7. Curve Tracing.	218 — 268
1. Curve tracing (cartesian equations). 2. Polar equations. 3. Parametric equations.	
UNIT - III	
1. Functions of a Real Variable, Limits, Continuity and Differentiability	1 — 37
1. Definitions. 2. Limits. 3. Continuity. 4. Differentiability.	
2. Rolle's Theorem, Mean Value Theorems, Taylor's and MacLaurin's Theorems.	38 — 59
1. Rolle's theorem. 2. Lagrange's mean value theorem. 3. Some important deductions from the mean value theorem. 4. Cauchy's mean value theorem. 5. Taylor's theorem with Lagrange's form of remainder after n terms. 6. Taylor's theorem with Cauchy's form of remainder.	

Note : The chapter 1 of Unit-I, chapter 1 of Unit-II and two chapters of Unit-III of this book are not in the syllabus of C.C.S. University, Meerut. The chapters 2 to 9 cover the syllabus of Unit I and the chapters 2 to 7 of Unit-II cover the syllabus of Unit II.

1

Differentiation

§ 1. Derivative (or differential coefficient) of a function.

Let $y = f(x)$ be a function of x defined in the interval (a, b) . For a small increment δx in x , let the corresponding increment in the value of y be δy . The quotient $\delta y/\delta x$ is called the *average rate of change* of y with respect to x in the interval $(x, x + \delta x)$. If the increment ratio $\delta y/\delta x$ tends to a finite limit as $\delta x \rightarrow 0$, then this limit is called the **differential coefficient** of y with respect to x and is denoted by

$$\frac{dy}{dx} \text{ or } \frac{d}{dx}f(x) \text{ or } f'(x) \text{ or } Df(x) \text{ or } Dy \text{ etc.}$$

Thus $\frac{dy}{dx}$ = the diff. coeff. of y w.r.t. x = $\lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x}$.

The diff. coeff. dy/dx is also called the *rate of change* of y with respect to x .

Now if $y = f(x)$ and $y + \delta y = f(x + \delta x)$, we have

$$\delta y = f(x + \delta x) - f(x).$$

$\therefore \frac{dy}{dx} = \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} = \lim_{\delta x \rightarrow 0} \frac{f(x + \delta x) - f(x)}{\delta x}$, provided the

limit exists. The process of finding the diff. coeff. of a given function $f(x)$ is called **differentiation**.

The differential coefficient of $f(x)$ for $x = a$ is defined as $\lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$, provided the limit exists. It is denoted by

$$\left(\frac{dy}{dx}\right)_{x=a} \text{ or } (y')_a \text{ or } f'(a).$$

§ 2. Some standard results.

To find the differential coefficients of some standard functions from the first principles.

I. Differential coefficient of a constant. Let $f(x) = c$, where c is a constant. Then $f(x + \delta x) = c$, because $f(x)$ takes the same value c for every value of x .

Now by the definition of a diff. coeff., we have

$$\begin{aligned} \frac{d}{dx}f(x) &= \lim_{\delta x \rightarrow 0} \frac{f(x + \delta x) - f(x)}{\delta x} = \lim_{\delta x \rightarrow 0} \frac{c - c}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \frac{0}{\delta x} = \lim_{\delta x \rightarrow 0} 0 = 0. \end{aligned}$$

Thus, the diff. coeff. of a constant is zero.

2. Diff. coeff. of the product of a constant and a function.

$$\begin{aligned} \text{We have, } \frac{d}{dx} \{cf(x)\} &= \lim_{\delta x \rightarrow 0} \frac{cf(x + \delta x) - cf(x)}{\delta x} \\ &= c \lim_{\delta x \rightarrow 0} \frac{f(x + \delta x) - f(x)}{\delta x} = c \frac{d}{dx} f(x). \end{aligned}$$

3. Derivative of a sum or a difference of two functions.

Let $f(x) = f_1(x) \pm f_2(x)$.

Then $f(x + \delta x) = f_1(x + \delta x) \pm f_2(x + \delta x)$.

$$\begin{aligned} \therefore \frac{d}{dx} f(x) &= \lim_{\delta x \rightarrow 0} \frac{\{f_1(x + \delta x) \pm f_2(x + \delta x)\} - \{f_1(x) \pm f_2(x)\}}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \frac{f_1(x + \delta x) - f_1(x)}{\delta x} \pm \lim_{\delta x \rightarrow 0} \frac{f_2(x + \delta x) - f_2(x)}{\delta x} \\ &= \frac{d}{dx} f_1(x) \pm \frac{d}{dx} f_2(x). \end{aligned}$$

4. Differential coefficient of the product of two functions.

(Delhi 1982)

Let $y = f_1(x)f_2(x)$. Then $y + \delta y = f_1(x + \delta x) \cdot f_2(x + \delta x)$.

$$\begin{aligned} \therefore \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{f_1(x + \delta x)f_2(x + \delta x) - f_1(x) \cdot f_2(x)}{\delta x}, \quad [\text{by the def. of a diff. coeff.}] \\ &= \lim_{\delta x \rightarrow 0} \left[f_1(x + \delta x) \cdot \frac{f_2(x + \delta x) - f_2(x)}{\delta x} + f_2(x) \frac{f_1(x + \delta x) - f_1(x)}{\delta x} \right] \\ &= \lim_{\delta x \rightarrow 0} f_1(x + \delta x) \frac{f_2(x + \delta x) - f_2(x)}{\delta x} \\ &\quad + \lim_{\delta x \rightarrow 0} f_2(x) \cdot \frac{f_1(x + \delta x) - f_1(x)}{\delta x} \\ &= f_1(x) \cdot \frac{d}{dx} f_2(x) + f_2(x) \cdot \frac{d}{dx} f_1(x). \end{aligned}$$

\therefore the diff. coeff. of the product of two functions

= (first function) \times (diff. coeff. of the second function)
+ (second function) \times (diff. coeff. of the first function).

5. Differential coefficient of the quotient of two functions.

Let $y = f_1(x)/f_2(x)$. Then $y + \delta y = f_1(x + \delta x)/f_2(x + \delta x)$.

$$\begin{aligned} \therefore \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{\left[\frac{f_1(x + \delta x)}{f_2(x + \delta x)} - \frac{f_1(x)}{f_2(x)} \right]}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \left[\frac{f_1(x + \delta x) \cdot f_2(x) - f_1(x) \cdot f_2(x + \delta x)}{\delta x \cdot f_2(x + \delta x) \cdot f_2(x)} \right] \end{aligned}$$

$$\begin{aligned}
 &= \lim_{\delta x \rightarrow 0} \left[\frac{f_1(x + \delta x) f_2(x) - f_1(x) f_2(x) - f_1(x) f_2(x + \delta x) + f_1(x) \cdot f_2(x)}{\delta x \cdot f_2(x + \delta x) \cdot f_2(x)} \right] \\
 &= \lim_{\delta x \rightarrow 0} \left[\frac{f_2(x) \cdot \left\{ \frac{f_1(x + \delta x) - f_1(x)}{\delta x} \right\} - f_1(x) \cdot \left\{ \frac{f_2(x + \delta x) - f_2(x)}{\delta x} \right\}}{f_2(x + \delta x) f_2(x)} \right] \\
 &= \frac{f_2(x) \cdot \frac{d}{dx} f_1(x) - f_1(x) \cdot \frac{d}{dx} f_2(x)}{[f_2(x)]^2}.
 \end{aligned}$$

\therefore the diff. coeff. of the quotient of two functions
 $= \frac{(\text{diff. coeff. of Nr.})(\text{Dr.}) - (\text{Nr.})(\text{diff. coeff. of Dr.})}{\text{square of the denominator}}$.

6. $(d/dx) x^n = nx^{n-1}$.

Let $y = x^n$. Then $y + \delta y = (x + \delta x)^n$.

$$\begin{aligned}
 \therefore \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{(x + \delta x)^n - x^n}{\delta x} = \lim_{\delta x \rightarrow 0} x^n \left[\frac{\left(1 + \frac{\delta x}{x}\right)^n - 1}{\delta x} \right] \\
 &= \lim_{\delta x \rightarrow 0} x^n \left[\frac{1 + n \frac{\delta x}{x} + \frac{n(n-1)}{2!} \frac{(\delta x)^2}{x^2} + \dots - 1}{\delta x} \right] \\
 &= \lim_{\delta x \rightarrow 0} x^n \left[\frac{n}{x} + \frac{n(n-1)}{2!} \frac{\delta x}{x^2} + \text{terms containing higher powers of } \delta x \right] \\
 &= x^n (n/x) = nx^{n-1}.
 \end{aligned}$$

7. $(d/dx) e^x = e^x$.

Let $y = e^x$. Then $y + \delta y = e^x + \delta x$.

$$\begin{aligned}
 \therefore \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{e^{x+\delta x} - e^x}{\delta x} \\
 &= \lim_{\delta x \rightarrow 0} e^x \left[\frac{e^{\delta x} - 1}{\delta x} \right] = \lim_{\delta x \rightarrow 0} e^x \left[\frac{1 + \delta x + (\delta x)^2/2! + \dots - 1}{\delta x} \right] \\
 &= \lim_{\delta x \rightarrow 0} e^x \left[1 + \frac{\delta x}{2!} + \text{terms containing higher powers of } \delta x \right] = e^x.
 \end{aligned}$$

8. $(d/dx) a^x = a^x \log a$.

Let $y = a^x$. Then $y + \delta y = a^{x+\delta x}$.

$$\begin{aligned}
 \therefore \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{a^{x+\delta x} - a^x}{\delta x} \\
 &= \lim_{\delta x \rightarrow 0} a^x \left[\frac{a^{\delta x} - 1}{\delta x} \right] \\
 &= \lim_{\delta x \rightarrow 0} a^x \left[1 + \delta x \log a + \frac{(\delta x)^2 (\log a)^2}{2!} + \dots - 1 \right]
 \end{aligned}$$

$$= \lim_{\delta x \rightarrow 0} a^x [\log a + (\delta x/2!) (\log a)^2 + \text{terms containing higher powers of } \delta x]$$

$$= a^x \log a.$$

9. $(d/dx) \log_e x = 1/x.$

Let $y = \log x$. Then $y + \delta y = \log(x + \delta x)$.

$$\begin{aligned}\therefore \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{\log(x + \delta x) - \log x}{\delta x} = \lim_{\delta x \rightarrow 0} \frac{\log \{(x + \delta x)/x\}}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \frac{\log \left(1 + \frac{\delta x}{x}\right)}{\delta x} = \lim_{\delta x \rightarrow 0} \frac{\frac{\delta x}{x} - \frac{(\delta x)^2}{2x^2} + \dots}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \left[\frac{1}{x} - \frac{\delta x}{2x^2} + \dots \right] = \frac{1}{x}.\end{aligned}$$

10. $\frac{d}{dx} \log_a x = \frac{1}{x \log_e a}.$

We have $\log_a x = \log_e x \cdot \log_a e = \log_a e \cdot \log_e x$, where $\log_a e$ is simply a constant.

$$\begin{aligned}\therefore \frac{d}{dx} (\log_a x) &= (\log_a e) \frac{d}{dx} (\log_e x) = (\log_a e) \frac{1}{x} = \frac{1}{x} \cdot \log_a e \\ &= \frac{1}{x \log_e a}, \text{ since } \log_a e \cdot \log_e a = 1.\end{aligned}$$

11. $(d/dx) \sin x = \cos x.$

Let $y = \sin x$. Then $y + \delta y = \sin(x + \delta x)$.

$$\begin{aligned}\therefore \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{\sin(x + \delta x) - \sin x}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \frac{2 \cos \{x + (\delta x/2)\} \sin(\delta x/2)}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \frac{\cos \{x + (\delta x/2)\} \sin(\delta x/2)}{(\delta x/2)} \\ &= \lim_{\delta x \rightarrow 0} \cos \left(x + \frac{\delta x}{2}\right) \cdot \lim_{\delta x \rightarrow 0} \frac{\sin(\delta x/2)}{(\delta x/2)} \\ &= (\cos x) \cdot 1 = \cos x.\end{aligned}$$

12. $(d/dx) \cos x = -\sin x.$

Let $y = \cos x$. Then $y + \delta y = \cos(x + \delta x)$.

$$\begin{aligned}\therefore \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{\cos(x + \delta x) - \cos x}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \frac{-2 \sin \{x + (\delta x/2)\} \sin(\delta x/2)}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} -\sin \left(x + \frac{\delta x}{2}\right) \cdot \frac{\sin(\delta x/2)}{(\delta x/2)} = (-\sin x) \cdot 1 = -\sin x.\end{aligned}$$

13. $(d/dx) \tan x = \sec^2 x.$

(Delhi 1982)

Let $y = \tan x$. Then $y + \delta y = \tan(x + \delta x)$.

$$\begin{aligned}\therefore \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{\tan(x + \delta x) - \tan x}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \frac{\left[\frac{\sin(x + \delta x)}{\cos(x + \delta x)} - \frac{\sin x}{\cos x} \right]}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \frac{\sin(x + \delta x) \cos x - \sin x \cos(x + \delta x)}{\delta x \cos(x + \delta x) \cdot \cos x} \\ &= \lim_{\delta x \rightarrow 0} \frac{\sin((x + \delta x) - x)}{\delta x \cdot \cos(x + \delta x) \cos x} \\ &= \lim_{\delta x \rightarrow 0} \left[\frac{\sin \delta x}{\delta x} \cdot \frac{1}{\cos(x + \delta x) \cos x} \right] = \frac{1}{\cos^2 x} = \sec^2 x.\end{aligned}$$

14. $(d/dx) \cot x = -\operatorname{cosec}^2 x.$

For proof proceed as in the case of $\tan x$.

15. $(d/dx) \sec x = \sec x \tan x.$

Let $y = \sec x$. Then $y + \delta y = \sec(x + \delta x)$.

$$\begin{aligned}\therefore \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{\sec(x + \delta x) - \sec x}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \frac{\frac{1}{\cos(x + \delta x)} - \frac{1}{\cos x}}{\delta x} \\ &= \lim_{\delta x \rightarrow 0} \frac{\cos x - \cos(x + \delta x)}{\delta x \cdot \cos(x + \delta x) \cos x} \\ &= \lim_{\delta x \rightarrow 0} \frac{2 \sin(x + \frac{1}{2} \delta x) \sin(\delta x/2)}{\delta x \cdot \cos(x + \delta x) \cos x} \\ &= \lim_{\delta x \rightarrow 0} \left[\frac{\sin(x + \frac{1}{2} \delta x)}{\cos(x + \delta x) \cos x} \cdot \frac{\sin(\delta x/2)}{(\delta x/2)} \right] \\ &= \frac{\sin x}{\cos x \cos x} = \sec x \tan x.\end{aligned}$$

16. $(d/dx) \operatorname{cosec} x = -\operatorname{cosec} x \cot x.$

For proof proceed as in the case of $\sec x$.

17. $\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}.$

Let $y = \sin^{-1} x$. Then $x = \sin y$ and so $x + \delta x = \sin(y + \delta y)$. As $\delta x \rightarrow 0$, δy also $\rightarrow 0$.

Now $\delta x = \sin(y + \delta y) - \sin y$.

$$\therefore 1 = \frac{\sin(y + \delta y) - \sin y}{\delta x}, \quad [\text{on dividing both sides by } \delta x]$$

$$\text{or } 1 = \frac{\sin(y + \delta y) - \sin y}{\delta y} \cdot \frac{\delta y}{\delta x}.$$

Taking limits of both sides when $\delta x \rightarrow 0$, we get

$$\begin{aligned} 1 &= \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y) - \sin y}{\delta y} \cdot \lim_{\delta y \rightarrow 0} \frac{\delta y}{\delta x}, \\ &= \left[\lim_{\delta y \rightarrow 0} \frac{2 \cos(y + \frac{1}{2}\delta y) \sin(\frac{1}{2}\delta y)}{\delta y} \right] \cdot \frac{dy}{dx} \quad [\because \delta y \rightarrow 0 \text{ when } \delta x \rightarrow 0] \\ &= \left[\lim_{\delta y \rightarrow 0} \cos(y + \frac{1}{2}\delta y) \cdot \frac{\sin(\frac{1}{2}\delta y)}{\delta y/2} \right] \frac{dy}{dx} \\ &= (\cos y) \cdot (dy/dx). \\ \therefore \frac{dy}{dx} &= \frac{1}{\cos y} = \frac{1}{\sqrt{1 - \sin^2 y}} = \frac{1}{\sqrt{1 - x^2}}. \end{aligned}$$

18. $\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1 - x^2}}$.

For proof proceed exactly as in the case of $\sin^{-1} x$.

19. $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1 + x^2}$.

Let $y = \tan^{-1} x$. Then $x = \tan y$ and so $x + \delta x = \tan(y + \delta y)$. As $\delta x \rightarrow 0$, δy also $\rightarrow 0$.

Now $\delta x = \tan(y + \delta y) - \tan y$.

$$\therefore 1 = \frac{\tan(y + \delta y) - \tan y}{\delta y} \cdot \frac{\delta y}{\delta x}.$$

Taking limits of both sides when $\delta x \rightarrow 0$, we get

$$\begin{aligned} 1 &= \lim_{\delta y \rightarrow 0} \frac{\tan(y + \delta y) - \tan y}{\delta y} \cdot \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} \\ &= \left[\lim_{\delta y \rightarrow 0} \left\{ \frac{\sin(y + \delta y)}{\cos(y + \delta y)} - \frac{\sin y}{\cos y} \right\} \right] \cdot \frac{dy}{dx} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y) \cos y - \cos(y + \delta y) \sin y}{\delta y \cos(y + \delta y) \cos y} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y - y)}{\delta y \cdot \cos(y + \delta y) \cos y} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \left[\frac{\sin \delta y}{\delta y} \cdot \frac{1}{\cos(y + \delta y) \cos y} \right] \\ &= \frac{dy}{dx} \cdot \frac{1}{\cos^2 y} = \frac{dy}{dx} \cdot \sec^2 y. \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{1}{\sec^2 y} = \frac{1}{1 + \tan^2 y} = \frac{1}{1 + x^2}.$$

20. $\frac{d}{dx}(\cot^{-1} x) = -\frac{1}{1 + x^2}$.

For proof proceed exactly as in the case of $\tan^{-1} x$.



Taking limits of both sides when $\delta x \rightarrow 0$, we get

$$1 = \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y) - \sin y}{\delta y} \cdot \lim_{\delta y \rightarrow 0} \frac{\delta y}{\delta x},$$

[$\because \delta y \rightarrow 0$ when $\delta x \rightarrow 0$]

$$\begin{aligned} &= \left[\lim_{\delta y \rightarrow 0} \frac{2 \cos(y + \frac{1}{2}\delta y) \sin(\frac{1}{2}\delta y)}{\delta y} \right] \cdot \frac{dy}{dx} \\ &= \left[\lim_{\delta y \rightarrow 0} \cos(y + \frac{1}{2}\delta y) \cdot \frac{\sin(\frac{1}{2}\delta y)}{\delta y/2} \right] \frac{dy}{dx} \\ &= (\cos y) \cdot (dy/dx). \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{1}{\cos y} = \frac{1}{\sqrt{1 - \sin^2 y}} = \frac{1}{\sqrt{1 - x^2}}.$$

18. $\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}}$.

For proof proceed exactly as in the case of $\sin^{-1} x$.

19. $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$.

Let $y = \tan^{-1} x$. Then $x = \tan y$ and so $x + \delta x = \tan(y + \delta y)$.
As $\delta x \rightarrow 0$, δy also $\rightarrow 0$.

Now $\delta x = \tan(y + \delta y) - \tan y$.

$$\therefore 1 = \frac{\tan(y + \delta y) - \tan y}{\delta y} \cdot \frac{\delta y}{\delta x}.$$

Taking limits of both sides when $\delta x \rightarrow 0$, we get

$$\begin{aligned} 1 &= \lim_{\delta y \rightarrow 0} \frac{\tan(y + \delta y) - \tan y}{\delta y} \cdot \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} \\ &= \left[\lim_{\delta y \rightarrow 0} \left\{ \frac{\sin(y + \delta y)}{\cos(y + \delta y)} - \frac{\sin y}{\cos y} \right\} \right] / \frac{dy}{dx} \cdot \frac{dy}{dx} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y) \cos y - \cos(y + \delta y) \sin y}{\delta y \cos(y + \delta y) \cos y} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y - y)}{\delta y \cos(y + \delta y) \cos y} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \left[\frac{\sin \delta y}{\delta y} \cdot \frac{1}{\cos(y + \delta y) \cos y} \right] \\ &= \frac{dy}{dx} \cdot \frac{1}{\cos^2 y} = \frac{dy}{dx} \cdot \sec^2 y. \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{1}{\sec^2 y} = \frac{1}{1 + \tan^2 y} = \frac{1}{1 + x^2}.$$

20. $\frac{d}{dx}(\cot^{-1} x) = -\frac{1}{1+x^2}$.

For proof proceed exactly as in the case of $\tan^{-1} x$.

21. $\frac{d}{dx}(\sec^{-1} x) = \frac{1}{x\sqrt{x^2 - 1}}$.

Let $y = \sec^{-1} x$. Then $x = \sec y$ and so $x + \delta x = \sec(y + \delta y)$.
As $\delta x \rightarrow 0$, δy also $\rightarrow 0$.

Now $\delta x = \sec(y + \delta y) - \sec y$.

$$\sec(y + \delta y) - \sec y = \frac{\sec(y + \delta y) - \sec y}{\sec y} \cdot \sec y$$

 δy δx

Copyrighted material

6

DIFFERENTIAL CALCULUS

Taking limits of both sides when $\delta x \rightarrow 0$, we get

$$1 = \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y) - \sin y}{\delta y} \cdot \lim_{\delta y \rightarrow 0} \frac{\delta y}{\delta x},$$

[$\because \delta y \rightarrow 0$ when $\delta x \rightarrow 0$]

$$\begin{aligned} &= \left[\lim_{\delta y \rightarrow 0} \frac{2 \cos(y + \frac{1}{2}\delta y) \sin(\frac{1}{2}\delta y)}{\delta y} \right] \cdot \frac{dy}{dx} \\ &= \left[\lim_{\delta y \rightarrow 0} \cos(y + \frac{1}{2}\delta y) \cdot \frac{\sin(\frac{1}{2}\delta y)}{\delta y/2} \right] \frac{dy}{dx} \\ &= (\cos y) \cdot (dy/dx). \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{1}{\cos y} = \frac{1}{\sqrt{1 - \sin^2 y}} = \frac{1}{\sqrt{1 - x^2}}.$$

18. $\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}}$.

For proof proceed exactly as in the case of $\sin^{-1} x$.

19. $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$.

Let $y = \tan^{-1} x$. Then $x = \tan y$ and so $x + \delta x = \tan(y + \delta y)$.
As $\delta x \rightarrow 0$, δy also $\rightarrow 0$.

Now $\delta x = \tan(y + \delta y) - \tan y$.

$$\therefore 1 = \frac{\tan(y + \delta y) - \tan y}{\delta y} \cdot \frac{\delta y}{\delta x}.$$

Taking limits of both sides when $\delta x \rightarrow 0$, we get

$$\begin{aligned} 1 &= \lim_{\delta y \rightarrow 0} \frac{\tan(y + \delta y) - \tan y}{\delta y} \cdot \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} \\ &= \left[\lim_{\delta y \rightarrow 0} \left\{ \frac{\sin(y + \delta y)}{\cos(y + \delta y)} - \frac{\sin y}{\cos y} \right\} \right] \cdot \frac{dy}{dx} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y) \cos y - \cos(y + \delta y) \sin y}{\delta y \cos(y + \delta y) \cos y} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y - y)}{\delta y \cdot \cos(y + \delta y) \cos y} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \left[\frac{\sin \delta y}{\delta y} \cdot \frac{1}{\cos(y + \delta y) \cos y} \right] \\ &= \frac{dy}{dx} \cdot \frac{1}{\cos^2 y} = \frac{dy}{dx} \cdot \sec^2 y. \\ \therefore \frac{dy}{dx} &= \frac{1}{\sec^2 y} = \frac{1}{1 + \tan^2 y} = \frac{1}{1 + x^2}. \end{aligned}$$

20. $\frac{d}{dx}(\cot^{-1} x) = -\frac{1}{1+x^2}$.

For proof proceed exactly as in the case of $\tan^{-1} x$.

Copyrighted material



Taking limits of both sides when $\delta x \rightarrow 0$, we get

$$1 = \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y) - \sin y}{\delta y} \cdot \lim_{\delta y \rightarrow 0} \frac{\delta y}{\delta x},$$

$$= \left[\lim_{\delta y \rightarrow 0} \frac{2 \cos(y + \frac{1}{2}\delta y) \sin(\frac{1}{2}\delta y)}{\delta y} \right] \cdot \frac{dy}{dx}$$

$$= \left[\lim_{\delta y \rightarrow 0} \cos(y + \frac{1}{2}\delta y) \cdot \frac{\sin(\frac{1}{2}\delta y)}{\delta y/2} \right] \frac{dy}{dx}$$

$$= (\cos y) \cdot (dy/dx).$$

$$\therefore \frac{dy}{dx} = \frac{1}{\cos y} = \frac{1}{\sqrt{(1 - \sin^2 y)}} = \frac{1}{\sqrt{(1 - x^2)}}.$$

18. $\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{(1 - x^2)}}.$

For proof proceed exactly as in the case of $\sin^{-1} x$.

19. $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1 + x^2}.$

Let $y = \tan^{-1} x$. Then $x = \tan y$ and so $x + \delta x = \tan(y + \delta y)$.

As $\delta x \rightarrow 0$, δy also $\rightarrow 0$.

Now $\delta x = \tan(y + \delta y) - \tan y$.

$$\therefore 1 = \frac{\tan(y + \delta y) - \tan y}{\delta y} \cdot \frac{\delta y}{\delta x}.$$

Taking limits of both sides when $\delta x \rightarrow 0$, we get

$$1 = \lim_{\delta y \rightarrow 0} \frac{\tan(y + \delta y) - \tan y}{\delta y} \cdot \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x}$$

$$= \left[\lim_{\delta y \rightarrow 0} \frac{\left\{ \frac{\sin(y + \delta y)}{\cos(y + \delta y)} - \frac{\sin y}{\cos y} \right\}}{\delta y} \right] \cdot \frac{dy}{dx}$$

$$= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y) \cos y - \cos(y + \delta y) \sin y}{\delta y \cos(y + \delta y) \cos y}$$

$$= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \frac{\sin(y + \delta y - y)}{\delta y \cdot \cos(y + \delta y) \cos y}$$

$$= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \left[\frac{\sin \delta y}{\delta y} \cdot \frac{1}{\cos(y + \delta y) \cos y} \right]$$

$$= \frac{dy}{dx} \cdot \frac{1}{\cos^2 y} = \frac{dy}{dx} \cdot \sec^2 y.$$

$$\therefore \frac{dy}{dx} = \frac{1}{\sec^2 y} = \frac{1}{1 + \tan^2 y} = \frac{1}{1 + x^2}.$$

20. $\frac{d}{dx}(\cot^{-1} x) = -\frac{1}{1 + x^2}.$

For proof proceed exactly as in the case of $\tan^{-1} x$.

Copyrighted material

21. $\frac{d}{dx}(\sec^{-1} x) = \frac{1}{x\sqrt{(x^2 - 1)}}.$

Let $y = \sec^{-1} x$. Then $x = \sec y$ and so $x + \delta x = \sec(y + \delta y)$.
As $\delta x \rightarrow 0$, δy also $\rightarrow 0$.

Now $\delta x = \sec(y + \delta y) - \sec y$.



$$21. \frac{d}{dx} (\sec^{-1} x) = \frac{1}{x\sqrt{x^2 - 1}}.$$

Let $y = \sec^{-1} x$. Then $x = \sec y$ and so $x + \delta x = \sec(y + \delta y)$. As $\delta x \rightarrow 0$, δy also $\rightarrow 0$.

Now $\delta x = \sec(y + \delta y) - \sec y$.

$$\therefore 1 = \frac{\sec(y + \delta y) - \sec y}{\delta y} \cdot \frac{\delta y}{\delta x}.$$

Taking limits of both sides when $\delta x \rightarrow 0$, we get

$$\begin{aligned} 1 &= \lim_{\delta y \rightarrow 0} \frac{\sec(y + \delta y) - \sec y}{\delta y} \cdot \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \left[\frac{1}{\cos(y + \delta y)} - \frac{1}{\cos y} \right] / \delta y \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \frac{\cos y - \cos(y + \delta y)}{\delta y \cdot \cos y \cos(y + \delta y)} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \frac{2 \sin(y + \frac{1}{2}\delta y) \sin(\frac{1}{2}\delta y)}{\delta y \cdot \cos y \cos(y + \delta y)} \\ &= \frac{dy}{dx} \cdot \lim_{\delta y \rightarrow 0} \left[\frac{\sin(y + \frac{1}{2}\delta y)}{\cos y \cos(y + \delta y)} \cdot \frac{\sin(\frac{1}{2}\delta y)}{\frac{1}{2}\delta y} \right] \\ &= \frac{dy}{dx} \cdot \frac{\sin y}{\cos y \cos y} = \frac{dy}{dx} \cdot \sec y \tan y. \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{1}{\sec y \tan y} = \frac{1}{\sec y \sqrt{\sec^2 y - 1}} = \frac{1}{x \sqrt{x^2 - 1}}.$$

$$22. \frac{d}{dx} (\operatorname{cosec}^{-1} x) = -\frac{1}{x\sqrt{x^2 - 1}}.$$

For proof proceed as in the case of $\sec^{-1} x$.

§ 3. Differential coefficient of a function of a function.

Consider the function $\log \sin x$. Here $\log(\sin x)$ is a function of $\sin x$ whereas $\sin x$ is itself a function of x . Thus we have the case of a function of a function.

Let $y = f\{\phi(x)\}$.

Put $t = \phi(x)$. Then $t + \delta t = \phi(x + \delta x)$. As $\delta x \rightarrow 0$, δt also $\rightarrow 0$.

We have

$$\begin{aligned} \frac{dy}{dx} &= \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} = \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta t} \cdot \frac{\delta t}{\delta x} = \left(\lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta t} \right) \cdot \left(\lim_{\delta x \rightarrow 0} \frac{\delta t}{\delta x} \right) \\ &= \left(\lim_{\delta t \rightarrow 0} \frac{\delta y}{\delta t} \right) \cdot \left(\lim_{\delta x \rightarrow 0} \frac{\delta t}{\delta x} \right), \text{ since } \delta t \rightarrow 0 \text{ when } \delta x \rightarrow 0 \\ &= \frac{dy}{dt} \cdot \frac{dt}{dx}. \end{aligned}$$

Thus, if y is a function of t and t is a function of x , then y is a function of x and we have

Copyrighted material

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}.$$

Similarly, if y is a function of u , u is a function of v and v is a function of x , then y is also a function of x , and we have

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx}.$$

This is known as the **chain rule of differentiation**. We can extend



$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}.$$

Similarly, if y is a function of u , u is a function of v and v is a function of x , then y is also a function of x , and we have

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx}.$$

This is known as the **chain rule of differentiation**. We can extend it still further.

Cor. To prove that $\frac{dy}{dx} \times \frac{dx}{dy} = 1$.

Let $y = f(x)$. Then $y + \delta y = f(x + \delta x)$. As $\delta x \rightarrow 0$, δy also $\rightarrow 0$. We have

$$\frac{\delta y}{\delta x} \cdot \frac{\delta x}{\delta y} = 1.$$

Taking limits of both sides when $\delta x \rightarrow 0$, we get

$$\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \cdot \frac{\delta x}{\delta y} \right) = 1 \text{ or } \left(\lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} \right) \left(\lim_{\delta x \rightarrow 0} \frac{\delta x}{\delta y} \right) = 1$$

$$\text{or } \left(\lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x} \right) \cdot \left(\lim_{\delta y \rightarrow 0} \frac{\delta x}{\delta y} \right) = 1, \text{ since } \delta y \rightarrow 0 \text{ as } \delta x \rightarrow 0$$

$$\text{or } \frac{dy}{dx} \cdot \frac{dx}{dy} = 1. \text{ Thus } \frac{dy}{dx} = \frac{1}{dx/dy}.$$

§ 4. Hyperbolic functions.

These are defined as follows :

$$\sinh x = \frac{1}{2}(e^x - e^{-x}); \cosh x = \frac{1}{2}(e^x + e^{-x});$$

$$\tanh x = \frac{\sinh x}{\cosh x} = \frac{e^x - e^{-x}}{e^x + e^{-x}}; \coth x = \frac{\cosh x}{\sinh x} = \frac{e^x + e^{-x}}{e^x - e^{-x}};$$

$$\operatorname{sech} x = \frac{1}{\cosh x}; \text{ and } \operatorname{cosech} x = \frac{1}{\sinh x}.$$

Remember the following relations between hyperbolic functions.

- (i) $\cosh^2 x - \sinh^2 x = 1$; (ii) $\sinh 2x = 2 \sinh x \cosh x$;
- (iii) $1 - \tanh^2 x = \operatorname{sech}^2 x$; (iv) $\coth^2 x - 1 = \operatorname{cosech}^2 x$;
- (v) $\cosh 2x = \cosh^2 x + \sinh^2 x$;
- (vi) $\tanh 2x = \frac{2 \tanh x}{1 + \tanh^2 x}$.

To get any of these relations first write the corresponding relation between circular functions. Then convert each circular function into the corresponding hyperbolic function and also change the sign of the term which contains the product of two sines.

Derivatives of hyperbolic functions.

We have, $\frac{d}{dx} \sinh x = \frac{d}{dx} \left\{ \frac{1}{2}(e^x - e^{-x}) \right\} = \frac{1}{2}(e^x + e^{-x}) = \cosh x$.

Similarly we can find the derivatives of other hyperbolic functions. Remember the following results.

$$\frac{d}{dx} \sinh x = \cosh x, \frac{d}{dx} \cosh x = \sinh x, \frac{d}{dx} \tanh x = \operatorname{sech}^2 x,$$



We have, $\frac{d}{dx} \sinh x = \frac{d}{dx} \left\{ \frac{1}{2} (e^x - e^{-x}) \right\} = \frac{1}{2} (e^x + e^{-x}) = \cosh x.$

Similarly we can find the derivatives of other hyperbolic functions. Remember the following results.

$$\frac{d}{dx} \sinh x = \cosh x, \quad \frac{d}{dx} \cosh x = \sinh x, \quad \frac{d}{dx} \tanh x = \operatorname{sech}^2 x,$$

$$\frac{d}{dx} \coth x = -\operatorname{cosech}^2 x, \quad \frac{d}{dx} \operatorname{sech} x = -\operatorname{sech} x \tanh x,$$

$$\frac{d}{dx} \operatorname{cosech} x = -\operatorname{cosech} x \coth x.$$

We note that these results are similar to the corresponding results on circular functions. The only change is that here the derivatives of $\sinh x$, $\cosh x$ and $\tanh x$ have +ive sign placed before them and the derivatives of the remaining three functions have -ive sign placed before them.

§ 5. Inverse hyperbolic functions.

Let $y = \sinh^{-1} x$. Then $\sinh y = x$. Therefore

$\cosh y = \sqrt{(\sinh^2 y + 1)} = \sqrt{(x^2 + 1)}$. Adding these, we get $\sinh y + \cosh y = x + \sqrt{(x^2 + 1)}$

or $\frac{1}{2} (e^y - e^{-y}) + \frac{1}{2} (e^y + e^{-y}) = x + \sqrt{(x^2 + 1)}$ or $e^y = x + \sqrt{(x^2 + 1)}$.

$$\therefore y = \log [x + \sqrt{(x^2 + 1)}].$$

Thus $\sinh^{-1} x = \log [x + \sqrt{(x^2 + 1)}]$.

Similarly, $\cosh^{-1} x = \log [x + \sqrt{(x^2 - 1)}]$;

$$\tanh^{-1} x = \frac{1}{2} \log \frac{1+x}{1-x}; \quad \coth^{-1} x = \frac{1}{2} \log \frac{x+1}{x-1};$$

$$\operatorname{sech}^{-1} x = \log \frac{1+\sqrt{(1-x^2)}}{x}; \quad \operatorname{cosech}^{-1} x = \log \frac{1+\sqrt{(1+x^2)}}{x}.$$

Derivatives of Inverse Hyperbolic functions.

Let $\sinh^{-1} x = y$. Then $x = \sinh y$.

$$\therefore \frac{d}{dx} (x) = \frac{d}{dy} (\sinh y) \text{ or } 1 = \left[\frac{d}{dy} (\sinh y) \right] \frac{dy}{dx} = \cosh y \frac{dy}{dx}.$$

$$\therefore \frac{dy}{dx} = \frac{1}{\cosh y} = \frac{1}{\sqrt{(\sinh^2 y + 1)}} = \frac{1}{\sqrt{(x^2 + 1)}}.$$

$$\therefore \frac{d}{dx} \sinh^{-1} x = \frac{1}{\sqrt{(x^2 + 1)}}.$$

$$\text{Similarly, } \frac{d}{dx} \cosh^{-1} x = \frac{1}{\sqrt{(x^2 - 1)}};$$

$$\frac{d}{dx} \tanh^{-1} x = \frac{1}{1-x^2}; \quad \frac{d}{dx} \coth^{-1} x = -\frac{1}{x^2-1}.$$

Copyrighted material

$$\frac{d}{dx} \operatorname{sech}^{-1} x = \frac{1}{x\sqrt{(1-x^2)}}; \quad \frac{d}{dx} \operatorname{cosech}^{-1} x = -\frac{1}{x\sqrt{(1+x^2)}}.$$

§ 6. Some more methods of differentiation.

(a) **Logarithmic differentiation.** When a function consists of the product or the quotient of a number of functions, we take the logarithm and differentiate. This process, called *logarithmic differentiation*, is also useful when a function of x is raised to a power which is itself a function of x .

(b) **Implicit functions.** If the relation between y and x is represented by an equation from which y cannot be easily expressed in terms of x then y is called an *implicit function* of x . But if y is given



Copyrighted material

Derivatives of hyperbolic functions.

We have, $\frac{d}{dx} \sinh x = \frac{d}{dx} \left\{ \frac{1}{2} (e^x - e^{-x}) \right\} = \frac{1}{2} (e^x + e^{-x}) = \cosh x.$

Similarly we can find the derivatives of other hyperbolic functions. Remember the following results.

$$\begin{aligned}\frac{d}{dx} \sinh x &= \cosh x, \quad \frac{d}{dx} \cosh x = \sinh x, \quad \frac{d}{dx} \tanh x = \operatorname{sech}^2 x, \\ \frac{d}{dx} \coth x &= -\operatorname{cosech}^2 x, \quad \frac{d}{dx} \operatorname{sech} x = -\operatorname{sech} x \tanh x, \\ \frac{d}{dx} \operatorname{cosech} x &= -\operatorname{cosech} x \coth x.\end{aligned}$$

We note that these results are similar to the corresponding results on circular functions. The only change is that here the derivatives of $\sinh x$, $\cosh x$ and $\tanh x$ have +ive sign placed before them and the derivatives of the remaining three functions have -ive sign placed before them.

§ 5. Inverse hyperbolic functions.

Let $y = \sinh^{-1} x$. Then $\sinh y = x$. Therefore

$\cosh y = \sqrt{(\sinh^2 y + 1)} = \sqrt{(x^2 + 1)}$. Adding these, we get $\sinh y + \cosh y = x + \sqrt{(x^2 + 1)}$

or $\frac{1}{2} (e^y - e^{-y}) + \frac{1}{2} (e^y + e^{-y}) = x + \sqrt{(x^2 + 1)}$ or $e^y = x + \sqrt{(x^2 + 1)}$.

$$\therefore y = \log [x + \sqrt{(x^2 + 1)}].$$

Thus $\sinh^{-1} x = \log [x + \sqrt{(x^2 + 1)}]$.

Similarly, $\cosh^{-1} x = \log [x + \sqrt{(x^2 - 1)}]$;

$$\tanh^{-1} x = \frac{1}{2} \log \frac{1+x}{1-x}; \quad \coth^{-1} x = \frac{1}{2} \log \frac{x+1}{x-1};$$

$$\operatorname{sech}^{-1} x = \log \frac{1+\sqrt{1-x^2}}{x}; \quad \operatorname{cosech}^{-1} x = \log \frac{1+\sqrt{1+x^2}}{x}.$$

Derivatives of Inverse Hyperbolic functions.

Let $\sinh^{-1} x = y$. Then $x = \sinh y$.

$$\therefore \frac{d}{dx} (x) = \frac{d}{dy} (\sinh y) \text{ or } 1 = \left[\frac{d}{dy} (\sinh y) \right] \frac{dy}{dx} = \cosh y \frac{dy}{dx}.$$

$$\therefore \frac{dy}{dx} = \frac{1}{\cosh y} = \frac{1}{\sqrt{(\sinh^2 y + 1)}} = \frac{1}{\sqrt{(x^2 + 1)}}.$$

$$\therefore \frac{d}{dx} \sinh^{-1} x = \frac{1}{\sqrt{(x^2 + 1)}}.$$

$$\text{Similarly, } \frac{d}{dx} \cosh^{-1} x = \frac{1}{\sqrt{(x^2 - 1)}};$$

$$\frac{d}{dx} \tanh^{-1} x = \frac{1}{1-x^2}; \quad \frac{d}{dx} \coth^{-1} x = -\frac{1}{x^2-1}.$$

Copyrighted material

$$\frac{d}{dx} \operatorname{sech}^{-1} x = \frac{1}{x\sqrt{(1-x^2)}}; \quad \frac{d}{dx} \operatorname{cosech}^{-1} x = -\frac{1}{x\sqrt{(1+x^2)}}.$$



$$\frac{d}{dx} \operatorname{sech}^{-1} x = \frac{1}{x\sqrt{(1-x^2)}}; \quad \frac{d}{dx} \operatorname{cosech}^{-1} x = -\frac{1}{x\sqrt{(1+x^2)}}.$$

§ 6. Some more methods of differentiation.

(a) **Logarithmic differentiation.** When a function consists of the product or the quotient of a number of functions, we take the logarithm and differentiate. This process, called *logarithmic differentiation*, is also useful when a function of x is raised to a power which is itself a function of x .

(b) **Implicit functions.** If the relation between y and x is represented by an equation from which y cannot be easily expressed in terms of x , then y is called an *implicit function* of x . But if x is given directly in terms of x , then y is called an *explicit function* of x .

To find dy/dx in the case of implicit functions, differentiate each term of the given equation w.r.t. 'x' and solve the resulting equation for dy/dx .

Note. Here the value of dy/dx shall usually contain both x and y .

(c) **Parametric equations.** Sometimes x and y are both expressed in terms of a third variable, say, t . This variable is called a parameter and the equations thus given are known as parametric equations. In the case of parametric equations $x = f_1(t)$ and $y = f_2(t)$, we find dy/dx in the following manner :

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = \frac{dy/dt}{dx/dt}.$$

(d) **Trigonometrical transformations.** Sometimes it is easy to differentiate after making some trigonometrical transformations.

Following formulae of trigonometry are of frequent use :

$$(i) \cos x = 2 \cos^2(x/2) - 1 = 1 - 2 \sin^2(x/2);$$

$$(ii) \sin x = 2 \tan \frac{1}{2}x / (1 + \tan^2 \frac{1}{2}x);$$

$$(iii) \cos x = (1 - \tan^2 \frac{1}{2}x) / (1 + \tan^2 \frac{1}{2}x);$$

$$(iv) \tan x = 2 \tan \frac{1}{2}x / (1 - \tan^2 \frac{1}{2}x);$$

$$(v) \tan^{-1} x + \tan^{-1} y = \tan^{-1} \frac{x+y}{1-xy};$$

$$(vi) \tan^{-1} x - \tan^{-1} y = \tan^{-1} \frac{x-y}{1+xy};$$

$$(vii) 2 \tan^{-1} x = \tan^{-1} 2x / (1 - x^2);$$

$$(viii) 3 \tan^{-1} x = \tan^{-1} \frac{3x - x^3}{1 - 3x^2};$$

$$(ix) \sin 3x = 3 \sin x - 4 \sin^3 x;$$

$$(x) \cos 3x = 4 \cos^3 x - 3 \cos x.$$

(e) Differentiation of a function w.r.t. a function.

Suppose we have to differentiate $f(x)$ w.r.t. $\phi(x)$. Put $\phi(x) = t$.

Copyrighted material

Then

$$\frac{df(x)}{d\phi(x)} = \frac{df(x)}{dt} = \frac{df(x)}{dx} \cdot \frac{dx}{dt} = \frac{df(x)}{dx} / \frac{dt}{dx}.$$

$$\text{Hence } \frac{df(x)}{d\phi(x)} = \frac{df(x)}{dx} / \frac{d\phi(x)}{dx}.$$

§ 7. List of Standard results to be Committed to Memory.



Then

$$\frac{df(x)}{d\phi(x)} = \frac{df(x)}{dt} = \frac{df(x)}{dx} \cdot \frac{dx}{dt} = \frac{df(x)}{dx} / \frac{dt}{dx}.$$

$$\text{Hence } \frac{df(x)}{d\phi(x)} = \frac{df(x)}{dx} / \frac{d\phi(x)}{dx}.$$

§ 7. List of Standard results to be Committed to Memory.

$$\frac{d}{dx} x^n = nx^{n-1}$$

$$\frac{d}{dx} \tan x = \sec^2 x$$

$$\frac{d}{dx} e^x = e^x$$

$$\frac{d}{dx} \cot x = -\operatorname{cosec}^2 x$$

$$\frac{d}{dx} \log_e x = \frac{1}{x}$$

$$\frac{d}{dx} \sec x = \sec x \tan x$$

$$\frac{d}{dx} \log_a x = \frac{1}{x} \log_a e$$

$$\frac{d}{dx} \operatorname{cosec} x = -\operatorname{cosec} x \cot x$$

$$\frac{d}{dx} a^x = a^x \log_e a$$

$$\frac{d}{dx} \sinh x = \cosh x$$

$$\frac{d}{dx} \sin x = \cos x$$

$$\frac{d}{dx} \cosh x = \sinh x$$

$$\frac{d}{dx} \cos x = -\sin x$$

$$\frac{d}{dx} \tanh x = \operatorname{sech}^2 x$$

$$\frac{d}{dx} \coth x = -\operatorname{cosech}^2 x$$

$$\frac{d}{dx} \cot^{-1} x = -\frac{1}{1+x^2}$$

$$\frac{d}{dx} \operatorname{sech} x = -\operatorname{sech} x \tanh x$$

$$\frac{d}{dx} \sec^{-1} x = \frac{1}{x\sqrt{x^2-1}}$$

$$\frac{d}{dx} \operatorname{cosech} x = -\operatorname{cosech} x \coth x$$

$$\frac{d}{dx} \operatorname{cosec}^{-1} x = -\frac{1}{x\sqrt{x^2-1}}$$

$$\frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \sinh^{-1} x = \frac{1}{\sqrt{x^2+1}}$$

$$\frac{d}{dx} \cos^{-1} x = -\frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \cosh^{-1} x = \frac{1}{\sqrt{x^2-1}}$$

$$\frac{d}{dx} \tan^{-1} x = \frac{1}{1+x^2}$$

$$\frac{d}{dx} \tanh^{-1} x = \frac{1}{1-x^2}$$

Solved Examples

Ex. 1. Find the differential coefficients of the following w.r.t. x.

$$(i) \sqrt{2x - x^{-2}} \quad (ii) \frac{\sqrt{5-2x}}{2x+1} \quad (iii) \log \frac{2}{x}$$

$$(iv) \sqrt{(1+\sin x)/(1-\sin x)} \quad (v) \frac{ax^2+b}{ax^2-b} + \frac{ax^2-b}{ax^2+b}$$

$$(vi) \log(\log x).$$

Copyrighted material

$$\begin{aligned} \text{Sol. } (i) \quad & \frac{d}{dx} (2x - x^{-2})^{1/2} = \frac{1}{2}(2x - x^{-2})^{-1/2} (2 + 2x^{-3}) \\ & = (2x - x^{-2})^{-1/2} (1 + x^{-3}) = \frac{1 + x^{-3}}{\sqrt{2x - x^{-2}}}. \end{aligned}$$

$$(ii) \quad \frac{d}{dx} \left\{ \frac{\sqrt{5-2x}}{2x+1} \right\} = \frac{d}{dx} \left\{ (5-2x)^{1/2} \cdot (2x+1)^{-1} \right\}$$



Copyrighted material

Sol. (i) $\frac{d}{dx} (2x - x^{-2})^{1/2} = \frac{1}{2}(2x - x^{-2})^{-1/2} (2 + 2x^{-3})$
 $= (2x - x^{-2})^{-1/2} (1 + x^{-3}) = \frac{1 + x^{-3}}{\sqrt{2x - x^{-2}}}.$

(ii) $\frac{d}{dx} \left\{ \frac{\sqrt{5-2x}}{2x+1} \right\} = \frac{d}{dx} \{(5-2x)^{1/2} \cdot (2x+1)^{-1}\}$
 $= \frac{1}{2}(5-2x)^{-1/2} (-2)(2x+1)^{-1} - (2x+1)^{-2} \cdot 2 \cdot (5-2x)^{1/2}$
 $= \frac{-1}{(2x+1)\sqrt{5-2x}} - \frac{2\sqrt{5-2x}}{(2x+1)^2}$
 $= \frac{-(2x+1)-2(5-2x)}{(2x+1)^2\sqrt{5-2x}} = \frac{2x-11}{(2x+1)^2\sqrt{5-2x}}.$

(iii) $\frac{d}{dx} \log \frac{2}{x} = \frac{d}{dx} (\log 2 - \log x) = -\frac{1}{x}.$

(iv) We have $\sqrt{\left(\frac{1+\sin x}{1-\sin x}\right)} = \sqrt{\left\{\frac{(1+\sin x)(1+\sin x)}{(1-\sin x)(1+\sin x)}\right\}}$
 $= \sqrt{\left\{\frac{(1+\sin x)^2}{1-\sin^2 x}\right\}} = \frac{1+\sin x}{\cos x} = \sec x + \tan x.$

∴ $\frac{d}{dx} \sqrt{\left(\frac{1+\sin x}{1-\sin x}\right)} = \frac{d}{dx} (\sec x + \tan x) = \sec x |\tan x + \sec^2 x.$

(v) $\frac{d}{dx} \left(\frac{ax^2+b}{ax^2-b} + \frac{ax^2-b}{ax^2+b} \right) = \frac{d}{dx} \left(\frac{2a^2x^4+2b^2}{a^2x^4-b^2} \right)$
 $= \frac{8a^2x^3(a^2x^4-b^2)-4a^2x^3(2a^2x^4+2b^2)}{(a^2x^4-b^2)^2}$
 $= -\frac{16a^2b^2x^3}{(a^2x^4-b^2)^2}.$

(vi) $\frac{d}{dx} \{\log(\log x)\} = \frac{1}{\log x} \frac{d}{dx} (\log x) = \frac{1}{x \log x}.$

Ex. 2. Find the differential coefficients of the following w.r.t. x.

(i) $7 \sin x + 2 \log x - e^x + (x^2 - 7x + 4)$

(ii) $(\cos x) \cdot (\log x)$ (iii) $e^{\sin^{-1} x}$

(iv) $e^{ax} \cos(bx+c)$ (v) $\log \sec x$

(vi) $a^{2x} \sinh 2x$ (vii) $\log \frac{1+\sqrt{x}}{1-\sqrt{x}}$

(viii) $\frac{e^x}{1+x}$ (ix) $\frac{\tan x}{x+e^x}$

(x) $\log [\sqrt{1+\log x} - \sin x]$ (xi) $\tan(\log \tan^{-1} \sqrt{x}).$

Sol. (i) Let $y = 7 \sin x + 2 \log x + e^x + (x^2 - 7x + 4).$

Then $\frac{dy}{dx} = 7 \cdot \frac{d}{dx} (\sin x) + 2 \frac{d}{dx} (\log x) - \frac{d}{dx} (e^x) + \frac{d}{dx} (x^2 - 7x + 4)$
 $= 7 \cos x + 2/x - e^x + 2x - 7.$

Copyrighted material



Copyrighted material

DIFFERENTIATION

13

(ii) Let $y = (\cos x) \cdot (\log x)$.

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= (\cos x) \frac{d}{dx} (\log x) + (\log x) \frac{d}{dx} (\cos x) \\ &= (\cos x) (1/x) + (\log x) (-\sin x) \\ &= (1/x) \cos x - (\sin x) \log x. \end{aligned}$$

$$\begin{aligned} \text{(iii) Let } y &= e^{\sin^{-1} x}. \text{ Then } \frac{dy}{dx} = e^{\sin^{-1} x} \frac{d}{dx} \sin^{-1} x \\ &= e^{\sin^{-1} x} \cdot \frac{1}{\sqrt{1-x^2}}. \end{aligned}$$

(iv) Let $y = e^{ax} \cos(bx + c)$.

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= e^{ax} \cdot \frac{d}{dx} \cos(bx + c) + \cos(bx + c) \cdot \frac{d}{dx} e^{ax} \\ &= e^{ax} [(-\sin(bx + c))b] + \cos(bx + c) \cdot e^{ax} \cdot a \\ &= e^{ax} [a \cos(bx + c) - b \sin(bx + c)]. \end{aligned}$$

$$\begin{aligned} \text{(v) Let } y &= \log \sec x. \text{ Then } \frac{dy}{dx} = \frac{1}{\sec x} \cdot \frac{d}{dx} (\sec x) \\ &= \frac{1}{\sec x} \sec x \tan x = \tan x. \end{aligned}$$

$$\begin{aligned} \text{(vi) Let } y &= a^{2x} \sinh 2x. \text{ Then } \frac{dy}{dx} = a^{2x} \cdot \frac{d}{dx} \sinh 2x + \sinh 2x \cdot \frac{d}{dx} a^{2x} \\ &= 2a^{2x} \cosh 2x + (\sinh 2x) \cdot (a^{2x} \log a) \cdot 2 \\ &= 2a^{2x} [\cosh 2x + (\log a) \sinh 2x]. \end{aligned}$$

$$\text{(vii) Let } y = \log \frac{1+\sqrt{x}}{1-\sqrt{x}} = \log(1+\sqrt{x}) - \log(1-\sqrt{x}).$$

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= \frac{1}{1+\sqrt{x}} \cdot \frac{d}{dx} (1+\sqrt{x}) - \frac{1}{1-\sqrt{x}} \cdot \frac{d}{dx} (1-\sqrt{x}) \\ &= \frac{1}{2\sqrt{x}} \left[\frac{1}{1+\sqrt{x}} + \frac{1}{1-\sqrt{x}} \right] \\ &= \frac{1}{2\sqrt{x}} \cdot \frac{2}{1-x} = \frac{1}{(1-x)\sqrt{x}}. \end{aligned}$$

(viii) Let $y = e^x/(1+x)$.

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= \frac{(1+x) \cdot \frac{d}{dx} e^x - e^x \cdot \frac{d}{dx} (1+x)}{(1+x)^2} \\ &= \frac{(1+x)e^x - e^x \cdot 1}{(1+x)^2} = \frac{x e^x}{(1+x)^2}. \end{aligned}$$

(ix) Let $y = \tan x/(x + e^x)$.

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= \frac{(x+e^x) \sec^2 x - (\tan x)(1+e^x)}{(x+e^x)^2} \\ &= \frac{x \sec^2 x + e^x (\sec^2 x - \tan x) - \tan x}{(x+e^x)^2} \end{aligned}$$

Copyrighted material



Copyrighted material

(x) Let $y = \log [\sqrt{1 + \log x} - \sin x]$.

$$\text{Then } \frac{dy}{dx} = \frac{1}{\sqrt{1 + \log x} - \sin x} \cdot \left[\frac{1}{2\sqrt{1 + \log x}} - \cos x \right]$$

(xi) Let $y = \tan(\log \tan^{-1} \sqrt{x})$.

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= \sec^2(\log \tan^{-1} \sqrt{x}) \left[\frac{1}{\tan^{-1}(\sqrt{x})} \cdot \frac{1}{1+x} \cdot \frac{1}{2\sqrt{x}} \right] \\ &= \frac{\sec^2(\log \tan^{-1} \sqrt{x})}{(2\sqrt{x})(1+x)\tan^{-1}(\sqrt{x})} \end{aligned}$$

Problems involving logarithmic differentiation, implicit functions and parametric equations.

Ex. 3. Find (dy/dx) , when

(i) $y = x^{(x)}$

(Delhi 1983, 79)

(ii) $y = (\tan x)^{\log x} + (\cot x)^{\sin x}$,

(iii) $y = (\cot x)^{\cot x} + (\cosh x)^{\cosh x}$, (iv) $y = x^x + x^{\sin x}$.

Sol. (i) Taking logarithm of both sides, we get

$\log y = (x^x) \log x$.

Now differentiating both sides w.r.t. x , we get

$$\frac{1}{y} \frac{dy}{dx} = x^x \cdot \frac{1}{x} + \log x \cdot \frac{d}{dx}(x^x). \quad \dots(1)$$

Let $z = x^x$. Then $\log z = x \log x$.Differentiating w.r.t. x , we have

$$\frac{1}{z} \frac{dz}{dx} = x \frac{1}{x} + \log x. \quad \therefore \frac{dz}{dx} = x^x (1 + \log x).$$

Now (1) gives, $\frac{dy}{dx} = x^{(x)} \cdot [x^{x-1} + x^x (\log x) \cdot (1 + \log x)]$.(ii) We have $y = (\tan x)^{\log x} + (\cot x)^{\sin x}$.Let $(\tan x)^{\log x} = u$ and $(\cot x)^{\sin x} = v$.

Then $y = u + v. \quad \therefore \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$.

Now $\log u = (\log x)(\log \tan x)$.

$$\therefore \frac{1}{u} \cdot \frac{du}{dx} = (\log x) \cdot \frac{\sec^2 x}{\tan x} + \frac{1}{x} \log \tan x$$

or $\frac{du}{dx} = (\tan x)^{\log x} [\cot x \sec^2 x \log x + (1/x) \log \tan x]$.

Again $\log v = (\sin x)(\log \cot x)$.

$$\therefore \frac{1}{v} \frac{dv}{dx} = \sin x \cdot \frac{1}{\cot x} (-\operatorname{cosec}^2 x) + \cos x \log \cot x$$

or $\frac{dv}{dx} = (\cot x)^{\sin x} [\cos x \log \cot x - \sec x]$.

$$\begin{aligned} \text{Hence } \frac{dy}{dx} &= (\tan x)^{\log x} [\cot x \sec^2 x \log x + (1/x) \log \tan x] \\ &\quad + (\cot x)^{\sin x} [\cos x \log \cot x - \sec x]. \end{aligned}$$

Copyrighted material

(iii) Here $y = (\cot x)^{\cot x} + (\cosh x)^{\cosh x}$.Let $(\cot x)^{\cot x} = u$ and $(\cosh x)^{\cosh x} = v$.Then $y = u + v. \quad \text{Therefore } dy/dx = du/dx + dv/dx$.



Copyrighted material

DIFFERENTIATION

15

(iii) Here $y = (\cot x)^{\cot x} + (\cosh x)^{\cosh x}$.Let $(\cot x)^{\cot x} = u$ and $(\cosh x)^{\cosh x} = v$.Then $y = u + v$. Therefore $dy/dx = du/dx + dv/dx$.Now $\log u = \cot x \cdot \log \cot x$.

$$\therefore \frac{1}{u} \frac{du}{dx} = \cot x \cdot \frac{1}{\cot x} (-\operatorname{cosec}^2 x) - \operatorname{cosec}^2 x \cdot \log \cot x$$

$$\text{or } \frac{du}{dx} = -\operatorname{cosec}^2 x (1 + \log \cot x).$$

Again $\log v = \cosh x \log \cosh x$.

$$\therefore \frac{1}{v} \frac{dv}{dx} = \cosh x \cdot \frac{1}{\cosh x} \cdot \sinh x + \sinh x \log \cosh x$$

$$\text{or } \frac{dv}{dx} = v \sinh x (1 + \log \cosh x).$$

Hence

$$\begin{aligned} \frac{dy}{dx} &= -u \operatorname{cosec}^2 x (1 + \log \cot x) + v \sinh x (1 + \log \cosh x) \\ &= -(\cot x)^{\cot x} \operatorname{cosec}^2 x (1 + \log \cot x) \\ &\quad + (\cosh x)^{\cosh x} \sinh x (1 + \log \cosh x). \end{aligned}$$

(iv) We have $y = x^x + (x)^{\sin x}$.Let $x^x = u$ and $(x)^{\sin x} = v$.

$$\text{Then } y = u + v \text{ and } \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}.$$

Now $\log u = x \log x$.

$$\therefore \frac{1}{u} \frac{du}{dx} = x \frac{1}{x} + \log x \text{ or } \frac{du}{dx} = x^x (1 + \log x).$$

Again $\log v = (\sin x) \log x$.

$$\therefore \frac{1}{v} \frac{dv}{dx} = (\sin x) \cdot \frac{1}{x} + (\cos x) \log x$$

$$\text{or } \frac{dv}{dx} = x^{\sin x} \cdot \left[\frac{1}{x} \sin x + (\cos x) \log x \right].$$

$$\text{Hence } \frac{dy}{dx} = x^x \cdot (1 + \log x) + (x)^{\sin x} \left[\frac{1}{x} \sin x + (\cos x) \log x \right].$$

Ex. 4. Find $\frac{dy}{dx}$ if(i) $y = (x)^{\tan x} + (\sin x)^{\cos x}$.

(Delhi 1980)

(ii) $y = (\sin x)^{\cos x} + (\cos x)^{\sin x}$.Sol. (i) Let $u = (x)^{\tan x}$ and $v = (\sin x)^{\cos x}$.

$$\therefore y = u + v \Rightarrow (dy/dx) = (du/dx) + (dv/dx). \quad \dots(1)$$

Now $u = (x)^{\tan x} \Rightarrow \log u = \tan x \log x$.∴ differentiating w.r.t. x , we have

$$(1/u) (du/dx) = (\sec^2 x) \cdot \log x + (\tan x)/x$$

Copyrighted material

$$\text{or } \frac{du}{dx} = (x)^{\tan x} \left[\sec^2 x \log x + \frac{\tan x}{x} \right].$$



Copyrighted material

$$\text{or } \frac{du}{dx} = (x)^{\tan x} \left[\sec^2 x \log x + \frac{\tan x}{x} \right].$$

$$\text{Similarly } \frac{dv}{dx} = (\sin x)^{\cos x} [(-\sin x) \cdot \log \sin x + (\cos x) \cdot \cot x].$$

∴ from (1), we get

$$\begin{aligned} (dy/dx) &= (du/dx) + (dv/dx) \\ &= (x)^{\tan x} [\sec^2 x \log x + (\tan x)/x] \\ &\quad + (\sin x)^{\cos x} [\cos x \cot x - \sin x \cdot \log \sin x]. \end{aligned}$$

(ii) Proceed exactly as in part (i).

Ex. 5. Find (dy/dx) if $\sin y = \log_{\sin x} \cos x$. (Delhi 1980)

Sol. We have

$$\sin y = \log_{\sin x} \cos x = \frac{\log_e \cos x}{\log_e \sin x}. \quad [\text{Note}]$$

Differentiating both sides w.r.t. x , we have

$$(\cos y) \cdot \frac{dy}{dx} = \frac{(-\tan x) \cdot \log \sin x - (\cot x) \cdot \log \cos x}{(\log_e \sin x)^2}.$$

$$\therefore \frac{dy}{dx} = -\frac{(\tan x) \log \sin x + (\cot x) \log \cos x}{(\cos y) (\log \sin x)^2}.$$

Ex. 6. Find (dy/dx) , when

(i) $x = a(t - \sin t), y = a(1 - \cos t)$.

(ii) $x = a(\cos t + \log \tan t/2), y = a \sin t$.

(iii) $y = \tan^{-1} \frac{2t}{1-t^2}, x = \sin^{-1} \frac{2t}{1+t^2}$.

(iv) $x = a \sqrt{\left(\frac{t^2-1}{t^2+1}\right)}, y = at \sqrt{\left(\frac{t^2-1}{t^2+1}\right)}$.

Sol. (i) We have $x = a(t - \sin t), y = a(1 - \cos t)$.

$$\therefore \frac{dx}{dt} = a(1 - \cos t) \text{ and } \frac{dy}{dt} = a \sin t.$$

$$\text{Now } \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{a \sin t}{a(1 - \cos t)} = \frac{2 \sin t / 2 \cos t / 2}{2 \sin^2 t / 2} = \cot t / 2.$$

(ii) Here $x = a(\cos t + \log \tan t/2), y = a \sin t$.

$$\begin{aligned} \therefore \frac{dx}{dt} &= a \left[-\sin t + \frac{1}{\tan t / 2} (\sec^2 t / 2) \cdot \frac{1}{2} \right] \\ &= a \left(-\sin t + \frac{1}{2 \sin t / 2 \cos t / 2} \right) \\ &= a \left[\frac{1}{\sin t} - \sin t \right] = a \left[\frac{1 - \sin^2 t}{\sin t} \right] = \frac{a \cos^2 t}{\sin t}. \end{aligned}$$

Again $(dy/dt) = a \cos t$.

$$\text{Now } \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = a \cos t \cdot \frac{\sin t}{a \cos^2 t} = \tan t.$$

Copyrighted material

(iii) Let $t = \tan \theta$. Then

$$y = \tan^{-1} \frac{2 \tan \theta}{1 - \tan^2 \theta} = \tan^{-1} \tan 2\theta = 2\theta = 2 \tan^{-1} t.$$

$$\therefore \frac{dy}{dt} = 2/(1 + t^2)$$



Copyrighted material

DIFFERENTIATION

17

(iii) Let $t = \tan \theta$. Then

$$y = \tan^{-1} \frac{2 \tan \theta}{1 - \tan^2 \theta} = \tan^{-1} \tan 2\theta = 2\theta = 2 \tan^{-1} t.$$

$$\therefore (dy/dt) = 2/(1+t^2).$$

$$\begin{aligned} \text{Again } x &= \sin^{-1} \frac{2t}{1+t^2} = \sin^{-1} \frac{2 \tan \theta}{1 + \tan^2 \theta} \\ &= \sin^{-1} (\sin 2\theta) = 2\theta = 2 \tan^{-1} t. \end{aligned}$$

$$\therefore (dx/dt) = 2/(1+t^2).$$

$$\text{Now } \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2/(1+t^2)}{2/(1+t^2)} = 1.$$

(iv) Here $x = a \sqrt{\frac{t^2 - 1}{t^2 + 1}}$ and $y = at \sqrt{\frac{t^2 - 1}{t^2 + 1}}$.We have, $\log y = \log a + \log t + \frac{1}{2} \log(t^2 - 1) - \frac{1}{2} \log(t^2 + 1)$.

$$\therefore \frac{1}{y} \frac{dy}{dt} = \frac{1}{t} + \frac{t}{t^2 - 1} - \frac{t}{t^2 + 1} = \frac{t^4 + 2t^2 - 1}{t(t^4 - 1)}$$

$$\text{or } \frac{dy}{dt} = a \sqrt{\frac{t^2 - 1}{t^2 + 1}} \cdot \frac{t^4 + 2t^2 - 1}{(t^4 - 1)}.$$

Again $\log x = \log a + \frac{1}{2} \log(t^2 - 1) - \frac{1}{2} \log(t^2 + 1)$.

$$\therefore \frac{1}{x} \frac{dx}{dt} = \frac{t}{t^2 - 1} - \frac{t}{t^2 + 1} = \frac{2t}{t^4 - 1}$$

$$\text{or } \frac{dx}{dt} = a \sqrt{\frac{t^2 - 1}{t^2 + 1}} \cdot \frac{2t}{t^4 - 1}.$$

$$\text{Now } \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{t^4 + 2t^2 - 1}{2t}.$$

**Ex. 7 (a). If $\sin y = x \sin(a+y)$, prove that

$$\frac{dy}{dx} = \frac{\sin^2(a+y)}{\sin a} \quad (\text{Delhi 1982, 75, 72})$$

(b) If $x^y = e^{x-y}$, prove that

$$\frac{dy}{dx} = \frac{\log x}{(1+\log x)^2} \quad (\text{Delhi 1981, 76; Meerut 81S})$$

(c) If $x = e^{\tan^{-1}((y-x^2)/x^2)}$, find (dy/dx) , expressing it as a function of x only:(d) If $x/(x-y) = \log \{a/(x-y)\}$, prove that

$$(dy/dx) = 2 - (x/y). \quad (\text{Delhi 1983, 74})$$

Sol. (a) We have $x = \sin y / \sin(a+y)$.Differentiating both sides w.r.t. x , we get

$$1 = \frac{\cos y \sin(a+y) - \sin y \cos(a+y)}{\sin^2(a+y)} \cdot \frac{dy}{dx}$$

Copyrighted material

$$\text{or } 1 = \frac{\sin \{(a+y)-y\}}{\sin^2(a+y)} \frac{dy}{dx}. \quad \text{Hence } \frac{dy}{dx} = \frac{\sin^2(a+y)}{\sin a}.$$



Copyrighted material

18

DIFFERENTIAL CALCULUS

or $1 = \frac{\sin \{(a+y)-y\}}{\sin^2(a+y)} \frac{dy}{dx}$. Hence $\frac{dy}{dx} = \frac{\sin^2(a+y)}{\sin a}$.

(b) We have, $x^y = e^{x-y}$.

$$\therefore y \log x = (x-y) \log e = x-y \text{ or } y + y \log x = x$$

or $y = x/(1+\log x)$. Hence $\frac{dy}{dx} = \frac{1(1+\log x) - x \cdot (1/x)}{(1+\log x)^2}$
 $= \log x/(1+\log x)^2$.

(c) We have $x = e^{\tan^{-1}((y-x^2)/x^2)}$.

Therefore $\log x = \tan^{-1} \frac{y-x^2}{x^2} \log e$

or $\frac{y-x^2}{x^2} = \tan \log x \text{ or } y = x^2 + x^2 \tan \log x$.

$\therefore \frac{dy}{dx} = 2x + x^2 (\sec^2 \log x) \cdot \frac{1}{x} + 2x \cdot \tan \log x$
 $= x [2 + \sec^2 \log x + 2 \tan \log x]$.

(d) We have $x/(x-y) = \log \{a/(x-y)\}$.

Differentiating w.r.t. x , we have

$$\frac{d}{dx} \left(\frac{x}{x-y} \right) = \frac{d}{dx} \left(\log \frac{a}{x-y} \right) = \frac{d}{dx} [\log a - \log(x-y)]$$

or $\frac{1 \cdot (x-y) - \{1 - (dy/dx)\} \cdot x}{(x-y)^2} = \left[0 - \frac{1}{(x-y)} \left(1 - \frac{dy}{dx} \right) \right]$

or $(x-y) - x + x(dy/dx) = -(x-y) + (x-y)(dy/dx)$

or $y(dy/dx) = 2y - x \text{ i.e., } dy/dx = 2 - (x/y)$. Proved.

Ex. 8. Find (dy/dx) if $y = \tan^{-1} \{(ax-b)/(bx+a)\}$.

(Delhi 1980)

Sol. We have $y = \tan^{-1} \left(\frac{ax-b}{bx+a} \right)$.

$\therefore \frac{dy}{dx} = \frac{1}{1 + (ax-b)^2/(bx+a)^2} \cdot \frac{d}{dx} \left(\frac{ax-b}{bx+a} \right)$

or $\frac{dy}{dx} = \frac{(bx+a)^2}{(bx+a)^2 + (ax-b)^2} \cdot \frac{a(bx+a) - b(ax-b)}{(bx+a)^2}$
 $= \frac{a^2 + b^2}{(a^2 + b^2)x^2 + (a^2 + b^2)} = \frac{1}{1+x^2}$.



Copyrighted material



3

Expansions of Functions

§ 1. Taylor's series.

Suppose $f(x)$ possesses continuous derivatives of all orders in the interval $[a, a+h]$. Then for every positive integral value of n , we have

$$f(a+h) = f(a) + hf'(a) + \frac{h^2}{2!}f''(a) + \dots + \frac{h^{n-1}}{(n-1)!}f^{(n-1)}(a) + R_n, \quad \dots(1)$$

where $R_n = \frac{h^n}{n!}f^{(n)}(a+\theta h)$, $(0 < \theta < 1)$.

Suppose $R_n \rightarrow 0$, as $n \rightarrow \infty$. Then taking limits of both sides of (1) when $n \rightarrow \infty$, we get

$$f(a+h) = f(a) + hf'(a) + \frac{h^2}{2!}f''(a) + \dots + \frac{h^n}{n!}f^{(n)}(a) + \dots \quad \dots(2)$$

The series given in (2) is known as **Taylor's infinite series** for the expansion of $f(a+h)$ as a **power series** in h .

§ 2. Maclaurin's series.

Suppose $f(x)$ possesses continuous derivatives of all orders in the interval $[0, x]$. Then for every positive integral value of n , we have

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2!}f''(0) + \dots + \frac{x^{n-1}}{(n-1)!}f^{(n-1)}(0) + R_n, \quad \dots(1)$$

where $R_n = \frac{x^n}{n!}f^{(n)}(\theta x)$, $(0 < \theta < 1)$.

Suppose $R_n \rightarrow 0$, as $n \rightarrow \infty$. Then taking limits of both sides of (1) when $n \rightarrow \infty$, we get

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2!}f''(0) + \dots + \frac{x^n}{n!}f^{(n)}(0) + \dots \quad \dots(2)$$

The series given in (2) is known as **Maclaurin's infinite series** for the expansion of $f(x)$ as a **power series** in x . Maclaurin's series is a particular case of Taylor's series. If in Taylor's series we put $a = 0$ and $h = x$ we get Maclaurin's series.

Maclaurin's expansion of $f(x)$ fails if any of the functions $f(x)$, $f'(x)$, $f''(x)$, ... becomes infinite or discontinuous at any point of the interval $[0, x]$ or if R_n does not tend to zero as $n \rightarrow \infty$.

Copyrighted material

§ 3. Formal expansions of functions.

We have seen that for the validity of the expansion of a function $f(x)$ as an infinite Maclaurin's series, it is necessary that $R_n \rightarrow 0$ as $n \rightarrow \infty$. But to examine the behaviour of R_n as $n \rightarrow \infty$ is a difficult job because in many cases it is not possible to find a general expression for the n th derivatives of the functions to be expanded. So in this



Copyrighted material

§ 3. Formal expansions of functions.

We have seen that for the validity of the expansion of a function $f(x)$ as an infinite Maclaurin's series, it is necessary that $R_n \rightarrow 0$ as $n \rightarrow \infty$. But to examine the behaviour of R_n as $n \rightarrow \infty$ is a difficult job because in many cases it is not possible to find a general expression for the n th derivatives of the functions to be expanded. So in this chapter we shall simply obtain *formal expansion* of a function $f(x)$ without showing that $R_n \rightarrow 0$ as $n \rightarrow \infty$.

Such an expansion will not give us any idea of the range of values of x for which the expansion is valid. To obtain such an expansion of $f(x)$ we have only to calculate the values of its derivatives for $x = 0$ and substitute them in the infinite Maclaurin's series

$$f(x) = f(0) + \frac{x}{1!} f'(0) + \dots + \frac{x^n}{n!} f^{(n)}(0) + \dots$$

For the convenience of the students we shall now give *formal proofs* of Maclaurin's and Taylor's theorems without bothering about the nature of R_n as $n \rightarrow \infty$.

Maclaurin's theorem.

(Gorakhpur 87; Meerut 84; Magadh 84; Bihar 82; Kashmir 83)

Let $f(x)$ be a function of x which possesses continuous derivatives of all orders in the interval $[0, x]$. Assuming that $f(x)$ can be expanded as an infinite power series in x , we have

$$f(x) = f(0) + \frac{x}{1!} f'(0) + \frac{x^2}{2!} f''(0) + \dots + \frac{x^n}{n!} f^{(n)}(0) + \dots$$

Proof. Suppose $f(x) = A_0 + A_1 x + A_2 x^2 + A_3 x^3 + \dots$... (1)

Let the expansion (1) be differentiable term by term any number of times. Then by successive differentiation, we have

$$f'(x) = A_1 + 2A_2 x + 3A_3 x^2 + 4A_4 x^3 + \dots,$$

$$f''(x) = 2A_2 + 3 \cdot 2 A_3 x + 4 \cdot 3 A_4 x^2 + \dots,$$

$$f'''(x) = 3 \cdot 2 \cdot 1 A_3 + 4 \cdot 3 \cdot 2 A_4 x + \dots, \text{ and so on.}$$

Putting $x = 0$ in each of these relations, we get

$$f(0) = A_0, f'(0) = A_1, f''(0) = 2! A_2, f'''(0) = 3! A_3, \dots$$

Substituting these values of A_0, A_1, A_2, \dots in (1), we get

$$f(x) = f(0) + x f'(0) + \frac{x^2}{2!} f''(0) + \dots + \frac{x^n}{n!} f^{(n)}(0) + \dots$$

This is Maclaurin's theorem. If we denote $f(x)$ by y , then Maclaurin's theorem can be written in either of the following ways :

$$y = (y)_0 + \frac{x}{1!} (y_1)_0 + \frac{x^2}{2!} (y_2)_0 + \frac{x^3}{3!} (y_3)_0 + \dots + \frac{x^n}{n!} (y_n)_0 + \dots$$

Copyrighted material

or $y = y(0) + \frac{x}{1!} y_1(0) + \frac{x^2}{2!} y_2(0) + \dots + \frac{x^n}{n!} y_n(0) + \dots$

Taylor's theorem.

(Gorakhpur 1988; Bihar 82; Vikram 88;
Meerut 85; Magadh 87; Jiwaji 90)



Copyrighted material

$$\text{or } y = y(0) + \frac{x}{1!} y_1(0) + \frac{x^2}{2!} y_2(0) + \dots + \frac{x^n}{n!} y_n(0) + \dots$$

Taylor's theorem.(Gorakhpur 1988; Bihar 82; Vikram 88;
Meerut 85; Magadh 87; Jiwaji 90)

Let $f(x)$ be a function of x which possesses continuous derivatives of all orders in the interval $[a, a+h]$. Assuming that $f(a+h)$ can be expanded as an infinite power series in h , we have

$$f(a+h) = f(a) + hf'(a) + \frac{h^2}{2!} f''(a) + \dots + \frac{h^n}{n!} f^{(n)}(a) + \dots$$

Proof. Suppose $f(a+h) = A_0 + A_1 h + A_2 h^2 + A_3 h^3 + \dots \quad \dots(1)$ Let the expansion (1) be differentiable term by term any number of times w.r.t. h . Then by successive differentiation w.r.t. h , we have

$$f'(a+h) = A_1 + 2A_2 h + 3A_3 h^2 + \dots,$$

$$f''(a+h) = 2A_2 + 3 \cdot 2A_3 h + \dots,$$

$$f'''(a+h) = 3 \cdot 2 \cdot 1 A_3 + \dots, \text{ and so on.}$$

Putting $h=0$ in each of the above relations, we get

$$f(a) = A_0, f'(a) = A_1, f''(a) = 2! A_2, f'''(a) = 3! A_3, \text{ and so on.}$$

$$\therefore A_0 = f(a), A_1 = f'(a), A_2 = \frac{1}{2!} f''(a), A_3 = \frac{1}{3!} f'''(a),$$

and so on.

Substituting these values of $A_0, A_1, A_2, A_3, \dots$ in (1), we get

$$f(a+h) = f(a) + hf'(a) + \frac{h^2}{2!} f''(a) + \dots + \frac{h^n}{n!} f^{(n)}(a) + \dots$$

This is Taylor's theorem. Another useful form is obtained on replacing h by $(x-a)$. Thus

$$f(x) = f(a) + (x-a)f'(a) + \frac{(x-a)^2}{2!} f''(a) + \dots + \frac{(x-a)^n}{n!} f^{(n)}(a) + \dots,$$

which is an expansion of $f(x)$ as a power series in $(x-a)$.Note. If we expand $f(x+h)$, by Taylor's theorem, as a power series in h , then the result is as follows :

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2!} f''(x) + \dots + \frac{h^n}{n!} f^{(n)}(x) + \dots$$

§ 4. Some Important Expansions.**1. Expansion of e^x . (Exponential series).**Let $f(x) = e^x$. Then $f(0) = e^0 = 1$; $f^{(n)}(x) = e^x$ so that $f^{(n)}(0) = e^0 = 1$, where $n = 1, 2, 3, \dots$

Substituting these values in Maclaurin's series

Copyrighted material

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2!} f''(0) + \frac{x^3}{3!} f'''(0) + \dots$$

$$\text{we set } e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots$$



Copyrighted material

EXPANSIONS OF FUNCTIONS

61

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2!}f''(0) + \frac{x^3}{3!}f'''(0) + \dots$$

we get $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots$

2. Expansion of $\sin x$. (Sine series)Let $f(x) = \sin x$. Then $f(0) = 0$,

$$f'(x) = \cos x \text{ so that } f'(0) = 1,$$

$$f''(x) = -\sin x \text{ so that } f''(0) = 0,$$

$$f'''(x) = -\cos x \text{ so that } f'''(0) = -1, \text{ and so on.}$$

In general, $f^n(x) = \sin(x + \frac{1}{2}n\pi)$ so that $f^n(x) = \sin(\frac{1}{2}n\pi)$.When $n = 2m$, $f^n(0) = \sin m\pi = 0$ and when $n = 2m + 1$,

$$\begin{aligned} f^n(0) &= \sin\{\frac{1}{2}(2m+1)\pi\} = \sin(m\pi + \frac{1}{2}\pi) \\ &= (-1)^m \sin(\frac{1}{2}\pi) = (-1)^m. \end{aligned}$$

Substituting these values in Maclaurin's series, we get

$$\sin x = 0 + x \cdot 1 + 0 + \frac{x^3}{3!}(-1) + 0 + \dots$$

$$+ 0 + (-1)^m \frac{x^{2m+1}}{(2m+1)!} + \dots$$

or $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + (-1)^m \frac{x^{2m+1}}{(2m+1)!} + \dots$ (Meerut 1985)

Similarly, we may obtain the Cosine series :

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots + (-1)^m \frac{x^{2m}}{(2m)!} + \dots$$

3. Expansion of $\log(1+x)$. (Meerut 1976, 89; Luck. 83)Let $f(x) = \log(1+x)$. Then $f(0) = \log 1 = 0$;

$$f^n(x) = \frac{(-1)^{n-1}(n-1)!}{(x+1)^n}$$

so that $f^n(0) = (-1)^{n-1}(n-1)!$, where $n = 1, 2, 3, \dots$

$$\therefore f'(0) = (-1)^{1-1}(1-1)! = 1,$$

$$f''(0) = (-1)^{2-1}(2-1)! = -1!,$$

$$f'''(0) = (-1)^{3-1}(3-1)! = 2!,$$

$$f^{iv}(0) = (-1)^{4-1}(4-1)! = 3!, \text{ and so on,}$$

Substituting the values of $f(0), f'(0), f''(0)$, etc. in Maclaurin's series

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2!}f''(0) + \dots + \frac{x^n}{n!}f^n(0) + \dots,$$

we get $\log(1+x) = 0 + x \cdot 1 - \frac{x^2}{2!} \cdot 1! + \frac{x^3}{3!} \cdot 2! - \frac{x^4}{4!} \cdot 3! + \dots$

$$+ \frac{x^n}{n!}(-1)^{n-1}(n-1)! + \dots$$

Copyrighted material

or $\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots + (-1)^{n-1} \frac{x^n}{n} + \dots$



Copyrighted material

$$\text{or } \log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots + (-1)^{n-1} \frac{x^n}{n} + \dots$$

Note. The function $\log x$ does not possess Maclaurin's series expansion because it is not defined at $x = 0$.

4. Expansion of $(1+x)^n$. (Binomial series).

Let $f(x) = (1+x)^n$. Then $f(0) = 1$;
 $f'(x) = n(n-1)(n-2)\dots(n-m+1)(1+x)^{n-m}$
 so that $f'(0) = n(n-1)\dots(n-m+1)$, where $m = 1, 2, 3, \dots$
 $\therefore f'(0) = n, f''(0) = n(n-1), f'''(0) = n(n-1)(n-2)$
 and so on.

Substituting the values of $f(0), f'(0), f''(0)$ etc. in Maclaurin's series for $f(x)$, we get

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!} x^2 + \dots + \frac{n(n-1)\dots(n-m+1)}{m!} x^m + \dots$$

Solved Examples

Ex. 1. Expand the following by Maclaurin's theorem :

- (i) a^x
- (ii) $\tan x$ (Meerut 1981; Vikram 87; Agra 81; Kashmir 84)
- (iii) $\log \sec x$ (Rohilkhand 1991; Meerut 82, 86, 88, 95, 98)
- (iv) $\log \cos x$ (v) $\sec x$
- (vi) $\log(\sec x + \tan x)$.

Sol. (i) Let $f(x) = a^x$. Then $f(0) = a^0 = 1$,

$f'(x) = a^x \log a$ so that $f'(0) = \log a$,

$f''(x) = a^x (\log a)^2$ so that $f''(0) = (\log a)^2$,

$f'''(x) = a^x (\log a)^3$ so that $f'''(0) = (\log a)^3$, and so on.

In general, $f^n(x) = a^x (\log a)^n$ so that $f^n(0) = (\log a)^n$.

Now by Maclaurin's theorem, we have

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2!} f''(0) + \frac{x^3}{3!} f'''(0) + \dots + \frac{x^n}{n!} f^n(0) + \dots$$

$$\therefore a^x = 1 + x \log a + \frac{x^2}{2!} (\log a)^2 + \frac{x^3}{3!} (\log a)^3 + \dots + \frac{x^n}{n!} (\log a)^n + \dots$$

(ii) Let $y = \tan x$. Then $(y)_0 = \tan 0 = 0$,

$y_1 = \sec^2 x = 1 + \tan^2 x = 1 + y^2$ so that

$$(y_1)_0 = 1 + (y)_0^2 = 1 + 0 = 1,$$

Copyrighted material

$y_2 = 2yy_1$ so that $(y_2)_0 = 2(y)_0 (y_1)_0 = 2 \times 0 \times 1 = 0$,

$y_3 = 2y_1y_1 + 2yy_2 = 2y_1^2 + 2yy_2$ so that $(y_3)_0 = 2 \times 1^2 + 0 = 2$,



$y_2 = 2yy_1$ so that $(y_2)_0 = 2(y_0)(y_1)_0 = 2 \times 0 \times 1 = 0$,

$y_3 = 2y_1y_1 + 2yy_2 = 2y_1^2 + 2yy_2$ so that $(y_3)_0 = 2 \times 1^2 + 0 = 2$,

$y_4 = 4y_1y_2 + 2y_1y_2 + 2yy_3 = 6y_1y_2 + 2yy_3$ so that

$$(y_4)_0 = 6 \times 1 \times 0 + 2 \times 0 \times 2 = 0,$$

$y_5 = 6y_2^2 + 6y_1y_3 + 2y_1y_3 + 2yy_4 = 6y_2^2 + 8y_1y_3 + 2yy_4$ so that

$$(y_5)_0 = 0 + 8 \times 1 \times 2 + 0 = 16, \text{ and so on.}$$

Now by Maclaurin's theorem, we have

$$y = (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \frac{x^3}{3!}(y_3)_0 + \frac{x^4}{4!}(y_4)_0 + \frac{x^5}{5!}(y_5)_0 + \dots$$

$$\therefore \tan x = 0 + x \cdot 1 + \frac{x^2}{2!} \cdot 0 + \frac{x^3}{3!} \cdot 2 + \frac{x^4}{4!} \cdot 0 + \frac{x^5}{5!} \cdot 16 + \dots$$

$$= x + \frac{x^3}{3} + \frac{2}{15}x^5 + \dots$$

(iii) Let $y = \log \sec x$. Then $(y)_0 = \log \sec 0 = \log 1 = 0$,

$$y_1 = \frac{1}{\sec x} \cdot \sec x \tan x = \tan x \text{ so that } (y_1)_0 = 0,$$

$$y_2 = \sec^2 x = 1 + \tan^2 x = 1 + y_1^2 \text{ so that } (y_2)_0 = 1 + (y_1)_0^2 = 1,$$

$$y_3 = 2y_1y_2 \text{ so that } (y_3)_0 = 2(y_1)_0(y_2)_0 = 0,$$

$$y_4 = 2y_2^2 + 2y_1y_3 \text{ so that } (y_4)_0 = 2 \times 1^2 + 0 = 2,$$

$$y_5 = 4y_2y_3 + 2y_2y_4 + 2y_1y_4 = 6y_2y_3 + 2y_1y_4 \text{ so that}$$

$$(y_5)_0 = 6 \times 1 \times 0 + 2 \times 0 \times 2 = 0,$$

$$y_6 = 6y_3^2 + 6y_2y_4 + 2y_2y_4 + 2y_1y_5 = 6y_3^2 + 8y_2y_4 + 2y_1y_5 \text{ so that}$$

$$(y_6)_0 = 0 + 8 \times 1 \times 2 + 0 = 16, \text{ and so on.}$$

Now by Maclaurin's theorem, we have

$$y = (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \frac{x^3}{3!}(y_3)_0 + \frac{x^4}{4!}(y_4)_0 + \dots$$

$$\therefore \log \sec x = 0 + x \cdot 0 + \frac{x^2}{2!} \cdot 1 + \frac{x^3}{3!} \cdot 0 + \frac{x^4}{4!} \cdot 2 + \frac{x^5}{5!} \cdot 0$$

$$+ \frac{x^5}{6!} \cdot 16 + \dots$$

$$= \frac{x^2}{2} + \frac{x^4}{12} + \frac{x^6}{45} + \dots$$

(iv) Let $y = \log \cos x$. Then $(y)_0 = 0$,

$$y_1 = \frac{-\sin x}{\cos x} = -\tan x \text{ so that } (y_1)_0 = 0,$$

$$y_2 = -\sec^2 x = -(1 + \tan^2 x) = -(1 + y_1^2) = -1 - y_1^2$$

$$\text{so that } (y_2)_0 = -1 - 0 = -1,$$

$$y_3 = -2y_1y_2 \text{ so that } (y_3)_0 = 0,$$

Copyrighted material

$$y_4 = -2y_2^2 - 2y_1y_3 \text{ so that } (y_4)_0 = -2(-1)^2 - 0 = -2,$$

$$(y_5)_0 = 0, (y_6)_0 = -16, \text{ and so on.}$$

[For calculation of $(y_5)_0$ and $(y_6)_0$ proceed as in part (iii)]

Now substituting these values in Maclaurin's theorem, we get



Copyrighted material

$y_4 = -2y_2^2 - 2y_1y_3$ so that $(y_4)_0 = -2(-1)^2 - 0 = -2$,
 $(y_5)_0 = 0$, $(y_6)_0 = -16$, and so on.

[For calculation of $(y_5)_0$ and $(y_6)_0$ proceed as in part (iii)]

Now substituting these values in Maclaurin's theorem, we get

$$\begin{aligned}\log \cos x &= 0 + x \cdot 0 + \frac{x^2}{2!} \cdot (-1) + \frac{x^3}{3!} \cdot 0 + \frac{x^4}{4!} \cdot (-2) + \frac{x^5}{5!} \cdot 0 \\ &\quad + (x^6/6!) \cdot (-16) + \dots \\ &= -(x^2/2) - (x^4/12) - (x^6/45) + \dots\end{aligned}$$

(v) Let $y = \sec x$. Then $(y)_0 = \sec 0 = 1$,

$y_1 = \sec x \tan x$ so that $(y_1)_0 = 1 \times 0 = 0$,

$$\begin{aligned}y_2 &= \sec x \sec^2 x + \sec x \tan x \tan x = \sec^3 x + \sec x \tan^2 x \\ &= \sec^3 x + \sec x (\sec^2 x - 1) = 2 \sec^3 x - \sec x = 2y^3 - y\end{aligned}$$

so that $(y_2)_0 = 2 \times 1^3 - 1 = 1$,

$y_3 = 6y^2y_1 - y_1$ so that $(y_3)_0 = 0 - 0 = 0$,

$$\begin{aligned}y_4 &= 6y^2y_2 + 12yy_1^2 - y_2 \text{ so that } (y_4)_0 = 6 \times 1^2 \times 1 + 0 - 1 = 5, \\ &\quad \text{and so on.}\end{aligned}$$

Now by Maclaurin's theorem, we have

$$\begin{aligned}y &= (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \frac{x^3}{3!}(y_3)_0 + \frac{x^4}{4!}(y_4)_0 + \dots \\ \therefore \sec x &= 1 + x \cdot 0 + \frac{x^2}{2!} \cdot 1 + \frac{x^3}{3!} \cdot 0 + \frac{x^4}{4!} \cdot 5 + \dots \\ &= 1 + (x^2/2!) + (5x^4/4!) + \dots\end{aligned}$$

(vi) Let $y = \log(\sec x + \tan x)$. Then $(y)_0 = \log(1 + 0) = 0$,

$$y_1 = \frac{\sec x \tan x + \sec^2 x}{\sec x + \tan x} = \sec x \text{ so that } (y_1)_0 = \sec 0 = 1,$$

$y_2 = \sec x \tan x$ so that $(y_2)_0 = 1 \times 0 = 0$,

$$y_3 = \sec x \sec^2 x + \sec x \tan^2 x = \sec^3 x + \sec x (\sec^2 x - 1)$$

$$= 2 \sec^3 x - \sec x = 2y^3 - y_1 \text{ so that } (y_3)_0 = 2 \times 1^3 - 1 = 1,$$

$y_4 = 6y^2y_2 - y_2$ so that $(y_4)_0 = 0$,

$$\begin{aligned}y_5 &= 6y^2y_3 + 12y_1y_2^2 - y_3 \text{ so that } (y_5)_0 = 6 \times 1^2 \times 1 + 0 - 1 = 5, \\ &\quad \text{and so on.}\end{aligned}$$

Substituting these values in Maclaurin's theorem, we get

$$\begin{aligned}\log(\sec x + \tan x) &= 0 + x \cdot 1 + \frac{x^2}{2!} \cdot 0 + \frac{x^3}{3!} \cdot 1 + \frac{x^4}{4!} \cdot 0 \\ &\quad + \frac{x^5}{5!} \cdot 5 + \dots \\ &= x + (x^3/6) + (x^5/24) + \dots\end{aligned}$$

**Ex. 2. Use Maclaurin's formula to show that

Copyrighted material

$$e^x \sec x = 1 + x + \frac{2x^2}{2!} + \frac{4x^3}{3!} + \dots$$

(Garhwal 1983; Kanpur 80; Gorakhpur 83)



Copyrighted material

EXPANSIONS OF FUNCTIONS

65

$$e^x \sec x = 1 + x + \frac{2x^2}{2!} + \frac{4x^3}{3!} + \dots$$

(Garhwal 1983; Kanpur 80; Gorakhpur 83)

Sol. Let $y = e^x \sec x$. Then $(y)_0 = 1$, $y_1 = e^x \sec x + e^x \sec x \tan x = y + y \tan x$ so that $(y_1)_0 = 1$, $y_2 = y_1 + y_1 \tan x + y \sec^2 x$ so that $(y_2)_0 = 1 + 0 + 1 = 2$, $y_3 = y_2 + y_2 \tan x + 2y_1 \sec^2 x + 2y \sec^2 x \tan x$ so that $(y_3)_0 = 2 + 2 = 4$, and so on.

Substituting these values in Maclaurin's theorem, we get

$$e^x \sec x = 1 + x + \frac{2x^2}{2!} + \frac{4x^3}{3!} + \dots$$

Ex. 3. Expand the following functions by Maclaurin's theorem :

*(i) $e^{\sin x}$ (Meerut 1990S; Gorakhpur 86; G.N.U. 81; K.U. 89)(ii) $e^x \cos x$ (iii) $e^x \log(1+x)$ (iv) $\log(1+\sin x)$ (Gorakhpur 1989; Kanpur 86, Meerut 92)Sol. (i) Let $y = e^{\sin x}$. Then $(y)_0 = e^{\sin 0} = e^0 = 1$, $y_1 = e^{\sin x} \cos x = y \cos x$ so that $(y_1)_0 = (y)_0 \cos 0 = 1 \times 1 = 1$, $y_2 = y_1 \cos x - y \sin x$ so that $(y_2)_0 = 1 \times 1 - 1 \times 0 = 1$, $y_3 = y_2 \cos x - y_1 \sin x - y_1 \sin x - y \cos x$ $= y_2 \cos x - 2y_1 \sin x - y \cos x$ so that $(y_3)_0 = 1 - 0 - 1 = 0$, $y_4 = y_3 \cos x - 3y_2 \sin x - 3y_1 \cos x + y \sin x$ so that $(y_4)_0 = -3$,

and so on.

Now by Maclaurin's theorem, we have

$$y = (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \frac{x^3}{3!}(y_3)_0 + \frac{x^4}{4!}(y_4)_0 + \dots$$

$$\therefore e^{\sin x} = 1 + x \cdot 1 + \frac{x^2}{2!} \cdot 1 + \frac{x^3}{3!} \cdot 0 + \frac{x^4}{4!} \cdot (-3) + \dots$$

$$= 1 + x + \frac{x^2}{2} - \frac{x^4}{8} + \dots$$

(ii) Let $y = e^x \cos x$. Then $(y)_0 = e^0 = 1$,

$$y_1 = e^x \cos x (\cos x - x \sin x) = y(\cos x - x \sin x)$$

so that $(y_1)_0 = 1 \cdot (1 - 0) = 1$,

$$y_2 = y_1 (\cos x - x \sin x) + y(-2 \sin x - x \cos x) \text{ giving}$$

$$(y_2)_0 = 1 \cdot (1 - 0) + 1 \cdot 0 = 1,$$

$$y_3 = y_2 (\cos x - x \sin x) + y_1 (-2 \sin x - x \cos x)$$

$$+ y_1 (-2 \sin x - x \cos x) + y(-3 \cos x + x \sin x)$$

Copyrighted material

$$= y_2 (\cos x - x \sin x) + 2y_1 (-2 \sin x - x \cos x)$$

$$+ y_1 (-2 \sin x - x \cos x) + y(-3 \cos x + x \sin x)$$



$$= y_2(\cos x - x \sin x) + 2y_1(-2 \sin x - x \cos x) \\ + y(-3 \cos x + x \sin x)$$

so that $(y_3)_0 = 1 \cdot (1 - 0) + 0 + 1 \cdot (-3) = -2$,

$$y_4 = y_3(\cos x - x \sin x) + 3y_2(-2 \sin x - x \cos x) \\ + 3y_1(-3 \cos x + x \sin x) + y(4 \sin x + x \cos x)$$

giving $(y_4)_0 = -2 \cdot (1 - 0) + 0 + 3 \cdot 1 \cdot (-3 + 0) + 0 = -11$,

$$y_5 = y_4(\cos x - x \sin x) + 4y_3(-2 \sin x - x \cos x) \\ + 6y_2(-3 \cos x + x \sin x) + 4y_1(4 \sin x + x \cos x) \\ + y(5 \cos x - x \sin x)$$

so that $(y_5)_0 = -11 + 6 \cdot 1 \cdot (-3) + 5 = -24$, and so on.

Substituting these values in Maclaurin's theorem, we get

$$e^{x \cos x} = 1 + x \cdot 1 + \frac{x^2}{2!} \cdot 1 + \frac{x^3}{3!} \cdot (-2) + \frac{x^4}{4!} \cdot (-11) \\ + \frac{x^5}{5!} \cdot (-24) + \dots$$

$$= 1 + x + \frac{x^2}{2} - \frac{x^3}{3} - \frac{11x^4}{24} - \frac{x^5}{5} + \dots$$

(iii) Let $y = e^x \log(1+x)$. Then $(y)_0 = e^0 \log 1 = 1 \times 0 = 0$,

$$y_1 = e^x \log(1+x) + e^x(1+x)^{-1} = y + e^x(1+x)^{-1}$$

so that $(y_1)_0 = 0 + 1 = 1$,

$$y_2 = y_1 + e^x(1+x)^{-1} - e^x(1+x)^{-2} \text{ so that } (y_2)_0 = 1 + 1 - 1 = 1,$$

$$y_3 = y_2 + e^x(1+x)^{-1} - e^x(1+x)^{-2} - e^x(1+x)^{-3}$$

$$+ 2e^x(1+x)^{-3}$$

$$= y_2 + e^x(1+x)^{-1} - 2e^x(1+x)^{-2} + 2e^x(1+x)^{-3}$$

so that $(y_3)_0 = 1 + 1 - 2 + 2 = 2$,

$$y_4 = y_3 + e^x(1+x)^{-1} - 3e^x(1+x)^{-2} + 6e^x(1+x)^{-3}$$

$$- 6e^x(1+x)^{-4} \text{ so that } (y_4)_0 = 2 + 1 - 3 + 6 - 6 = 0,$$

$$y_5 = y_4 + e^x(1+x)^{-1} - 4e^x(1+x)^{-2} + 12e^x(1+x)^{-3}$$

$$- 24e^x(1+x)^{-4} + 24e^x(1+x)^{-5} \text{ so that }$$

$$(y_5)_0 = 0 + 1 - 4 + 12 - 24 + 24 = 9, \text{ and so on.}$$

Substituting these values in Maclaurin's theorem, we get

$$e^x \log(1+x) = 0 + x \cdot 1 + \frac{x^2}{2!} \cdot 1 + \frac{x^3}{3!} \cdot 2 + \frac{x^4}{4!} \cdot 0 + \frac{x^5}{5!} \cdot 9 + \dots \\ = x + \frac{x^2}{2!} + \frac{2x^3}{3!} + \frac{9x^5}{5!} + \dots$$

(iv) Let $y = \log(1+\sin x)$. Then $(y)_0 = 0$,

$$y_1 = \frac{\cos x}{1 + \sin x} \text{ so that } (y_1)_0 = 1,$$

Copyrighted material

$$y_2 = \frac{-\sin x(1 + \sin x) - \cos^2 x}{(1 + \sin x)^2} = \frac{-(1 + \sin x)}{(1 + \sin x)^2} \\ = -\frac{1}{1 + \sin x} \text{ so that } (y_2)_0 = -1,$$



$$\begin{aligned}
 y_2 &= \frac{-\sin x (1 + \sin x) - \cos^2 x}{(1 + \sin x)^2} = \frac{-(1 + \sin x)}{(1 + \sin x)^2} \\
 &= -\frac{1}{1 + \sin x} \text{ so that } (y_2)_0 = -1, \\
 y_3 &= \frac{\cos x}{(1 + \sin x)^2} = \frac{\cos x}{1 + \sin x} \cdot \frac{1}{1 + \sin x} = -y_1 y_2 \\
 &\quad \text{so that } (y_3)_0 = -1 \cdot (-1) = 1, \\
 y_4 &= -y_1 y_3 - y_2^2 \text{ so that } (y_4)_0 = -1 \cdot 1 - (-1)^2 = -1 - 1 = -2, \\
 y_5 &= -y_1 y_4 - y_2 y_3 - 2y_2 y_3 = -y_1 y_4 - 3y_2 y_3 \text{ so that} \\
 (y_5)_0 &= -1 \cdot (-2) - 3 \cdot (-1) \cdot 1 = 2 + 3 = 5, \text{ and so on.} \\
 \text{Substituting these values in Maclaurin's theorem, we get} \\
 \log(1 + \sin x) &= 0 + x \cdot 1 + (x^2/2!) \cdot (-1) + (x^3/3!) \cdot 1 \\
 &\quad + (x^4/4!) \cdot (-2) + (x^5/5!) \cdot 5 + \dots \\
 &= x - \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{12} + \frac{x^5}{24} + \dots
 \end{aligned}$$

Ex. 4. Expand by Maclaurin's theorem $e^x/(1 + e^x)$ as far as the term x^3 .

Sol. Let $y = \frac{e^x}{1 + e^x} = \frac{1 + e^x - 1}{1 + e^x} = 1 - \frac{1}{1 + e^x}$.

Then $(y)_0 = \frac{e^0}{1 + e^0} = \frac{1}{2}$,

$$\begin{aligned}
 y_1 &= 0 + \frac{e^x}{(1 + e^x)^2} = \frac{e^x}{1 + e^x} \cdot \frac{1}{1 + e^x} = y(1 - y) = y - y^2 \text{ so that} \\
 (y_1)_0 &= \frac{1}{2} - \frac{1}{4} = \frac{1}{4},
 \end{aligned}$$

$$y_2 = y_1 - 2yy_1 \text{ so that } (y_2)_0 = \frac{1}{4} - 2 \cdot \frac{1}{2} \cdot \frac{1}{4} = 0,$$

$$y_3 = y_2 - 2y_1^2 - 2yy_2 \text{ so that } (y_3)_0 = 0 - 2 \cdot (1/4)^2 - 0 = -1/8,$$

and so on.

Substituting these values in Maclaurin's theorem, we get

$$\begin{aligned}
 \frac{e^x}{1 + e^x} &= \frac{1}{2} + x \cdot \frac{1}{4} + \frac{x^2}{2!} \cdot 0 + \frac{x^3}{3!} \cdot (-1/8) + \dots \\
 &= \frac{1}{2} + \frac{x}{4} - \frac{1}{48}x^3 + \dots
 \end{aligned}$$

Ex. 5. Apply Maclaurin's theorem to find the expansion in ascending powers of x of $\log_e(1 + e^x)$ to the term containing x^4 .

(Gorakhpur 1981; Kanpur 88; Ranchi 84)

Sol. Let $y = \log_e(1 + e^x)$. Then $(y)_0 = \log_e(1 + e^0) = \log_e 2$,

$$y_1 = \frac{e^x}{1 + e^x} = \frac{(1 + e^x) - 1}{1 + e^x} = 1 - \frac{1}{1 + e^x} \text{ so that } (y_1)_0 = 1 - \frac{1}{2} = \frac{1}{2},$$

Copyrighted material

$$y_2 = 0 + \frac{e^x}{(1 + e^x)^2} = \frac{e^x}{(1 + e^x)} \cdot \frac{1}{1 + e^x} = y_1(1 - y_1) = y_1 - y_1^2$$

so that $(y_2)_0 = \frac{1}{2} - (\frac{1}{2})^2 = \frac{1}{4}$,

$$y_3 = y_2 - 2y_1 y_2 \text{ so that } (y_3)_0 = \frac{1}{4} - 2 \cdot \frac{1}{2} \cdot \frac{1}{4} = 0,$$

$$y_4 = y_3 - 2y_2^2 - 2y_1 y_3 \text{ so that } (y_4)_0 = 0 - 2 \cdot (\frac{1}{4})^2 - 0 = -1/8,$$



$$y_1 = \frac{e^x}{1+e^x} = \frac{(1+e^x)-1}{1+e^x} = 1 - \frac{1}{1+e^x} \text{ so that } (y_1)_0 = 1 - \frac{1}{2} = \frac{1}{2},$$

Copyrighted material

$$y_2 = 0 + \frac{e^x}{(1+e^x)^2} = \frac{e^x}{(1+e^x)} \cdot \frac{1}{1+e^x} = y_1(1-y_1) = y_1 - y_1^2$$

$$\text{so that } (y_2)_0 = \frac{1}{2} - (\frac{1}{2})^2 = \frac{1}{4},$$

$$y_3 = y_2 - 2y_1y_2 \text{ so that } (y_3)_0 = \frac{1}{4} - 2 \cdot \frac{1}{2} \cdot \frac{1}{4} = 0,$$

$$y_4 = y_3 - 2y_2^2 - 2y_1y_3 \text{ so that } (y_4)_0 = 0 - 2 \cdot (\frac{1}{4})^2 - 0 = -1/8,$$

and so on.

Now by Maclaurin's theorem, we have

$$y = (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \frac{x^3}{3!}(y_3)_0 + \frac{x^4}{4!}(y_4)_0 + \dots$$

$$\therefore \log(1+e^x)$$

$$= \log 2 + x \cdot \frac{1}{2} + \frac{x^2}{2!} \cdot \frac{1}{4} + \frac{x^3}{3!} \cdot 0 + \frac{x^4}{4!} \cdot (-1/8) + \dots$$

$$= \log 2 + \frac{x}{2} + \frac{x^2}{8} - \frac{x^4}{192} + \dots$$

Ex. 6. Find the first four terms in the expansion of $\log(1+\tan x)$ in powers of x . (Magadh 1986)

Sol. Let $y = \log(1+\tan x)$. Then $(y)_0 = \log(1+\tan 0) = 0$.

Now $e^y = 1 + \tan x$. Differentiating both sides w.r.t. x , we get

$$e^y y_1 = \sec^2 x. \quad \dots(1)$$

Putting $x = 0$ on both sides of (1), we get

$$e^0 (y_1)_0 = 1 \text{ or } (y_1)_0 = 1.$$

Differentiating (1), we get

$$e^y y_1^2 + e^y y_2 = 2 \sec^2 x \tan x$$

$$\text{or } e^y (y_1^2 + y_2) = 2 \sec^2 x \tan x. \quad \dots(2)$$

Putting $x = 0$ in (2), we get

$$1 + (y_2)_0 = 0 \text{ or } (y_2)_0 = -1.$$

Differentiating (2), we get

$$e^y y_1 (y_1^2 + y_2) + e^y (2y_1 y_2 + y_3) = 2 \sec^4 x + 4 \sec^2 x \tan^2 x$$

$$\text{or } e^y (y_1^3 + 3y_1 y_2 + y_3) = 2 \sec^4 x + 4 \sec^2 x \tan^2 x. \quad \dots(3)$$

Putting $x = 0$ in (3), we get

$$1 + 3 \cdot 1 \cdot (-1) + (y_3)_0 = 2 \text{ or } (y_3)_0 = 4.$$

Differentiating (3), we get

$$e^y y_1 (y_1^3 + 3y_1 y_2 + y_3) + e^y (3y_1^2 y_2 + 3y_2^2 + 3y_1 y_3 + y_4)$$

$$= 8 \sec^4 x \tan x + 8 \sec^2 x \tan^3 x + 8 \sec^4 x \tan x$$

$$\text{or } e^y (y_1^4 + 6y_1^2 y_2 + 4y_1 y_3 + 3y_2^2 + y_4) = 16 \sec^4 x \tan x + 8 \sec^2 x \tan^3 x. \quad \dots(4)$$

Putting $x = 0$ in (4), we get

$$1 + 6 \cdot 1 \cdot (-1) + 4 \cdot 1 \cdot 4 + 3 \cdot (-1)^2 + (y_4)_0 = 0$$

Copyrighted material



$$1 + 0 \cdot 1 \cdot (-1) + 4 \cdot 1 \cdot 4 + 5 \cdot (-1)^2 + (y_4)_0 = 0$$

Copyrighted material

EXPANSIONS OF FUNCTIONS

69

or $(y_4)_0 + 14 = 0$ or $(y_4)_0 = -14.$

Now substituting these values in Maclaurin's theorem, we get

$$\begin{aligned} \log(1 + \tan x) &= 0 + x \cdot 1 + \frac{x^2}{2!} \cdot (-1) + \frac{x^3}{3!} \cdot 4 + \frac{x^4}{4!} \cdot (-14) + \dots \\ &= x - \frac{1}{2}x^2 + \frac{2}{3}x^3 - \frac{7}{12}x^4 + \dots \end{aligned}$$

Ex. 7. Apply Maclaurin's theorem to obtain terms upto x^4 in the expansion of $\log(1 + \sin^2 x).$ (Rohilkhand 1979)

Sol. Let $y = \log(1 + \sin^2 x).$ Then $(y)_0 = 0.$

Now $e^y = 1 + \sin^2 x.$

Differentiating, we get

$$e^y y_1 = 2 \sin x \cos x = \sin 2x. \quad \dots(1)$$

Putting $x = 0$ in (1), we get

$$e^0 (y_1)_0 = 0 \text{ or } (y_1)_0 = 0.$$

Differentiating (1), we get

$$e^y (y_1^2 + y_2) = 2 \cos 2x. \quad \dots(2)$$

Putting $x = 0$ in (2), we get

$$(y_2)_0 = 2.$$

Differentiating (2), we get

$$e^y [(y_1^2 + y_2)y_1 + 2y_1y_2 + y_3] = -4 \sin 2x$$

or $e^y (y_1^3 + 3y_1y_2 + y_3) = -4 \sin 2x. \quad \dots(3)$

Putting $x = 0$ in (3), we get

$$(y_3)_0 = 0.$$

Differentiating (3), we get

$$e^y (y_1^3 + 3y_1y_2 + y_3) + e^y (3y_1^2y_2 + 3y_2^2 + 3y_1y_3 + y_4) = -8 \cos 2x$$

or $e^y (y_1^4 + 6y_1^2y_2 + 4y_1y_3 + 3y_2^2 + y_4) = -8 \cos 2x. \quad \dots(4)$

Putting $x = 0$ in (4), we get

$$3 \cdot 2^2 + (y_4)_0 = -8 \text{ or } (y_4)_0 = -20.$$

Now substituting these values in Maclaurin's theorem, we get

$$\begin{aligned} \log(1 + \sin^2 x) &= 0 + x \cdot 0 + \frac{x^2}{2!} \cdot 2 + \frac{x^3}{3!} \cdot 0 + \frac{x^4}{4!} (-20) + \dots \\ &= x^2 - \frac{5}{6}x^4 + \dots \end{aligned}$$

****Ex. 8.** Show that

(i) $e^x \cos x = 1 + x - \frac{2x^3}{3!} - \frac{2^2x^4}{4!} - \frac{2^2x^5}{5!} + \frac{2^3x^7}{7!} + \dots$

$$+ 2^{n/2} \cos \frac{n\pi}{4} \cdot \frac{x^n}{n!} + \dots$$

(Meerut 1974, 96; Kanpur 85; Rohilkhand 88, 89; Jhansi 88)

Copyrighted material

(ii) $e^x \sin x = x + x^2 - \frac{2}{2!}x^3 - \frac{2^2}{4!}x^5 - \dots + \sin(\frac{1}{4}\pi) \frac{2^{n/2}}{n!} x^n + \dots$



$$(ii) e^x \sin x = x + x^2 - \frac{2}{3!} x^3 - \frac{2^2}{5!} x^5 - \dots + \sin\left(\frac{1}{4}n\pi\right) \frac{2^{n/2}}{n!} x^n + \dots$$

(Meerut 1983, 95, 96 BP; Avadh 87; Lucknow 82;
Gorakhpur 81; Agra 80, 84, 89)

Sol. (i) Let $y = e^x \cos x$. Then $(y)_0 = e^0 \cos 0 = 1$,

$$y_1 = e^x \cos x - e^x \sin x = e^x (\cos x - \sin x)$$

so that $(y_1)_0 = 1(1 - 0) = 1$,

$$y_2 = e^x (\cos x - \sin x) + e^x (-\sin x - \cos x) = -2e^x \sin x$$

so that $(y_2)_0 = 0$,

$$y_3 = -2e^x \sin x - 2e^x \cos x = -2e^x (\sin x + \cos x)$$

so that $(y_3)_0 = -2$,

$$y_4 = -2e^x (\sin x + \cos x) - 2e^x (\cos x - \sin x)$$

$$= -4e^x \cos x = -2^2 y_1 \text{ so that } (y_4)_0 = -2^2,$$

$$y_5 = -2^2 y_1 \text{ so that } (y_5)_0 = -2^2,$$

$$y_6 = -2^2 y_2 \text{ so that } (y_6)_0 = 0,$$

$$y_7 = -2^2 y_3 \text{ so that } (y_7)_0 = 2^3, \text{ and so on.}$$

In general

$$y_n = (1+1)^{n/2} \cos(x + n \tan^{-1} 1) = 2^{n/2} \cos(x + n\pi/4)$$

$$\text{so that } (y_n)_0 = 2^{n/2} \cos\left(\frac{1}{4}n\pi\right).$$

Now by Maclaurin's theorem, we have

$$\begin{aligned} y &= (y)_0 + x(y)_1 + \frac{x^2}{2!}(y)_2 + \dots + \frac{x^n}{n!}(y)_n + \dots \\ &= 1 + x \cdot 1 + \frac{x^2}{2!} \cdot 0 + \frac{x^3}{3!} \cdot (-2) + \frac{x^4}{4!} \cdot (-2^2) + \frac{x^5}{5!} \cdot (-2^2) \\ &\quad + \frac{x^6}{6!} \cdot 0 + \frac{x^7}{7!} \cdot 2^3 + \dots + \frac{x^n}{n!} 2^{n/2} \cos\left(\frac{1}{4}n\pi\right) + \dots \\ &= 1 + x - \frac{2x^3}{3!} - \frac{2^2x^4}{4!} - \frac{2^2x^5}{5!} + \frac{2^3x^7}{7!} + \dots + 2^{n/2} \cos\left(\frac{1}{4}n\pi\right) \frac{x^n}{n!} + \dots \end{aligned}$$

(ii) Proceed as in part (i).

**Ex. 9. Apply Maclaurin's theorem to prove that

$$(i) e^{ax} \sin bx = bx + abx^2 + \frac{3a^2b - b^3}{3!} x^3 + \dots$$

$$+ \frac{(a^2 + b^2)^{n/2}}{n!} x^n \sin\{n \tan^{-1}(b/a)\} + \dots$$

(Meerut 1984)

$$(ii) e^{ax} \cos bx = 1 + ax + \frac{a^2 - b^2}{2} x^2 + \frac{a(a^2 - 3b^2)}{3!} x^3 + \dots$$

$$+ \frac{(a^2 + b^2)^{n/2}}{n!} x^n \cos\{n \tan^{-1}(b/a)\} + \dots$$

Copyrighted material

(Meerut 1998, 84; 83S, 77; Gorakhpur 89; Allahabad 84;
Kurukshtera 82; Magadh 83; Delhi 88; G.N.U. 85)

Hence deduce that

$$e^{x \cos \alpha} \cos(x \sin \alpha) = 1 + x \cos \alpha + \frac{x^2}{2!} \cos 2\alpha + \frac{x^3}{3!} \cos 3\alpha + \dots$$

(Kanpur 1982)



(Meerut 1998, 84; 83S, 77; Gorakhpur 89; Allahabad 84;
Kurukshetra 82; Magadh 83; Delhi 88; G.N.U. 85)

Hence deduce that

$$e^{x \cos \alpha} \cos(x \sin \alpha) = 1 + x \cos \alpha + \frac{x^2}{2!} \cos 2\alpha + \frac{x^3}{3!} \cos 3\alpha + \dots$$

(Kanpur 1982)

Sol. (i) Let $y = e^{ax} \sin bx$. Then $(y)_0 = e^0 \sin 0 = 0$,

$$y_1 = ae^{ax} \sin bx + be^{ax} \cos bx = ay + be^{ax} \cos bx$$

so that $(y_1)_0 = b$,

$$y_2 = ay_1 + abe^{ax} \cos bx - b^2 e^{ax} \sin bx$$

$= ay_1 - b^2 y + abe^{ax} \cos bx$ so that $(y_2)_0 = ab - 0 + ab = 2ab$,

$$y_3 = ay_2 - b^2 y_1 + a^2 be^{ax} \cos bx - ab^2 e^{ax} \sin bx$$

$$= ay_2 - b^2 y_1 - ab^2 y + a^2 be^{ax} \cos bx$$

so that $(y_3)_0 = 2a^2 b - b^3 + a^2 b = 3a^2 b - b^3$, and so on.

In general,

$$y_n = (a^2 + b^2)^{n/2} \sin \{bx + n \tan^{-1}(b/a)\}$$

so that $(y_n)_0 = (a^2 + b^2)^{n/2} \sin \{n \tan^{-1}(b/a)\}$.

Now by Maclaurin's theorem, we have

$$\begin{aligned} y &= (y)_0 + \frac{x}{1!} \cdot (y_1)_0 + \frac{x^2}{2!} \cdot (y_2)_0 + \frac{x^3}{3!} \cdot (y_3)_0 + \dots + \frac{x^n}{n!} \cdot (y_n)_0 + \dots \\ &= 0 + \frac{x}{1!} \cdot b + \frac{x^2}{2!} \cdot (2ab) + \frac{x^3}{3!} \cdot (3a^2b - b^3) + \dots \\ &\quad + \frac{x^n}{n!} \cdot (a^2 + b^2)^{n/2} \sin \{n \tan^{-1}(b/a)\} + \dots \\ &= bx + abx^2 + \frac{3a^2b - b^3}{3!} x^3 + \dots \\ &\quad + \frac{(a^2 + b^2)^{n/2}}{n!} x^n \sin \{n \tan^{-1}(b/a)\} + \dots \end{aligned}$$

(ii) Let $y = e^{ax} \cos bx$. Then $(y)_0 = e^0 \cos 0 = 1$,

$$y_1 = ae^{ax} \cos bx - be^{ax} \sin bx = ay - be^{ax} \sin bx$$

so that $(y_1)_0 = a$,

$$y_2 = ay_1 - abe^{ax} \sin bx - b^2 e^{ax} \cos bx = ay_1 - b^2 y - abe^{ax} \sin bx$$

so that $(y_2)_0 = a^2 - b^2$,

$$y_3 = ay_2 - b^2 y_1 - a^2 be^{ax} \sin bx - ab^2 e^{ax} \cos bx$$

$$= ay_2 - b^2 y_1 - ab^2 y - a^2 be^{ax} \sin bx$$

so that $(y_3)_0 = a(a^2 - b^2) - b^2 a - ab^2 = a(a^2 - 3b^2)$, and so on.

In general, $y_n = (a^2 + b^2)^{n/2} \cos \{bx + n \tan^{-1}(b/a)\}$ so that

$$(y_n)_0 = (a^2 + b^2)^{n/2} \cos \{n \tan^{-1}(b/a)\}.$$

Substituting these values in Maclaurin's theorem, we get

Copyrighted material

$$\begin{aligned} e^{ax} \cos bx &= 1 + ax + \frac{a^2 - b^2}{2!} x^2 + \frac{a(a^2 - 3b^2)}{3!} x^3 + \dots \\ &\quad + \frac{(a^2 + b^2)^{n/2}}{n!} x^n \cos \{n \tan^{-1}(b/a)\} + \dots \end{aligned}$$

Deduction. Putting $a = \cos \alpha$ and $b = \sin \alpha$, we get

$$(y_n)_0 = (\cos^2 \alpha + \sin^2 \alpha)^{n/2} \cos(n \tan^{-1} \tan \alpha) = \cos n\alpha$$
 so that

$$(y_1)_0 = \cos \alpha, (y_2)_0 = \cos 2\alpha, (y_3)_0 = \cos 3\alpha, \text{ etc.}$$



$$e^{ax} \cos bx = 1 + ax + \frac{a^2 - b^2}{2!} x^2 + \frac{a(a^2 - 3b^2)}{3!} x^3 + \dots \\ + \frac{(a^2 + b^2)^{n/2}}{n!} x^n \cos \{n \tan^{-1}(b/a)\} + \dots$$

Deduction. Putting $a = \cos \alpha$ and $b = \sin \alpha$, we get

$$(y_n)_0 = (\cos^2 \alpha + \sin^2 \alpha)^{n/2} \cos(n \tan^{-1} \tan \alpha) = \cos n\alpha \text{ so that}$$

$$(y_1)_0 = \cos \alpha, (y_2)_0 = \cos 2\alpha, (y_3)_0 = \cos 3\alpha, \text{ etc.}$$

$$\therefore e^{x \cos \alpha} \cos(x \sin \alpha) \\ = 1 + x \cos \alpha + (x^2/2!) \cos 2\alpha + (x^3/3!) \cos 3\alpha + \dots$$

Ex. 10. Prove that

$$\sin^{-1} x = x + \frac{1}{2} \cdot \frac{x^3}{3} + \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{x^5}{5} + \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdot \frac{x^7}{7} + \dots$$

(Rohilkhand 1980, 87S)

Sol. Let $y = \sin^{-1} x$. Then $y_1 = 1/\sqrt{1-x^2}$

$$\text{or } (1-x^2)y_1^2 - 1 = 0.$$

Differentiating again, we get

$$(1-x^2)2y_1y_2 - 2xy_1^2 = 0$$

$$\text{or } (1-x^2)y_2 - xy_1 = 0, \text{ since } 2y_1 \neq 0.$$

Now differentiating n times by Leibnitz's theorem, we get

$$(1-x^2)y_{n+2} - (2n+1)xy_{n+1} - n^2y_n = 0.$$

Putting $x = 0$ in the above relations, we get

$$(y)_0 = 0, (y_1)_0 = 1, (y_2)_0 = 0$$

$$\text{and } (y_{n+2})_0 = n^2(y_n)_0. \quad \dots(1)$$

Putting $n = 1, 2, 3, \dots$ in (1), we get

$$(y_3)_0 = 1^2(y_1)_0 = 1^2, (y_4)_0 = 2^2(y_2)_0 = 0, (y_5)_0 = 3^2(y_3)_0 = 3^2 \cdot 1^2,$$

$$(y_6)_0 = 4^2(y_4)_0 = 0, (y_7)_0 = 5^2(y_5)_0 = 5^2 \cdot 3^2 \cdot 1^2, \text{ and so on.}$$

Now by Maclaurin's theorem, we have

$$y = (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \frac{x^3}{3!}(y_3)_0 + \dots \\ \therefore \sin^{-1} x = 0 + x \cdot 1 + \frac{x^2}{2!} \cdot 0 + \frac{x^3}{3!} \cdot 1^2 + \frac{x^4}{4!} \cdot 0 + \frac{x^5}{5!} \cdot 3^2 \cdot 1^2 \\ + \frac{x^6}{6!} \cdot 0 + \frac{x^7}{7!} \cdot 5^2 \cdot 3^2 \cdot 1^2 + \dots \\ = x + \frac{1}{2} \cdot \frac{x^3}{3} + \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{x^5}{5} + \frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdot \frac{x^7}{7} + \dots$$

Ex. 11. Expand $\tan^{-1} x$ by Maclaurin's theorem. Write also the general term. (Delhi 1981; Bundelkhand 82; Kashmir 87)

Sol. Let $y = \tan^{-1} x$. Proceeding as in Ex. 36 on page 50, we get

$$(y)_0 = 0, (y_1)_0 = 1, (y_2)_0 = 0,$$



and $(y_{n+2})_0 = -\{(n+1)n\}(y_n)_0$... (1)

Putting $n = 1, 2, 3, \dots$ in (1), we get

$$(y_3)_0 = -(2 \cdot 1)(y_1)_0 = -2!, (y_4)_0 = -(3 \cdot 2)(y_2)_0 = 0,$$

$$(y_5)_0 = -(4 \cdot 3)(y_3)_0 = -(4 \cdot 3) \cdot (-2!) = 4!, \text{ etc.}$$

Now by Maclaurin's theorem, we have

$$y = (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \frac{x^3}{3!}(y_3)_0 + \dots$$

$$\therefore \tan^{-1}x = 0 + x \cdot 1 + \frac{x^2}{2!} \cdot 0 + \frac{x^3}{3!} \cdot (-2!) + \frac{x^4}{4!} \cdot 0$$

$$+ \frac{x^5}{5!} \cdot (4!) + \dots$$

$$= x - (x^3/3) + (x^5/5) - \dots$$

The general term in this expansion is $(x^n/n!)(y_n)_0$.

So we need the value of $(y_n)_0$. Putting $(n-2)$ in place of n in (1), we get

$$(y_n)_0 = -\{(n-1)(n-2)\}(y_{n-2})_0 \\ = [-\{(n-1)(n-2)\}] [-\{(n-3)(n-4)\}] (y_{n-4})_0.$$

Now there arise two cases :

Case I. When n is even, we have

$$(y_n)_0 = [-\{(n-1)(n-2)\}] [-\{(n-3)(n-4)\}] \dots [-\{(3)(2)\}](y_2)_0 \\ = 0, \text{ since } (y_2)_0 = 0.$$

Case II. When n is odd, we have

$$(y_n)_0 = [-\{(n-1)(n-2)\}] [-\{(n-3)(n-4)\}] \dots \\ [-\{(4)(3)\}] [-\{(2)(1)\}](y_1)_0 \\ = (-1)^{(n-1)/2}(n-1)!, \text{ since } (y_1)_0 = 1.$$

Thus in the expansion of $\tan^{-1}x$, the coefficient of x^n is 0 if n is even and is $\frac{(-1)^{(n-1)/2}(n-1)!}{n!}$ i.e., $\frac{(-1)^{(n-1)/2}}{n}$ if n is odd.

$$\therefore \tan^{-1}x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots + \frac{(-1)^{[(2n-1)-1]/2}}{2n-1} x^{2n-1} + \dots \\ = x - \frac{x^3}{3} + \dots + (-1)^{n-1} \frac{x^{2n-1}}{2n-1} + \dots$$

Ex. 12. Expand $\log\{1 - \log(1-x)\}$ in powers of x by Maclaurin's theorem as far as the term x^3 .

By substituting $x/(1+x)$ for x deduce the expansion of $\log\{1 + \log(1+x)\}$ as far as the term in x^3 .

Sol. Let $y = \log\{1 - \log(1-x)\}$. Then $(y)_0 = 0$.

Now $e^y = 1 - \log(1-x)$. Differentiating, we get

$$e^y y_1 = (1-x)^{-1}. \quad \dots(1)$$

Putting $x = 0$ in (1), we get $(y_1)_0 = 1$.

Differentiating (1), we get

$$e^y y_1^2 + e^y y_2 = (1-x)^{-2}$$



Copyrighted material

Putting $x = 0$ in (1), we get $(y_1)_0 = 1$.

Differentiating (1), we get

$$e^y y_1^2 + e^y y_2 = (1-x)^{-2}$$

or $e^y (y_1^2 + y_2) = (1-x)^{-2} \quad \dots(2)$

Putting $x = 0$ in (2), we get

$$1 + (y_2)_0 = 1 \quad \text{or} \quad (y_2)_0 = 0.$$

Differentiating (2), we get

$$e^y (y_1^3 + 3y_1 y_2 + y_3) = 2(1-x)^{-3} \quad \dots(3)$$

Putting $x = 0$ in (3), we get

$$1 + (y_3)_0 = 2 \quad \text{or} \quad (y_3)_0 = 1.$$

Substituting these values in Maclaurin's theorem, we get

$$\begin{aligned} \log\{1 - \log(1-x)\} &= 0 + x \cdot 1 + \frac{x^2}{2!} \cdot 0 + \frac{x^3}{3!} \cdot 1 + \dots \\ &= x + (x^3/6) + \dots \quad \dots(A) \end{aligned}$$

Now substituting $x/(1+x)$ for x on both sides of (A), we get

$$\log\left\{1 - \log\left(1 - \frac{x}{1+x}\right)\right\} = \frac{x}{1+x} + \frac{1}{6}\left(\frac{x}{1+x}\right)^3 + \dots$$

or $\log\{1 + \log(1+x)\} = x(1+x)^{-1} + (1/6)x^3(1+x)^{-3} + \dots$
 $= x\left\{1 + (-1)x + \frac{(-1)(-2)}{1 \cdot 2}x^2 + \dots\right\} + \frac{1}{6}x^3\{1 + (-3)x + \dots\},$

on expanding by binomial theorem

$$= (x - x^2 + x^3 + \dots) + (\frac{1}{6}x^3 + \dots)$$

$$= x - x^2 + \frac{7}{6}x^3 + \dots$$

Ex. 13. Expand $\{x + \sqrt{1+x^2}\}^m$ in ascending powers of x and find the general term also. (Agra 1978; Kanpur 78)

Sol. Let $y = \{x + \sqrt{1+x^2}\}^m$. Proceeding as in Ex. 39 (a) on page 52, we get

$$(y)_0 = 1, (y_1)_0 = m, (y_2)_0 = m^2,$$

and $(y_{n+2})_0 = (m^2 - n^2)(y_n)_0. \quad \dots(1)$

Now putting $n = 1, 2, 3, 4, \dots$ in (1), we get

$$(y_3)_0 = (m^2 - 1^2)(y_1)_0 = (m^2 - 1^2)m,$$

$$(y_4)_0 = (m^2 - 2^2)(y_2)_0 = (m^2 - 2^2)m^2,$$

$$(y_5)_0 = (m^2 - 3^2)(y_3)_0 = (m^2 - 3^2)(m^2 - 1^2)m,$$

$$(y_6)_0 = (m^2 - 4^2)(y_4)_0 = (m^2 - 4^2)(m^2 - 2^2)m^2, \text{ etc.}$$

In general,

$$\text{if } n \text{ is odd, } (y_n)_0 = \{m^2 - (n-2)^2\} \dots (m^2 - 3^2)(m^2 - 1^2)m$$

and if n is even, $(y_n)_0 = \{m^2 - (n-2)^2\} \dots (m^2 - 4^2)(m^2 - 2^2)m^2 \quad \dots(2)$

Copyrighted material

Now by Maclaurin's theorem, we have

$$y = (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \dots + \frac{x^n}{n!}(y_n)_0 + \dots$$

$$\therefore \{x + \sqrt{1+x^2}\}^m = 1 + mx + \frac{m^2}{2!}x^2 + \frac{m(m^2 - 1^2)}{3!}x^3$$



Copyrighted material

Now by Maclaurin's theorem, we have

$$\begin{aligned}y &= (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \dots + \frac{x^n}{n!}(y_n)_0 + \dots \\ \therefore \quad \{x + \sqrt{1+x^2}\}^m &= 1 + mx + \frac{m^2}{2!}x^2 + \frac{m(m^2-1^2)}{3!}x^3 \\ &\quad + \frac{m^2(m^2-2^2)}{4!}x^4 + \frac{m(m^2-1^2)(m^2-3^2)}{5!}x^5 + \dots\end{aligned}$$

The general term = $\frac{x^n}{n!}(y_n)_0$, where $(y_n)_0$ is given by (2).

Ex. 14. Expand $\log \{x + \sqrt{1+x^2}\}$ in ascending powers of x and find the general term. (Meerut 1991; K.U. 85; Bihar 84)

Sol. Let $y = \log \{x + \sqrt{1+x^2}\}$. Then

$$y_1 = \frac{1}{x + \sqrt{1+x^2}} \cdot \left\{ 1 + \frac{2x}{2\sqrt{1+x^2}} \right\} = \frac{1}{\sqrt{1+x^2}}$$

$$\text{Therefore } y_1^2(x^2+1) - 1 = 0.$$

Differentiating again, we get

$$(x^2+1)2y_1y_2 + 2xy_1^2 = 0$$

$$\text{or } (x^2+1)y_2 + xy_1 = 0, \text{ since } 2y_1 \neq 0.$$

Now differentiating n times by Leibnitz's theorem, we get

$$(x^2+1)y_{n+2} + ny_{n+1} \cdot 2x + \frac{n(n-1)}{2!}y_n \cdot 2 + xy_{n+1} + ny_n = 0$$

$$\text{or } (x^2+1)y_{n+2} + (2n+1)xy_{n+1} + n^2y_n = 0.$$

Putting $x = 0$ in the above relations, we get

$$(y)_0 = 0, (y_1)_0 = 1, (y_2)_0 = 0, \text{ and}$$

$$(y_{n+2})_0 = -n^2(y_n)_0. \quad \dots(1)$$

Now putting $n = 1, 3, 5, \dots$ in (1), we get

$$(y_3)_0 = -1^2(y_1)_0 = -1^2, (y_5)_0 = (-3^2)(y_3)_0 = (-3^2)(-1^2) = 3^2 \cdot 1^2,$$

$$(y_7)_0 = (-5^2)(y_5)_0 = (-5^2)(-3^2)(-1^2) = -5^2 \cdot 3^2 \cdot 1^2, \text{ etc.}$$

In general, if n is odd, we have

$$\begin{aligned}(y_n)_0 &= \{- (n-2)^2\} \{- (n-4)^2\} \dots (-5^2)(-3^2)(-1^2) \\ &= (-1)^{(n-1)/2} (n-2)^2 (n-4)^2 \dots 5^2 \cdot 3^2 \cdot 1^2. \quad \dots(2)\end{aligned}$$

Again, putting $n = 2, 4, 6, \dots$ in (1), we get

$$(y_4)_0 = -2^2(y_2)_0 = 0, (y_6)_0 = -4^2(y_4)_0 = 0, \text{ etc.}$$

Thus, if n is even, we have $(y_n)_0 = 0$.

Now by Maclaurin's theorem, we have

$$\begin{aligned}\log [x + \sqrt{1+x^2}] &= (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \frac{x^3}{3!}(y_3)_0 + \dots \\ &= 0 + x \cdot 1 + \frac{x^2}{2!} \cdot 0 + \frac{x^3}{3!}(-1^2) + \frac{x^4}{4!} \cdot 0 + \frac{x^5}{5!}(3^2 \cdot 1^2) + \dots\end{aligned}$$

Copyrighted material

$$= x - \frac{x^3}{3!} 1^2 + \frac{x^5}{5!} (3^2 \cdot 1^2) - \frac{x^7}{7!} (5^2 \cdot 3^2 \cdot 1^2) + \dots$$

The general term = $(x^n/n!) (y_n)_0$, where $(y_n)_0$ is given by (2) when



$$= x - \frac{x^3}{3!} 1^2 + \frac{x^5}{5!} (3^2 \cdot 1^2) - \frac{x^7}{7!} (5^2 \cdot 3^2 \cdot 1^2) + \dots$$

The general term = $(x^n/n!)(y_n)_0$, where $(y_n)_0$ is given by (2) when n is odd and $(y_n)_0 = 0$, when n is even.

Putting $2n - 1$ in place of n in (2), we find that

$$(y_{2n-1})_0 = (-1)^{n-1} (2n-3)^2 \dots 5^2 \cdot 3^2 \cdot 1^2.$$

Thus $\log [x + \sqrt{(1+x^2)}]$

$$= x - 1^2 \cdot \frac{x^3}{3!} + 1^2 \cdot 3^2 \cdot \frac{x^5}{5!} - 1^2 \cdot 3^2 \cdot 5^2 \cdot \frac{x^7}{7!} + \dots$$

$$+ (-1)^{(n-1)} 1^2 \cdot 3^2 \cdot 5^2 \dots (2n-3)^2 \cdot \frac{x^{2n-1}}{(2n-1)!} + \dots$$

Ex. 15. Expand $e^a \sin^{-1} x$ by Maclaurin's theorem and find the general term. Hence show that

$$e^\theta = 1 + \sin \theta + \frac{1}{2!} \sin^2 \theta + \frac{2}{3!} \sin^3 \theta + \dots$$

(Meerut 1982, 84, 94; Allahabad 87; Rohilkhand 78; Agra 77;

Magadh 77; Ranchi 75; Indore 73;

Lucknow 83, 79; Kanpur 87, 89)

Sol. Let $y = e^a \sin^{-1} x$. Proceeding as in Ex. 37 on page 51, we get

$$y(0) = 1, y_1(0) = a, y_2(0) = a^2,$$

$$\text{and } y_{n+2}(0) = (n^2 + a^2)y_n(0). \quad \dots(1)$$

Putting $n = 1, 2, 3, 4, \dots$ in (1), we get

$$y_3(0) = (1^2 + a^2)y_1(0) = (1^2 + a^2)a, y_4(0) = (2^2 + a^2)y_2(0)$$

$$= (2^2 + a^2)a^2, y_5(0) = (3^2 + a^2)y_3(0) = (3^2 + a^2)(1^2 + a^2)a,$$

$$y_6(0) = (4^2 + a^2)y_4(0) = (4^2 + a^2)(2^2 + a^2)a^2, \text{ etc.}$$

In general,

$$y_n(0) = \begin{cases} a(1^2 + a^2)(3^2 + a^2) \dots [(n-2)^2 + a^2] & \text{if } n \text{ is odd} \\ a^2(2^2 + a^2)(4^2 + a^2) \dots [(n-2)^2 + a^2] & \text{if } n \text{ is even.} \end{cases}$$

Substituting these values in Maclaurin's expansion

$$y = y(0) + xy_1(0) + \frac{x^2}{2!} y_2(0) + \dots + \frac{x^n}{n!} y_n(0) + \dots,$$

we get

$$e^a \sin^{-1} x = 1 + ax + \frac{a^2}{2!} x^2 + \frac{a(1^2 + a^2)}{3!} x^3 + \frac{a^2(2^2 + a^2)}{4!} x^4 + \dots \quad \dots(2)$$

The general term is $(x^n/n!)y_n(0)$, where $y_n(0)$ is as given above.

Now putting $x = \sin \theta$ and $a = 1$ in (2), we get

$$e^\theta = 1 + \sin \theta + \frac{1}{2!} \sin^2 \theta + \frac{2}{3!} \sin^3 \theta + \dots$$

Ex. 16. Use Maclaurin's theorem to show that

$$e^{m \cos^{-1} x} = e^{m\pi/2} \left[1 - mx + \frac{m^2}{2!} x^2 - \frac{m(1^2 + m^2)}{3!} x^3 + \frac{m^2(2^2 + m^2)}{4!} x^4 - \dots \right]$$



Copyrighted material

EXPANSIONS OF FUNCTIONS

77

$$e^{m \cos^{-1} x} = e^{m\pi/2} \left[1 - mx + \frac{m^2}{2!} x^2 - \frac{m(1^2 + m^2)}{3!} x^3 + \frac{m^2(2^2 + m^2)}{4!} x^4 - \dots \right]$$

(Agra 1983; Delhi 82)

Sol. Proceed as in Ex. 15.**Ex. 17.** Expand $\sin(m \sin^{-1} x)$ by Maclaurin's theorem as far as x^5 . Hence expand $\sin m\theta$ in powers of $\sin \theta$.(Meerut 1978, 93; G.N.U. 72; Kanpur 88;
Rohilkhand 83, 85, 90; Lucknow 75)**Sol.** Let $y = \sin(m \sin^{-1} x)$. Proceeding as in Ex. 34 on page 49, we get

$$\begin{aligned} y(0) &= 0, y_1(0) = m, y_2(0) = 0, \\ \text{and } y_{n+2}(0) &= (n^2 - m^2)y_n(0). \end{aligned} \quad \dots(1)$$

Putting $n = 1, 2, 3, \dots$ in (1), we get

$$y_3(0) = (1^2 - m^2)y_1(0) = (1^2 - m^2)m,$$

$$y_4(0) = (2^2 - m^2)y_2(0) = 0,$$

$$y_5(0) = (3^2 - m^2)y_3(0) = (3^2 - m^2)(1^2 - m^2)m, \text{ etc.}$$

Substituting these values in Maclaurin's theorem, we get

$$\begin{aligned} \sin(m \sin^{-1} x) &= y(0) + xy_1(0) + \frac{x^2}{2!}y_2(0) + \dots \\ &= mx + \frac{m(1^2 - m^2)}{3!}x^3 + \frac{m(1^2 - m^2)(3^2 - m^2)}{5!}x^5 + \dots \end{aligned}$$

Putting $x = \sin \theta$ on both sides, we get

$$\begin{aligned} \sin m\theta &= m \sin \theta + \frac{m(1^2 - m^2)}{3!} \sin^3 \theta \\ &\quad + \frac{m(1^2 - m^2)(3^2 - m^2)}{5!} \sin^5 \theta + \dots \end{aligned}$$

Ex. 18. Show that

$$(\sin^{-1} x)^2 = \frac{2}{2!}x^2 + \frac{2 \cdot 2^2}{4!}x^4 + \frac{2 \cdot 2^2 \cdot 4^2}{6!}x^6 + \dots$$

(Meerut 1983, 84, 86)

Deduce that

$$\theta^2 = 2 \cdot \frac{\sin^2 \theta}{2!} + 2^2 \cdot \frac{2 \sin^4 \theta}{4!} + 2^2 \cdot 4^2 \frac{2 \sin^6 \theta}{6!} + \dots$$

(Rajasthan 1987; Meerut 86 P)

Sol. Let $y = (\sin^{-1} x)^2$. Proceeding as in Ex. 32 on page 47, we get

$$\begin{aligned} y(0) &= 0, y_1(0) = 0, y_2(0) = 2, \\ \text{and } y_{n+2}(0) &= n^2 y_n(0). \end{aligned} \quad \dots(1)$$

Putting $n = 1, 2, 3, 4, \dots$ in (1), we get

Copyrighted material

$$y_3(0) = 1^2 y_1(0) = 0, y_4(0) = 2^2 y_2(0) = 2^2 \cdot 2,$$

$$\dots$$



$$\begin{aligned}y_3(0) &= 1^2 y_1(0) = 0, y_4(0) = 2^2 y_2(0) = 2^2 \cdot 2, \\y_5(0) &= 3^2 y_3(0) = 0, y_6(0) = 4^2 y_4(0) = 4^2 \cdot 2^2 \cdot 2, \text{ etc.}\end{aligned}$$

Hence by Maclaurin's theorem, we get

$$\begin{aligned}(\sin^{-1} x)^2 &= y(0) + xy_1(0) + \frac{x^2}{2!} y_2(0) + \dots \\&= \frac{2}{2!} x^2 + \frac{2 \cdot 2^2}{4!} x^4 + \frac{2 \cdot 2^2 \cdot 4^2}{6!} x^6 + \dots\end{aligned}$$

Now putting $\sin^{-1} x = \theta$ or $x = \sin \theta$ on both sides, we get the required expansion for θ^2 .

Ex. 19. (a) By Maclaurin's theorem or otherwise find the expansion of $y = \sin(e^x - 1)$ upto and including the term in x^4 .

(Gorakhpur 1982, 88)

(b) Also show that $x = y - \frac{1}{2}y^2 + \dots$ (Allahabad 1973)

Sol. (a) Let $y = \sin(e^x - 1)$. Then $(y)_0 = \sin 0 = 0$,

$$y_1 = [\cos(e^x - 1)] \cdot e^x \text{ so that } (y_1)_0 = (\cos 0) \cdot e^0 = 1,$$

$$y_2 = [\cos(e^x - 1)] \cdot e^x - [\sin(e^x - 1)] \cdot e^{2x} = y_1 - ye^{2x}$$

$$\text{so that } (y_2)_0 = (y_1)_0 - (y)_0 e^0 = 1 - 0 = 1,$$

$$y_3 = y_2 - y_1 e^{2x} - 2y e^{2x} \text{ so that } (y_3)_0 = 1 - 1 - 0 = 0,$$

$$y_4 = y_3 - y_2 e^{2x} - 4y_1 e^{2x} - 4y e^{2x} \text{ so that } (y_4)_0 = -5, \text{ etc.}$$

Hence by Maclaurin's theorem, we get

$$\begin{aligned}\sin(e^x - 1) &= (y)_0 + x(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \frac{x^3}{3!}(y_3)_0 + \dots \\&= 0 + x \cdot 1 + \frac{x^2}{2!} \cdot 1 + \frac{x^3}{3!} \cdot 0 + \frac{x^4}{4!} \cdot (-5) + \dots \\&= x + \frac{1}{2}x^2 - \frac{5}{24}x^4 + \dots\end{aligned}$$

(b) We have

$$y = \sin(e^x - 1) \Rightarrow e^x - 1 = \sin^{-1} y \Rightarrow e^x = 1 + \sin^{-1} y. \quad \dots(1)$$

Differentiating (1) w.r.t. 'y', we get

$$e^x \cdot x_1 = 1/\sqrt{1-y^2}, \text{ where } x_1 = dx/dy. \quad \dots(2)$$

From (2), we get $(1-y^2)x_1^2 = e^{-2x}$.

Differentiating it w.r.t. 'y', we get

$$(1-y^2)2x_1x_2 - 2yx_1^2 = e^{-2x}(-2x_1)$$

$$\text{or } (1-y^2)x_2 - yx_1 = -e^{-2x}, \text{ since } 2x_1 \neq 0. \quad \dots(3)$$

$$\text{From (1), we have } x = \log(1 + \sin^{-1} y). \quad \dots(4)$$

Now putting $y = 0$ in (4), (2) and (3), we get

$$(x)_0 = \log(1+0) = 0, e^0 \cdot (x_1)_0 = 1/\sqrt{1-0} \text{ giving } (x_1)_0 = 1,$$

$$(x_2)_0 = -e^0 = -1. \quad [\text{Note that } (x)_y=0 = 0]$$

Hence by Maclaurin's theorem, we get

Copyrighted material

$$\begin{aligned}x &= (x)_0 + y(x_1)_0 + (y^2/2!) (x_2)_0 + \dots \\&= 0 + y \cdot 1 + (y^2/2!)(-1) + \dots = y - \frac{1}{2}y^2 + \dots\end{aligned}$$

***Ex. 20. (a)** If $y = (\sin^{-1} x)/\sqrt{1-x^2}$, where $-1 < x < 1$ and $-\pi/2 < \sin^{-1} x < \pi/2$, prove that $(1-x^2)y_{n+1} - (2n+1)xy_n - n^2y_{n-1} = 0$.

Also if $v = a_0 + a_1x + a_2x^2 + \dots + a_nx^n + \dots$



Copyrighted material

EXPANSIONS OF FUNCTIONS

79

$$\begin{aligned}x &= (x)_0 + y(x_1)_0 + (y^2/2!) (x_2)_0 + \dots \\&= 0 + y \cdot 1 + (y^2/2!) (-1) + \dots = y - \frac{1}{2}y^2 + \dots\end{aligned}$$

*Ex. 20. (a) If $y = (\sin^{-1}x)/\sqrt{1-x^2}$, where $-1 < x < 1$
and $-\pi/2 < \sin^{-1}x < \pi/2$,
prove that $(1-x^2)y_{n+1} - (2n+1)xy_n - n^2y_{n-1} = 0$.

Also if $y = a_0 + a_1x + a_2x^2 + \dots + a_nx^n + \dots$,
prove that $(n+1)a_{n+1} = n a_{n-1}$ and hence obtain the general term of
the expansion. (Lucknow 1976)

(b) Expand $(\sin^{-1}x)/\sqrt{1-x^2}$ in powers of x upto three terms.

Sol. (a) Here $y = (\sin^{-1}x)/\sqrt{1-x^2}$ (1)
 $\therefore y^2(1-x^2) = (\sin^{-1}x)^2$.

Differentiating w.r.t. x , we get

$$2yy_1(1-x^2) - 2xy^2 = 2(\sin^{-1}x)/\sqrt{1-x^2} = 2y.$$

Since $2y \neq 0$, therefore $y_1(1-x^2) - xy = 1$

$$\text{i.e. } y_1(1-x^2) - xy - 1 = 0. \quad \dots (2)$$

Differentiating (2) n times by Leibnitz's theorem, we get

$$\begin{aligned}y_{n+1}(1-x^2) + ny_n(-2x) + \frac{n(n-1)}{2!}y_{n-1} \cdot (-2) \\- xy_n - ny_{n-1} = 0\end{aligned}$$

$$\text{or } (1-x^2)y_{n+1} - (2n+1)xy_n - n^2y_{n-1} = 0. \quad \dots (3)$$

Now putting $x = 0$ in (1), (2) and (3), we get

$$(y)_0 = 0, (y_1)_0 = 1 \quad \text{and} \quad (y_{n+1})_0 = n^2(y_{n-1})_0. \quad \dots (4)$$

By Maclaurin's theorem, we have

$$y = (y)_0 + x(y_1)_0 + (x^2/2!)(y_2)_0 + \dots + (x^n/n!)(y_n)_0 + \dots$$

Also we are given that

$$y = a_0 + a_1x + a_2x^2 + \dots + a_nx^n + \dots$$

Comparing the coefficients of x^n in the two expansions for y , we
get

$$a_n = (y_n)_0/n!$$

$$\begin{aligned}\therefore \frac{a_{n+1}}{a_{n-1}} &= \frac{(y_{n+1})_0}{(n+1)!} + \frac{(y_{n-1})_0}{(n-1)!} = \frac{(y_{n+1})_0}{(y_{n-1})_0} \cdot \frac{(n-1)!}{(n+1)!} \\&= n^2 \cdot \frac{1}{n(n+1)}, \quad \text{from (4)} \\&= \frac{n}{n+1}.\end{aligned}$$

$$\therefore (n+1)a_{n+1} = n a_{n-1} \quad \text{Proved.}$$

$$\text{or } a_{n+1} = \frac{n}{n+1} a_{n-1}. \quad \dots (5)$$

Copyrighted material

Now $a_0 = (y)_0 = 0$, $a_1 = (y_1)_0 = 1$. Putting $n = 1, 3, 5, \dots$ in (5),
we get $a_2 = \frac{1}{2}a_0 = 0$, $a_4 = \frac{3}{4}a_2 = 0$, $a_6 = \frac{5}{6}a_4 = 0$, etc.



Now $a_0 = (y)_0 = 0$, $a_1 = (y_1)_0 = 1$. Putting $n = 1, 3, 5, \dots$ in (5), we get $a_2 = \frac{1}{2}a_0 = 0$, $a_4 = \frac{3}{4}a_2 = 0$, $a_6 = \frac{5}{6}a_4 = 0$, etc.

Thus $a_n = 0$ if n is even i.e., $a_{2n} = 0$.

Again putting $n = 2, 4, 6, \dots$ in (5), we get

$$a_3 = \frac{2}{3}a_1 = \frac{2}{3}, a_5 = \frac{4}{5} \cdot a_3 = \frac{4}{5} \cdot \frac{2}{3}, a_7 = \frac{6}{7}a_5 = \frac{6}{7} \cdot \frac{4}{5} \cdot \frac{2}{3}, \text{ etc.}$$

In general, if n is odd, we have

$$a_n = \frac{n-1}{n} \cdot \frac{n-3}{n-2} \cdots \frac{4}{5} \cdot \frac{2}{3}.$$

$$\text{Thus } a_{2n+1} = \frac{2n}{2n+1} \cdot \frac{2n-2}{2n-1} \cdots \frac{4}{5} \cdot \frac{2}{3}.$$

(b) As found in part (a), we have $a_0 = 0$, $a_1 = 1$, $a_2 = 0$, $a_3 = 2/3$, $a_4 = 0$, $a_5 = \frac{2}{3} \cdot \frac{4}{5}$, etc.

$$\therefore \frac{\sin^{-1}x}{\sqrt{1-x^2}} = x + \frac{2}{3}x^3 + \frac{2.4}{3.5}x^5 + \dots$$

Ex. 20. (c) If $y = \sin^{-1}x = a_0 + a_1x + a_2x^2 + \dots$, prove that

$$(n+1)(n+2)a_{n+2} = n^2a_n. \quad (\text{Meerut 1990 P})$$

Sol. Let $y = \sin^{-1}x$ (1)

$$\text{Then } y_1 = \frac{1}{\sqrt{1-x^2}}. \quad \dots (2)$$

$$\therefore (1-x^2)y_1^2 - 1 = 0.$$

Differentiating again, we get

$$(1-x^2)2y_1y_2 - 2xy_1^2 = 0 \text{ or } 2y_1[(1-x^2)y_2 - xy_1] = 0$$

$$\text{or } (1-x^2)y_2 - xy_1 = 0, \quad \dots (3)$$

since $2y_1 \neq 0$.

Now differentiating (3) n times by Leibnitz's theorem, we get

$$(1-x^2)y_{n+2} + n \cdot y_{n+1} \cdot (-2x) + \frac{n(n-1)}{1 \cdot 2}y_n \cdot (-2) \\ - y_{n+1} \cdot x - ny_n \cdot 1 = 0$$

$$\text{or } (1-x^2)y_{n+2} - (2n+1)xy_{n+1} - n^2y_n = 0. \quad \dots (4)$$

Putting $x = 0$ in (4), we get

$$(y_{n+2})_0 = n^2(y_n)_0. \quad \dots (5)$$

By Maclaurin's theorem, we have

$$y = (y)_0 + \frac{x}{1!}(y_1)_0 + \frac{x^2}{2!}(y_2)_0 + \frac{x^3}{3!}(y_3)_0 + \dots + \frac{x^n}{n!}(y_n)_0 + \dots$$

Also we are given that

$$y = \sin^{-1}x = a_0 + a_1x + a_2x^2 + \dots + a_nx^n + \dots$$

Equating the coefficients of x^n in the two expansions for y , we get

Copyrighted material

$$a_n = \frac{(y_n)_0}{n!}.$$

$$\therefore \frac{a_{n+2}}{(y_{n+2})_0} = \frac{1}{n!} = \frac{(y_{n+2})_0}{\dots} \cdot \frac{1}{\dots}$$



$$\begin{aligned} a_n &= \frac{(y_n)_0}{n!} \\ \therefore \frac{a_{n+2}}{a_n} &= \frac{(y_{n+2})_0}{(n+2)!} \cdot \frac{n!}{(y_n)_0} = \frac{(y_{n+2})_0}{(y_n)_0} \cdot \frac{1}{(n+2)(n+1)} \\ &= \frac{n^2}{(n+2)(n+1)}, \text{ substituting for } \frac{(y_{n+2})_0}{(y_n)_0} \text{ from (5).} \end{aligned}$$

Hence $(n+1)(n+2)a_{n+2} = n^2 a_n$.

Ex. 20 (d). If $y = \sin \log(x^2 + 2x + 1)$, prove that

$$(x+1)^2 y_{n+2} + (2n+1)(x+1)y_{n+1} + (n^2+4)y_n = 0.$$

Hence or otherwise expand y in ascending powers of x as far as x^6 .

(Allahabad 1989; Agra 85)

Sol. Here $y = \sin \log(x^2 + 2x + 1) = \sin \log(x+1)^2 \dots (1)$

$$\begin{aligned} \therefore y_1 &= [\cos \log(x+1)^2] \cdot \frac{1}{(x+1)^2} \cdot 2(x+1) \\ &= [\cos \log(x+1)^2] \cdot \frac{2}{x+1}. \end{aligned} \dots (2)$$

Squaring both sides of (2), we get

$$(x+1)^2 y_1^2 = 4 \cos^2 \log(x+1)^2 = 4[1 - \sin^2 \log(x+1)^2]$$

$$= 4(1 - y^2)$$

$$\text{or } (x+1)^2 y_1^2 + 4y^2 - 4 = 0. \dots (3)$$

Differentiating (3), we get

$$(x+1)^2 2y_1 y_2 + 2(x+1)y_1^2 + 8yy_1 = 0$$

$$\text{or } 2y_1 [(x+1)^2 y_2 + (x+1)y_1 + 4y] = 0$$

$$\text{or } (x+1)^2 y_2 + (x+1)y_1 + 4y = 0, \dots (4)$$

since $2y_1 \neq 0$.

Differentiating (4) n times by Leibnitz's theorem, we get

$$\begin{aligned} (x+1)^2 y_{n+2} + {}^n C_1 \cdot y_{n+1} \cdot 2(x+1) + {}^n C_2 \cdot y_n \cdot 2 \\ + (x+1)y_{n+1} + {}^n C_1 \cdot y_n \cdot 1 + 4y_n = 0 \end{aligned}$$

$$\text{or } (x+1)^2 y_{n+2} + (2n+1)(x+1)y_{n+1} + (n^2+4)y_n = 0. \dots (5)$$

Putting $x = 0$ in (1), (2) and (4), we get

$$(y)_0 = 0, (y_1)_0 = 2, (y_2)_0 + (y_1)_0 + 4(y)_0 = 0 \text{ or } (y_2)_0 = -2.$$

Also putting $x = 0$ in (5), we get

$$(y_{n+2})_0 + (2n+1)(y_{n+1})_0 + (n^2+4)(y_n)_0 = 0$$

$$\text{or } (y_{n+2})_0 = -[(2n+1)(y_{n+1})_0 + (n^2+4)(y_n)_0]. \dots (6)$$

Now putting $n = 1, 2, 3, 4$ in (6), we get

$$(y_3)_0 = -[3(y_2)_0 + 5(y_1)_0] = -[3 \cdot (-2) + 5 \cdot 2] = -4,$$

$$(y_4)_0 = -[5(y_3)_0 + 8(y_2)_0] = -[5 \cdot (-4) + 8 \cdot (-2)] = 36,$$

Copyrighted material

Some pages are omitted from this book preview.

