

# Assignment 3

Kavya Kamal

Download all python codes from

<https://github.com/kavyakamal66/IITH-INTERNSHIP/blob/main/Assignment3/code3.py>

and latex-tikz codes from

<https://github.com/kavyakamal66/IITH-INTERNSHIP/blob/main/Assignment3/assignment3.tex>

## 1 QUESTION NO. 2.1 - QUADRATIC FORMS

Find the equation of circle passing through  $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$  making intercepts  $a$  and  $b$  on the co-ordinate axis.

### 2 SOLUTION

The general equation of circle is,

$$\mathbf{x}^T \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (2.0.1)$$

Since the circle passes through  $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$ , the equation of given circle is,

$$\mathbf{x}^T \mathbf{x} + 2\mathbf{u}^T \mathbf{x} = 0 \quad (2.0.2)$$

Given intercepts are  $\begin{pmatrix} a \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 \\ b \end{pmatrix}$

Let x-intercept be  $\begin{pmatrix} 6 \\ 0 \end{pmatrix}$ , y-intercept be  $\begin{pmatrix} 0 \\ 8 \end{pmatrix}$  and  $\mathbf{u}$  be  $\begin{pmatrix} p \\ q \end{pmatrix}$

Substituting  $\begin{pmatrix} 6 \\ 0 \end{pmatrix}$  in 2.0.2

$$\begin{pmatrix} 6 & 0 \end{pmatrix} \begin{pmatrix} 6 \\ 0 \end{pmatrix} + 2\mathbf{u}^T \begin{pmatrix} 6 \\ 0 \end{pmatrix} = 0 \quad (2.0.3)$$

$$\Rightarrow 36 + 2(p \ q) \begin{pmatrix} 6 \\ 0 \end{pmatrix} = 0 \quad (2.0.4)$$

$$\Rightarrow (p \ q) \begin{pmatrix} 6 \\ 0 \end{pmatrix} = -18 \quad (2.0.5)$$

$$\Rightarrow p = -3 \quad (2.0.6)$$

Substituting  $\begin{pmatrix} 0 \\ 8 \end{pmatrix}$  in 2.0.2

$$\begin{pmatrix} 0 & 8 \end{pmatrix} \begin{pmatrix} 0 \\ 8 \end{pmatrix} + 2\mathbf{u}^T \begin{pmatrix} 0 \\ 8 \end{pmatrix} = 0 \quad (2.0.7)$$

$$\Rightarrow 64 + 2(p \ q) \begin{pmatrix} 0 \\ 8 \end{pmatrix} = 0 \quad (2.0.8)$$

$$\Rightarrow (-3 \ q) \begin{pmatrix} 0 \\ 8 \end{pmatrix} = -32 \quad (2.0.9)$$

$$\Rightarrow q = -4 \quad (2.0.10)$$

ie,

$$\mathbf{u} = \begin{pmatrix} -3 & -4 \end{pmatrix} \quad (2.0.11)$$

Centre of the circle,  $\mathbf{O} = -\mathbf{u}$

$$\mathbf{O} = \begin{pmatrix} 3 & 4 \end{pmatrix} \quad (2.0.12)$$

The radius of the circle can be found out using

$$r^2 = \mathbf{u}^T \mathbf{u} \quad (2.0.13)$$

$$\Rightarrow \begin{pmatrix} -3 \\ -4 \end{pmatrix} \begin{pmatrix} -3 & -4 \end{pmatrix} = 25 \quad (2.0.14)$$




Fig. 0: Plot of the required circle

$$\implies r = 5 \quad (2.0.15)$$

Hence , the equation of the given circle is,

$$\mathbf{x}^\top \mathbf{x} + 2 \begin{pmatrix} -3 & -4 \end{pmatrix} \mathbf{x} = 0 \quad (2.0.16)$$

$$\implies \mathbf{x}^\top \mathbf{x} - \begin{pmatrix} 6 & 8 \end{pmatrix} \mathbf{x} = 0 \quad (2.0.17)$$