

How (95)–(97) are obtained

a step-by-step companion to Sec. VI-E

every step below is checked by `analysis/vie_bound_constants.py`

0. What is being bounded, and why the rate

The estimator does not see the tilt angle. It fits the two-segment model to the *rate*

$$\omega_{\text{nom}}(\tau) = C_1 (\cosh C_2 \tau - 1), \quad C_1 = \frac{\dot{M}}{Wz_{CoM}}, \quad C_2 = \sqrt{\frac{Wz_{CoM}}{J_P}}, \quad (\text{D.1})$$

reconstructed from the gyro, and reports the onset as the argument that minimises the residual of that fit. Two questions follow, and (97) answers both:

- (i) how far can the *true* rate depart from (D.1) before the fit stops describing the data? — the quantity $|e_\omega|_{\text{end}}/\omega_{\text{nom, end}}$;
- (ii) how far does that departure move the *reported onset moment*? — the quantity $|\Delta M_{\text{crit}}|$.

These are different questions with different answers, which is why the two halves of (97) carry different constants ($\frac{1}{2}$ and 1 in the first, $\frac{1}{7}$ and $\frac{1}{5}$ in the second). Sections 5–6 do (i) and Section 7 does (ii). If only one line of this document is read, let it be that one.

The departure obeys the Duhamel integral (88),

$$e_\omega(\tau) = \int_0^\tau \dot{h}_\varphi(\tau - s) \rho(s) ds, \quad \dot{h}_\varphi(u) = \frac{\cosh C_2 u}{J_P}, \quad (\text{D.2})$$

with ρ the part of the true forcing that the estimator’s span $\{1, \tau, \delta\varphi\}$ cannot absorb — the gravity remainder ρ_φ and the bilinear ground-effect coupling ρ_{GE} of (89). Everything below is about turning (D.2) into a number.

1. The crude bound, and why it is not enough

Put $|\rho(s)| \leq \rho_{\text{max}}$ inside (D.2) and pull it out:

$$|e_\omega(\tau_{\text{end}})| \leq \rho_{\text{max}} \int_0^{\tau_{\text{end}}} \dot{h}_\varphi(\tau_{\text{end}} - s) ds = \rho_{\text{max}} \int_0^{\tau_{\text{end}}} \frac{\cosh C_2 u}{J_P} du = \frac{\rho_{\text{max}} \sinh x}{J_P C_2}, \quad x \triangleq C_2 \tau_{\text{end}}. \quad (\text{D.3})$$

The substitution $u = \tau_{\text{end}} - s$ was used in the middle step; the integral of $\cosh$ is $\sinh$. Using $J_P = Wz_{CoM}/C_2^2$, i.e. $J_P C_2 = Wz_{CoM}/C_2$, the inertia disappears: $|e_\omega(\tau_{\text{end}})| \leq \rho_{\text{max}} C_2 \sinh x / Wz_{CoM}$. This is (90).

(D.3) is honest but blunt. It treats ρ as though it sat at its maximum for the whole window, whereas both channels start at zero at the onset and grow like τ^6 and τ^4 . Evaluated over the identified constants it returns a deviation of 446% at the slowest ramp — bigger than the signal it is supposed to bound. Sections 2–4 recover the lost factor.

2. Chebyshev's integral inequality

Statement. Let f, g be integrable on $[a, b]$. If they are *oppositely* monotone (one non-decreasing, the other non-increasing), then

$$\int_a^b fg \leq \frac{1}{b-a} \int_a^b f \int_a^b g. \quad (\text{D.4})$$

If they are monotone in the *same* direction, the inequality reverses. (Textbooks usually print the same-direction version, with $\geq$; the one needed here is the reversed one, so the direction is worth checking rather than remembering.)

Proof. Opposite monotonicity means $(f(u) - f(v))(g(u) - g(v)) \leq 0$ for every $u, v \in [a, b]$: when $u > v$ one bracket is ≥ 0 and the other ≤ 0 . Integrate that product over the square $[a, b]^2$ and expand the four terms,

$$\underbrace{\int \int f(u)g(u)}_{(b-a) \int fg} - \underbrace{\int \int f(u)g(v)}_{\int f \int g} - \underbrace{\int \int f(v)g(u)}_{\int f \int g} + \underbrace{\int \int f(v)g(v)}_{(b-a) \int fg} = 2(b-a) \int fg - 2 \int f \int g \leq 0,$$

which is (D.4). $\square$

Application to (D.2). Take $[a, b] = [0, \tau_{\text{end}}]$ with $f(s) = \rho(s)$ and $g(s) = \dot{h}_\varphi(\tau_{\text{end}} - s)$.

- f is non-decreasing: both channels vanish at the onset and grow monotonically over the window ($\rho_\varphi \propto \varphi^2$ with φ rising, $\rho_{GE} \propto \tau \varphi$ likewise).
- g is non-increasing *in s* : as s grows the argument $\tau_{\text{end}} - s$ shrinks and cosh decreases. Note the kernel is increasing in its own argument; it is the reflection $s \mapsto \tau_{\text{end}} - s$ that makes it decreasing in s .

They are oppositely monotone, so (D.4) applies with $\leq$, which is the useful direction:

$$e_\omega(\tau_{\text{end}}) \leq \underbrace{\frac{1}{\tau_{\text{end}}} \int_0^{\tau_{\text{end}}} \rho}_{\bar{\rho}} \cdot \int_0^{\tau_{\text{end}}} \dot{h}_\varphi(\tau_{\text{end}} - s) ds = \frac{\bar{\rho} \sinh x}{J_P C_2}. \quad (\text{D.5})$$

(D.5) is (D.3) with ρ_{max} replaced by the *time average* $\bar{\rho}$. Nothing else changed. All that remains is to compute $\bar{\rho}/\rho_{\text{max}}$ for the two channels.

3. The reduction factors R_φ and R_{GE} : (95)

Both channels have a known shape, because φ does. From (D.1), $\varphi_{\text{nom}}(\tau) = (C_1/C_2)\Lambda(C_2\tau)$ with $\Lambda(u) = \sinh u - u$. Hence, writing $u = C_2\tau$,

$$\rho_\varphi(\tau) = \frac{1}{2}G''\varphi^2 = A\Lambda(u)^2, \quad \rho_{GE}(\tau) = \beta_M \dot{M}\tau \varphi = B u \Lambda(u),$$

with A, B constants whose values never matter — they cancel in the ratio below.

Gravity channel. The maximum is at the window end, $\rho_{\varphi, \text{max}} = A\Lambda(x)^2$. The average is, substituting $u = C_2\tau$ so that $d\tau = du/C_2$ and $\tau_{\text{end}} = x/C_2$,

$$\bar{\rho}_\varphi = \frac{1}{\tau_{\text{end}}} \int_0^{\tau_{\text{end}}} A\Lambda(C_2\tau)^2 d\tau = \frac{C_2}{x} \cdot \frac{A}{C_2} \int_0^x \Lambda(u)^2 du = \frac{A}{x} \int_0^x \Lambda(u)^2 du,$$

so the constant A cancels and

$$R_\varphi(x) \triangleq \frac{\bar{\rho}_\varphi}{\rho_{\varphi,\max}} = \frac{1}{x} \frac{\int_0^x \Lambda(u)^2 du}{\Lambda(x)^2}. \quad (\text{D.6})$$

Ground-effect channel. Identically, $\rho_{GE,\max} = B x \Lambda(x)$ and

$$R_{GE}(x) = \frac{1}{x} \frac{\int_0^x u \Lambda(u) du}{x \Lambda(x)}. \quad (\text{D.7})$$

Where $1/7$ and $1/5$ come from. Expand $\Lambda(u) = u^3/6 + u^5/120 + \dots$ and keep the leading term.

Small x . $\Lambda^2 \simeq u^6/36$, so $\int_0^x \Lambda^2 \simeq x^7/(7 \cdot 36)$ and

$$R_\varphi \simeq \frac{1}{x} \cdot \frac{x^7/(7 \cdot 36)}{x^6/36} = \frac{1}{7}.$$

Similarly $u\Lambda \simeq u^4/6$, so $\int_0^x u\Lambda \simeq x^5/30$ and

$$R_{GE} \simeq \frac{1}{x} \cdot \frac{x^5/30}{x \cdot x^3/6} = \frac{x^4/30}{x^4/6} = \frac{1}{5}.$$

The two numbers are just the orders of the channels: a τ^6 signal averages to $1/7$ of its endpoint value, a τ^4 signal to $1/5$. (For τ^n the average over $[0, T]$ is $T^n/(n+1)$ against the endpoint T^n .)

Large x . $\Lambda \rightarrow \sinh u \simeq e^u/2$, so $\int_0^x \Lambda^2 \simeq e^{2x}/8$ and $R_\varphi \simeq 1/(2x)$; $\int_0^x u\Lambda \simeq x e^x/2$ and $R_{GE} \simeq 1/x$. Both fall off like $1/x$.

Why the small- x value is the supremum. Both R 's decrease throughout, and this needs no numerics. Substituting $u = xt$ in the numerator rescales the window to $[0, 1]$:

$$R(x) = \frac{1}{x} \frac{\int_0^x f(u) du}{f(x)} = \int_0^1 \frac{f(xt)}{f(x)} dt. \quad (\text{D.6a})$$

In this form the small- x limit is immediate: if $f \simeq u^n$ near the origin then $f(xt)/f(x) \rightarrow t^n$ and $R \rightarrow \int_0^1 t^n dt = 1/(n+1)$. Now let

$$n_{\text{eff}}(u) \triangleq \frac{u f'(u)}{f(u)} = \frac{d \log f}{d \log u}$$

be the *local order* of f . Differentiating the integrand of (D.6a) logarithmically,

$$\frac{\partial}{\partial x} \log \frac{f(xt)}{f(x)} = t \frac{f'(xt)}{f(xt)} - \frac{f'(x)}{f(x)} = \frac{n_{\text{eff}}(xt) - n_{\text{eff}}(x)}{x}.$$

For $t \in (0, 1)$ we have $xt < x$, so if n_{eff} is increasing the bracket is ≤ 0 : every integrand of (D.6a) decreases in x , hence so does R . $\square$

Both channels satisfy the hypothesis, and with $\Lambda' = \cosh u - 1$ the local orders are expressible through the same $\Psi_\varphi(u) = u(\cosh u - 1)/(\sinh u - u) = u\Lambda'/\Lambda$ that Sec. 4 discarded as the wrong normaliser:

$$f = \Lambda^2 : \quad n_{\text{eff}} = 2\Psi_\varphi(u) \text{ from } 6 \text{ to } \infty; \quad f = u\Lambda : \quad n_{\text{eff}} = 1 + \Psi_\varphi(u) \text{ from } 4 \text{ to } \infty,$$

Ψ_φ being increasing with $\Psi_\varphi(0^+) = 3$. The mechanism is physical: Λ behaves as u^3 near the onset and grows exponentially later, so the larger the window the more sharply both channels are concentrated at its end, and the smaller their average is relative to the endpoint value. The small- x values are therefore suprema:

$$\bar{\rho} = R_\varphi(x) \rho_{\varphi,\max} + R_{GE}(x) \rho_{GE,\max}, \quad R_\varphi \leq \frac{1}{7}, \quad R_{GE} \leq \frac{1}{5}. \quad (95)$$

The suprema are approached but never attained: for $x > 0$ the inequalities are strict. Over the flown $x \in [1.79, 5.20]$ the factors run 0.089–0.133 and 0.145–0.191, so quoting $\frac{1}{7}$ and $\frac{1}{5}$ gives away 10–60% — while gaining a factor of 5 to 11 over (D.3).

4. Normalising on the rate: (96)

Divide (D.5) by the nominal rate at the same instant, $\omega_{\text{nom, end}} = C_1(\cosh x - 1)$ with $C_1 = \dot{M}/Wz_{CoM}$, and use $J_P C_2 = Wz_{CoM}/C_2$:

$$\frac{|e_\omega|_{\text{end}}}{\omega_{\text{nom, end}}} \leq \frac{\bar{\rho} C_2 \sinh x / Wz_{CoM}}{(\dot{M}/Wz_{CoM})(\cosh x - 1)} = \frac{\bar{\rho} C_2}{\dot{M}} \cdot \frac{\sinh x}{\cosh x - 1}.$$

Wz_{CoM} has cancelled. Now eliminate $\dot{M}$ in favour of the moment increment the window actually spans. Since $\tau_{\text{end}} = x/C_2$,

$$\Delta M_{\text{win}} = \dot{M} \tau_{\text{end}} = \frac{\dot{M} x}{C_2} \quad \Longleftrightarrow \quad \frac{C_2}{\dot{M}} = \frac{x}{\Delta M_{\text{win}}},$$

and substituting,

$$\frac{|e_\omega|_{\text{end}}}{\omega_{\text{nom, end}}} \leq \frac{\Psi(x) \bar{\rho}}{\Delta M_{\text{win}}}, \quad \Psi(x) = \frac{x \sinh x}{\cosh x - 1}. \quad (96)$$

The half-angle identities $\cosh x - 1 = 2 \sinh^2(x/2)$ and $\sinh x = 2 \sinh(x/2) \cosh(x/2)$ collapse the quotient:

$$\Psi(x) = x \frac{2 \sinh(x/2) \cosh(x/2)}{2 \sinh^2(x/2)} = x \coth \frac{x}{2} \geq 2.$$

Ψ rises from 2 at $x \rightarrow 0$ to x as $x \rightarrow \infty$. (Had the excursion been used as the normaliser instead of the rate, the factor would have been $x(\cosh x - 1)/(\sinh x - x)$, rising from 3 — a different function, and the wrong one, since the estimator never looks at φ .)

5. Why $\Psi \bar{\rho}$ stays bounded: first half of (97)

Ψ grows with x and the R 's shrink with it. They are not merely both bounded — they are inverse to each other, and the products converge:

$$\Psi R_\varphi : \quad x \rightarrow 0 : 2 \cdot \frac{1}{7} = \frac{2}{7}, \quad x \rightarrow \infty : x \cdot \frac{1}{2x} = \frac{1}{2};$$

$$\Psi R_{GE} : \quad x \rightarrow 0 : 2 \cdot \frac{1}{5} = \frac{2}{5}, \quad x \rightarrow \infty : x \cdot \frac{1}{x} = 1.$$

Both products increase monotonically between those limits (checked numerically over $x \in (0, 12]$), so the limits at $x \rightarrow \infty$ are suprema. Expanding $\bar{\rho}$ by (95),

$$\Psi \bar{\rho} = (\Psi R_\varphi) \rho_{\varphi, \text{max}} + (\Psi R_{GE}) \rho_{GE, \text{max}} \leq \frac{1}{2} \rho_{\varphi, \text{max}} + \rho_{GE, \text{max}},$$

which is the left half of (97). x has disappeared — and with it C_2 , J_P and Wz_{CoM} . This is the whole point of the exercise: the bound no longer depends on the window length, so it does not have to be solved for, only bounded well enough to keep $\rho_{GE, \text{max}}$ finite (which is what (92)–(93) do).

6. From the rate deviation to the threshold: second half of (97)

The reported quantity is not e_ω but the onset moment, so the deviation must be pushed one step further.

How the onset responds to a perturbation. The estimator picks t_0 to minimise $J(t_0) = \int (y(t) - \hat{\omega}(t; t_0))^2 dt$ with $\hat{\omega}(t; t_0) = C_1(\cosh C_2(t - t_0) - 1)$. Write the data as the nominal

solution plus the deviation, $y = \hat{\omega}(t; t_0^*) + e_\omega$, and the minimiser as $t_0 = t_0^* + \delta$. To first order $\hat{\omega}(t; t_0^* + \delta) \simeq \hat{\omega}(t; t_0^*) + \delta \partial_{t_0} \hat{\omega}$, so the residual is $e_\omega - \delta \partial_{t_0} \hat{\omega}$ and stationarity $\partial J / \partial t_0 = 0$ reads

$$\int (e_\omega - \delta \partial_{t_0} \hat{\omega}) \partial_{t_0} \hat{\omega} dt = 0 \quad \implies \quad \delta = \frac{\langle e_\omega, \partial_{t_0} \hat{\omega} \rangle}{\|\partial_{t_0} \hat{\omega}\|^2}.$$

With $\partial_{t_0} \hat{\omega} = -C_1 C_2 \sinh(C_2 \tau)$ and $\Delta M_{\text{crit}} = \dot{M} \delta$, and using $C_1 = \dot{M} / W z_{CoM}$,

$$\Delta M_{\text{crit}} = - \frac{W z_{CoM} \langle e_\omega, \sinh(C_2 \tau) \rangle}{C_2 \|\sinh(C_2 \tau)\|^2}. \quad (\text{D.8})$$

This is exactly the reduction `error_budget.py` implements.

It is an average of ρ with a non-negative weight. Substitute the Duhamel integral (D.2) into (D.8) and exchange the order of integration (Fubini; both integrands are continuous on a bounded domain):

$$\int_0^T \sinh(C_2 \tau) \int_0^\tau \dot{h}_\varphi(\tau - s) \rho(s) ds d\tau = \int_0^T \rho(s) \underbrace{\left[\int_s^T \sinh(C_2 \tau) \dot{h}_\varphi(\tau - s) d\tau \right]}_{\text{depends on } s \text{ only}} ds,$$

the inner limits changing from $s \in [0, \tau]$ to $\tau \in [s, T]$. Hence

$$\Delta M_{\text{crit}} = - \int_0^T \rho(s) w(s) ds, \quad w(s) = \frac{W z_{CoM} \int_s^T \sinh(C_2 \tau) \cosh(C_2(\tau - s)) d\tau}{J_P C_2 \|\sinh(C_2 \tau)\|^2}. \quad (\text{D.9})$$

Three properties of w finish the argument, and all three are checked in code (`error_budget.py --check`):

1. $w \geq 0$, because the integrand is a product of two hyperbolic functions of non-negative arguments;
2. $\int_0^T w = 1$. The check evaluates (D.9) at $\rho \equiv 1$ and obtains $\Delta M_{\text{crit}} = -1.0005$ to -1.0001 across five operating points. Physically: a uniform unmodelled moment displaces the identified threshold one-for-one, which is the only normalisation the reduction could have;
3. w is non-increasing in s — as s grows both the integration range and the cosh factor shrink (verified numerically over the window).

Property 1 plus 2 alone would give $|\Delta M_{\text{crit}}| \leq \rho_{\text{max}}$. But ρ is increasing and w is decreasing, so *Chebyshev applies a second time*, now to (D.9):

$$\left| \int_0^T \rho w \right| \leq \frac{1}{T} \int_0^T \rho \int_0^T w = \bar{\rho} \cdot 1 = \bar{\rho},$$

and substituting (95),

$$|\Delta M_{\text{crit}}| \leq \bar{\rho} \leq \frac{\rho_{\varphi, \text{max}}}{7} + \frac{\rho_{GE, \text{max}}}{5}, \quad (97b)$$

the right half of (97).

Why the constants differ from the left half. The left half asks how far the rate curve is displaced, so the kernel integral Ψ multiplies $\bar{\rho}$ and the products ΨR climb to $\frac{1}{2}$ and 1. The right half asks how far the *minimiser* moves, and the minimiser is found by correlating against $\sinh(C_2 \tau)$ — a normalised operation, $\int w = 1$, with no kernel gain left over. So there $\bar{\rho}$ enters bare, and the constants are the R 's themselves, $\frac{1}{7}$ and $\frac{1}{5}$. The threshold bound is the stronger of the two by roughly a factor of four, which is why it, and not the relative rate bound, is the number carried into the offset budget.

7. A worked example, checkable by hand

Take `case_02/Mx` at the fastest ramp: $C_2 = 5.428 \text{ s}^{-1}$, $K = 0.1503$ (so $Wz_{CoM} = 1/K = 6.653 \text{ N m}$, $C_1 = K\dot{M} = 0.1804$), $\dot{M} = 1.20 \text{ N m/s}$, $\varphi_{\max} = 10^\circ = 0.17453 \text{ rad}$, $\beta_M = 0.0345 \text{ rad}^{-1}$, $W(l_p + \lambda_{\text{off}}) = 31.59 \times 0.160 \text{ N m}$.

step	expression	value
tilt cap $\rightarrow \Lambda$	$\Lambda_{\max} = \varphi_{\max} Wz_{CoM} C_2 / \dot{M}$	5.253
(92)	$\bar{x} = (6\Lambda_{\max})^{1/3}$	3.159
exact	solve $\sinh x - x = \Lambda_{\max}$	$x = 2.781$
	$\tau_{\text{end}} = x/C_2$	0.512 s
	$\Delta M_{\text{win}} = \dot{M}\tau_{\text{end}}$	0.615 N m
channel	$\rho_{\varphi, \max} = \frac{1}{2}W(l_p + \lambda)\varphi_{\max}^2$	76.98 mN m
(93)	$\rho_{GE, \max} = \beta_M \Delta M_{\text{win}}\varphi_{\max}$	3.70 mN m
(D.6)	$R_\varphi(2.781)$	0.1217
(D.7)	$R_{GE}(2.781)$	0.1800
(95)	$\bar{\rho} = R_\varphi\rho_{\varphi, \max} + R_{GE}\rho_{GE, \max}$ (against $\rho_{\max} = 80.7 \text{ mN m}$)	10.04 mN m
(96)	$\Psi(x) = x \coth(x/2)$	3.148
(96)	$\Psi\bar{\rho}/\Delta M_{\text{win}}$	5.14%
(97)	$(\frac{1}{2}\rho_{\varphi, \max} + \rho_{GE, \max})/\Delta M_{\text{win}}$	6.86%
(97b)	$\rho_{\varphi, \max}/7 + \rho_{GE, \max}/5$	11.74 mN m
	$\div W$	0.372 mm
(D.3)	envelope only, $\Psi\rho_{\max}/\Delta M_{\text{win}}$	41.3%

Two readings. First, (97) costs about a third over (96) — the price of dropping x from the statement — whereas the envelope form costs a factor of eight. Second, this is the *worst admissible* case, evaluated at the 10° tilt cap; over the runs actually flown the excursions are $5\text{--}9^\circ$ and the exact per-run propagation, which convolves the measured ρ instead of any envelope, gives 0.04 mN m in the median and 0.15 at worst — 0.001 to 0.004 mm of offset.

8. Summary of the chain

step	what it buys
(D.2) $e_\omega = \int \dot{h}_\varphi(\tau_{\text{end}} - s)\rho(s)ds$	the exact deviation
(D.3) $ \rho \leq \rho_{\max}$, integrate the kernel	(90); too weak by $\sim 8\times$
(D.4) Chebyshev, $\rho \uparrow$ against $\dot{h}_\varphi(\tau_{\text{end}} - \cdot) \downarrow$	$\rho_{\max} \rightarrow \bar{\rho}$
(D.6–7) known shapes Λ^2 and $u\Lambda$	the ratios $\bar{\rho}/\rho_{\max}$
(D.6a) $R = \int_0^1 f(xt)/f(x)dt$, $n_{\text{eff}} \uparrow$	(95): $R_\varphi \leq \frac{1}{7}$, $R_{GE} \leq \frac{1}{5}$
(96) divide by $\omega_{\text{nom, end}}$, use $\Delta M_{\text{win}} = \dot{M}x/C_2$	$\Psi = x \coth(x/2)$; Wz_{CoM} , J_P gone
	$\Psi R_\varphi \uparrow \frac{1}{2}$, $\Psi R_{GE} \uparrow 1$
(D.8–9) linearise the onset, Fubini, $w \geq 0$, $\int w = 1$	$\Delta M_{\text{crit}} = -\langle \rho \rangle_w$
	Chebyshev again, $\rho \uparrow$ against $w \downarrow$
	(97) right half: $\frac{1}{7}, \frac{1}{5}$

Chebyshev is used twice, for the same reason both times: the forcing is concentrated at the end of the window and every kernel in the problem weights the end of the window least. The two uses are independent — the first converts a kernel integral, the second a normalised correlation — which is why the constants they leave behind differ.