

Question 1 | Correct Mark 1.00 out of 1.00

Given an array `nums` of size `n`, return *the majority element*.

The majority element is the element that appears more than $\lfloor n / 2 \rfloor$ times. You may assume that the majority element always exists in the array.

Example 1:

Input: `nums = [3,2,3]`

Output: 3

Example 2:

Input: `nums = [2,2,1,1,1,2,2]`

Output: 2

Constraints:

- `n == nums.length`
- `1 <= n <= 5 * 104`
- `-231 <= nums[i] <= 231 - 1`

For example:

Input	Result
3	3
3 2 3	

Input	Result
7 2 2 1 1 1 2 2	2

Answer: (penalty regime: 0 %)

```
1  #include<stdio.h>
2  int main(){
3      int i,n,count=0,ans;
4      scanf("%d",&n);
5      int a[n];
6      for(i=0;i<n;i++)
7          scanf("%d",&a[i]);
8      ans=a[0];
9      for(i=0;i<n;i++){
10         if(count==0)
11         {
12             ans=a[i];
13             if(a[i]==ans)
14                 count++;
15             else count--;
16         }
17     }
18     printf("%d", ans);
19
20 }
21
```

	Input	Expected	Got	
✓	3	3	3	✓
	3 2 3			

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[Back to Course](#)