Supplementary information for: Thermodynamic modeling tools for the interpretation of melt inclusions and volcanic gases

Kayla Iacovino *[α], Ery Hughes[β,γ], Penny Wieser[δ], Alain Burgisser[ε], Philippa Liggins[ζ], Shuo Ding[η], Chenguang Sun[θ], and Geoff Kilgour[γ]

[α] Carl Sagan Center for Research, SETI Institute, Mountain View, CA, 94043, US.
[β] Department of Earth Sciences, University College London, London, WC1E 6BT, UK.
[γ] Earth Sciences New Zealand – Te Whatu Ao, Wairakei Research Centre, Taupō, New Zealand
[δ] Department of Earth and Planetary Science, University of California Berkeley, Berkeley, CA, US.
[ε] Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IRD, Univ. Gustave Eiffel, ISTerre, Grenoble, 38000, France.
[ζ] Oxford Research Software Engineering Group, MPLS Doctoral Training Centre, University of Oxford, Oxford OX1 2JD, UK.
[η] Department of Geological Sciences, University of Florida, Gainesville, FL, 32611, US.
[θ] Department of Earth and Planetary Sciences, Jackson School of Geosciences, University of Texas, Austin, TX, 78712, US.

1 List of Supplementary Materials

All of these materials are available through the journal's repository and in the GitHub repo: `https://github.com/kaylai/volcanic-gas-modeling-tool-comparison`.

1. This PDF
2. All modeling results from all tools generated in this work: in folder "`Model_Outputs`"
3. All modeling tool settings used to generate results: in folder "`Model_Outputs/resolved_inputs`"
4. Notebooks for creating every figure in this manuscript: in folder "`Figure_Notebooks`"
5. Full datasets used to choose starting compositions for our four basaltic systems: "`Melt_Inclusions/melt_inclusion_datasets.xlsx`"
6. Jupyter notebook for describing how starting compositions were chosen from the full datasets: "`Melt_Inclusions/Choosing_MI_Kilauea.ipynb`"

2 Description of Basaltic Systems

Mid-ocean ridge basalts are represented by whole rock lava samples and representative MORB volatile concentrations. Major element data are from Lamont Seamount Chain lavas and adjacent East Pacific Rise lavas originally reported in Allan et al. [1989]. This is the same composition used in the parametrization database for MagmaSat MELTS [Ghiorso and Gualda 2015] and in the review by Wieser et al. [2022]. Temperature and volatile contents were chosen from representative MORB samples reported in Cottrell et al. [2021]. $Fe^{3+}/\Sigma Fe$ was set at 0.1, reflective of the majority of mid-ocean ridge basalts recording fO_2 around FMQ [Cottrell et al. 2021]. MORB lavas represent a volatile-poor end-member and position MORB as the most relevant to magma ocean and extraterrestrial model scenarios compared to other basalts examined here.

Major and volatile element compositions for all other samples are from glassy MI and include CO_2 abundances in vapor bubbles within MI. CO_2 we report here and use in modeling is the sum of the dissolved CO_2 in the melt and the exsolved CO_2 in the vapor bubble. Because they record the state of the melt at various points during crystal growth, MI are commonly paired with volatile solubility models to assess magma evolution and degassing patterns, including estimations of pre-eruptive magma conditions. As such, MI provide evidence-based starting conditions for magmatic degassing scenarios.

Volatile-poor OIBs are represented by Kīlauea volcano, Hawai'i. The volatile and redox systematics of Kīlauea magmas have been extensively studied using both olivine-hosted MI [Sides et al. 2014; Moore et al. 2015; Moussallam et al. 2016; Tucker et al. 2019; Lerner et al. 2021a; b; Wieser et al. 2021] and submarine glasses [Dixon et al. 1991; Clague et al. 1995], making them an ideal target for model inter-comparison. It is noteworthy that Kīlauean melts show variable H_2O contents, with some eruptions mostly yielding 0.3 wt%, and others 0.5–1.0 wt%. To address these variable H_2O contents, we performed two different Kīlauea models, Kīlauea 'dry' with 0.3 wt%, and 'wet' with 0.6 wt%. No significant differences were observed between these two runs, and so only the 'dry' runs are presented here. The major elements were chosen to be representative of higher MgO (10.2 wt%) MI corrected for post-entrapment crystallization [Wieser et al. 2020; Lerner et al. 2021a; Wieser et al. 2021]. The chosen initial CO_2 concentration of 800 ppm represents the most CO_2-rich MI from the 2018 eruption. For S, we use a concentration of 1500 ppm after Wieser et al. [2020] and Sides et al. [2014]. $Fe^{3+}/\Sigma Fe$ was set at 0.18 following Lerner et al. [2021b] and T at 1200 °C using the Helz and Thornber [1987] thermometer.

Oxidized H_2O-rich arc basalts are represented by Fuego volcano, Guatemala. Fuego MI from the 1974 eruption have up to 4.5 wt% H_2O [Rasmussen et al. 2020], relatively elevated S contents (2650 ppm) with up to 0.7 $S^{6+}/\Sigma S$ [Taracsák et al. 2023], and experimentally reconstructed CO_2 concentrations up to 3500 ppm [Rasmussen et al. 2020]. The major element composition

Draft for review

representing Fuego is based on a compilation of data from Rasmussen et al. [2020], Lloyd et al. [2013], Moore et al. [2015], and Newcombe et al. [2020], whilst volatiles are from Rasmussen et al. [2020]. $Fe^{3+}/\Sigma Fe$ was set at 0.24 and T at 1030 °C following Lloyd et al. [2013] and Ding et al. [2023].

Volatile-rich OIBs are represented by Fogo Volcano, Cabo Verde, an end-member volcano representing one of the world's most volatile- and alkali-rich silica-undersaturated ocean island volcanoes. Fogo magmas have extremely high CO_2 and S contents (1.15 wt% and 4690 ppm, respectively) relative to other OIBs [DeVitre et al. 2023]. The major element composition was chosen as the average composition of 5 PEC-corrected melt inclusions from sample FG18 with the highest H_2O and CO_2 contents [DeVitre et al. 2023]. We take the redox conditions to be ΔNNO+0.7, as calculated by DeVitre et al. [2023] using the V partitioning oximeter of Canil [2002]. All values—majors, volatiles, redox, and T—are taken from DeVitre et al. [2023].

3 Abbreviations

All models are given by name and full citation in the first appearance in the text and thereafter referred to by a model name or abbreviation only: "Allison" [**Allison2019**], "Eguchi" [Eguchi and Dasgupta 2018], "Frost" [Frost 1991], "Gaillard" [Gaillard et al. 2003], "Iacono-Marziano" or "IM" [Iacono-Marziano et al. 2012], "Kress and Carmichael" or "KC" [Kress and Carmichael 1991], "Liu" [Liu et al. 2005], "Moore" [Moore et al. 1998], "Muth" [Muth and Wallace 2021], "Nash" [Nash et al. 2019], "Shishkina" [Shishkina et al. 2014], and "VolatileCalc" or "VC" [referring to both the original model equations and the Excel VisualBasic app: Dixon et al. 1995; Newman and Lowenstern 2002].

4 Model Options

All model options chosen for the calculations in this manuscript are listed in Table 1. Equilibrium constants are for the following reactions:

K_{H_2O}: $H_2O = H_2 + 0.5O_2$
K_{CO_2}: $CO + 0.5O_2 = CO_2$
K_{H_2S-1}: $0.5S_2 + H_2O = H_2S + 0.5O_2$
K_{H_2S-2}: $H_2S + 1.5O_2 = SO_2 + H_2O$
K_{SO_2}: $0.5S_2 + O_2 = SO_2$
K_{CH_4}: $CO_2 + 2H_2O = CH_4 + 2O_2$
K_{OCS}: $3CO + SO_2 = 2CO_2 + OCS$

Modeling tool settings used to generate our results are included in the Supplement in the folder "`Model_Outputs/resolved_inputs`". Note that for Sulfur_X, the user can define a value for the B constant term to tweak the relationship between Fe and S redox (e.g., to be consistent with XANES data for the system as performed here for Kīlauea). Because of a paucity of experimental data for C-O-H-S-bearing systems at $P < \sim$250 bars, instability in the computed combined K_D can trend to infinity in special or unusual cases [Ding et al. 2023]. To mitigate this, Sulfur_X allows the user to specify an arbitrary increase in the combined K_D value at each pressure for the lowest pressure steps. In our model runs, we iteratively adjusted this behavior as well as other computational settings that control solver convergence until obtaining a Sulfur_X run with no or minimal low-pressure instability.

draft for review

Table 1: Model parameterizations used by each tool. See notes and reference key below the table.

Parameter	VolFe	Sulfur_X	D-Compress	MAGEC	EVo
SOLUBILITY					
H_2O	H24 FigS2*	IM12 Eq11[1]*	IM12 Eq11[1]*	IM12 Eq11[2]*	B15 Table1-basalt*
CO_2	[ξ,ζ,θ]D95 [Ψ]TH94*	IM12 Eq7[1]*	IM12 Eq7[1]*	IM12 Eq7[1]*	B15 Table1-basalt*
S^{2-}	O21 Eq10.34[3]*	D23 Table3-RxnI	N/A	N99 Eq3*	O21 Eq10.34*
S^{6+}	OM22 Eq12a[3]*	D23 Table3-RxnII	N/A	N/A	N/A
SO_2	N/A	N/A	B15 Table1-basalt*	N/A	N/A
H_2S	H24 FigS6-basalt*	N/A	B15 Table1-basalt*	N/A	N/A
H_2	H24 TableS4-basalt*	N/A	B15 Table1-basalt*	G03 Sec5.4	G03 Sec5.4*
CH_4	A13 Eq7a	N/A	0*	A13 Eq7a	A13 Eq7a
CO	H24 TableS4	N/A	0*	A15 Eq10	A15 Eq10
O_2	N/A	N/A	0*	N/A	N/A
S_2	N/A	N/A	0*	N/A	N/A
MELT SPECIATION					
$Fe^{3+}/\Sigma Fe$	KC91 EqA5, A6	KC91 Eq7	KC91 Eq7[4]	SY24 Eq6-9, Table2*	KC91 Eq7
$S^{6+}/\Sigma S$	N/A*	[ξ,Ψ]MW21 [ζ]MW21[5] [θ]OM22 Eq13*	N/A	SY24 Eq10*	N19 Eq11*
FUGACITY COEFFICIENT					
H_2O	HP91 Eq4, 6, A1-3, Table1*	SS92*	HP91 EqA2-3	PR76 Eq15	HP91
CO_2	SS92*	N/A	SS92	SS92	HP91
SO_2	SS92[6]*	SS92	SS92	SS92	SS92
H_2S	SS92[7]*	SS92	SS92	SS92	SS92
O_2	SS92*	N/A	SS92	SS92	SS92
H_2	SW64 Eq4*	N/A	SW64 Eq4	SS92	SW64 Eq4
CO	SS92*	N/A	SS92	SS92	SS92
CH_4	SS92*	N/A	SS92	SS92	SS92
S_2	SS92*	N/A	SS92	SS92	SS92
OCS	SS92*	N/A	N/A	SS92	N/A

continued on next page

draft for review

Tablele 1, continued.

Parameter	VolFe	Sulfur_X	D-Compress	MAGEC	EVo
EQUILIBRIUM CONSTANTS					
K_{H_2O}	OK77 Table1-d	N/A	OK77 Table1-d	SL22[8]	EVo RTD[9]
K_{CO_2}	OK77 Table1-c	N/A	OK77 Table1-c	SL22[8]	EVo RTD[9]
$K_{H_2S\text{-}1}$	OK77 Table1-h*	N/A	OK77 Table1-h	SL22[8]	EVo RTD[9]
$K_{H_2S\text{-}2}$	N/A	D23 Eq4	N/A	N/A	N/A
K_{SO_2}	OK77 Table1-f*	N/A	OK77 Table1-f	SL22[8]	EVo RTD[9]
K_{CH_4}	OK77 Table1-e*	N/A	OK77 Table1-e	SL22[8]	EVo RTD[9]
K_{OCS}	M19 Eq8	N/A	N/A	SY24[8]	N/A
OXYGEN FUGACITY					
NNO	F91 Table1	N/A	F91 Table1	H08 Table7[10]	F91 Table1
FMQ	F91 Table1-FMβQ*	F91 Table1-FMβQ	F91 Table1-FMβQ[11]	H08 Table7[12]	F91 Table1-FMβQ
Redox input options	$Fe^{3+}/\Sigma Fe$ $FeO + Fe_2O_3$ ΔFMQ ΔNNO $\log_{10} fO_2$ $S^{6+}/\Sigma S$	ΔFMQ	ΔNNO	$\log_{10} fO_2$ $Fe^{3+}/\Sigma Fe$ $S^{6+}/\Sigma S$ ΔIW ΔFMQ ΔNNO	$Fe^{3+}/\Sigma Fe$ $FeO + Fe_2O_3$ ΔFMQ ΔNNO fO_2 (bar)

Notes: Parameterisations used for all calculations, except where specified for individual melt compositions: ξMORB, ζKīlauea, θFuego, and ΨFogo. * Indicates other parameterizations are available within the tool. [1]Hydrous parameters; [2]Hydrous web-app parameters; [3]Including H_2O dilution; [4]At 1 bar; [5]Using 6.5 as the last constant value; [6]As modified in Fig. S1 of [Hughes et al. 2023]; [7]As modified in Fig. S1 of [Hughes et al. 2024b]; [8]Based on NIST-JANAF tables [Chase 1996]; [9]Based on JANAF tables [Chase et al. 1998]; [10]References of Frost [1991] and O'Neill and Pownceby [1993]; [11]For this study only because not included in the original tool; [12]References of O'Neill [1987].

References: A13 [Ardia et al. 2013]; A15 [Armstrong et al. 2015]; B15 [Burgisser et al. 2015]; D95 [Dixon et al. 1995]; D23 [Ding et al. 2023]; EVo RTD (`https://evo-outgas.readthedocs.io`); F91 [Frost 1991]; G03 [Gaillard et al. 2003]; H08 [Hirschmann et al. 2008]; HP91 [Holland and Powell 1991]; H24 [Hughes et al. 2024b]; IM12 [Iacono-Marziano et al. 2012]; KC91 [Kress and Carmichael 1991]; M19 [Moussallam et al. 2019]; MW21 [Muth and Wallace 2021]; N99 [Nzotta et al. 1999]; N19 [Nash et al. 2019]; OK77 [Ohmoto and Kerrick 1977]; O21 [O'Neill 2021a]; OM22 [O'Neill and Mavrogenes 2022]; PR76 [Peng and Robinson 1976]; SW64 [Shaw and Wones 1964]; SS92 [Shi and Saxena 1992]; SL22 [Sun and Lee 2022]; SY24 [Sun and Yao 2024]; TH94 [Thibault and Holloway 1994].

draft for review

5 How to cite models

When using model results in a publication, the specific model used should be: 1. well justified (see main text section ??; and 2. properly cited. The version of the code should be included; model options used within the tool should be clearly recorded and cited appropriately (including if implemented via another tool); and a notebook or script to recreate the calculations and all model results should be included as supplementary material. Here we provide an example suitable for a publication:

> Closed- and open-system degassing curves were modeled using VolFe v1.0 [Hughes et al. 2025] using solubility functions from Dixon et al. [1995], Ardia et al. [2013], O'Neill [2021b], O'Neill and Mavrogenes [2022], and Hughes et al. [2024a], fugacity coefficients from Shaw and Wones [1964], Holland and Powell [1991], Shi and Saxena [1992], and Hughes et al. [2023, 2024b], homogeneous vapor equilibria constants from Ohmoto and Kerrick [1977] and Moussallam et al. [2019], Fe^{3+}/Fe_T from Kress and Carmichael [1991], FMQ and NNO buffers from Frost [1991], and sulfide and anhydrite saturation from Chowdhury and Dasgupta [2019], O'Neill [2021b], and Wieser and Gleeson [2023]. Specific equations used are detailed in Table 5. Parameterizations for model dependent variables were chosen where the calibration dataset most closely matched the composition of the MI of interest. Test calculations show that this selection of model dependent variables in VolFe is able to reproduce saturation pressures from the experiments of "X" to within 20%, which are close in composition to our MI. A full list of model configuration options and notebooks for reproducing test calculations and modeling outputs presented here are in the Supplementary Material.

6 Vapor

The vapor composition during degassing is quite different between different tools (Figure 1). Here we expand on potential differences in the tools that may be contributing to these changes.

6.1 Homogeneous vapor equilibrium constants

The equilibrium constants for homogeneous equilibria in the vapor (K, which are independent of P and melt composition but depend on T) control the vapor composition at a given set of conditions and therefore differences in K can cause variations between model outputs. These K can be calculated from model results using the vapor composition (x_i^v and f_{O_2}) at 1 bar when $\gamma_i = 1$ (i.e., $f_i = x_i^v P$ at 1 bar) according to:

$$H_2(v) + 0.5O_2(v) = H_2O(v) \tag{1}$$

$$K_H = \frac{f_{H_2O}}{f_{H_2} f_{O_2}^{0.5}} = \frac{\gamma_{H_2O} x^v_{H_2O} P}{\gamma_{H_2} x^v_{H_2} P f_{O_2}^{0.5}} \tag{2}$$

$$CO(v) + 0.5O_2(v) = CO_2(v) \tag{3}$$

$$K_C = \frac{f_{CO_2}}{f_{CO} f_{O_2}^{0.5}} = \frac{\gamma_{CO_2} x^v_{CO_2} P}{\gamma_{CO} x^v_{CO} P f_{O_2}^{0.5}} \tag{4}$$

$$0.5S_2(v) + O_2(v) = SO_2(v) \tag{5}$$

$$K_S = \frac{f_{SO_2}}{f_{S_2}^{0.5} f_{O_2}} = \frac{\gamma_{SO_2} x^v_{SO_2} P}{\left(\gamma_{S_2} x^v_{S_2} P\right)^{0.5} f_{O_2}} \tag{6}$$

$$CH_4(v) + 0.5O_2(v) = CO(v) + 2H_2(v) \tag{7}$$

$$K_{CH} = \frac{f_{CO} f_{H_2}^2}{f_{CH_4} f_{O_2}^{0.5}} = \frac{\gamma_{CO} x^v_{CO} P \left(\gamma_{H_2} x^v_{H_2} P\right)^2}{\gamma_{CH_4} x^v_{CH_4} P f_{O_2}^{0.5}} \tag{8}$$

$$SO_2(v) + H_2(v) = H_2S(v) + O_2(v) \tag{9}$$

$$K_{HS} = \frac{f_{H_2S} f_{O_2}}{f_{SO_2} f_{H_2}} = \frac{\gamma_{H_2S} x^v_{H_2S} P f_{O_2}}{\gamma_{SO_2} x^v_{SO_2} P \gamma_{H_2O} x^v_{H_2} P} \tag{10}$$

draft for review

Table 2: Parameterizations used for model dependent variables in degassing calculations using VolFe.

Variable	Reference
Fugacity coefficients	
O_2	Shi and Saxena [1992]
CO	Shi and Saxena [1992]
H_2	Shaw and Wones [1964]
S_2	Shi and Saxena [1992]
CO_2	Shi and Saxena [1992]
H_2O	Eq. (4, 6: T > 673 K, A1-3) & Table 1 in Holland and Powell [1991]
SO_2	Shi and Saxena [1992] as modified in Fig. S1 of Hughes et al. [2023]
CH_4	Shi and Saxena [1992]
H_2S	Shi and Saxena [1992] as modified in Fig. S1 of Hughes et al. [2024b]
OCS	Shi and Saxena [1992]
Homogeneous vapor equilibrium constants	
CO_2	Reaction (c) in Table 1 of Ohmoto and Kerrick [1977]
H_2O	Reaction (d) in Table 1 of Ohmoto and Kerrick [1977]
SO_2	Reaction (f) in Table 1 of Ohmoto and Kerrick [1977]
CH_4	Reaction (e) in Table 1 of Ohmoto and Kerrick [1977]
H_2S	Reaction (h) in Table 1 of Ohmoto and Kerrick [1977]
OCS	Eq. (8) in Moussallam et al. [2019]
Solubility functions	
H_2O_T	Fig. S2 in Hughes et al. [2024b]
$CO_{2,T}$	Bullet (5) in Dixon et al. [1995]
$H_{2,mol}$	Basalt from Table S4 in Hughes et al. [2024b]
CO_{mol}	Table S4 in Hughes et al. [2024b]
$CH_{4,mol}$	Eq. (7a) from Ardia et al. [2013]
$^*S^{2-}$	Eq. (10.43) from O'Neill [2021a] (including H_2O dilution)
SO_4^{2-}	Eq. (12a) from O'Neill and Mavrogenes [2022] (including H_2O dilution)
H_2S_{mol}	Basalt from Fig. S6 in Hughes et al. [2024b]
fO_2 related parameters	
Fe^{3+}/Fe_T	Eq. (A-5, 6) from Kress and Carmichael [1991] (no H_2O dilution)
FMQ	FMβQ in Table 1 of Frost [1991]
NNO	NiNiO in Table 1 of Frost [1991]
Other	
S^{2-}CSS	Eq. (10.34, 10.43, 10.45, 10.46) in O'Neill [2021a]
S^{6+}CAS	Eq. (8) in Chowdhury and Dasgupta [2019] using PySulfSat v1.0.2 [Wieser and Gleeson 2023]

under review




Figure 1: Vapor composition against normalized P for each system using the different tools.




Figure 2: Equilibrium constants for each tool and system (i.e., different temperatures). Calculated at P = 1 bar as described in the text.

$$OCS(v) = CO(v) + 0.5S_2(v) \tag{11}$$

$$K_{SC} = \frac{f_{CO} f_{S_2}^{0.5}}{f_{OCS}} = \frac{\gamma_{CO} x_{CO}^{v} P \left(\gamma_{S_2} x_{S_2}^{v} P\right)^{0.5}}{\gamma_{OCS} x_{OCS}^{v} P} \tag{12}$$

Although these specific reactions and K may not have been used in the tools directly, these are six independent reactions that describe a vapor containing O_2, H_2, CO, S_2, H_2O, CO_2, SO_2, CH_4, H_2S, and OCS (i.e., no reaction can be made from a combination of the other reactions). Hence, this represents a minimum number of K in the system. Equilibrium constants are only calculated if the tool outputs all required x_i^v for a specific reaction (e.g., Sulfur_X, D-Compress, and EVo do not include x_{OCS}^v and therefore K_{OCS} cannot be calculated) and if the system calculation reached $P \leq 1$ bar (e.g., VolFe calculations for Fuego and MORB did not reach $P = 1$ bar and therefore values are not calculated; Figure 2). D-Compress and VolFe use Ohmoto and Kerrick [1977], whereas Evo and MAGEC use in-house versions based on NIST/JANAF databases. Implications of these variations are discussed in the main text.

6.2 Fugacity coefficients

We can also use the K_i equations to investigate the difference between γ_i in the different tools, which only depend on P and T (i.e., not melt composition). Individual γ_i cannot be calculated, but ratios of various combinations can be. If the ratios are similar, this suggests the tools are using similar γ_i (although the γ_i could be perfectly off-setting each other, but that would be unlikely). As for calculating K_i, the ratios of γ_i are only calculated if all the necessary vapor species are calculated and the calculation reaches ≤ 1 bar as K_i must also be calculated.

Most tools are using very similar values for γ_{H_2}, γ_{S_2}, γ_{SO_2}, γ_{H_2S}, and γ_{OCS}, as can be seen by comparing $\gamma_{S_2}^{0.5}/\gamma_{SO_2}$, $[\gamma_{SO_2}\gamma_{H_2}]/\gamma_{H_2S}$, and $\gamma_{OCS}/[\gamma_{CO}\gamma_{S_2}^{0.5}]$ in Figure 3. This makes sense given all tools are using Shi and Saxena [1992] for γ_{S_2}, γ_{SO_2}, γ_{H_2S}, and γ_{OCS}. The differences for VolFe at low-P for γ_{SO_2} and γ_{H_2S} are due to modifications described in Hughes et al. [2023, 2024b] and are therefore expected. Most tools use Shaw and Wones [1964] for γ_{H_2}, except MAGEC that uses a modified version of Shi and Saxena [1992], and therefore the similarity in γ_{H_2} is good. Most tools use equations from Holland and Powell [1991] (although not the same specific equations), except MAGEC that uses eq. (15) from Peng and Robinson [1976], which are likely giving different values leading to the differences seen in $\gamma_{H_2}/\gamma_{H_2O}$ (Figure 3). Values for γ_{CO_2}, γ_{CO}, and γ_{CH_4} seem similar between tools (e.g., $\gamma_{CH_4}/[\gamma_{CO}(\gamma_{H_2})^2]$ and $\gamma_{CH_4}/[\gamma_{CO_2}(\gamma_{H_2})^2]$) in Figure 3). All tools use Shi and Saxena [1992] for γ_{CH_4} and γ_{CO}, and most also use this for γ_{CO_2}, except EVo that uses Holland and Powell [1991]. There appear to be some minor variations in $\gamma_{CO}/\gamma_{CO_2}$, suggesting minor differences in γ_{CO} and/or γ_{CO_2}.

draft for review




Figure 3: Ratios of fugacity coefficients (γ_i calculated using equilibrium constant equations in the text)




Figure 4: Non-ideal factor γ for O_2 vs. pressure for four models.

O_2 in the vapor is highly non-ideal, and the deviation from ideality, the fugacity coefficient (γ), is dependent upon P and T. We can test whether models are treating the non-ideality of O_2 in the same way by calculating γ_{O_2} from the definition of fugacity as a function of P:

$$fO_2 = \gamma_{O_2} \times x^v_{O_2} \times P. \quad (13)$$

γ_{O_2} is unity for an ideal gas, but deviates strongly for a real gas, particularly at high P [e.g., Shi and Saxena 1992]. Supplementary figure 4 shows nearly perfect overlap, with γ_{O_2} from unity at 1 bar up to ~1.5 at 2,000 bars, meaning that this is not a factor in the divergence of computed redox variables and is expected given all tools use the model of Shi and Saxena [1992] for γ_{O_2}. Hence, differences in γ_{O_2} are not driving differences between model outputs.

6.3 Molar mass balance in the vapor

One difficulty when using models from the literature is that some post-processing calculations cannot be carried out because they are at odds with either model assumptions or numerical limitations. The molar fraction of O_2 in the vapor, $x^v_{O_2}$, is a prime example of such limitations. Sulfur_X does not include O_2 in the vapor calculation and therefore is not included in the following discussion. All other models assume that the sum of the molar fractions of all species in the vapor is equal to one, and thus one should be able to deduce the fraction of one species by calculating one minus the sum of all the others species. $x^v_{O_2}$ is notoriously small (typically 10^{-15}–10^{-11}), whereas $x^v_{H_2O}$ is typically 10^{-1}. This means that numerical limitations may be encountered because most models report (and calculate) in double precision (~15 significant digits). Figure 5 shows $x^v_{O_2}$ calculated by difference versus $x^v_{O_2}$ values output by the models. The bottom row shows the residual of the closure calculation $\Sigma x^v_i - 1$ (the sum of all vapor species minus 1) at each P using Kīlauea as a representative example.

D-Compress uses Turbo Delphi 2006 with TuboPower SysTools BCD allowing for floating point precision to 26 significant digits. All other models use Python or MATLAB's default double precision (IEEE 754 binary64) which is precise to 15–17 significant digits. This matters when very small values are converted between decimal representation (as we read it) and binary (as the computer reads it). Our plotted $x^v_{O_2}$ by difference and residual values were likewise computed in binary64




Figure 5: Molar fraction of O_2 in the vapor as reported by modeling tools vs. calculated by difference for all models that report it: D-Compress, EVo, MAGEC, and VolFe. Some by difference values computed for MAGEC were negative and so we plot the absolute values in order to plot on a logarithmic axis. The bottom row shows the residual of the closure calculation (the departure from zero of the sum of all vapor species minus 1) for Kīlauea runs.

from each model's CSV output, meaning that residuals below this floor cannot be resolved regardless of how the values were computed internally. In other words, any trailing digits are meaningless noise from rounding error.

D-Compress closure shows symmetric noise at the $\sim 10^{-17}$ level scattered around zero, reflecting only float64 rounding caused by our by-difference calculation and indicating higher precision within the model itself. The closure plot in EVo reveals systematically negative residuals $\sim 10^{-16}$–10^{-17}, indicating a normalization step within the solver capping sums to one. At reducing conditions below $\log fO_2$ of -15 ($\sim$ ΔIW-3), $x^v_{O_2}$ cannot be recovered by-difference. Data by difference from MAGEC oscillate between negative and positive and, for that reason, we plot the absolute value in the 1:1 figure for MAGEC. The MAGEC output files used in this comparison were exported to 5 significant figures, resulting in the closure error of $\sim 10^{-5}$–10^{-6}. Each species is rounded independently at output time, so per-P step sign distributions are random and centered on zero (output-format rounding, not algorithmic bias). This does not indicate anything wrong with MAGEC's internal solver, but this output precision means that fO_2 cannot be reliably recovered by difference from the exported MAGEC outputs used here. Most VolFe outputs reflect the same expected double-precision behavior as EVo but show closure failure in certain cases, specifically for high-CO_2, low-H_2O points at the mid-P. This is likely the result of its iterative Newton-Raphson solver converging on a near-solution and exiting before normalization of gas species to one can be enforced. Adjusting the solver tolerance dependent upon composition could improve the recoverability of $x^v_{O_2}$ in certain cases.

High precision is not typically required to answer geologic questions, but the impact can be significant due to repeated computation within the code itself before a result gets returned to a user. Likewise, if precision is truncated when the code writes results to disk, any calculation those values are fed back into will never be better than the number of truncated digits, and this would almost certainly go undetected, since floating point error thereafter will mask the previous truncation. It is also worth flagging floating point rounding behavior in Microsoft Excel. Although computations within Excel adhere to IEEE 754 binary64, when any value is *displayed* in the user interface, it is truncated to 15 significant figures, replacing the final two with zeroes. Additionally, if a file with numbers exceeding 11 digits is *saved* from Excel, it's "General" format defaults to scientific notation and may silently drop several significant figures from all values [Griffiths 2022]. This effect has been noted as particularly damaging for data analysis in the genome biology literature [Ziemann et al. 2016].

Copyright notice

References

Allan, J. F., R. Batiza, M. R. Perfit, D. J. Fornari, and R. O. Sack (1989). "Petrology of Lavas from the Lamont Seamount Chain and Adjacent East Pacific Rise, 10° N". *Journal of Petrology* 30(5), pages 1245–1298. DOI: 10.1093/petrology/30.5.1245.

Ardia, P., M. Hirschmann, A. Withers, and B. Stanley (2013). "Solubility of CH4 in a synthetic basaltic melt, with applications to atmosphere–magma ocean–core partitioning of volatiles and to the evolution of the Martian atmosphere". *Geochimica et Cosmochimica Acta* 114, pages 52–71. DOI: 10.1016/j.gca.2013.03.028.

Draft for review




Figure 6: Volatile proportionality factors (eq. 15 with all tools for all basalts showing the change in H_2O, CO_2, reduced S (S^{red}), and oxidized S (S^{ox}) proportionality factors (H_i') as a function of pressure during degassing.

Armstrong, L. S., M. M. Hirschmann, B. D. Stanley, E. G. Falksen, and S. D. Jacobsen (2015). "Speciation and solubility of reduced C–O–H–N volatiles in mafic melt: Implications for volcanism, atmospheric evolution, and deep volatile cycles in the terrestrial planets". *Geochimica et Cosmochimica Acta* 171, pages 283–302. DOI: 10.1016/j.gca.2015.07.007.

Burgisser, A., M. Alletti, and B. Scaillet (2015). "Simulating the behavior of volatiles belonging to the C–O–H–S system in silicate melts under magmatic conditions with the software D-Compress". *Computers & Geosciences* 79, pages 1–14. DOI: 10.1016/j.cageo.2015.03.002.

Canil, D. (2002). "Vanadium in peridotites, mantle redox and tectonic environments: Archean to present". *Earth and Planetary Science Letters* 195(1-2), pages 75–90.

Chase, M. W. (1996). "NIST-JANAF thermochemical tables for the iodine oxides". *Journal of physical and chemical reference data* 25, pages 1229–1339.

Chase, M. W. et al. (1998). "NIST-JANAF thermochemical tables". *Journal of physical and chemical reference data* 28, page 1951.

Chowdhury, P. and R. Dasgupta (2019). "Effect of sulfate on the basaltic liquidus and sulfur Concentration at Anhydrite Saturation (SCAS) of hydrous basalts–Implications for sulfur cycle in subduction zones". *Chemical Geology* 522, pages 162–174.

Clague, D. A., J. G. Moore, J. E. DIXON, and W. B. Friesen (1995). "Petrology of submarine lavas from Kilauea's Puna Ridge, Hawaii". *Journal of Petrology* 36(2), pages 299–349.

Cottrell, E., S. K. Birner, M. Brounce, F. A. Davis, L. E. Waters, and K. A. Kelley (2021). "Oxygen Fugacity Across Tectonic Settings". In: American Geophysical Union (AGU). Chapter 3, pages 33–61.

DeVitre, C. L., E. Gazel, R. S. Ramalho, S. Venugopal, M. Steele-MacInnis, J. Hua, C. M. Allison, L. R. Moore, J. C. Carracedo, and B. Monteleone (2023). "Oceanic intraplate explosive eruptions fed directly from the mantle". *Proceedings of the National Academy of Sciences* 120(33), e2302093120.

Ding, S., T. Plank, P. J. Wallace, and D. J. Rasmussen (2023). "Sulfur_X: A Model of Sulfur Degassing During Magma Ascent". *Geochemistry, Geophysics, Geosystems* 24(4). DOI: 10.1029/2022gc010552.

Dixon, J. E., D. A. Clague, and E. M. Stolper (1991). "Degassing history of water, sulfur, and carbon in submarine lavas from Kilauea Volcano, Hawaii". *The Journal of Geology* 99(3), pages 371–394.

Dixon, J. E., E. M. Stolper, and J. R. Holloway (1995). "An Experimental Study of Water and Carbon Dioxide Solubilities in Mid-Ocean Ridge Basaltic Liquids. Part I: Calibration and Solubility Models". *Journal of Petrology*.

Eguchi, J. and R. Dasgupta (2018). "A CO2 solubility model for silicate melts from fluid saturation to graphite or diamond saturation". *Chemical Geology* 487, pages 23–38. DOI: 10.1016/j.chemgeo.2018.04.012.

Frost, B. R. (1991). "Introduction to oxygen fugacity and its petrologic importance". *Reviews in Mineralogy and Geochemistry* 25(1), pages 1–9.

Gaillard, F., M. Pichavant, S. Mackwell, R. Champallier, B. Scaillet, and C. Mccammon (2003). "Chemical transfer during redox exchanges between H2 and Fe-bearing silicate melts". *American Mineralogist* 88(2-3), pages 308–315. DOI: 10.2138/am-2003-2-308.

Ghiorso, M. S. and G. A. R. Gualda (2015). "An H2O–CO2 mixed fluid saturation model compatible with rhyolite-MELTS". *Contributions to Mineralogy and Petrology* 169(6), page 53. DOI: 10.1007/s00410-015-1141-8.

Griffiths, I. (2022). *Excel, data loss, IEEE754, and precision*. endjin Blog. Accessed: 2026-05-20.

Helz, R. T. and C. R. Thornber (1987). "Geothermometry of Kilauea Iki lava lake, Hawaii". *Bulletin of volcanology* 49(5), pages 651–668.

Hirschmann, M. M., M. S. Ghiorso, F. A. Davis, S. M. Gordon, S. Mukherjee, T. L. Grove, M. Krawczynski, E. Medard, and C. B. Till (2008). "Library of Experimental Phase Relations (LEPR): A database and Web portal for experimental magmatic phase equilibria data: LIBRARY OF EXPERIMENTAL PHASE RELATIONS". *Geochemistry, Geophysics, Geosystems* 9(3). DOI: 10.1029/2007gc001894.

Holland, T. and R. Powell (1991). "A Compensated-Redlich-Kwong (CORK) equation for volumes and fugacities of CO2 and H2O in the range 1 bar to 50 kbar and 100–1600°C". *Contributions to Mineralogy and Petrology* 109(2), pages 265–273. DOI: 10.1007/{BF}00306484.

Hughes, E., P. Liggins, P. Wieser, and E. Stolper (2025). "VolFe: an open-source Python package for calculating melt-vapour equilibria including silicate melt, carbon, hydrogen, sulfur, and noble gases". *Volcanica* 8(2), pages 457–481. DOI: 10.30909/vol/imvc1781.

Hughes, E. C., J. Biasi, I. Fendley, K. Rahilly, T. D. Schlieder, H. Winslow, T. P. Fischer, and P. J. Wallace (2024a). "Modeling the behavior of sulfur in magmatic systems from source to surface: Application to Whakaari/White Island, Aotearoa New Zealand, and Etna, Italy". *Journal of Volcanology and Geothermal Research* 446, page 107939. DOI: 10.1016/j.jvolgeores.2023.107939.

Hughes, E. C., P. Liggins, L. Saper, and E. M. Stolper (2024b). "The effects of oxygen fugacity and sulfur on the pressure of vapor-saturation of magma". *American Mineralogist* 109(3), pages 422–438. DOI: 10.2138/am-2022-8739.

Hughes, E. C., L. M. Saper, P. Liggins, H. S. C. O'Neill, and E. M. Stolper (2023). "The sulfur solubility minimum and maximum in silicate melt". *Journal of the Geological Society* 180(3), jgs2021–125. DOI: 10.1144/jgs2021-125.

Draft for review

Iacono-Marziano, G., Y. Morizet, E. L. Trong, and F. Gaillard (2012). "New experimental data and semi-empirical parameterization of H2O–CO2 solubility in mafic melts". *Geochimica et Cosmochimica Acta* 97, pages 1–23. DOI: 10.1016/j.gca.2012.08.035.

Kress, V. C. and I. S. E. Carmichael (1991). "The compressibility of silicate liquids containing Fe2O3 and the effect of composition, temperature, oxygen fugacity and pressure on their redox states". *Contributions to Mineralogy and Petrology* 108(1-2), pages 82–92. DOI: 10.1007/bf00307328.

Lerner, A. H., M. J. Muth, P. J. Wallace, A. Lanzirotti, M. Newville, G. A. Gaetani, P. Chowdhury, and R. Dasgupta (2021a). "Improving the reliability of Fe-and S-XANES measurements in silicate glasses: correcting beam damage and identifying Fe-oxide nanolites in hydrous and anhydrous melt inclusions". *Chemical Geology* 586, page 120610.

Lerner, A. H., P. J. Wallace, T. Shea, A. J. Mourey, P. J. Kelly, P. A. Nadeau, T. Elias, C. Kern, L. E. Clor, C. Gansecki, et al. (2021b). "The petrologic and degassing behavior of sulfur and other magmatic volatiles from the 2018 eruption of Kılauea, Hawaii: melt concentrations, magma storage depths, and magma recycling". *Bulletin of Volcanology* 83(6), page 43.

Liu, Y., Y. Zhang, and H. Behrens (2005). "Solubility of H2O in rhyolitic melts at low pressures and a new empirical model for mixed H2O–CO2 solubility in rhyolitic melts". *Journal of Volcanology and Geothermal Research* 143(1-3), pages 219–235. DOI: 10.1016/j.jvolgeores.2004.09.019.

Lloyd, A. S., T. Plank, P. Ruprecht, E. H. Hauri, and W. Rose (2013). "Volatile loss from melt inclusions in pyroclasts of differing sizes". *Contributions to Mineralogy and Petrology* 165(1), pages 129–153. DOI: 10.1007/s00410-012-0800-2.

Moore, G. M., T. Vennemann, and I. Carmichael (1998). "An empirical model for the solubility of H2O in magmas to 3 kilobars". *American Mineralogist* 83, pages 36–42.

Moore, L. R., E. Gazel, R. Tuohy, A. S. Lloyd, R. Esposito, M. Steele-MacInnis, E. H. Hauri, P. J. Wallace, T. Plank, and R. J. Bodnar (2015). "Bubbles matter: An assessment of the contribution of vapor bubbles to melt inclusion volatile budgets". *American Mineralogist* 100(4), pages 806–823.

Moussallam, Y., M. Edmonds, B. Scaillet, N. Peters, E. Gennaro, I. Sides, and C. Oppenheimer (2016). "The impact of degassing on the oxidation state of basaltic magmas: a case study of Kılauea volcano". *Earth and Planetary Science Letters* 450, pages 317–325.

Moussallam, Y., C. Oppenheimer, and B. Scaillet (2019). "On the relationship between oxidation state and temperature of volcanic gas emissions". *Earth and Planetary Science Letters* 520, pages 260–267. DOI: 10.1016/j.epsl.2019.05.036.

Muth, M. J. and P. J. Wallace (2021). "Slab-derived sulfate generates oxidized basaltic magmas in the southern Cascade arc (California, USA)". *Geology*. DOI: 10.1130/g48759.1.

Nash, W. M., D. J. Smythe, and B. J. Wood (2019). "Compositional and temperature effects on sulfur speciation and solubility in silicate melts". *Earth and Planetary Science Letters* 507. DOI: 10.1016/j.epsl.2018.12.006.

Newcombe, M. E., T. Plank, Y. Zhang, M. Holycross, A. Barth, A. S. Lloyd, D. Ferguson, B. F. Houghton, and E. Hauri (2020). "Magma Pressure-Temperature-Time Paths During Mafic Explosive Eruptions". *Frontiers in Earth Science* 8, page 531911. DOI: 10.3389/feart.2020.531911.

Newman, S. and J. B. Lowenstern (2002). "VolatileCalc: a silicate melt–H2O–CO2 solution model written in Visual Basic for excel". *Computers & Geosciences* 28(5), pages 597–604. DOI: 10.1016/s0098-3004(01)00081-4.

Nzotta, M. M., D. Sichen, and S. Seetharaman (1999). "A study of the sulfide capacities of iron-oxide containing slags". *Metallurgical and Materials Transactions B* 30(5), pages 909–920. DOI: 10.1007/s11663-999-0096-4.

O'Neill, H. S. C. (1987). "Quartz-fayalite-iron and quartz-fayalite-magnetite equilibria and the free energy of formation of fayalite (Fe 2 SiO 4) and magnetite (Fe 3 O 4)". *American Mineralogist* 72(1-2), pages 67–75.

O'Neill, H. S. C. (2021a). "The thermodynamic controls on sulfide saturation in silicate melts with application to ocean floor basalts". *Magma Redox Geochemistry*, pages 177–213.

O'Neill, H. S. C. and M. I. Pownceby (1993). "Thermodynamic data from redox reactions at high temperatures. I. An experimental and theoretical assessment of the electrochemical method using stabilized zirconia electrolytes, with revised values for the Fe-"FeO", Co-CoO, Ni-NiO and Cu-Cu2O oxygen buffers, and new data for the W-WO2 buffer". *Contributions to Mineralogy and Petrology* 114(3), pages 296–314. DOI: 10.1007/BF01046533.

O'Neill, H. S. (2021b). "Magma Redox Geochemistry". *Geophysical Monograph Series*, pages 177–213. DOI: 10.1002/9781119473206.ch10.

O'Neill, H. S. and J. A. Mavrogenes (2022). "The sulfate capacities of silicate melts". *Geochimica et Cosmochimica Acta* 334, pages 368–382. DOI: 10.1016/j.gca.2022.06.020.

Ohmoto, H. and D. M. Kerrick (1977). "Devolatilization equilibria in graphitic systems". *American Journal of Science* 277(8), pages 1013–1044. DOI: 10.2475/ajs.277.8.1013.

Peng, D.-Y. and D. B. Robinson (1976). "A New Two-Constant Equation of State". *Industrial & Engineering Chemistry Fundamentals* 15(1), pages 59–64. DOI: 10.1021/i160057a011.

Rasmussen, D. J., T. A. Plank, P. J. Wallace, M. E. Newcombe, and J. B. Lowenstern (2020). "Vapor-bubble growth in olivine-hosted melt inclusions". *American Mineralogist* 105(12), pages 1898–1919.

Shaw, H. R. and D. R. Wones (1964). "Fugacity coefficients for hydrogen gas between O degrees and 1000 degrees C, for pressures to 3000 atm". *American Journal of Science* 262(7), pages 918–929. DOI: 10.2475/ajs.262.7.918.

UNDER PEER REVIEW

Shi, P. and S. K. Saxena (1992). "Thermodynamic modeling of the C-H-O-S fluid system". *American Mineralogist* 77, pages 1038–1049.

Shishkina, T. A., R. E. Botcharnikov, F. Holtz, R. R. Almeev, A. M. Jazwa, and A. A. Jakubiak (2014). "Compositional and pressure effects on the solubility of H2O and CO2 in mafic melts". *Chemical Geology* 388, pages 112–129. DOI: 10.1016/j.chemgeo.2014.09.001.

Sides, I., M. Edmonds, J. Maclennan, B. F. Houghton, D. Swanson, and M. J. Steele-MacInnis (2014). "Magma mixing and high fountaining during the 1959 Kılauea Iki eruption, Hawai 'i". *Earth and Planetary Science Letters* 400, pages 102–112.

Sun, C. and C.-T. A. Lee (2022). "Redox evolution of crystallizing magmas with C-H-O-S volatiles and its implications for atmospheric oxygenation". *Geochimica et Cosmochimica Acta* 338, pages 302–321. DOI: https://doi.org/10.1016/j.gca.2022.09.044.

Sun, C. and L. Yao (2024). "Redox equilibria of iron in low- to high-silica melts: A simple model and its applications to C-H-O-S degassing". *Earth and Planetary Science Letters* 638, page 118742. DOI: https://doi.org/10.1016/j.epsl.2024.118742.

Taracsák, Z., T. Mather, S. Ding, T. Plank, M. Brounce, D. Pyle, A. Aiuppa, et al. (2023). "Sulfur from the subducted slab dominates the sulfur budget of the mantle wedge under volcanic arcs". *Earth and Planetary Science Letters* 602, page 117948. DOI: 10.1016/j.epsl.2022.117948.

Thibault, Y. and J. R. Holloway (1994). "Solubility of CO2 in a Ca-rich leucitite: effects of pressure, temperature, and oxygen fugacity". *Contributions to Mineralogy and Petrology* 116(1-2), pages 216–224. DOI: 10.1007/bf00310701.

Tucker, J. M., E. H. Hauri, A. J. Pietruszka, M. O. Garcia, J. P. Marske, and F. A. Trusdell (2019). "A high carbon content of the Hawaiian mantle from olivine-hosted melt inclusions". *Geochimica et Cosmochimica Acta* 254, pages 156–172.

Wieser, P. E., K. Iacovino, S. Matthews, G. Moore, and C. M. Allison (2022). "VESIcal Part II: A critical approach to volatile solubility modelling using an open-source Python3 engine". *Earth and Space Science*. DOI: 10.1029/2021ea001932.

Wieser, P. and M. Gleeson (2023). "PySulfSat: An open-source Python3 tool for modeling sulfide and sulfate saturation". *Volcanica* 6(1), pages 107–127.

Wieser, P. E., F. Jenner, M. Edmonds, J. Maclennan, and B. E. Kunz (2020). "Chalcophile elements track the fate of sulfur at Kılauea Volcano, Hawai'i". *Geochimica et Cosmochimica Acta* 282, pages 245–275.

Wieser, P. E., H. Lamadrid, J. Maclennan, M. Edmonds, S. Matthews, K. Iacovino, F. E. Jenner, C. Gansecki, F. Trusdell, R. L. Lee, et al. (2021). "Reconstructing magma storage depths for the 2018 Kilauean eruption from melt inclusion CO2 contents: the importance of vapor bubbles". *Geochemistry, Geophysics, Geosystems* 22(2), e2020GC009364.

Ziemann, M., Y. Eren, and A. El-Osta (2016). "Gene name errors are widespread in the scientific literature". *Genome Biology* 17(1), page 177. DOI: 10.1186/s13059-016-1044-7.

draft for review