

# Assignment 7: GLMs (Linear Regressios, ANOVA, & t-tests)

Kayla Emerson

Fall 2024

## OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on generalized linear models.

## Directions

1. Rename this file `<FirstLast>_A07_GLMs.Rmd` (replacing `<FirstLast>` with your first and last name).
2. Change “Student Name” on line 3 (above) with your name.
3. Work through the steps, **creating code and output** that fulfill each instruction.
4. Be sure to **answer the questions** in this assignment document.
5. When you have completed the assignment, **Knit** the text and code into a single PDF file.

## Set up your session

1. Set up your session. Check your working directory. Load the tidyverse, agricolae and other needed packages. Import the *raw* NTL-LTER raw data file for chemistry/physics (NTL-LTER\_Lake\_ChemistryPhysics\_Raw.csv). Set date columns to date objects.
2. Build a ggplot theme and set it as your default theme.

```
#1
#load packages
library(tidyverse)
```

```
## -- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
## v dplyr      1.1.4      v readr      2.1.5
## v forcats    1.0.0      v stringr   1.5.1
## v ggplot2    3.5.1      v tibble    3.2.1
## v lubridate  1.9.3      v tidyr     1.3.1
## v purrr      1.0.2
## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()     masks stats::lag()
## i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to become errors
```

```
library(agricolae)
library(here)
```

```
## here() starts at /home/guest/EDE_Fall2024
```

```
library(lubridate)
here()
```

```
## [1] "/home/guest/EDE_Fall2024"
```

```
#import raw dataset
lake_chem <- read.csv(here("Data/Raw/NTL-LTER_Lake_ChemistryPhysics_Raw.csv"),
                      stringsAsFactors = TRUE)

#set date cols to dates
lake_chem$sampldate <- mdy(lake_chem$sampldate)
class(lake_chem$sampldate)
```

```
## [1] "Date"
```

```
#2
#create theme
mytheme <- theme_classic(base_size = 14) +
  theme(axis.text = element_text(color = "darkblue"),
        legend.position = "right")
theme_set(mytheme)
```

## Simple regression

Our first research question is: Does mean lake temperature recorded during July change with depth across all lakes?

3. State the null and alternative hypotheses for this question: > Answer: H0: Mean lake temperature recorded during July does not change with depth across all lakes Ha: Mean lake temperature recorded during July does change with depth across all lakes.
4. Wrangle your NTL-LTER dataset with a pipe function so that the records meet the following criteria:
  - Only dates in July.
  - Only the columns: lakename, year4, daynum, depth, temperature\_C
  - Only complete cases (i.e., remove NAs)
5. Visualize the relationship among the two continuous variables with a scatter plot of temperature by depth. Add a smoothed line showing the linear model, and limit temperature values from 0 to 35 °C. Make this plot look pretty and easy to read.

```
#4
#wrangle dataset
lake_chem_wrangled <- lake_chem %>%
  filter(month(sampldate) == 7) %>%
  select(lakename, year4, daynum, depth, temperature_C) %>%
  filter(!is.na(depth)) %>%
  filter(!is.na(temperature_C))
```


```
#5
```

```
#visualize relationship of continuous vars
plot_one <- lake_chem_wrangled %>%
  ggplot(aes(y = temperature_C, x = depth)) +
  geom_point() +
  labs(x = "Depth",
       y = "Temperature_C") +
  ylim(0, 35) +
  geom_smooth(method = "lm") +
  mytheme

print(plot_one)
```

```
## 'geom_smooth()' using formula = 'y ~ x'
```

```
## Warning: Removed 24 rows containing missing values or values outside the scale range
## ('geom_smooth()').
```



6. Interpret the figure. What does it suggest with regards to the response of temperature to depth? Do the distribution of points suggest anything about the linearity of this trend?

Answer: The figure suggests that as depth increases, temperature decreases. The points appear to be somewhat linearly related, but their distribution is mainly in the mid-range of both variables.

7. Perform a linear regression to test the relationship and display the results.

```
#7
#perform linear regression
regression_one <- lm(
  data = lake_chem_wrangled,
  temperature_C ~ depth)
print(summary(regression_one))

##
## Call:
## lm(formula = temperature_C ~ depth, data = lake_chem_wrangled)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.5173 -3.0192  0.0633  2.9365 13.5834
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) 21.95597    0.06792   323.3  <2e-16 ***
## depth      -1.94621    0.01174  -165.8  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.835 on 9726 degrees of freedom
## Multiple R-squared:  0.7387, Adjusted R-squared:  0.7387
## F-statistic: 2.75e+04 on 1 and 9726 DF,  p-value: < 2.2e-16
```

8. Interpret your model results in words. Include how much of the variability in temperature is explained by changes in depth, the degrees of freedom on which this finding is based, and the statistical significance of the result. Also mention how much temperature is predicted to change for every 1m change in depth.

Answer: A linear regression was run on the dataset wherein temperature was the dependent variable and depth was the independent variable. The coefficients in the output give us the equation for the line,  $y = -1.94621(x) + 21.95597$ . For every 1m increase in depth, temperature is expected to drop 1.94621 degrees. Because the slope is negative, we know that depth and temperature have a negative relationship (when one gets larger, the other gets smaller). The residual standard error is 3.835 on 9726 degrees of freedom which means that the actual values are on average 3.835 points from the estimated values based on the linear model, in units of Y. The multiple R-squared value is 0.7387 which means 73.87% of the variation in depth, temperature is explaining. It describes the fit of the data and 0.7387 is not too bad. Lastly, the p-value is lower than 0.05, so we can reject the null-hypothesis and we can conclude there is a statistically significant relationship between depth and temperature.

---

## Multiple regression

Let's tackle a similar question from a different approach. Here, we want to explore what might the best set of predictors for lake temperature in July across the monitoring period at the North Temperate Lakes LTER.

9. Run an AIC to determine what set of explanatory variables (year4, daynum, depth) is best suited to predict temperature.
10. Run a multiple regression on the recommended set of variables.

```
#9
#see corr plot
library(corrplot)
```

```
## corrplot 0.94 loaded
```

```
lake_chem_wrangled.test <-
  lake_chem_wrangled %>%
  select(temperature_C, depth, year4, daynum) %>%
  na.omit()
```


```
lake_chem_Corr <- cor(lake_chem_wrangled.test)
```

```
corrplot(lake_chem_Corr, upper = "ellipse")
```

```
## Warning in text.default(pos.xlabel[, 1], pos.xlabel[, 2], newcolnames, srt =
## tl.srt, : "upper" is not a graphical parameter
```

```
## Warning in text.default(pos.ylabel[, 1], pos.ylabel[, 2], newrownames, col =
## tl.col, : "upper" is not a graphical parameter
```

```
## Warning in title(title, ...): "upper" is not a graphical parameter
```



```
#run AIC
lake_chem_all <- lm(data = lake_chem_wrangled,
                    temperature_C ~ depth + year4 + daynum)

summary(lake_chem_all)

##
## Call:
## lm(formula = temperature_C ~ depth + year4 + daynum, data = lake_chem_wrangled)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.6536 -3.0000  0.0902  2.9658 13.6123
##
## Coefficients:
##              Estimate Std. Error  t value Pr(>|t|)
## (Intercept) -8.575564   8.630715  -0.994   0.32044
## depth       -1.946437   0.011683 -166.611 < 2e-16 ***
## year4        0.011345   0.004299   2.639   0.00833 **
## daynum       0.039780   0.004317   9.215   < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.817 on 9724 degrees of freedom
## Multiple R-squared:  0.7412, Adjusted R-squared:  0.7411
## F-statistic: 9283 on 3 and 9724 DF,  p-value: < 2.2e-16
```

```
step(lake_chem_all)
```

```
## Start:  AIC=26065.53
## temperature_C ~ depth + year4 + daynum
##
##           Df Sum of Sq  RSS   AIC
## <none>                 141687 26066
## - year4    1         101 141788 26070
## - daynum   1        1237 142924 26148
## - depth    1       404475 546161 39189
##
##
## Call:
## lm(formula = temperature_C ~ depth + year4 + daynum, data = lake_chem_wrangled)
##
## Coefficients:
## (Intercept)      depth      year4      daynum
##    -8.57556    -1.94644     0.01134     0.03978
```

```
#10
#run multiple regression on recommended set of vars
```

```
regression.two <- lm(
  data = lake_chem_wrangled,
  temperature_C ~ depth + daynum)
summary(regression.two)
```

```
##
## Call:
## lm(formula = temperature_C ~ depth + daynum, data = lake_chem_wrangled)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.6174 -2.9809  0.0845  2.9681 13.4406
##
## Coefficients:
##              Estimate Std. Error  t value Pr(>|t|)
## (Intercept) 14.088588   0.855505   16.468  <2e-16 ***
## depth       -1.946111   0.011685  -166.541 <2e-16 ***
## daynum        0.039836   0.004318    9.225  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.818 on 9725 degrees of freedom
## Multiple R-squared:  0.741, Adjusted R-squared:  0.741
## F-statistic: 1.391e+04 on 2 and 9725 DF, p-value: < 2.2e-16
```

11. What is the final set of explanatory variables that the AIC method suggests we use to predict temperature in our multiple regression? How much of the observed variance does this model explain? Is this an improvement over the model using only depth as the explanatory variable?

Answer: The AIC method suggests we use depth and daynum to predict the temperature in the multiple regression. This model explains 74.12% of the the observed variance, which is a very small improvement using only depth as the explanatory variable.

---

## Analysis of Variance

12. Now we want to see whether the different lakes have, on average, different temperatures in the month of July. Run an ANOVA test to complete this analysis. (No need to test assumptions of normality or similar variances.) Create two sets of models: one expressed as an ANOVA models and another expressed as a linear model (as done in our lessons).

```
#12
#run a one-way ANOVA
lake_temps.anova <- aov(data = lake_chem_wrangled, temperature_C ~ lakename)
summary(lake_temps.anova)
```

```
##              Df Sum Sq Mean Sq F value Pr(>F)
## lakename        8  21642   2705.2      50 <2e-16 ***
## Residuals     9719 525813     54.1
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
#run an ANOVA expressed as a lm
lake_temps.anova2 <- lm(data = lake_chem_wrangled, temperature_C ~ lakename)
summary(lake_temps.anova2)
```

```
##
## Call:
## lm(formula = temperature_C ~ lakename, data = lake_chem_wrangled)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -10.769  -6.614  -2.679   7.684  23.832
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)      17.6664     0.6501  27.174 < 2e-16 ***
## lakenameCrampton Lake    -2.3145     0.7699  -3.006 0.002653 **
## lakenameEast Long Lake   -7.3987     0.6918 -10.695 < 2e-16 ***
## lakenameHummingbird Lake  -6.8931     0.9429  -7.311 2.87e-13 ***
## lakenamePaul Lake        -3.8522     0.6656  -5.788 7.36e-09 ***
## lakenamePeter Lake       -4.3501     0.6645  -6.547 6.17e-11 ***
## lakenameTuesday Lake    -6.5972     0.6769  -9.746 < 2e-16 ***
## lakenameWard Lake        -3.2078     0.9429  -3.402 0.000672 ***
## lakenameWest Long Lake   -6.0878     0.6895  -8.829 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 7.355 on 9719 degrees of freedom
## Multiple R-squared:  0.03953,    Adjusted R-squared:  0.03874
## F-statistic:    50 on 8 and 9719 DF,  p-value: < 2.2e-16
```

13. Is there a significant difference in mean temperature among the lakes? Report your findings.

Answer: There is a significant difference in mean temperature among the lakes. Both the one-way ANOVA and the ANOVA as a linear model resulted in a p-value of less than 0.05, so we can reject the null hypothesis.

14. Create a graph that depicts temperature by depth, with a separate color for each lake. Add a `geom_smooth` (method = "lm", se = FALSE) for each lake. Make your points 50 % transparent. Adjust your y axis limits to go from 0 to 35 degrees. Clean up your graph to make it pretty.


```
#14.
plot_two <- lake_chem_wrangled %>%
  ggplot(aes(y = temperature_C, x = depth, color = lakename)) +
  geom_point(alpha = 0.5) +
  geom_smooth(method = "lm", se = FALSE) +
  labs(x = "Depth",
       y = "Temperature_C",
       color = "Lake Name") +
  scale_y_continuous(limits= c(0, 35), expand = c(0,0)) +
  mytheme

print(plot_two)
```

```
## 'geom_smooth()' using formula = 'y ~ x'
```

```
## Warning: Removed 73 rows containing missing values or values outside the scale range
## ('geom_smooth()').
```





15. Use the Tukey's HSD test to determine which lakes have different means.

```
#15
#run tukey test
TukeyHSD(lake_temps.anova)
```

```
## Tukey multiple comparisons of means
## 95% family-wise confidence level
##
## Fit: aov(formula = temperature_C ~ lakename, data = lake_chem_wrangled)
##
## $lakename
##
```

	diff	lwr	upr	p adj
Crampton Lake-Central Long Lake	-2.3145195	-4.7031913	0.0741524	0.0661566
East Long Lake-Central Long Lake	-7.3987410	-9.5449411	-5.2525408	0.0000000
Hummingbird Lake-Central Long Lake	-6.8931304	-9.8184178	-3.9678430	0.0000000
Paul Lake-Central Long Lake	-3.8521506	-5.9170942	-1.7872070	0.0000003
Peter Lake-Central Long Lake	-4.3501458	-6.4115874	-2.2887042	0.0000000
Tuesday Lake-Central Long Lake	-6.5971805	-8.6971605	-4.4972005	0.0000000
Ward Lake-Central Long Lake	-3.2077856	-6.1330730	-0.2824982	0.0193405
West Long Lake-Central Long Lake	-6.0877513	-8.2268550	-3.9486475	0.0000000
East Long Lake-Crampton Lake	-5.0842215	-6.5591700	-3.6092730	0.0000000
Hummingbird Lake-Crampton Lake	-4.5786109	-7.0538088	-2.1034131	0.0000004
Paul Lake-Crampton Lake	-1.5376312	-2.8916215	-0.1836408	0.0127491
Peter Lake-Crampton Lake	-2.0356263	-3.3842699	-0.6869828	0.0000999

## Tuesday Lake-Crampton Lake	-4.2826611	-5.6895065	-2.8758157	0.0000000
## Ward Lake-Crampton Lake	-0.8932661	-3.3684639	1.5819317	0.9714459
## West Long Lake-Crampton Lake	-3.7732318	-5.2378351	-2.3086285	0.0000000
## Hummingbird Lake-East Long Lake	0.5056106	-1.7364925	2.7477137	0.9988050
## Paul Lake-East Long Lake	3.5465903	2.6900206	4.4031601	0.0000000
## Peter Lake-East Long Lake	3.0485952	2.2005025	3.8966879	0.0000000
## Tuesday Lake-East Long Lake	0.8015604	-0.1363286	1.7394495	0.1657485
## Ward Lake-East Long Lake	4.1909554	1.9488523	6.4330585	0.0000002
## West Long Lake-East Long Lake	1.3109897	0.2885003	2.3334791	0.0022805
## Paul Lake-Hummingbird Lake	3.0409798	0.8765299	5.2054296	0.0004495
## Peter Lake-Hummingbird Lake	2.5429846	0.3818755	4.7040937	0.0080666
## Tuesday Lake-Hummingbird Lake	0.2959499	-1.9019508	2.4938505	0.9999752
## Ward Lake-Hummingbird Lake	3.6853448	0.6889874	6.6817022	0.0043297
## West Long Lake-Hummingbird Lake	0.8053791	-1.4299320	3.0406903	0.9717297
## Peter Lake-Paul Lake	-0.4979952	-1.1120620	0.1160717	0.2241586
## Tuesday Lake-Paul Lake	-2.7450299	-3.4781416	-2.0119182	0.0000000
## Ward Lake-Paul Lake	0.6443651	-1.5200848	2.8088149	0.9916978
## West Long Lake-Paul Lake	-2.2356007	-3.0742314	-1.3969699	0.0000000
## Tuesday Lake-Peter Lake	-2.2470347	-2.9702236	-1.5238458	0.0000000
## Ward Lake-Peter Lake	1.1423602	-1.0187489	3.3034693	0.7827037
## West Long Lake-Peter Lake	-1.7376055	-2.5675759	-0.9076350	0.0000000
## Ward Lake-Tuesday Lake	3.3893950	1.1914943	5.5872956	0.0000609
## West Long Lake-Tuesday Lake	0.5094292	-0.4121051	1.4309636	0.7374387
## West Long Lake-Ward Lake	-2.8799657	-5.1152769	-0.6446546	0.0021080

16. From the findings above, which lakes have the same mean temperature, statistically speaking, as Peter Lake? Does any lake have a mean temperature that is statistically distinct from all the other lakes?

Answer: From the Tukey test, the lakes that have the same mean temperature, statistically, as Peter Lake are Paul Lake and Ward Lake. None of the lakes have a mean temperature that is statistically distinct from all of the other lakes.

17. If we were just looking at Peter Lake and Paul Lake. What's another test we might explore to see whether they have distinct mean temperatures?

Answer: When just looking at Peter and Paul lakes, another test we can use to explore whether they have distinct mean temperatures is the T-test.

18. Wrangle the July data to include only records for Crampton Lake and Ward Lake. Run the two-sample T-test on these data to determine whether their July temperature are same or different. What does the test say? Are the mean temperatures for the lakes equal? Does that match you answer for part 16?

```
#Filter for only Crampton Lake and Ward Lake
lake_chem_super_wrangled <- lake_chem_wrangled %>%
  filter(lakename == "Crampton Lake" | lakename == "Ward Lake")

#Run a two-sample T-test
lake_two_ttest <- t.test(lake_chem_super_wrangled$temperature_C ~ lake_chem_super_wrangled$lakename)
lake_two_ttest
```

##

```
## Welch Two Sample t-test
##
## data: lake_chem_super_wrangled$temperature_C by lake_chem_super_wrangled$lakename
## t = 1.1181, df = 200.37, p-value = 0.2649
## alternative hypothesis: true difference in means between group Crampton Lake and group Ward Lake is not equal to 0
## 95 percent confidence interval:
## -0.6821129 2.4686451
## sample estimates:
## mean in group Crampton Lake      mean in group Ward Lake
##                15.35189                14.45862
```

Answer: The two-sample T-test says that the mean temperatures at Crampton and Ward Lakes in the month of July are not statistically different, because the p-value is greater than 0.05. The mean temperatures are not equal, but they are not statistically different enough to reject the null hypothesis. This does match my answer for part 16 because the Tukey test that was run between Crampton and Ward lakes did not result in a p-value below 0.05.