Search

home / study / business / economics / economics questions and answers / the tensile strength of a synthetic fiber was of interest to the manufact...

Question: The tensile strength of a synthetic fiber was of interest to the ...

(2 bookmarks)

2. (30 points) The tensile strength of a synthetic fiber was of interest to the manufacturer. It is suspected that strength is related to the percentage of cotton in the fiber. Five levels of cotton percentage were used, and five replicates were run in random order, resulting in the following data.

	1	2	3	4	5	
15	7	7	15	11	9	
20	12	17	12	18	18	
25	14	18	18	19	19	
30	19	25	22	19	23	
35	7	10	11	15	11	

c) Estimate σ^2 .

Show transcribed image text

Expert Answer (



Anonymous answered this

609 answers

Was this answer helpful?















View all solutions

Algorithms

1st Edition

Answer:

In the given data, the tensile strength of synthetic fibre under 5 different percentages of cotton are measured.

Test the data whether there is significant difference between the means or not at $\alpha = 0.05$ level of significance.

State the null and alternative hypotheses for the given scenario as below.


$$H_0: \mu_1 = \mu_2 = \mu_3 = \mu_4 = \mu_5$$

$$H_1$$
: At least one μ_i is different $(i = 1, 2, 3, 4, 5)$


Test the hypothesis at $\alpha=0.05$ level of significance.

Use the following MINITAB instructions to conduct ANOVA for the given data.


- 1) Import the data into MINITAB worksheet.
- 2) From the menu bar, select $Stat \rightarrow ANOVA \rightarrow One-Way...$
- 3) Select the data columns into Responses field.
- 4) Define the Confidence level as "95.0".
- 5) Click on Comparisons.
- 6) Check the option Fisher's, individual error rate and define the value as 5.
- 7) Click on Graphs.
- 8) Check the option Boxplots of data.
- 9) Click on OK and again on OK.



Snap a photo from your phone to post a question We'll send you a one-time download



Thus, the resultant ANOVA is summarized as below.



From the output, the F-statistic is obtained as F = 14.76.

The corresponding ρ -value of the test statistic at (4,20) degrees of freedom is obtained as p=0.000.

As the p-value is less than $\alpha = 0.05$ level of significance, so reject the null hypothesis.

Therefore, the mean tensile strength of synthetic fibre under 5 different percentages will differ significantly.

(a)

From the results of ANOVA, one can conclude that the mean tensile strength of synthetic fibre under 5 different percentages of cotton will differ significantly.

The Fisher's LSD method is applied when there is significant difference between the means of the treatments.

State the null and alternative hypotheses for pairwise comparisons of means as follows:

$$\begin{array}{lll} H_{01}: \mu_1 = \mu_2 & H_{11}: \mu_1 \neq \mu_2 \\ H_{02}: \mu_1 = \mu_3 & H_{12}: \mu_1 \neq \mu_3 \\ H_{03}: \mu_1 = \mu_4 & H_{13}: \mu_1 \neq \mu_4 \\ H_{04}: \mu_1 = \mu_5 & H_{14}: \mu_1 \neq \mu_5 \\ H_{05}: \mu_2 = \mu_3 & \text{Versus} \\ H_{06}: \mu_2 = \mu_4 & H_{16}: \mu_2 \neq \mu_4 \\ H_{07}: \mu_2 = \mu_5 & H_{17}: \mu_2 \neq \mu_5 \\ H_{08}: \mu_3 = \mu_4 & H_{18}: \mu_3 \neq \mu_4 \\ H_{09}: \mu_3 = \mu_5 & H_{19}: \mu_3 \neq \mu_5 \\ H_{10}: \mu_4 = \mu_5 & H_{110}: \mu_4 \neq \mu_5 \end{array}$$

Test the hypotheses at $\alpha = 0.05$ level of significance.



Chegg Study Textbook Solutions Expert Q&A Practice Percentage N Mean Grouping

```
Percentage N Mean Groups
30 5 21.600 A
25 5 17.600 B
20 5 15.400 B
35 5 10.800 C
15 5 9.800 C
```

Means that do not share a letter are significantly different.

Fisher 95% Individual Confidence Intervals All Pairwise Comparisons among Levels of Percentage Simultaneous confidence level = 73.57%

Percentage = 15 subtracted from:

Percentage	Lower	Center	Upper		+	+	+-
20	1.855	5.600	9.345		(*)	
25	4.055	7.800	11.545			(*)
30	8.055	11.800	15.545			(-*)
35	-2.745	1.000	4.745		(*)	
					+	+	+-
				-8.0	0.0	8.0	16.0

Percentage = 20 subtracted from:

Percentage	Lower	Center	Upper	
25	-1.545	2.200	5.945	(*)
30	2.455	6.200	9.945	(*)
35	-8.345	-4.600	-0.855	(*)
				0 0 0 0 0 16 0

Percentage = 25 subtracted from:

Percentage	Lower	Center	Upper		+	+	+-
30	0.255	4.000	7.745		(-*)	
35	-10.545	-6.800	-3.055	(*)			
					+		
							16.0

Percentage = 30 subtracted from:




From the above boxplot, it can be observed that the mean tensile strength of synthetic fibre under cotton percentages 15% and 35%, and the mean tensile strength of synthetic fibre under cotton percentages 20% and 25% are not significantly different and the mean tensile strength of synthetic fibre under cotton percentage 30% is significantly different from others.



(h)

The above MINITAB instructions (7) and (8) will plot a comparative boxplots with mean tensile strengths at each level of cotton percentages as follows:



From the above boxplot, it can be observed that the mean tensile strength of synthetic fibre under cotton percentages 15% and 35% are slightly closer, the mean tensile strength of synthetic fibre under cotton percentages 20% and 25% are slightly closer and the mean tensile strength of synthetic fibre under cotton percentage 30% is comparatively more than the others.

Therefore, the comparison through boxplot is similar to the Fisher's LSD method.

Comment >

Practice with similar questions

Q: The tensile strength of a synthetic fiber was of interest to the manufacturer. It is suspected that strength is related to the percentage of cotton in the fiber. Five levels of cotton percentage were used, and five replicates were run in random order, resulting in the following data. a) Test the hypothesis that the percentage of cotton affect the tensile strength. Use alpha = 0.0...

A: See answer

Questions viewed by other students

Q: A manufacturer is interested in testing the tensile strength of a synthetic fiber. We suspect strength is related to the percentage of cotton in the fiber. Five levels of cotton percentage were tested, and five data points were collected at each percentage. Observations cotton % 1 2 3 4 5 15 7 7 15 11 9 20 12 17 12 18 18 25 14 18 18 19 19 30 19 25 22 19 23 35 7 10 1...

A: See answer 100% (1 rating)

Q: Tensile strength of synthetic fiber, which contains different amount of cotton. It is suspected that strength is related to the percentage of cotton in the fiber. Five levels of cotton percentage were used (denoted as rows in table), and measurements took place in five days (denoted as columns). day cotton percentage 1 2 3 4 5 15 20 25 30 35 7 7 15 11 9 12 17 12 18 18 14 18 18 19 1...

A: See answer

Show more v

CO	М	PA	N	Υ

About Chegg Chegg For Good College Marketing Corporate Development

Investor Relations Jobs

Join Our Affiliate Program Media Center Site Map

LEGAL & POLICIES

Honor Shield

Advertising Choices Cookie Notice General Policies Intellectual Property Rights Terms of Use Global Privacy Policy DO NOT SELL MY INFO Honor Code

Cheap Textbooks Chegg Coupon Chegg Play Chegg Study Help College Textbooks eTextbooks Flashcards Learn Chegg Math Solver

CHEGG PRODUCTS AND SERVICES

Mobile Apps Sell Textbooks Solutions Manual Study 101 Textbook Rental Used Textbooks Digital Access Codes Chegg Money

CHEGG NETWORK

EasyBib Internships.com Thinkful

CUSTOMER SERVICE

Customer Service Give Us Feedback Help with eTextbooks Help to use EasyBib Plus . Manage Chegg Study Subscription Return Your Books Textbook Return Policy





© 2003-2021 Chegg Inc. All rights reserved.