Instructions for preparing the solution script:

- Write your name, ID#, and Section number clearly in the very front page.
- Write all answers sequentially.
- Start answering a question (not the pat of the question) from the top of a new page.
- Write legibly and in orderly fashion maintaining all mathematical norms and rules. Prepare a single solution file and rename it "ID#\_FirstName\_TheorySection#.pdf".
- Start working right away. There is no late submission form. If you miss the deadline, you need to use the make-up assignment to cover up the marks.
- 1. Consider a set of data points (0, 1), (0.5, 1.6487) and (1, 2.7183). Now answer the following:
  - (a) (6 marks) Compute the Lagrange bases:  $l_0(x)$ ,  $l_1(x)$  and  $l_2(x)$ .
  - (b) (3 marks) Write the interpolating polynomial and find the value of the interpolating polynomial at x = 0.2.
  - (c) (3 marks) If the data are given by the function  $f(x) = e^x$ , what will be the maximum error of the Lagrange polynomial function at 0.2? (use Weierstrass Approximation)
  - (d) (2 marks) Why is the Lagrange method better than the Vandermonde matrix method to find an interpolating polynomial?
- 2. Consider the data points (-1, 8), (0, 4) and (1, 16) and . Using these data and Newton's divided/difference method, answer the following:
  - (a) (6 marks) Compute the coefficients  $a_k$  for k = 0, 1, 2. Also write the interpolating polynomial.
  - (b) (2 marks) Using the polynomial you found in the previous part, find  $p_n(0.5)$  and  $p_n(-0.9)$ .
- 3. (a) (6 marks) Compute the upper bound of the error using Cauchy's Theorem for  $f(x) = \sin^2(x/2)$  with nodes  $\{-\pi/3, 0, \pi/3\}$  within the interval [-1.2, 1.2].
  - (b) (2 marks) Why are Chebyshev nodes an optimal choice in Interpolation?
- 4. Consider the following data set:

	$\overline{x}$	f(x)	f'(x)
(	).1	-0.62050	3.58502
(	0.2	-0.28340	3.14033

Answer the following based on the above data:

- (a) (8 marks) Compute the Hermite bases:  $h_0(x)$ ,  $h_1(x)$ ,  $\hat{h}_0(x)$  and  $\hat{h}_1(x)$ .
- (b) (2 marks) Write the Hermite polynomial and find the value at x = 0.15.