

1. Information about the indicator	
Goal	Eliminate hunger, ensure food security and improve nutrition, and promote sustainable agricultural development.
Task	End all forms of malnutrition by 2030, including achieving the internationally agreed targets for stunting and wasting in children under the age of five by 2025, and meeting the nutritional needs of adolescent girls, pregnant and lactating women, and the elderly.
The indicator	Prevalence of malnutrition (weight for height $>+2$ or <-2 standard deviation from the median of the WHO Child Growth Standards) among children under 5 years of age, by type (wasting and overweight)
2. Information about the organization	
Organization	Bureau of National Statistics of the Agency for Strategic Planning and Reforms of the Republic of Kazakhstan
3. Определения и понятия	
Definition	Percentage of children under the age of 5 who have a weight-to-height ratio: a) by two standard deviations (moderate and severe deviation from the norm); b) three standard deviations (a strong degree of deviation from the norm) below the median value of this indicator for children of the same age, established by WHO.
Basic concepts:	Exhaustion refers to a child who is too thin for his height. Exhaustion or acute malnutrition, the result of recent rapid weight loss or inability to gain weight. Overweight refers to a child who is too heavy for his height.
Justification and interpretation	Child growth is an internationally accepted characteristic of child development, depending on the level of infant nutrition. Child exhaustion carries an increased risk of death, but treatment is possible. Overweight due to improper nutrition occurs as a result of the fact that a person spends too few calories relative to the amount consumed from food and drinks, and increases the risk of non-communicable diseases in the future.
4. Data sources and collection methods	
Data sources Data	Sample household survey "Multi-indicator Cluster Survey" (MIX)
Collection Methods	Weighing the height and weight of children under 5 years of age in households selected by random sampling
Unit of measurement	Percent
Periodicity	Once every five years
5. Calculation method and other methodological considerations	
Calculation method:	The percentage of children under the age of 5 who have a weight-to-height ratio below/above the median value of this indicator. It is based on a standardized methodology using Child Growth Standards developed by WHO. The weight-to-height ratio is expressed in units of standard deviation (z-value)

	based on the median of the reference population.
Comments and restrictions:	-
6. Доступность данных и дезагрегация	
Data availability and gaps:	The data is available on the website of the Bureau of National Statistics of the Agency for Strategic Planning and Reforms of the Republic of Kazakhstan.
The level of disaggregation:	By gender, type of locality, region, age in months, mother's education level
7. Comparability with international data/standards	The MIX is a UNICEF international survey that uses international standards in methodology.
8. References and documentation	www.stat.gov.kz