$\tilde{\chi}_1^{\pm}(100) \to W^{\pm} \tilde{\chi}_1^0(0)$ (ATLAS_2014_I1286761 (1403.5294))

• Process: $\tilde{\chi}_1^+ \tilde{\chi}_1^- : \tilde{\chi}_1^{\pm} \to W^{\pm} \tilde{\chi}_1^0$.

• The number of events: $5 \cdot 10^4$.

• Event Generator: Herwig++ 2.5.2.

#	cut name	$\epsilon_{ m Exp}$	$\epsilon_{ ext{Atom}}$	Atom Exp	$\frac{\text{(Exp-Atom)}}{\text{Error}}$	#/?	$R_{\rm Exp}$	R_{Atom}	Atom Exp	(Exp-Atom) Error
0	$= 2 \text{ OSlep } p_T > 35, 20: \text{ SF}$	100.0	100.0							
1	Jet Veto: SF	49.5 ± 1.96	68.02 ± 3.24	1.37	4.88	0	0.49 ± 0.02	0.68 ± 0.03	1.37	4.88
2	Z Veto: SF	40.81 ± 1.78	53.67 ± 2.88	1.31	3.79	1	0.82 ± 0.04	0.79 ± 0.04	0.96	-0.64
3	WWa: $p_T(\ell\ell) > 80$: SF	6.85 ± 0.73	7.96 ± 1.11	1.16	0.83	2	0.17 ± 0.02	0.15 ± 0.02	0.88	-0.72
4	WWa: METrel > 80 : SF	4.06 ± 0.56	5.46 ± 0.92	1.34	1.3	3	0.59 ± 0.08	0.69 ± 0.12	1.16	0.66
5	WWa: $m_{\ell\ell} < 120$: SF	2.77 ± 0.46	4.21 ± 0.81	1.52	1.54	4	0.68 ± 0.11	0.77 ± 0.15	1.13	0.47

Table 1: The cut-flow table for the same flavour channel.

#	cut name	$\epsilon_{ m Exp}$	$\epsilon_{ ext{Atom}}$	Atom Exp	$\frac{\text{(Exp-Atom)}}{\text{Error}}$	#/?	R_{Exp}	R_{Atom}	Atom Exp	(Exp-Atom) Error
0	$= 2 \text{ OSlep } p_T > 35, 20: \text{ DF}$	100.0	100.0							
1	Jet Veto: DF	49.93 ± 1.96	65.23 ± 3.15	1.31	4.12	0	0.5 ± 0.02	0.65 ± 0.03	1.31	4.12
2	Z Veto: DF	49.93 ± 1.96	65.23 ± 3.15	1.31	4.12	1	1.0 ± 0.04	1.0 ± 0.05	1.0	0.0
3	WWa: $p_T(\ell\ell) > 80$: DF	7.69 ± 0.77	6.46 ± 1.0	0.84	-0.98	2	0.15 ± 0.02	0.1 ± 0.02	0.64	-2.53
4	WWa: METrel > 80 : DF	4.82 ± 0.61	3.69 ± 0.75	0.77	-1.16	3	0.63 ± 0.08	0.57 ± 0.12	0.91	-0.39
5	WWa: $m_{\ell\ell} < 120$: DF	3.29 ± 0.5	3.08 ± 0.69	0.93	-0.25	4	0.68 ± 0.1	0.83 ± 0.19	1.22	0.7

Table 2: The cut-flow table for the different flavour channel.