

Stéphane Lehericy  
Malgorzata Marjanska  
Lilia Mesrob  
Marie Sarazin  
Serge Kinkingnehun

## Magnetic resonance imaging of Alzheimer's disease

Received: 24 February 2006  
Revised: 12 May 2006  
Accepted: 22 May 2006  
Published online: 25 July 2006  
© Springer-Verlag 2006

S. Lehericy (✉)  
Department of Neuroradiology,  
Université Pierre et Marie Curie–Paris 6,  
Groupe Hospitalier, Pitié-Salpêtrière,  
47-83 boulevard de l'Hôpital,  
Paris 75651, Cedex 13, France  
Tel.: +33-1-46163510  
Fax: +33-142-163515  
e-mail: stephane.lehericy@psl.aphp.fr

S. Lehericy · L. Mesrob ·  
S. Kinkingnehun  
Inserm U610,  
Université Pierre et Marie Curie–Paris 6,  
Groupe Hospitalier, Pitié-Salpêtrière,  
47-83 boulevard de l'Hôpital,  
Paris 75651, Cedex 13, France  
e-mail: lmesrob@yahoo.fr  
e-mail: serge@skinkin.com

M. Marjanska  
Center for Magnetic Resonance  
Research and Department of  
Radiology, University of Minnesota,  
2021 6th Street SE,  
Minneapolis, MN 55455, USA  
e-mail: gosia@cmrr.umn.edu

M. Sarazin  
Department of Neurology,  
Université Pierre et Marie Curie–Paris 6,  
Groupe Hospitalier Pitié-Salpêtrière,  
47-83 boulevard de l'Hôpital,  
Paris 75651, Cedex 13, France  
e-mail: marie.sarazin@psl.aphp.fr

**Abstract** A modern challenge for neuroimaging techniques is to contribute to the early diagnosis of neurodegenerative diseases, such as Alzheimer's disease (AD). Early diagnosis includes recognition of pre-demented conditions, such as mild cognitive impairment (MCI) or having a high risk of developing AD. The role of neuroimaging therefore extends beyond its traditional role of excluding other conditions such as neurosurgical lesions. In addition, early diagnosis would allow early treatment using currently available therapies or new therapies in the future. Structural imaging can detect and follow the time course of subtle brain atrophy as a surrogate marker for pathological processes. New MR techniques and image analysis software can detect

subtle brain microstructural, perfusion or metabolic changes that provide new tools to study the pathological processes and detect pre-demented conditions. This review focuses on markers of macro- and microstructural, perfusion, diffusion and metabolic MR imaging and spectroscopy in AD.

**Keywords** Alzheimer's disease · MRI · Volumetry · Diffusion · Perfusion · Spectroscopy

### Introduction

Alzheimer's disease and neurodegenerative dementias are growing health problems, since populations are constantly aging. The diagnosis of neurodegenerative diseases is therefore a challenge for modern neuroimaging techniques that extends them beyond their traditional role of excluding other conditions such as neurosurgical lesions. In addition, early diagnosis would allow early treatment using currently available therapies, such as cholinesterase inhibitors that

improve or stabilize cognition more efficiently when administered in the early stages of the disease, or using new therapies in the future. Early diagnosis includes recognition of the pre-demented conditions, such as the identification of people with mild cognitive impairment (MCI) or with a high risk of developing AD.

The hallmark of Alzheimer's disease is the presence of extracellular deposits of  $\beta$ -amyloid plaques, intraneuronal neurofibrillary tangles (NFT) and neuropil threads (paired helical filaments containing abnormally phosphorylated

tau protein). Histological studies have shown that the progression of neurofibrillary changes in Alzheimer's disease follows a specific pattern in the brain [1, 2]. Lesion deposition begins in the entorhinal area, then the hippocampal formation, and extends sequentially toward the temporal neocortex, multimodal associative temporal areas, and other neocortical areas. Therefore, lesion burden and neuronal loss are higher in the medial temporal lobe (MTL) than in other brain regions [1, 2].

#### CT in Alzheimer's disease

Computed tomography (CT) is mainly used to rule out surgical causes of dementia. To obtain the best data, the CT scan should be performed using thin slices parallel to the long axis of the hippocampus. Helical acquisitions allow coronal reconstruction perpendicular to the long axis of the hippocampus and therefore a better visualization of medial temporal lobe atrophy. CT shows an enlargement of cerebrospinal fluid spaces, lateral ventricles, particularly the temporal horn of the lateral ventricle, and cortical sulci. CT studies showed that the rate of brain atrophy is larger than the annual rate observed in normal aging [3] and that the hippocampal fissure is wider than in age-matched nondemented people [4]. CT is less efficient than MR in identifying early positive signs of neurodegenerative dementias, i.e., medial temporal lobe atrophy in Alzheimer's disease. Therefore, MR is preferred over CT whenever possible.

#### MRI in Alzheimer's disease

##### *Brain atrophy in Alzheimer's disease*

MR acquisitions that are useful in Alzheimer's disease include high resolution T<sub>1</sub>-weighted images perpendicular

to the long axis of the hippocampus to evaluate medial temporal lobe atrophy (Figs. 1 and 2), axial or coronal T<sub>2</sub>-weighted or FLAIR images to evaluate cerebrovascular pathology and white matter hyperintensities, and axial gradient echo T<sub>2</sub>-weighted images to detect microbleeds in vascular dementia.


Rapid evolution of MTL atrophy on successive MR examinations is a highly evocative feature of AD (Fig. 3). For more than a decade, researchers have documented MTL atrophy using MRI morphometry. In the early 1990s, MRI replaced CT as a preferred method for the detection and quantification of brain atrophy. Initially, MRI measurements were performed in the MTL, including the hippocampus and the amygdala, then the entorhinal cortex using region-of-interest (ROI) methods. A few early studies used linear measurements [5]. This method is easy and quick, but provides an indirect estimate of MTL volumes. When rigorously applied, it can give a reliable estimate of hippocampal atrophy [6]. MR volumetry is theoretically the best method to assess atrophy when the intra- and inter-rater reliability has been thoroughly evaluated. MR volumetry is time consuming, requiring the manual outline of the ROI on each slice (Fig. 1); visual assessment of MTL atrophy was proposed as an alternative to volumetry [7, 8]. Visual assessment of MTL atrophy is easily applicable in clinical practice, but is subjective, highly observer dependent and does not provide a true quantitative assessment of the volumes. Therefore, most studies interested in MTL structures included MR volumetry [9]. New areas of research include the development of dedicated software for MR volumetry and computer-assisted diagnosis of AD.

The volume of the hippocampus is significantly reduced in Alzheimer's disease compared to control subjects, by 30–40% in moderate [5, 9–11], 20–30% in mild (MMSE greater than 20) [12–18] and about 10–12% in early Alzheimer's disease (MMSE about 27) [19]. Measurements of the entorhinal cortex are more difficult because



**Fig. 1** MR anatomy of the medial temporal lobe. Oblique coronal slices perpendicular to the long axis of the hippocampus passing at the level of the head (*left*), the body (*middle*) and the tail of the

hippocampus (*right*). The contours of the hippocampus (*solid line*) and the amygdala (*dashed line*) are shown. The *arrow* points to the entorhinal cortex



**Fig. 2** Histological MR comparison of the hippocampus. **a** Coronal MR slice passing at the level of the body of the hippocampus. The hippocampal formation is outlined using *white dots*. **b** Corresponding histological slice (courtesy of Michel Baulac): (1) Ammon's horn,


(2) dentate gyrus, (3) subiculum, (4) fimbria, (5) parahippocampal gyrus, (6) transverse fissure, (7) choroids plexus, (8) temporal horn, (9) lateral geniculate body, (10) collateral sulcus, and (11) hippocampal sulcus

the anatomical boundaries of the structure are not as easy to determine. According to models of disease progression in AD [2], atrophy of the entorhinal cortex should be more severe than hippocampal atrophy in early stages of Alzheimer's disease. This was confirmed in some studies, which reported higher atrophy of the entorhinal cortex than the hippocampus in early AD [19]. Some studies indicated that entorhinal cortex measurements have a higher sensitivity to differentiate AD patients from control subjects [19–23], whereas other studies did not [24–26]. Overall, the sensitivity and specificity of hippocampal and entorhinal cortex measurements to distinguish patients from age-matched control subjects are in the range of 80–100% and 80–95%, respectively.

Aside from ROI approaches, computerized morphometry methods have gained great interest because they allow whole brain analysis and do not depend on the examiner's abilities. Voxel-based morphometry (VBM) has become a tool of predilection to study the anatomical differences between two populations [27]. VBM is an automatic whole brain method that compares signal intensity voxel by voxel. VBM has been widely used to compare AD populations to healthy elderly subjects [28–31]. Using VBM, several studies have reported differences in the grey

matter between controls and mild to moderate AD in the posterior and temporal areas (Fig. 4) [28–35]. Patients with early onset (age  $\leq 65$  years) showed greater neocortical atrophy at the temporoparietal junction than patients with late onset [36]. VBM may be more accurate than hippocampal volumetry to detect medial temporal lobe atrophy in Alzheimer's disease, but this needs to be confirmed in larger series [37, 38].


Several imaging methods have been developed to follow disease progression and understand the effects of disease in AD. Earlier studies used serial manual measurements of the hippocampus [39]. More recent studies have used specific software that normalizes brain structures into a common reference space, allowing statistical comparisons and group difference mapping [40, 41]. Some software also match cortical features across individuals, by warping one brain surface onto another [42, 43]. Iterative principal component analysis by considering voxel intensity pairs from co-registered MR images has also been successfully used to map sequential volume changes [44]. Atlas-based automatic segmentation of the hippocampus has been reported [45]. Overall, the annual rate of hippocampal atrophy in AD was calculated at 2.2% [46] to 3.6–5.9% [20, 39, 47] compared to 0.24% [46] to 1.4–1.73% [39, 47] in age-



**Fig. 3** Progression of hippocampal atrophy in Alzheimer's disease. From *left to right*, the yearly examination of the same subject is shown. Increasing hippocampal and parahippocampal atrophy is

observed over time as well as ventricular enlargement and widening of cortical sulci. The *arrows* point to the hippocampus





**Fig. 4** VBM comparison of patients with Alzheimer's disease and age-matched control subjects. Maps of significantly lower grey matter density in 15 patients with Alzheimer's disease ( $\text{MMSE} = 20.9 \pm 3.7$ ) compared to 15 age-matched control subjects. Maps are superimposed on a three-dimensional surface rendering (*upper row*) and on axial

slices of a brain template (*lower row*). Clusters were significant at  $P < 0.05$  corrected for multiple comparisons. Grey matter density was decreased in the medial temporal lobe structures, the lateral temporal and inferior parietal areas, the posterior cingulum, and the cerebellum

matched control subjects. The annual rate of entorhinal cortex atrophy in AD was about 7 to 8% [20, 47, 48]. Dynamic maps of disease progression in AD show a spreading wave of grey matter loss extending from limbic to frontal cortices, whereas the sensorimotor cortex is relatively spared [43], in agreement with postmortem observations [1, 2].

#### *Basal forebrain atrophy in Alzheimer's disease*

Degeneration of cholinergic neurons in the basal forebrain, including the nucleus basalis of Meynert, is another feature of AD [49, 50]. The cholinergic deficit partially contributes to the cognitive deficits observed in AD and is the basis for the use of cholinesterase inhibitors. Atrophy of the substantia innominata was observed in AD patients and was more pronounced in patients who responded to cholinesterase inhibitors [51, 52].

#### *Brain atrophy in patients with risk factors of Alzheimer's disease*

In line with histological findings, MR studies have shown that MTL atrophy was present before subjects developed

the first symptoms. Longitudinal studies of subjects with a family history of Alzheimer's disease [46, 53–55], carriers of apolipoprotein E4 (apo E4) [56, 57] and normal elderly subjects [47, 58] have detected MTL atrophy in subjects who developed Alzheimer's disease. In subjects who developed dementia during the study, the hippocampal volume at entry was smaller than in subjects who did not develop dementia. The annual rate of atrophy was either similar for the two groups [58] or higher in subjects with familial risk factors who became demented (1.5% per year in demented and 0.2% in non-demented subjects) [46, 53, 54] or in carriers of apo E4 (2.86% in apo E4+ and 0.85% in apo E4– subjects) [59]. Similarly, the rate of temporal atrophy was higher in normal subjects who converted to dementia during the follow-up [47, 58]. The annual rate of hippocampal atrophy was increased in presymptomatic subjects and to a greater extent when the first cognitive symptoms have occurred [46].

#### *Medial temporal lobe atrophy in mild cognitive impairment*


MCI defines a state of cognition where the deficiency is greater than expected for subjects of the same age and socio-cultural background, but not severe enough for

dementia. The concept of MCI is not grounded on a causal mechanism and encompasses a heterogeneous population of patients, including those with other degenerative or vascular diseases, and with attention deficits such as those observed in metabolic or iatrogenic disorders, depression or anxiety. Although all patients with MCI do not develop Alzheimer's disease, they present an increased risk of progression to dementia [60–63]. A fundamental question is thus to determine whether the diagnosis of Alzheimer's disease can be made at this stage. This is particularly important in the selection of patients that can benefit from future treatment.

Given the pattern of progression of histological lesions in Alzheimer's disease, the medial temporal lobe, and particularly the hippocampus, is the structure where early lesions are expected in MCI patients who will develop Alzheimer's disease. Early studies have shown abnormalities in hippocampal volumes in patients with mild

cognitive impairment [64, 65]. A 10 to 20% reduction in hippocampal volume was reported in MCI patients [21, 26, 65–72]. Using VBM, reduced grey matter density was observed in the medial temporal lobe and the posterior association cortex of MCI patients [73–75]. Moreover, greater grey matter loss was observed in the medial, inferior and middle temporal areas, posterior cingulate, and precuneus in MCI patients who developed Alzheimer's disease relative to non-converters [74] (Fig. 5). In AD patients, the hippocampal region's atrophy observed in MCI patients extends to the neighboring temporal association neocortex.

Longitudinal MRI studies using an individual's first scan as the reference have shown that MTL volumes can predict which MCI subject will develop AD. Atrophy of the hippocampus [67] and the entorhinal cortex [68, 76, 77], as well as an increased rate of atrophy [47, 78, 79], measured using MRI, were associated with an increased risk of



**Fig. 5** VBM differences in MCI patients who developed Alzheimer's disease compared to non-converters. *Top*: surface rendering of the maps of percent annual gray-matter loss in converters and in non-converters (from left, right, and medial viewpoints). *Bottom*: significant clusters of

greater grey-matter loss in converters relative to non-converters as projected onto the 'glass-brain' representation of SPM (*left*) and onto coronal sections of a whole-brain template (*right*) (courtesy of Gaël Chetelat, Ph.D.)

Alzheimer's disease. Similar results were obtained using visual scaling [80]. The annual rate of hippocampal atrophy was higher in MCI patients whose cognitive impairment increases over time (3.69%) compared to stable patients (2.55%) and control subjects (1.73%) [78]. Measurements of the volume of the entorhinal cortex, the superior temporal sulcus and the anterior cingulate cortex differentiate control subjects from patients with AD, as well as 93% of MCI patients who will develop AD, and 85% of MCI patients who will not develop AD [68]. However, the goal is not to define structural predictive factors of Alzheimer's disease, but to diagnose patients with a pre-demented stage of Alzheimer's disease among MCI patients. Recently, a longitudinal study of 205 aged subjects and MCI patients followed over 5.5 to 8.5 years has shown that MR volumetric measurements of the hippocampus were able to help predict the conversion of MCI patients to AD [61]. In this study, the best predictive model was obtained when genetic susceptibility factors, specific memory tests, and volumetric measurements of the hippocampus were combined. A large multi-site longitudinal MRI and FDG-PET study, the Alzheimer's Disease Neuroimaging Initiative (ADNI <http://www.ADNI-info.org>), has recently started to develop improved imaging methods to measure longitudinal changes of the brain in normal aging, during the transition to early Alzheimer's disease, and in Alzheimer's disease patients [81].

It should be noted that MTL structures decrease in size with normal aging, mainly the hippocampal formations [39, 47, 53, 58, 82, 83], as well as the temporal cortex [83] and temporal gray matter [82]. However, atrophy in MCI patients is more marked than in normal aging.

#### *Medial temporal lobe atrophy in patients with other degenerative diseases*

MTL atrophy has been reported in other dementias, such as Parkinson's disease (with or without dementia) [84, 85], dementia with Lewy bodies [86, 87] and vascular dementia [88]. MTL atrophy is more severe in Alzheimer's disease than in these diseases, however. The rate of progression of MTL atrophy over time is also higher in patients with Lewy body dementia and vascular dementia than in age-matched controls [89]. Therefore, MTL atrophy is apparently not specific for Alzheimer's disease. Other neurodegenerative diseases are associated with atrophy in other brain regions that depends on the type of neurodegenerative process. This suggests that whole brain analysis of the profiles of grey matter loss has better potential to differentiate neurodegenerative dementias [34, 90, 91].

#### *Correlations between structural lesions and neuro-psychological deficits*

Early CT studies have reported correlations between performances in global measurements of cognitive functions, such as the MMSE [92] or the Mattis Dementia Rating Scale, and global measurements of atrophy such as ventricular enlargement [3].

A deficit in recent memory is generally predominant at the early stage of Alzheimer's disease [93, 94]. Classically, this deficit relates to lesions of medial temporal lobe structures that are affected early in AD [1, 2, 95]. Several studies have confirmed that volumes of MTL structures, and particularly the hippocampus, were correlated with memory performances [20, 84, 96–101]. Correlations were also reported with the entorhinal cortex [20, 47].

The correlation between MTL structures was specific to memory performances [96, 97, 102], in line with the specific role of the MTL structures in early memory impairment observed in the disease. Overall, the correlation was higher for delayed than for immediate memory tasks [96, 97]. Performances at verbal memory tests correlated best with left MTL volumes, whereas performances at non-verbal memory tests correlated best with right medial temporal lobe volumes [98–100]. This specificity indicates that hippocampal atrophy does not correlate with global indices of cognitive impairment, but is specifically associated with memory function deficits. Memory scores were less correlated with volumes of the para-hippocampal gyrus or the temporal lobe [99, 100]. As already mentioned, these structures are more difficult to measure and therefore correlations may be more difficult to determine. Indeed, a recent study reported that the Delayed List Verbal Recall test correlated significantly with atrophy rates of the entorhinal cortex [20].

Other studies have reported correlations between non-memory tests and various brain regions. Anterior and inferior structures of the left temporal lobe were related to semantic memory deficits, as suggested by the difference in patterns of atrophy between semantic dementia (predominance in the left infero-anterior temporal regions) and Alzheimer's disease (symmetric temporal atrophy without antero-posterior gradient) [103]. Impaired naming difficulties correlated with grey matter atrophy in the left anterior-lateral temporal and anterior cingulate regions [33]. A selective correlation between performances at verbal fluency tests and the frontal lobe volumes was observed, in line with the role of the frontal lobe in the generation of verbal and non-verbal material, whereas this correlation was not observed with MTL structures [104].

In contrast to ROI studies, VBM or other whole brain techniques allow the correlation of neuropsychological variables and grey matter loss in the entire brain. Indeed, significant positive correlations were observed among the MMSE scores and temporal, posterior cingulate gyri, and precuneus regions in AD patients [31, 90], as well as



hippocampal atrophy and enlargement of the temporal horn of the lateral ventricle [105].

#### *White matter pathology in Alzheimer's disease*

Vascular lesions can occur in AD. Vascular risk factors such as hypertension, hypercholesterolemia, and apolipoprotein E4 allele may predispose to AD [106, 107]. Changes in the endothelium and amyloid angiopathy have been described in AD [108]. Over 60% of patients with Alzheimer's disease present white matter lesions [109]. The clinical differentiation of vascular dementia from Alzheimer's disease with cerebrovascular disease can therefore be difficult [110]. Vascular lesions are milder in AD as compared with vascular dementias, however. The

recognition of white matter and other vascular lesions in AD is important, since vascular pathology can lower the threshold for dementia or increase its severity [111–113]. Vascular lesions may represent a target for treatment.


#### *Other MRI techniques*

Beside MR morphometry, several more recent imaging techniques have shown a potential to detect early abnormalities in AD that may be useful in the clinical diagnosis of dementias. These techniques include perfusion, diffusion, spectroscopy and microscopy.

##### – Perfusion MRI

Perfusion MRI techniques can assess cerebral perfusion (cerebral blood volume and cerebral blood flow).

**Fig. 6** Voxel based-DTI comparison of patients with Alzheimer's disease and age-matched control subjects. Maps of significantly higher FA in ten patients with Alzheimer's disease ( $MMSE = 24.0 \pm 3.5$ ) compared to ten age-matched control subjects ( $MMSE = 28.0 \pm 1.6$ ). Maps are superimposed on a three dimensional surface rendering of a template brain (upper left: anterior view; upper right: posterior view, middle: lateral views, lower left: inferior view, lower right: superior view). Clusters were significant at  $P < 0.05$  corrected for multiple comparisons. FA was decreased in frontal, parietal and temporal structures and the cerebellum



They therefore have the potential to depict functional deficiencies in a way similar to HMPAO SPECT [114–116]. Techniques using radiotracers have consistently shown a reduction in cerebral blood flow in patients with AD, in the temporo-parietal association areas, the posterior cingulate cortex, and, to a lesser extent, in frontal association areas with relative sparing of the primary motor and sensory areas [114–118].


Cerebral perfusion can be assessed using contrast-enhanced techniques that require the injection of a paramagnetic contrast agent. In Alzheimer's disease, reduced perfusion was reported in parietal areas using contrast-enhanced perfusion imaging [119]. Arterial spin labeling (ASL) techniques use electromagnetic labeling of water in the blood to acquire images sensitive to flow without any external contrast agents [120]. Initially limited to the acquisition of a few slices only, recent technical advances have allowed the acquisition of multiple slices [121] and absolute quantification [122]. In Alzheimer's disease, cerebral perfusion assessed using ASL techniques was altered in the parietal, temporal and cingulate regions, similar to those seen in HMPAO SPECT [119, 123–125]. Perfusion in posterior parietal and cingulate areas correlated with neuropsychological performances such as memory scores [123] or the MMSE [124]. By the assessment of cerebral blood flow in AD with spin labeling techniques, MRI has the potential to assist in the diagnosis of neurodegenerative dementias.

#### – Diffusion imaging

Diffusion-weighted imaging (DWI) is sensitive to the random motion of water molecules in the brain. Measures of water diffusivity from DWI provide an estimate of the microstructural integrity of the brain parenchyma. Diffusion tensor imaging (DTI) can also detect the directionality of molecular diffusion in the white matter and allows the demonstration of fiber tracts in vivo in humans [126–128]. The directionality dependence of diffusion is called anisotropy and is measured using indices such as fractional anisotropy (FA) [129].

Increased diffusivity has been reported in the temporal lobe [130, 131] and the posterior white matter [132] of patients with Alzheimer's disease, as well as MCI subjects [133, 134]. Increased diffusivity can reflect a change in the ultrastructural organization of brain tissue or represent an index of neuronal atrophy. In contrast, brain anisotropy was decreased in several white matter regions including the corpus callosum, the cingulum, the superior longitudinal fasciculus [135–138] or the area of the perforant pathway [139], reflecting a change in the white matter bundles connecting the brain areas affected in AD (Fig. 6). In MCI subjects, higher baseline hippocampal diffusivity was associated with a greater risk of progression

to AD [140]. Similarly to voxel-based morphometry, voxel-based comparison of anisotropy maps showed reduced FA in posterior brain regions [141]. DTI abnormalities showed some correlation with measures of cognitive performances. Left hippocampal volumes were associated with poor verbal memory performance, particularly when associated with high diffusivity values [133]. Performance in the delayed verbal recall test correlated significantly with posterior cingulate bundle anisotropy and diffusivity [136]. Although the neurobiological abnormalities that result in abnormal DTI are not completely understood, DTI is a useful tool for the in vivo detection of microstructural abnormalities in AD. Improvements in diffusion techniques should further enhance the information that can be derived from DTI.



**Fig. 7**  $^1\text{H}$  MR spectra comparison of a patient with Alzheimer's disease and age-matched subject. MR spectra obtained using single voxel spectroscopy with short echo time (30 ms) from the posterior cingulate in an age-matched control subject (top) and in a patient with Alzheimer's disease (bottom). NAA/Cr ratio is lower in the patient with AD than in the control subject, and mIns/Cr ratio is higher in the AD patient than in the control subject (courtesy of Kejal Kantarci, M.D.)




In Alzheimer's disease, white matter abnormalities had already been reported in the corpus callosum using measurements on T<sub>1</sub>-weighted images [142, 143]. Atrophy predominated in the posterior part of the structure, whereas the anterior part was atrophic in fronto-temporal dementia.

– Magnetic resonance spectroscopy

Proton magnetic resonance spectroscopy (<sup>1</sup>H MRS) provides information about the levels of metabolites in the brain. Metabolite levels are sensitive to different in vivo pathological processes at the molecular or cellular level; consequently, they can provide both qualitative and quantitative information about the degenerative process. Different metabolites provide information about different processes; for example, *N*-acetylaspartate (NAA) is believed to be a marker for neuronal number and health; choline-containing compounds (Cho) are markers of demyelination and cell proliferation; *myo*-inositol (mIns) is thought to be a marker for osmotic stress or astrogliosis; lactate is frequently associated with the presence of pathology.

In most AD studies, the metabolite levels are reported as their ratios over the sum of creatine and phosphocreatine (Cr). Cr is used as a concentration reference, since its concentration is regarded as being relatively stable. Two methods are generally used to obtain information about the levels of metabolites in the brain: chemical shift imaging (CSI is also referred to as spectroscopic imaging) and single voxel spectroscopy. CSI can provide information about the spatial distributions of metabolite levels from multiple slices simultaneously (3D-CSI), with the drawback of providing information mostly about NAA, Cho and Cr, especially when run at a long echo time, and with problems with implementation of efficient water suppression. There is usually not enough signal to quantify the concentration of mIns or other *J*-coupled metabolites. The metabolite levels cannot be obtained from the entire brain due to the problems with magnetic field homogeneity when optimizing large volumes of interest and with lipid artifact signals when the borders of the VOI are too close to the skull. The

**Fig. 8** MR microscopy Four-way correlation in a 24-month-old AD mouse. In vivo MR microscopy (a, b), ex vivo MR microscopy (c, d), thioflavin-S-stained (e, f), and iron-stained (g, h) images have been precisely spatially registered over a circumscribed area of the cortex, indicated by the box. The boxes in the right column represent 3× magnified portions of the adjacent parent image in the left column. The numbered arrows indicate individual plaques visualized in each of the four different image types (reprinted with permission from [174], copyright 2005 by the Society for Neuroscience)



single voxel spectroscopy run at shorter echo times allows a good quantification of *J*-coupled metabolites such as mIns, but the information is coming from one spatial location in the brain. In AD, both spectroscopy methods have been used.

The proton spectrum is drastically different in AD patients from normal subjects with reduced NAA/Cr and with elevated mIns/Cr [144] (Fig. 7). The ratios of NAA/Cr, NAA/Cho and NAA/mIns are consistently decreased [145–149]. A larger decrease in the NAA level has been reported in the temporal lobe tissue in patients with AD compared with controls than in the rest of the brain, which showed a seemingly uniform decrease [150]. The decrease in NAA ratios accelerates with the disease progression [151], and changes in NAA concentration are detectable early in the progression of the disease [152]. Cognitive dysfunction of demented subjects was significantly correlated with a reduction of NAA in some studies [153], but not in others [149]. Note that in healthy elderly people NAA does not change significantly with further aging [154] and that those individuals with a higher white matter Cho/Cr ratio were shown to have a higher risk of developing dementia or AD within 4 years [155]. The raised mIns/Cr levels allow for the distinction between AD patients and cognitively normal elderly subjects. The mIns levels were raised uniformly throughout the brain [150]. Elevated levels of mIns were also observed in frontotemporal lobar degeneration (FTLB) [156] and mild cognitive impairment (MCI) patients [157]. NAA/mIns ratios distinguished clinically diagnosed patients with AD from cognitively normal elderly with a sensitivity of 83% and specificity of 98% in one cohort [146], and a sensitivity of 82% and specificity of 80% in another [158]. There are conflicting reports about Cho levels in AD. Some studies identified elevated Cho levels [156], and others did not [150, 159]. Decreased levels of glutamate + glutamine (Glx) have been reported using single voxel

spectroscopy [160, 161]. In spite of the fact that lactate was found in a significant portion of elderly persons [162] and was observed in the CSF in Alzheimer's disease [163], no studies have reported increased brain tissue lactate levels in Alzheimer's disease.

#### MR microscopy

Amyloid plaques in patients with AD range in size from 2 to 200  $\mu\text{m}$ . The direct imaging of plaques requires very high spatial resolution. Although scintigraphic markers (such as the "Pittsburgh-B" compound) have been developed that can visualize the plaque burden with positron emission tomography (PET) in living Alzheimer's patients [164–166], individual plaques are beyond the resolution of PET. Very high field magnets ( $\geq 7$  T) allow very high spatial resolution and therefore MR microscopy. Individual plaques were first evidenced at 7 T using a gradient echo  $T_2$ -weighted sequence and a spatial resolution of 40  $\mu\text{m}$  cubic isovoxel size [167]. Plaques appeared as dark round spots, probably because they contain metals such as iron. This finding was not replicated in another study that reported the dark spots to be vascular structures [168]. Later studies in ex-vivo transgenic mouse brain specimens confirmed that plaques could be imaged at 7 T [169–172] (Fig. 8). These methods are not yet applicable to human studies as the imaging time was greater than 10 h. Recently, the imaging time was reduced to 1 h–1 h 30 m using optimized sequences at 9.4 T, and images were obtained for the first time in a transgenic mouse in vivo [173, 174]. Scanning time is thus approaching a reasonable time for in vivo human studies. Additionally, the amyloid plaques have been detected in mice with  $^{19}\text{F}$  MRI using a  $^{19}\text{F}$ -containing amyloidophilic probe [175]. The images were obtained in 2 h with virtually zero background noise since  $^{19}\text{F}$  is absent in living beings. The used resolution did not allow the detection of the individual plaques in vivo.

## References

1. Delacourte A, David JP, Sergeant N, Buee L, Wattez A, Vermersch P, Ghazali F, Fallet-Bianco C, Pasquier F, Lebert F, Petit H, Di Menza C (1999) The biochemical pathway of neurofibrillary degeneration in aging and Alzheimer's disease. *Neurology* 52 (6):1158–1165
2. Braak H, Braak E (1995) Staging of Alzheimer's disease-related neurofibrillary changes. *Neurobiol Aging* 16 (3):271–278, discussion 8–84
3. DeCarli C, Kaye JA, Horwitz B, Rapoport SI (1990) Critical analysis of the use of computer-assisted transverse axial tomography to study human brain in aging and dementia of the Alzheimer type. *Neurology* 40(6):872–883
4. de Leon MJ, George AE, Stylopoulos LA, Smith G, Miller DC (1989) Early marker for Alzheimer's disease: the atrophic hippocampus. *Lancet* 2 (8664):672–673

5. Seab JP, Jagust WJ, Wong ST, Roos MS, Reed BR, Budinger TF (1988) Quantitative NMR measurements of hippocampal atrophy in Alzheimer's disease. *Magn Reson Med* 8(2):200–208
6. Meguro K, LeMestric C, Landeau B, Desgranges B, Eustache F, Baron JC (2001) Relations between hypometabolism in the posterior association neocortex and hippocampal atrophy in Alzheimer's disease: a PET/MRI correlative study. *J Neurol Neurosurg Psychiatry* 71(3):315–321
7. Scheltens P, Fox N, Barkhof F, De Carli C (2002) Structural magnetic resonance imaging in the practical assessment of dementia: beyond exclusion. *Lancet Neurol* 1(1):13–21
8. Wahlund LO, Julin P, Johansson SE, Scheltens P (2000) Visual rating and volumetry of the medial temporal lobe on magnetic resonance imaging in dementia: a comparative study. *J Neurol Neurosurg Psychiatry* 69(5):630–635
9. Jack CR Jr, Petersen RC, O'Brien PC, Tangalos EG (1992) MR-based hippocampal volumetry in the diagnosis of Alzheimer's disease. *Neurology* 42(1):183–188
10. Hampel H, Teipel SJ, Bayer W, Alexander GE, Schwarz R, Schapiro MB, Rapoport SI, Moller HJ (2002) Age transformation of combined hippocampus and amygdala volume improves diagnostic accuracy in Alzheimer's disease. *J Neurol Sci* 194(1):15–19
11. Kesslak JP, Nalcioglu O, Cotman CW (1991) Quantification of magnetic resonance scans for hippocampal and parahippocampal atrophy in Alzheimer's disease. *Neurology* 41(1):51–54
12. Killiany RJ, Moss MB, Albert MS, Sandor T, Tieman J, Jolesz F (1993) Temporal lobe regions on magnetic resonance imaging identify patients with early Alzheimer's disease. *Arch Neurol* 50(9):949–954
13. Cuenod CA, Denys A, Michot JL, Jehenson P, Forette F, Kaplan D, Syrota A, Boller F (1993) Amygdala atrophy in Alzheimer's disease. An in vivo magnetic resonance imaging study. *Arch Neurol* 50(9):941–945
14. Lehericy S, Baulac M, Chiras J, Pierot L, Martin N, Pillon B, Deweer B, Dubois B, Marsault C (1994) Amygdalohippocampal MR volume measurements in the early stages of Alzheimer disease. *Am J Neuroradiol* 15(5):929–937
15. Pitkanen A, Laakso M, Kalviainen R, Partanen K, Vainio P, Lehtovirta M, Riekkinen P, Soininen H (1996) Severity of hippocampal atrophy correlates with the prolongation of MRI T2 relaxation time in temporal lobe epilepsy but not in Alzheimer's disease. *Neurology* 46(6):1724–1730
16. Frisoni GB, Beltramello A, Weiss C, Geroldi C, Bianchetti A, Trabucchi M (1996) Linear measures of atrophy in mild Alzheimer disease. *Am J Neuroradiol* 17(5):913–923
17. Jack CR Jr, Petersen RC, Xu YC, Waring SC, O'Brien PC, Tangalos EG, Smith GE, Ivnik RJ, Kokmen E (1997) Medial temporal atrophy on MRI in normal aging and very mild Alzheimer's disease. *Neurology* 49(3):786–794
18. Laakso MP, Lehtovirta M, Partanen K, Riekkinen PJ, Soininen H (2000) Hippocampus in Alzheimer's disease: a 3-year follow-up MRI study. *Biol Psychiatry* 47(6):557–561
19. Bobinski M, de Leon MJ, Convit A, De Santi S, Wegiel J, Tarshish CY, Saint Louis LA, Wisniewski HM (1999) MRI of entorhinal cortex in mild Alzheimer's disease. *Lancet* 353(9146):38–40
20. Du AT, Schuff N, Kramer JH, Ganzer S, Zhu XP, Jagust WJ, Miller BL, Reed BR, Mungas D, Yaffe K, Chui HC, Weiner MW (2004) Higher atrophy rate of entorhinal cortex than hippocampus in AD. *Neurology* 62(3):422–427
21. Du AT, Schuff N, Amend D, Laakso MP, Hsu YY, Jagust WJ, Yaffe K, Kramer JH, Reed B, Norman D, Chui HC, Weiner MW (2001) Magnetic resonance imaging of the entorhinal cortex and hippocampus in mild cognitive impairment and Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 71(4):441–447
22. Killiany RJ, Hyman BT, Gomez-Isla T, Moss MB, Kikinis R, Jolesz F, Tanzi R, Jones K, Albert MS (2002) MRI measures of entorhinal cortex vs hippocampus in preclinical AD. *Neurology* 58(8):1188–1196
23. Juottonen K, Laakso MP, Insausti R, Lehtovirta M, Pitkanen A, Partanen K, Soininen H (1998) Volumes of the entorhinal and perirhinal cortices in Alzheimer's disease. *Neurobiol Aging* 19(1):15–22
24. Frisoni GB, Laakso MP, Beltramello A, Geroldi C, Bianchetti A, Soininen H, Trabucchi M (1999) Hippocampal and entorhinal cortex atrophy in frontotemporal dementia and Alzheimer's disease. *Neurology* 52(1):91–100
25. Juottonen K, Laakso MP, Partanen K, Soininen H (1999) Comparative MR analysis of the entorhinal cortex and hippocampus in diagnosing Alzheimer disease. *Am J Neuroradiol* 20(1):139–144
26. Xu Y, Jack CR Jr, O'Brien PC, Kokmen E, Smith GE, Ivnik RJ, Boeve BF, Tangalos RG, Petersen RC (2000) Usefulness of MRI measures of entorhinal cortex versus hippocampus in AD. *Neurology* 54(9):1760–1767
27. Ashburner J, Friston KJ (2000) Voxel-based morphometry—the methods. *Neuroimage* 11(6 Pt 1):805–821
28. Baron JC, Chetelat G, Desgranges B, Percey G, Landeau B, de la Sayette V, Eustache F (2001) In vivo mapping of gray matter loss with voxel-based morphometry in mild Alzheimer's disease. *Neuroimage* 14(2):298–309
29. Karas GB, Burton EJ, Rombouts SA, van Schijndel RA, O'Brien JT, Scheltens P, McKeith IG, Williams D, Ballard C, Barkhof F (2003) A comprehensive study of gray matter loss in patients with Alzheimer's disease using optimized voxel-based morphometry. *Neuroimage* 18(4):895–907
30. Garrido GE, Furuie SS, Buchpiguel CA, Bottino CM, Almeida OP, Cid CG, Camargo CH, Castro CC, Glabus MF, Busatto GF (2002) Relation between medial temporal atrophy and functional brain activity during memory processing in Alzheimer's disease: a combined MRI and SPECT study. *J Neurol Neurosurg Psychiatry* 73(5):508–516
31. Frisoni GB, Testa C, Zorzan A, Sabatoli F, Beltramello A, Soininen H, Laakso MP (2002) Detection of grey matter loss in mild Alzheimer's disease with voxel based morphometry. *J Neurol Neurosurg Psychiatry* 73(6):657–664
32. Hirata Y, Matsuda H, Nemoto K, Ohnishi T, Hirao K, Yamashita F, Asada T, Iwabuchi S, Samejima H (2005) Voxel-based morphometry to discriminate early Alzheimer's disease from controls. *Neurosci Lett* 382(3):269–274



33. Grossman M, McMillan C, Moore P, Ding L, Glosser G, Work M, Gee J (2004) What's in a name: voxel-based morphometric analyses of MRI and naming difficulty in Alzheimer's disease, frontotemporal dementia and corticobasal degeneration. *Brain* 127(Pt 3):628–649
34. Burton EJ, McKeith IG, Burn DJ, Williams ED, JT OB (2004) Cerebral atrophy in Parkinson's disease with and without dementia: a comparison with Alzheimer's disease, dementia with Lewy bodies and controls. *Brain* 127 (Pt 4):791–800
35. Busatto GF, Garrido GE, Almeida OP, Castro CC, Camargo CH, Cid CG, Buchpiguel CA, Furuie S, Bottino CM (2003) A voxel-based morphometry study of temporal lobe gray matter reductions in Alzheimer's disease. *Neurobiol Aging* 24(2):221–231
36. Frisoni GB, Testa C, Sabatoli F, Beltramello A, Soininen H, Laakso MP (2005) Structural correlates of early and late onset Alzheimer's disease: voxel based morphometric study. *J Neurol Neurosurg Psychiatry* 76(1):112–114
37. Good CD, Scahill RI, Fox NC, Ashburner J, Friston KJ, Chan D, Crum WR, Rossor MN, Frackowiak RS (2002) Automatic differentiation of anatomical patterns in the human brain: validation with studies of degenerative dementias. *Neuroimage* 17(1):29–46
38. Testa C, Laakso MP, Sabatoli F, Rossi R, Beltramello A, Soininen H, Frisoni GB (2004) A comparison between the accuracy of voxel-based morphometry and hippocampal volumetry in Alzheimer's disease. *J Magn Reson Imaging* 19(3):274–282
39. Jack CR Jr, Petersen RC, Xu Y, O'Brien PC, Smith GE, Ivnik RJ, Tangalos EG, Kokmen E (1998) Rate of medial temporal lobe atrophy in typical aging and Alzheimer's disease. *Neurology* 51(4):993–999
40. Good CD, Johnsrude IS, Ashburner J, Henson RN, Friston KJ, Frackowiak RS (2001) A voxel-based morphometric study of ageing in 465 normal adult human brains. *Neuroimage* 14(1 Pt 1):21–36
41. Thompson PM, Hayashi KM, Sowell ER, Gogtay N, Giedd JN, Rapoport JL, de Zubicaray GI, Janke AL, Rose SE, Semple J, Doddrell DM, Wang Y, van Erp TG, Cannon TD, Toga AW (2004) Mapping cortical change in Alzheimer's disease, brain development, and schizophrenia. *Neuroimage* 23 (Suppl 1):S2–S18
42. Thompson PM, Mega MS, Woods RP, Zoumalan CI, Lindshield CJ, Blanton RE, Moussai J, Holmes CJ, Cummings JL, Toga AW (2001) Cortical change in Alzheimer's disease detected with a disease-specific population-based brain atlas. *Cereb Cortex* 11(1):1–16
43. Thompson PM, Hayashi KM, de Zubicaray G, Janke AL, Rose SE, Semple J, Herman D, Hong MS, Dittmer SS, Doddrell DM, Toga AW (2003) Dynamics of gray matter loss in Alzheimer's disease. *J Neurosci* 23 (3):994–1005
44. Chen K, Reiman EM, Alexander GE, Bandy D, Renaut R, Crum WR, Fox NC, Rossor MN (2004) An automated algorithm for the computation of brain volume change from sequential MRIs using an iterative principal component analysis and its evaluation for the assessment of whole-brain atrophy rates in patients with probable Alzheimer's disease. *Neuroimage* 22 (1):134–143
45. Carmichael OT, Aizenstein HA, Davis SW, Becker JT, Thompson PM, Meltzer CC, Liu Y (2005) Atlas-based hippocampus segmentation in Alzheimer's disease and mild cognitive impairment. *Neuroimage* 27(4):979–990
46. Fox NC, Crum WR, Scahill RI, Stevens JM, Janssen JC, Rossor MN (2001) Imaging of onset and progression of Alzheimer's disease with voxel-compression mapping of serial magnetic resonance images. *Lancet* 358 (9277):201–205
47. Jack CR Jr, Shiung MM, Gunter JL, O'Brien PC, Weigand SD, Knopman DS, Boeve BF, Ivnik RJ, Smith GE, Cha RH, Tangalos EG, Petersen RC (2004) Comparison of different MRI brain atrophy rate measures with clinical disease progression in AD. *Neurology* 62(4):591–600
48. Du AT, Schuff N, Zhu XP, Jagust WJ, Miller BL, Reed BR, Kramer JH, Mungas D, Yaffe K, Chui HC, Weiner MW (2003) Atrophy rates of entorhinal cortex in AD and normal aging. *Neurology* 60(3):481–486
49. Whitehouse PJ, Price DL, Clark AW, Coyle JT, DeLong MR (1981) Alzheimer disease: evidence for selective loss of cholinergic neurons in the nucleus basalis. *Ann Neurol* 10 (2):122–126
50. Lehericy S, Hirsch EC, Cervera-Pierot P, Hersch LB, Bakchine S, Piette F, Duyckaerts C, Hauw JJ, Javoy-Agid F, Agid Y (1993) Heterogeneity and selectivity of the degeneration of cholinergic neurons in the basal forebrain of patients with Alzheimer's disease. *J Comp Neurol* 330(1):15–31
51. Sasaki M, Ehara S, Tamakawa Y, Takahashi S, Tohgi H, Sakai A, Mita T (1995) MR anatomy of the substantia innominata and findings in Alzheimer disease: a preliminary report. *Am J Neuroradiol* 16(10):2001–2007
52. Tanaka Y, Hanyu H, Sakurai H, Takasaki M, Abe K (2003) Atrophy of the substantia innominata on magnetic resonance imaging predicts response to donepezil treatment in Alzheimer's disease patients. *Dement Geriatr Cogn Disord* 16(3):119–125
53. Fox NC, Warrington EK, Rossor MN (1999) Serial magnetic resonance imaging of cerebral atrophy in preclinical Alzheimer's disease. *Lancet* 353 (9170):2125
54. Scahill RI, Schott JM, Stevens JM, Rossor MN, Fox NC (2002) Mapping the evolution of regional atrophy in Alzheimer's disease: unbiased analysis of fluid-registered serial MRI. *Proc Natl Acad Sci USA* 99(7):4703–4707
55. Fox NC, Scahill RI, Crum WR, Rossor MN (1999) Correlation between rates of brain atrophy and cognitive decline in AD. *Neurology* 52(8):1687–1689
56. Bigler ED, Lowry CM, Anderson CV, Johnson SC, Terry J, Steed M (2000) Dementia, quantitative neuroimaging, and apolipoprotein E genotype. *Am J Neuroradiol* 21(10):1857–1868
57. Hashimoto M, Yasuda M, Tanimukai S, Matsui M, Hirano N, Kazui H, Mori E (2001) Apolipoprotein E epsilon 4 and the pattern of regional brain atrophy in Alzheimer's disease. *Neurology* 57 (8):1461–1466
58. Kaye JA, Swihart T, Howieson D, Dame A, Moore MM, Karnos T, Camicioli R, Ball M, Oken B, Sexton G (1997) Volume loss of the hippocampus and temporal lobe in healthy elderly persons destined to develop dementia. *Neurology* 48(5):1297–1304
59. Moffat SD, Szekely CA, Zonderman AB, Kabani NJ, Resnick SM (2000) Longitudinal change in hippocampal volume as a function of apolipoprotein E genotype. *Neurology* 55(1):134–136
60. Masur DM, Sliwinski M, Lipton RB, Blau AD, Crystal HA (1994) Neuropsychological prediction of dementia and the absence of dementia in healthy elderly persons. *Neurology* 44 (8):1427–1432
61. Petersen RC, Knopman D, Boeve BF et al (2005) Role of hippocampal volumes in predicting progression to mild cognitive impairment. *Neurology* 64(6, Suppl 1):A126–A127

62. Linn RT, Wolf PA, Bachman DL, Knoefel JE, Cobb JL, Belanger AJ, Kaplan EF, D'Agostino RB (1995) The 'preclinical phase' of probable Alzheimer's disease. A 13-year prospective study of the Framingham cohort. *Arch Neurol* 52(5):485-490
63. Bowen J, Teri L, Kukull W, McCormick W, McCurry SM, Larson EB (1997) Progression to dementia in patients with isolated memory loss. *Lancet* 349(9054):763-765
64. Soininen HS, Partanen K, Pitkanen A, Vainio P, Hanninen T, Hallikainen M, Koivisto K, Riekkinen PJ Sr (1994) Volumetric MRI analysis of the amygdala and the hippocampus in subjects with age-associated memory impairment: correlation to visual and verbal memory. *Neurology* 44(9):1660-1668
65. Jack CR Jr, Petersen RC, Xu YC, O'Brien PC, Smith GE, Ivnik RJ, Boeve BF, Waring SC, Tangalos EG, Kokmen E (1999) Prediction of AD with MRI-based hippocampal volume in mild cognitive impairment. *Neurology* 52(7):1397-1403
66. De Santi S, de Leon MJ, Rusinek H, Convit A, Tarshish CY, Roche A, Tsui WH, Kandil E, Boppana M, Daisley K, Wang GJ, Schlyer D, Fowler J (2001) Hippocampal formation glucose metabolism and volume losses in MCI and AD. *Neurobiol Aging* 22(4):529-539
67. Visser PJ, Verhey FR, Hofman PA, Scheltens P, Jolles J (2002) Medial temporal lobe atrophy predicts Alzheimer's disease in patients with minor cognitive impairment. *J Neurol Neurosurg Psychiatry* 72(4):491-497
68. Killiany RJ, Gomez-Isla T, Moss M, Kikinis R, Sandor T, Jolesz F, Tanzi R, Jones K, Hyman BT, Albert MS (2000) Use of structural magnetic resonance imaging to predict who will get Alzheimer's disease. *Ann Neurol* 47(4):430-439
69. Wolf H, Grunwald M, Kruggel F, Riedel-Heller SG, Angerhofer S, Hojjatoleslami A, Hensel A, Arendt T, Gertz H (2001) Hippocampal volume discriminates between normal cognition; questionable and mild dementia in the elderly. *Neurobiol Aging* 22(2):177-186
70. Visser PJ, Scheltens P, Verhey FR, Schmand B, Launer LJ, Jolles J, Jonker C (1999) Medial temporal lobe atrophy and memory dysfunction as predictors for dementia in subjects with mild cognitive impairment. *J Neurol* 246(6):477-485
71. Convit A, de Asis J, de Leon MJ, Tarshish CY, De Santi S, Rusinek H (2000) Atrophy of the medial occipitotemporal, inferior, and middle temporal gyri in non-demented elderly predict decline to Alzheimer's disease. *Neurobiol Aging* 21(1):19-26
72. Convit A, De Leon MJ, Tarshish C, De Santi S, Tsui W, Rusinek H, George A (1997) Specific hippocampal volume reductions in individuals at risk for Alzheimer's disease. *Neurobiol Aging* 18(2):131-138
73. Pennanen C, Testa C, Laakso MP, Hallikainen M, Helkala EL, Hanninen T, Kivipelto M, Kononen M, Nissinen A, Tervo S, Vanhanen M, Vanninen R, Frisoni GB, Soininen H (2005) A voxel based morphometry study on mild cognitive impairment. *J Neurol Neurosurg Psychiatry* 76(1):11-14
74. Chetelat G, Landeau B, Eustache F, Mezege F, Viader F, de la Sayette V, Desgranges B, Baron JC (2005) Using voxel-based morphometry to map the structural changes associated with rapid conversion in MCI: a longitudinal MRI study. *Neuroimage* 27(4):934-946
75. Karas GB, Scheltens P, Rombouts SA, Visser PJ, van Schijndel RA, Fox NC, Barkhof F (2004) Global and local gray matter loss in mild cognitive impairment and Alzheimer's disease. *Neuroimage* 23(2):708-716
76. deToledo-Morrell L, Stoub TR, Bulgakova M, Wilson RS, Bennett DA, Leurgans S, Wu J, Turner DA (2004) MRI-derived entorhinal volume is a good predictor of conversion from MCI to AD. *Neurobiol Aging* 25(9):1197-1203
77. Dickerson BC, Goncharova I, Sullivan MP, Forchetti C, Wilson RS, Bennett DA, Beckett LA, deToledo-Morrell L (2001) MRI-derived entorhinal and hippocampal atrophy in incipient and very mild Alzheimer's disease. *Neurobiol Aging* 22(5):747-754
78. Jack CR Jr, Petersen RC, Xu YJ, O'Brien PC, Smith GE, Ivnik RJ, Boeve BF, Tangalos EG, Kokmen E (2000) Rates of hippocampal atrophy correlate with change in clinical status in aging and AD. *Neurology* 55(4):484-489
79. Stoub TR, Bulkakova M, Leurgans S, Bennett DA, Fleischman D, Turner DA, deToledo-Morrell L (2005) MRI predictors of risks of incident Alzheimer disease: a longitudinal study. *Neurology* 64(9):1520-1524
80. Korf ES, Wahlund LO, Visser PJ, Scheltens P (2004) Medial temporal lobe atrophy on MRI predicts dementia in patients with mild cognitive impairment. *Neurology* 63(1):94-100
81. Mueller SG, Weiner MW, Thal LJ, Petersen RC, Jack C, Jagust W, Trojanowski JQ, Toga AW, Beckett L (2005) The Alzheimer's disease neuroimaging initiative. *Neuroimaging Clin N Am* 15(4):869-877
82. Sullivan EV, Marsh L, Mathalon DH, Lim KO, Pfefferbaum A (1995) Age-related decline in MRI volumes of temporal lobe gray matter but not hippocampus. *Neurobiol Aging* 16(4):591-606
83. Insausti R, Juottonen K, Soininen H, Insausti AM, Partanen K, Vainio P, Laakso MP, Pitkanen A (1998) MR volumetric analysis of the human entorhinal, perirhinal, and temporopolar cortices. *Am J Neuroradiol* 19(4):659-671
84. Laakso MP, Partanen K, Riekkinen P, Lehtovirta M, Helkala EL, Hallikainen M, Hanninen T, Vainio P, Soininen H (1996) Hippocampal volumes in Alzheimer's disease, Parkinson's disease with and without dementia, and in vascular dementia: An MRI study. *Neurology* 46(3):678-681
85. Tam CW, Burton EJ, McKeith IG, Burn DJ, O'Brien JT (2005) Temporal lobe atrophy on MRI in Parkinson disease with dementia: a comparison with Alzheimer disease and dementia with Lewy bodies. *Neurology* 64(5):861-865
86. Hashimoto M, Kitagaki H, Imamura T, Hirono N, Shimomura T, Kazui H, Tanimukai S, Hanihara T, Mori E (1998) Medial temporal and whole-brain atrophy in dementia with Lewy bodies: a volumetric MRI study. *Neurology* 51(2):357-362
87. Barber R, Gholkar A, Scheltens P, Ballard C, McKeith IG, O'Brien JT (1999) Medial temporal lobe atrophy on MRI in dementia with Lewy bodies. *Neurology* 52(6):1153-1158
88. Barber R, Ballard C, McKeith IG, Gholkar A, O'Brien JT (2000) MRI volumetric study of dementia with Lewy bodies: a comparison with AD and vascular dementia. *Neurology* 54(6):1304-1309
89. O'Brien JT, Paling S, Barber R, Williams ED, Ballard C, McKeith IG, Gholkar A, Crum WR, Rossor MN, Fox NC (2001) Progressive brain atrophy on serial MRI in dementia with Lewy bodies, AD, and vascular dementia. *Neurology* 56(10):1386-1388

90. Ballmaier M, O'Brien JT, Burton EJ, Thompson PM, Rex DE, Narr KL, McKeith IG, DeLuca H, Toga AW (2004) Comparing gray matter loss profiles between dementia with Lewy bodies and Alzheimer's disease using cortical pattern matching: diagnosis and gender effects. *Neuroimage* 23(1):325–335
91. Burton EJ, Karas G, Paling SM, Barber R, Williams ED, Ballard CG, McKeith IG, Scheltens P, Barkhof F, O'Brien JT (2002) Patterns of cerebral atrophy in dementia with Lewy bodies using voxel-based morphometry. *Neuroimage* 17(2):618–630
92. Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 12(3):189–198
93. Petersen RC, Smith GE, Ivnik RJ, Kokmen E, Tangalos EG (1994) Memory function in very early Alzheimer's disease. *Neurology* 44(5):867–872
94. Jacobs DM, Sano M, Dooneief G, Marder K, Bell KL, Stern Y (1995) Neuropsychological detection and characterization of preclinical Alzheimer's disease. *Neurology* 45(5):957–962
95. Hyman BT, Van Hoesen GW, Damasio AR, Barnes CL (1984) Alzheimer's disease: cell-specific pathology isolates the hippocampal formation. *Science* 225(4667):1168–1170
96. Deweer B, Lehericy S, Pillon B, Baulac M, Chiras J, Marsault C, Agid Y, Dubois B (1995) Memory disorders in probable Alzheimer's disease: the role of hippocampal atrophy as shown with MRI. *J Neurol Neurosurg Psychiatry* 58(5):590–597
97. Fama R, Sullivan EV, Shear PK, Marsh L, Yesavage JA, Tinklenberg JR, Lim KO, Pfefferbaum A (1997) Selective cortical and hippocampal volume correlates of Mattis Dementia Rating Scale in Alzheimer disease. *Arch Neurol* 54(6):719–728
98. Cahn DA, Sullivan EV, Shear PK, Marsh L, Fama R, Lim KO, Yesavage JA, Tinklenberg JR, Pfefferbaum A (1998) Structural MRI correlates of recognition memory in Alzheimer's disease. *J Int Neuropsychol Soc* 4(2):106–114
99. de Toledo-Morrell L, Dickerson B, Sullivan MP, Spanovic C, Wilson R, Bennett DA (2000) Hemispheric differences in hippocampal volume predict verbal and spatial memory performance in patients with Alzheimer's disease. *Hippocampus* 10(2):136–142
100. Petersen RC, Jack CR Jr, Xu YC, Waring SC, O'Brien PC, Smith GE, Ivnik RJ, Tangalos EG, Boeve BF, Kokmen E (2000) Memory and MRI-based hippocampal volumes in aging and AD. *Neurology* 54(3):581–587
101. Sencakova D, Graff-Radford NR, Willis FB, Lucas JA, Parfitt F, Cha RH, O'Brien PC, Petersen RC, Jack CR Jr (2001) Hippocampal atrophy correlates with clinical features of Alzheimer disease in African Americans. *Arch Neurol* 58(10):1593–1597
102. Mungas D, Harvey D, Reed BR, Jagust WJ, DeCarli C, Beckett L, Mack WJ, Kramer JH, Weiner MW, Schuff N, Chui HC (2005) Longitudinal volumetric MRI change and rate of cognitive decline. *Neurology* 65(4):565–571
103. Chan D, Fox NC, Scallan RI, Crum WR, Whitwell JL, Leschziner G, Rossor AM, Stevens JM, Cipolotti L, Rossor MN (2001) Patterns of temporal lobe atrophy in semantic dementia and Alzheimer's disease. *Ann Neurol* 49(4):433–442
104. Fama R, Sullivan EV, Shear PK, Cahn-Weiner DA, Marsh L, Lim KO, Yesavage JA, Tinklenberg JR, Pfefferbaum A (2000) Structural brain correlates of verbal and nonverbal fluency measures in Alzheimer's disease. *Neuropsychology* 14(1):29–40
105. Thompson PM, Hayashi KM, De Zubicaray GI, Janke AL, Rose SE, Semple J, Hong MS, Herman DH, Gravano D, Doddrell DM, Toga AW (2004) Mapping hippocampal and ventricular change in Alzheimer disease. *Neuroimage* 22(4):1754–1766
106. de la Torre JC (2002) Alzheimer disease as a vascular disorder: nosological evidence. *Stroke* 33(4):1152–1162
107. Skoog I, Kalaria RN, Breteler MM (1999) Vascular factors and Alzheimer disease. *Alzheimer Dis Assoc Disord* 13(Suppl 3):S106–S114
108. Hachinski V, Munoz DG (1997) Cerebrovascular pathology in Alzheimer's disease: cause, effect or epiphenomenon? *Ann N Y Acad Sci* 826:1–6
109. de Groot JC, de Leeuw FE, Oudkerk M, Hofman A, Jolles J, Breteler MM (2001) Cerebral white matter lesions and subjective cognitive dysfunction: the Rotterdam Scan Study. *Neurology* 56(11):1539–1545
110. Roman GC, Erkinjuntti T, Wallin A, Pantoni L, Chui HC (2002) Subcortical ischaemic vascular dementia. *Lancet Neurol* 1(7):426–436
111. Bastos Leite AJ, Scheltens P, Barkhof F (2004) Pathological aging of the brain: an overview. *Top Magn Reson Imaging* 15(6):369–389
112. Nagy Z, Esiri MM, Jobst KA, Morris JH, King EM, McDonald B, Joachim C, Litchfield S, Barnettson L, Smith AD (1997) The effects of additional pathology on the cognitive deficit in Alzheimer disease. *J Neuropathol Exp Neurol* 56(2):165–170
113. Goulding JM, Signorini DF, Chatterjee S, Nicoll JA, Stewart J, Morris R, Lammie GA (1999) Inverse relation between Braak stage and cerebrovascular pathology in Alzheimer predominant dementia. *J Neurol Neurosurg Psychiatry* 67(5):654–657
114. Jagust W, Thisted R, Devous MD Sr, Van Heertum R, Mayberg H, Jobst K, Smith AD, Borys N (2001) SPECT perfusion imaging in the diagnosis of Alzheimer's disease: a clinical-pathologic study. *Neurology* 56(7):950–956
115. Jobst KA, Barnettson LP, Shepstone BJ (1998) Accurate prediction of histologically confirmed Alzheimer's disease and the differential diagnosis of dementia: the use of NINCDS-ADRDA and DSM-III-R criteria, SPECT, X-ray CT, and Apo E4 in medial temporal lobe dementias. Oxford Project to Investigate Memory and Aging. *Int Psychogeriatr* 10(3):271–302
116. Scheltens P, Launer LJ, Barkhof F, Weinstein HC, Jonker C (1997) The diagnostic value of magnetic resonance imaging and technetium 99 m-HMPAO single-photon-emission computed tomography for the diagnosis of Alzheimer disease in a community-dwelling elderly population. *Alzheimer Dis Assoc Disord* 11(2):63–70
117. Jagust WJ (1996) Functional imaging patterns in Alzheimer's disease. Relationships to neurobiology. *Ann N Y Acad Sci* 777:30–36
118. Bartenstein P, Minoshima S, Hirsch C, Buch K, Willoch F, Mosch D, Schad D, Schwaiger M, Kurz A (1997) Quantitative assessment of cerebral blood flow in patients with Alzheimer's disease by SPECT. *J Nucl Med* 38(7):1095–1101
119. Bozzao A, Floris R, Baviera ME, Apruzzese A, Simonetti G (2001) Diffusion and perfusion MR imaging in cases of Alzheimer's disease: correlations with cortical atrophy and lesion load. *Am J Neuroradiol* 22(6):1030–1036



120. Detre JA, Alsop DC (1999) Perfusion magnetic resonance imaging with continuous arterial spin labeling: methods and clinical applications in the central nervous system. *Eur J Radiol* 30 (2):115–124
121. Alsop DC, Detre JA (1998) Multisection cerebral blood flow MR imaging with continuous arterial spin labeling. *Radiology* 208(2):410–416
122. Alsop DC, Detre JA (1996) Reduced transit-time sensitivity in noninvasive magnetic resonance imaging of human cerebral blood flow. *J Cereb Blood Flow Metab* 16(6):1236–1249
123. Sandson TA, O'Connor M, Sperling RA, Edelman RR, Warach S (1996) Noninvasive perfusion MRI in Alzheimer's disease: a preliminary report. *Neurology* 47(5):1339–1342
124. Alsop DC, Detre JA, Grossman M (2000) Assessment of cerebral blood flow in Alzheimer's disease by spin-labeled magnetic resonance imaging. *Ann Neurol* 47(1):93–100
125. Johnson NA, Jahng GH, Weiner MW, Miller BL, Chui HC, Jagust WJ, Gorno-Tempini ML, Schuff N (2005) Pattern of cerebral hypoperfusion in Alzheimer disease and mild cognitive impairment measured with arterial spin-labeling MR imaging: initial experience. *Radiology* 234(3):851–859
126. Bassar PJ, Pajevic S, Pierpaoli C, Dud-a J, Aldroubi A (2000) In vivo fiber tractography using DT-MRI data. *Magn Reson Med* 44(4):625–632
127. Mori S, Crain BJ, Chacko VP, van Zijl PC (1999) Three-dimensional tracking of axonal projections in the brain by magnetic resonance imaging. *Ann Neurol* 45(2):265–269
128. Poupon C, Clark CA, Frouin V, Regis J, Bloch I, Le Bihan D, Mangin J (2000) Regularization of diffusion-based direction maps for the tracking of brain white matter fascicles. *Neuroimage* 12(2):184–195
129. Le Bihan D, Mangin JF, Poupon C, Clark CA, Pappata S, Molko N, Chabriet H (2001) Diffusion tensor imaging: concepts and applications. *J Magn Reson Imaging* 13(4):534–546
130. Hanyu H, Sakurai H, Iwamoto T, Takasaki M, Shindo H, Abe K (1998) Diffusion-weighted MR imaging of the hippocampus and temporal white matter in Alzheimer's disease. *J Neurol Sci* 156(2):195–200
131. Sandson TA, Felician O, Edelman RR, Warach S (1999) Diffusion-weighted magnetic resonance imaging in Alzheimer's disease. *Dement Geriatr Cogn Disord* 10(2):166–171
132. Yoshiura T, Mihara F, Tanaka A, Ogomori K, Ohayagi Y, Taniwaki T, Yamada T, Yamasaki T, Ichimiya A, Kinukawa N, Kuwabara Y, Honda H (2003) High b value diffusion-weighted imaging is more sensitive to white matter degeneration in Alzheimer's disease. *Neuroimage* 20(1):413–419
133. Muller MJ, Greverus D, Dellani PR, Weibrich C, Wille PR, Scheurich A, Stoeter P, Fellgiebel A (2005) Functional implications of hippocampal volume and diffusivity in mild cognitive impairment. *Neuroimage* 28 (4):1033–1042
134. Kantarci K, Jack CR Jr, Xu YC, Campeau NG, O'Brien PC, Smith GE, Ivnik RJ, Boeve BF, Kokmen E, Tangalos EG, Petersen RC (2001) Mild cognitive impairment and Alzheimer disease: regional diffusivity of water. *Radiology* 219(1):101–107
135. Rose SE, Chen F, Chalk JB, Zelaya FO, Strugnell WE, Benson M, Semple J, Doddrell DM (2000) Loss of connectivity in Alzheimer's disease: an evaluation of white matter tract integrity with colour coded MR diffusion tensor imaging. *J Neurol Neurosurg Psychiatry* 69(4):528–530
136. Fellgiebel A, Muller MJ, Wille P, Dellani PR, Scheurich A, Schmidt LG, Stoeter P (2005) Color-coded diffusion-tensor-imaging of posterior cingulate fiber tracts in mild cognitive impairment. *Neurobiol Aging* 26(8):1193–1198
137. Sugihara S, Kinoshita T, Matsusue E, Fujii S, Ogawa T (2004) Usefulness of diffusion tensor imaging of white matter in Alzheimer disease and vascular dementia. *Acta Radiol* 45(6):658–663
138. Takahashi S, Yonezawa H, Takahashi J, Kudo M, Inoue T, Tohgi H (2002) Selective reduction of diffusion anisotropy in white matter of Alzheimer disease brains measured by 3.0 Tesla magnetic resonance imaging. *Neurosci Lett* 332(1):45–48
139. Kalus P, Slotboom J, Gallinat J, Mahlberg R, Cattapan-Ludewig K, Wiest R, Nyffeler T, Buri C, Federspiel A, Kunz D, Schroth G, Kiefer C (2005) Examining the gateway to the limbic system with diffusion tensor imaging: The perforant pathway in dementia. *Neuroimage*
140. Kantarci K, Petersen RC, Boeve BF, Knopman DS, Weigand SD, O'Brien PC, Shiung MM, Smith GE, Ivnik RJ, Tangalos EG, Jack CR Jr (2005) DWI predicts future progression to Alzheimer disease in amnesic mild cognitive impairment. *Neurology* 64 (5):902–904
141. Medina D, Detoledo-Morrell L, Urresta F, Gabrieli JD, Moseley M, Fleischman D, Bennett DA, Leurgans S, Turner DA, Stebbins GT (2005) White matter changes in mild cognitive impairment and AD: A diffusion tensor imaging study. *Neurobiol Aging*
142. Kaufer DI, Miller BL, Itti L, Fairbanks LA, Li J, Fishman J, Kushi J, Cummings JL (1997) Midline cerebral morphometry distinguishes frontotemporal dementia and Alzheimer's disease. *Neurology* 48(4):978–985
143. Yamauchi H, Fukuyama H, Nagahama Y, Katsumi Y, Hayashi T, Oyanagi C, Konishi J, Shio H (2000) Comparison of the pattern of atrophy of the corpus callosum in frontotemporal dementia, progressive supranuclear palsy, and Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 69(5):623–629
144. Ross BD, Bluml S, Cowan R, Danielsen E, Farrow N, Tan J (1998) In vivo MR spectroscopy of human dementia. *Neuroimaging Clin N Am* 8(4):809–822
145. Miller BL, Moats RA, Shonk T, Ernst T, Woolley S, Ross BD (1993) Alzheimer disease: depiction of increased cerebral myo-inositol with proton MR spectroscopy. *Radiology* 187(2):433–437
146. Shonk TK, Moats RA, Gifford P, Michaelis T, Mandigo JC, Izumi J, Ross BD (1995) Probable Alzheimer disease: diagnosis with proton MR spectroscopy. *Radiology* 195(1):65–72
147. Moats RA, Ernst T, Shonk TK, Ross BD (1994) Abnormal cerebral metabolite concentrations in patients with probable Alzheimer disease. *Magn Reson Med* 32(1):110–115
148. Jessen F, Block W, Traber F, Keller E, Flacke S, Papassotiropoulos A, Lamerichs R, Heun R, Schild HH (2000) Proton MR spectroscopy detects a relative decrease of N-acetylaspartate in the medial temporal lobe of patients with AD. *Neurology* 55(5):684–688
149. Jessen F, Traeber F, Freymann N, Maier W, Schild HH, Heun R, Block W (2005) A comparative study of the different N-acetylaspartate measures of the medial temporal lobe in Alzheimer's disease. *Dement Geriatr Cogn Disord* 20(2–3):178–183
150. Firbank MJ, Harrison RM, O'Brien JT (2002) A comprehensive review of proton magnetic resonance spectroscopy studies in dementia and Parkinson's disease. *Dement Geriatr Cogn Disord* 14(2):64–76

151. Adalsteinsson E, Sullivan EV, Kleinmans N, Spielman DM, Pfefferbaum A (2000) Longitudinal decline of the neuronal marker N-acetyl aspartate in Alzheimer's disease. *Lancet* 355(9216):1696–1697
152. Schuff N, Amend D, Ezekiel F, Steinman SK, Tanabe J, Norman D, Jagust W, Kramer JH, Mastrianni JA, Fein G, Weiner MW (1997) Changes of hippocampal N-acetyl aspartate and volume in Alzheimer's disease. A proton MR spectroscopic imaging and MRI study. *Neurology* 49(6):1513–1521
153. Heun R, Schlegel S, Graf-Morgenstern M, Tintera J, Gawehn J, Stoeter P (1997) Proton magnetic resonance spectroscopy in dementia of Alzheimer type. *Int J Geriatr Psychiatry* 12(3):349–358
154. Sijens PE, den Heijer T, Origgi D, Vermeer SE, Breteler MM, Hofman A, Oudkerk M (2003) Brain changes with aging: MR spectroscopy at supraventricular plane shows differences between women and men. *Radiology* 226(3):889–896
155. den Heijer T, Sijens PE, Prins ND, Hofman A, Koudstaal PJ, Oudkerk M, Breteler MM (2006) MR spectroscopy of brain white matter in the prediction of dementia. *Neurology* 66(4):540–544
156. Kantarci K, Petersen RC, Boeve BF, Knopman DS, Tang-Wai DF, O'Brien PC, Weigand SD, Edland SD, Smith GE, Ivnik RJ, Ferman TJ, Tangalos EG, Jack CR Jr (2004) 1 H MR spectroscopy in common dementias. *Neurology* 63(8):1393–1398
157. Kantarci K, Reynolds G, Petersen RC, Boeve BF, Knopman DS, Edland SD, Smith GE, Ivnik RJ, Tangalos EG, Jack CR Jr (2003) Proton MR spectroscopy in mild cognitive impairment and Alzheimer disease: comparison of 1.5 and 3 T. *Am J Neuroradiol* 24(5):843–849
158. Kantarci K, Jack CR Jr, Xu YC, Campeau NG, O'Brien PC, Smith GE, Ivnik RJ, Boeve BF, Kokmen E, Tangalos EG, Petersen RC (2000) Regional metabolic patterns in mild cognitive impairment and Alzheimer's disease: A 1 H MRS study. *Neurology* 55(2):210–217
159. Valenzuela MJ, Sachdev P (2001) Magnetic resonance spectroscopy in AD. *Neurology* 56(5):592–598
160. Antuono PG, Jones JL, Wang Y, Li SJ (2001) Decreased glutamate+glutamine in Alzheimer's disease detected in vivo with 1 H-MRS at 0.5 T. *Neurology* 56(6):737–742
161. Hattori N, Abe K, Sakoda S, Sawada T (2002) Proton MR spectroscopic study at 3 Tesla on glutamate/glutamine in Alzheimer's disease. *Neuroreport* 13(1):183–186
162. Sijens PE, den Heijer T, de Leeuw FE, de Groot JC, Achten E, Heijboer RJ, Hofman A, Breteler MM, Oudkerk M (2001) MR spectroscopy detection of lactate and lipid signals in the brains of healthy elderly people. *Eur Radiol* 11(8):1495–1501
163. Pugliese M, Carrasco JL, Andrade C, Mas E, Mascort J, Mahy N (2005) Severe cognitive impairment correlates with higher cerebrospinal fluid levels of lactate and pyruvate in a canine model of senile dementia. *Prog Neuropsychopharmacol Biol Psychiatry* 29(4):603–610
164. Klunk WE, Engler H, Nordberg A, Wang Y, Blomqvist G, Holt DP, Bergstrom M, Savitcheva I, Huang GF, Estrada S, Ausen B, Debnath ML, Barletta J, Price JC, Sandell J, Lopresti BJ, Wall A, Koivisto P, Antoni G, Mathis CA, Langstrom B (2004) Imaging brain amyloid in Alzheimer's disease with Pittsburgh Compound-B. *Ann Neurol* 55(3):306–319
165. Sair HI, Doraiswamy PM, Petrella JR (2004) In vivo amyloid imaging in Alzheimer's disease. *Neuroradiology* 46(2):93–104
166. Shoghi-Jadid K, Small GW, Agdeppa ED, Kepe V, Ercole LM, Siddarth P, Read S, Satyamurthy N, Petric A, Huang SC, Barrio JR (2002) Localization of neurofibrillary tangles and beta-amyloid plaques in the brains of living patients with Alzheimer disease. *Am J Geriatr Psychiatry* 10(1):24–35
167. Benveniste H, Einstein G, Kim KR, Hulette C, Johnson GA (1999) Detection of neuritic plaques in Alzheimer's disease by magnetic resonance microscopy. *Proc Natl Acad Sci USA* 96(24):14079–14084
168. Dhenain M, Privat N, Duyckaerts C, Jacobs RE (2002) Senile plaques do not induce susceptibility effects in T2\*-weighted MR microscopic images. *NMR Biomed* 15(3):197–203
169. Poduslo JF, Wengenack TM, Curran GL, Wisniewski T, Sigurdsson EM, Macura SI, Borowski BJ, Jack CR Jr (2002) Molecular targeting of Alzheimer's amyloid plaques for contrast-enhanced magnetic resonance imaging. *Neurobiol Dis* 11(2):315–329
170. Wadghiri YZ, Sigurdsson EM, Wisniewski T, Turnbull DH (2005) Magnetic resonance imaging of amyloid plaques in transgenic mice. *Methods Mol Biol* 299:365–379
171. Helpert JA, Lee SP, Falangola MF, Dyakin VV, Bogart A, Ardekani B, Duff K, Branch C, Wisniewski T, de Leon MJ, Wolf O, O'Shea J, Nixon RA (2004) MRI assessment of neuropathology in a transgenic mouse model of Alzheimer's disease. *Magn Reson Med* 51(4):794–798
172. Zhang J, Yarowsky P, Gordon MN, Di Carlo G, Munireddy S, van Zijl PC, Mori S (2004) Detection of amyloid plaques in mouse models of Alzheimer's disease by magnetic resonance imaging. *Magn Reson Med* 51(3):452–457
173. Jack CR Jr, Garwood M, Wengenack TM, Borowski B, Curran GL, Lin J, Adriany G, Grohn OH, Grimm R, Poduslo JF (2004) In vivo visualization of Alzheimer's amyloid plaques by magnetic resonance imaging in transgenic mice without a contrast agent. *Magn Reson Med* 52(6):1263–1271
174. Jack CR Jr, Wengenack TM, Reyes DA, Garwood M, Curran GL, Borowski BJ, Lin J, Preboske GM, Holasek SS, Adriany G, Poduslo JF (2005) In vivo magnetic resonance microimaging of individual amyloid plaques in Alzheimer's transgenic mice. *J Neurosci* 25(43):10041–10048
175. Higuchi M, Iwata N, Matsuba Y, Sato K, Sasamoto K, Saido TC (2005) 19F and 1 H MRI detection of amyloid beta plaques in vivo. *Nat Neurosci* 8(4):527–533