

MEASURE THEORY

CONTENTS

1. Lebesgue integral	1
1.1. Definition, basic properties and monotone convergence	1
1.2. Additivity of the integral	5
1.3. Integrable functions and convergence	7
2. Outer measures	8
3. Monotone classes	10
4. Product measures and Fubini-Tonelli theorem	11
5. Jensen's Inequality	15
6. Hausdorff measure	18
6.1. Definition and Borel measurability	18
6.2. Equality of Hausdorff and Lebesgue measure when $s = d$.	20
7. Area formula and change of variables in integration	25
7.1. Area formula for Linear maps	26
7.2. Proof of the Area formula when $Jf(x) > 0$ for all $x \in X$.	26
7.3. Proof of the area formula without assuming $Jf(x) > 0$	28
7.4. Change of variables	29
7.5. Volume and surface area of the unit ball	31

1. LEBESGUE INTEGRAL

1.1. Definition, basic properties and monotone convergence.

Definition 1.1. Let X be a measurable space. We say that $f : X \rightarrow \mathbb{C}$ is a simple function if it is measurable and the range $f(X) = \{f(x) \mid x \in X\}$ is a finite set.

Lemma 1.2. Let $f : X \rightarrow \mathbb{C}$ be measurable. Then f is a simple function if and only if

$$f = \sum_{i=1}^n c_i \mathbb{1}_{A_i}$$

for some positive integer n and measurable $A_1, \dots, A_n \subset X$ and $c_i \in \mathbb{C}$.

Proof. If f is simple then the range of f is a finite set $\{c_1, \dots, c_n\}$. So we can just take $A_i = f^{-1}(c_i)$.

Conversely, suppose f takes such a form. Define now $q : X \rightarrow \{0, 1\}^n$ by the map

$$f(x) = (\mathbb{1}_{A_1}(x), \dots, \mathbb{1}_{A_n}(x)).$$

Thus $f(x) = \langle q(x), c \rangle$ where $c = (c_1, \dots, c_n)$. So clearly f takes finitely many values (at most 2^n), as it only depends on $q(x)$. $\square$

Thus, the space of simple functions form a vector space.

We say that $A_1, \dots, A_n$ is a measurable partition of X if each $A_i \subset X$ is measurable, $A_i \cap A_j = \emptyset$ for $i \neq j$ and

$$\bigcup_{i=1}^n A_i = X.$$

Lemma 1.3. Let μ be a measure on a space X and suppose $f : X \rightarrow \mathbb{C}$ is measurable. Suppose that

$$f = \sum_{i=1}^n c_i \mathbb{1}_{A_i},$$

for some positive integer n , some measurable partition $A_1, \dots, A_n \subset X$ of X and some $c_i \in \mathbb{R}$. Then the quantity

$$I_\mu(f) := \sum_{i=1}^n c_i \mu(A_i)$$

does not depend on n and the $(A_1, c_1), \dots, (A_n, c_n)$.

Proof. Let $q_1, \dots, q_m$ be all the possible distinct values of c_i , for $i \in \{1, \dots, n\}$ such that $A_i \neq \emptyset$. Let

$$I_j = \{i \in \{1, \dots, n\} \mid c_i = q_j\}.$$

Let

$$B_j = \bigsqcup_{i \in I_j} A_i.$$

Note that the B_j form a measurable partition of X into non-empty subsets. Thus

$$f = \sum_{j=1}^m q_j \sum_{i \in I_j} \mathbb{1}_{A_i} = \sum_{j=1}^m q_j \mathbb{1}_{B_j}.$$

Likewise,

$$\sum_{i=1}^n c_i \mu(A_i) = \sum_{j=1}^m q_j \sum_{i \in I_j} \mu(A_i) = \sum_{j=1}^m q_j \mu(B_j).$$

If $x \in B_j$ then $f(x) = q_j$ and so as the B_j are disjoint and non-empty, we thus have that

$$f^{-1}(q_j) = B_j.$$

This shows that

$$\sum_{i=1}^n c_i \mu(A_i) = \sum_{y \in \mathbb{R}} y \mu(f^{-1}(y))$$

and is thus not dependent on the choice of $A_1, \dots, A_n$ nor the $c_1, \dots, c_n$. $\square$

Lemma 1.4. Suppose that $f, g : X \rightarrow \mathbb{C}$ are simple functions. Then

$$I_\mu(f + g) = I_\mu(f) + I_\mu(g).$$

Proof. We can write

$$f = \sum_{i=1}^n c_i \mathbb{1}_{A_i}$$

and

$$g = \sum_{i=1}^m d_i \mathbb{1}_{B_i}.$$

Now let

$$Q_{i,j} = A_i \cap B_j.$$

Thus

$$f = \sum_{i=1}^n c_i \mathbb{1}_{A_i} = \sum_{i=1}^n c_i \sum_{j=1}^m \mathbb{1}_{Q_{i,j}} = \sum_{i,j} c_i \mathbb{1}_{Q_{i,j}}.$$

Likewise

$$g = \sum_{i,j} d_j \mathbb{1}_{Q_{i,j}}.$$

Thus by Lemma 1.3 we have

$$I_\mu(f + g) = \sum_{i,j} (c_i + d_j) \mu(Q_{i,j}) = \sum_{i,j} c_i \mu(Q_{i,j}) + \sum_{i,j} d_j \mu(Q_{i,j}) = I_\mu(f) + I_\mu(g).$$

□

Definition 1.5. Let μ be a measure on a space X . Let $f : X \rightarrow [0, \infty]$ be a positive measurable function. Define the Lebesgue integral to be

$$\int_X f d\mu = \sup\{I_\mu(f) \mid g : X \rightarrow [0, \infty) \text{ is a simple function with } 0 \leq g \leq f\}.$$

Lemma 1.6. If $f_1, f_2 : X \rightarrow [0, \infty]$ are such that $0 \leq f_1 \leq f_2$ then

$$0 \leq \int_X f_1 d\mu \leq \int_X f_2 d\mu.$$

Proof. Obvious. □

Lemma 1.7. If $f : X \rightarrow [0, \infty)$ is simple then $I_\mu(f) = \int_X f d\mu$.

Proof. Clearly $I_\mu(f) \leq \int_X f d\mu$ since we can take $f = g$. Now we prove the other inequality. Thus suppose $g : X \rightarrow [0, \infty)$ is simple such that $0 \leq g \leq f$. Then $0 \leq f - g$ and $f - g$ is simple so $I_\mu(f - g) \geq 0$. However

$$I_\mu(f) = I_\mu(g) + I_\mu(f - g) \geq I_\mu(g).$$

Thus $I_\mu(f) \geq \int_X f d\mu$. □

Theorem 1.8 (Monotone convergence theorem). Suppose that $f_1, f_2, \dots : X \rightarrow [0, \infty]$ be a sequence of measurable functions that is pointwise increasing, i.e., such that $f_n(x) \leq f_{n+1}(x)$ for all $x \in X$ and $n \in \mathbb{Z}_{>0}$. Then

$$\int_X f_i d\mu \rightarrow \int_X \lim_{n \rightarrow \infty} f_n(x).$$

Proof. Let $f(x) = \lim_{n \rightarrow \infty} f_n(x)$. Let $I = \int_X f_n d\mu$. If $I = 0$ then the theorem is true, so assume $I > 0$. Hence let $0 < I_1 < I_2 < \dots < I$ be a sequence such that $I_n \rightarrow I$, i.e., if $I < \infty$ then we can let $I_n = I - n^{-1}$ otherwise we can let $I_n = n$. Fix an integer $Q > 0$. Now there exists a simple function $g : X \rightarrow [0, \infty)$ such that $0 \leq g \leq f$ and

$$(1) \quad \int g d\mu > I_Q.$$

Now write

$$g = \sum_{i=1}^m c_i \mathbb{1}_{A_i}$$

for some measurable disjoint $A_1, \dots, A_m \subset X$ and $0 < c_i < \infty$. For convenience, let us consider two cases.

Case 1: $\mu(A_i) = \infty$ for some A_i .

Then in this case $I = \infty$ as $\int_X g d\mu \geq c_i \mu(A_i) \infty$. Now let

$$B_n = \{x \in A_i \mid f_n(x) > c_i/2\}.$$

Note that

$$\bigcup_n B_n = A_i,$$

and thus $\mu(B_n) \rightarrow \mu(A_i) = \infty$. Now $f_n(x) \geq \frac{c_i}{2} \mathbb{1}_{B_n}$ and hence

$$\int_X f_n d\mu \geq \frac{c_i}{2} \mu(B_n) \rightarrow \infty,$$

which completes the proof of the monotone convergence theorem in this case.

Case 2: $\mu(A_i) < \infty$ for each A_i .

From (1), we can now fix $0 < \epsilon < 1$ such that $(1 - \epsilon) \int_X g d\mu > I_Q$. Let

$$B_{n,i} = \{x \in A_i \mid f_n(x) > (1 - \epsilon)c_i\}.$$

Now

$$\int_X f_n d\mu \geq \sum_{i=1}^m (1 - \epsilon)c_i \mu(B_{n,i}) \rightarrow \sum_{i=1}^m (1 - \epsilon)c_i \mu(A_i) \quad \text{as } n \rightarrow \infty.$$

But

$$\sum_{i=1}^m (1 - \epsilon)c_i \mu(A_i) = (1 - \epsilon) \int_X g d\mu > I_Q.$$

Thus we have shown that for large enough n we have

$$\int_X f_n d\mu > I_Q,$$

which completes the proof. $\square$

Proposition 1.9. Let $f : X \rightarrow [0, \infty]$ be measurable. Then there exists measurable $A_1, A_2, \dots \subset X$ and such that

$$f(x) = \sum_{n=1}^{\infty} \frac{1}{n} \mathbb{1}_{A_n}.$$

Thus f is a monotonic limit of simple functions.

Proof. We define $A_1, A_2, \dots$ recursively as follows. Let

$$f_n = \sum_{i=1}^n \frac{1}{i} \mathbb{1}_{A_i}.$$

We let $f_0 = 0$. Let $n \geq 0$ and suppose $0 \leq f_n \leq f$ has already been constructed (more precisely, we mean $A_1, \dots, A_n$ have already been constructed). Let

$$A_{n+1} = \{x \in X \mid f_n(x) + \frac{1}{n+1} \leq f(x)\}.$$

Thus

$$f_{n+1} = f_n + \frac{1}{n+1} \mathbb{1}_{A_{n+1}} \leq f$$

still holds.

We must now show that

$$f_n \rightarrow f.$$

First, if $f(x) = \infty$ then $x \in A_n$ for all n and thus

$$\sum_{n=1}^{\infty} \frac{1}{n} \mathbb{1}_{A_n}(x) = \sum_{n=1}^{\infty} \frac{1}{n} = \infty = f(x).$$

Now suppose that $f(x) < \infty$. Suppose for contradiction that

$$\lim_{k \rightarrow \infty} f_k(x) < f(x) - 1/N$$

for some $N > 0$. As f_n is increasing, it now follows that for $n > N$ we have

$$f_n(x) \leq \lim_{k \rightarrow \infty} f_k(x) \leq f(x) - \frac{1}{N} \leq f(x) - \frac{1}{n+1}.$$

But this precisely means that $x \in A_n$ for all $n > N$. Thus

$$f(x) \geq \sum_{n=N+1}^{\infty} \frac{1}{n} = \infty,$$

which is a contradiction. □

1.2. Additivity of the integral.

Theorem 1.10. Suppose that $f, g : X \rightarrow [0, \infty]$ are measurable. Then

$$\int_X (f + g) d\mu = \int_X f d\mu + \int_X g d\mu.$$

Proof. We know this is true for f, g simple by Lemma 1.4 and Lemma 1.7. Now we can write f and g as monotonic limits of simple functions f_n, g_n . But $f_n + g_n \rightarrow f + g$ monotonically. So the monotone convergence theorem completes the proof. □

Definition 1.11. We let $\int_A f d\mu = \int_X \mathbb{1}_A f d\mu$. We now define

$$\int_X f d\mu = \int_{f \geq 0} f d\mu - \int_{f < 0} |f| d\mu,$$

in case it is not the case that these integrals are not both ∞ (in which case, it is undefined).

Observe that in the definition above, we can replace strict inequality with non-strict inequality and vice versa (this is because $\int_X 0 d\mu = 0$ and the integral is additive on non-negative functions. Also, let us define the positive part $f^+ = \mathbf{1}_{f \geq 0} f$ and negative part $f^- = -\mathbf{1}_{f < 0} f$ of f . Thus by definition we have f^+ and f^- are both ≥ 0 and

$$\int_X f d\mu = \int_X f^+ d\mu - \int_X f^- d\mu$$

under the assumption that at least one of these integrals is not infinite.

Lemma 1.12. Suppose that $f, g : X \rightarrow [0, \infty]$ are measurable with $\int_X g d\mu < \infty$ or $\int_X f d\mu < \infty$. Then

$$\int_X (f - g) d\mu = \int_X f d\mu - \int_X g d\mu$$

with the left hand side being well defined (either the positive or negative part has finite integral).

Proof. By definition

$$\begin{aligned} \int_X (f - g) d\mu &= \int_X (f - g)^+ d\mu - \int_X (g - f)^+ d\mu \\ &= \int_{f \geq g} (f - g) d\mu - \int_{g > f} (g - f) d\mu \end{aligned}$$

but $(f - g)^+ \leq f$ and $(g - f)^+ \leq g$ so at least one of these integrals on the right hand side is non-infinite. Thus the integral is well defined. Now by Theorem 1.10, we have

$$(2) \quad \int_{f \geq g} (f - g) d\mu + \int_{f \geq g} g d\mu = \int_{f \geq g} f d\mu$$

and

$$(3) \quad \int_{g > f} (g - f) d\mu + \int_{g > f} f d\mu = \int_{g > f} g d\mu.$$

We will now assume $\int_X f d\mu < \infty$ (the other case will then follow by symmetry). Thus the right hand side of (2) is non-infinite and so all terms are non-infinite. So we can safely rearrange this equation. Likewise, the term $\int_{g > f} d\mu < \infty$ so it can safely be moved to the right hand side in the other equation (3). After rearranging and subtracting these two equations, we obtain that

$$\int_X (f - g) d\mu = \int_{f \geq g} f d\mu - \int_{f \geq g} g d\mu - \left(\int_{g > f} g d\mu - \int_{g > f} f d\mu \right).$$

Only the third integral can possibly be infinite as argued above, thus the terms can be safely grouped to give

$$\int_X (f - g) d\mu = \int_{f \geq g} f d\mu + \int_{g > f} d\mu - \left(\int_{g > f} g d\mu + \int_{f \geq g} g d\mu \right).$$

Finally, the proof is complete using Theorem 1.10. □

Theorem 1.13. Suppose that $f, g : X \rightarrow (-\infty, \infty]$ are such that either

$$\int_{f \geq 0} f d\mu + \int_{g \geq 0} g d\mu < \infty$$

or

$$\int_{f < 0} f d\mu + \int_{g < 0} g d\mu > -\infty.$$

Then the following holds and is well defined (no $\infty - \infty$ terms and all integrals exist)

$$\int_X (f + g) d\mu = \int_X f d\mu + \int_X g d\mu.$$

Proof. Write $f = f^+ - f^-$ and $g = g^+ - g^-$ as above. We assume the first condition, thus

$$\int_X f^+ + g^+ d\mu < \infty.$$

Now

$$f + g = f^+ + g^+ - (f^- + g^-),$$

so the assumption of the previous theorem must apply (likewise, they also apply for the other case). After applying this theorem, rearranging and using additivity of the integral for positive functions the proof is complete. $\square$

1.3. Integrable functions and convergence. We say that a measurable $f : X \rightarrow \mathbb{C}$ is integrable if $\int_X |f| d\mu < \infty$.

Lemma 1.14. Suppose that $f : X \rightarrow \mathbb{C}$ is integrable. Then for $\epsilon > 0$ there exists a measurable $A \subset X$ such that $\mu(A) < \infty$ and

$$\int_{X \setminus A} |f| d\mu < \epsilon.$$

Proof. Let $0 \leq h_1 \leq h_2 \leq \dots$ be a sequence of simple functions converging monotonically to $|f|$. Thus $\int_X h_n d\mu < \infty$ and so, as h_n is simple, there is a measurable set A_n such that $\mu(A_n) < \infty$ and $h_n(x) = 0$ for $x \in X \setminus A_n$. By the monotone convergence theorem, there is a large enough n so that

$$\int_X (|f| - h_n) d\mu < \epsilon.$$

However

$$\epsilon > \int_X (|f| - h_n) d\mu = \int_{A_n} (|f| - h_n) + \int_{X \setminus A_n} |f| d\mu$$

as $h_n = 0$ on $X \setminus A_n$. So we can take $A = A_n$ as all integrands are non-negative. $\square$

Theorem 1.15 (Lebesgue dominated convergence theorem). Suppose that $f_1, f_2, \dots : X \rightarrow \mathbb{C}$ are measurable and $g : X \rightarrow [0, \infty]$ is integrable, i.e., $\int_X g d\mu < \infty$ such that $|f_n| \leq g$ for all f_n . Suppose that $f_n \rightarrow f$ pointwise, then f is integrable and

$$\int_X |f_n - f| d\mu \rightarrow 0,$$

thus

$$\int_X f_n d\mu \rightarrow \int_X f d\mu.$$

Proof. Note that $|f| \leq |g|$ thus $|f|$ is integrable. Fix $\epsilon > 0$ and use the previous lemma to find $A \subset X$ with $\mu(A) < \infty$ and $\int_{X \setminus A} |g| d\mu < \epsilon$. Thus

$$\int_{X \setminus A} |f - f_n| \leq \int_{X \setminus A} 2g d\mu < 2\epsilon.$$

Now as $\mu(A) < \infty$, we can choose $\delta > 0$ so that

$$\delta \mu(A) < \epsilon.$$

Let

$$A_n = \{x \in A \mid |f_m(x) - f(x)| < \delta \text{ for all } m \geq n\}.$$

Notice that $A_n \subset A_{n+1}$ and $\bigcup_n A_n = A$. Thus by the monotone convergence theorem, we have

$$\left| \int_{A_n} g - \int_A g \right| < \epsilon$$

for all large enough $n > N$. This means that

$$\int_{A \setminus A_n} g < \epsilon$$

for all $n > N$ and consequently

$$\int_{A \setminus A_n} |f - f_n| \leq \int_{A \setminus A_n} 2g < 2\epsilon.$$

Finally, $|f(x) - f_n(x)| < \delta$ for all $x \in A_n$, by definition of A_n . Thus

$$\int_{A_n} |f(x) - f_n(x)| \leq \mu(A)\delta < \epsilon.$$

By splitting up $\int_X = \int_{X \setminus A} + \int_{A \setminus A_n} + \int_{A_n}$ and combining the estimates gathered, we obtain

$$\int_X |f - f_n| d\mu < 2\epsilon + 2\epsilon + \epsilon = 5\epsilon$$

for $n > N$. Thus completing the proof. $\square$

2. OUTER MEASURES

Definition 2.1. An *outer measure* on a set X is a function $\mu : 2^X \rightarrow [0, \infty]$ which satisfies the following conditions:

(1) Countably sub-additivity: if $A_1, A_2, \dots \subset X$ then

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) \leq \sum_{i=1}^{\infty} \mu(A_i).$$

(2) Monotonicity: If $A \subset B \subset X$ then $\mu(A) \leq \mu(B)$.

(3) Empty set is null: $\mu(\emptyset) = 0$.

We say that $A \subset X$ is μ -measurable if

$$\mu(B) = \mu(B \cap A) + \mu(B \cap A^c) \quad \text{for all } B \subset X.$$

Note that, by sub-additivity, this is equivalent to

$$\mu(B) \geq \mu(B \cap A) + \mu(B \cap A^c) \quad \text{for all } B \subset X.$$

Theorem 2.2. The set of all measurable subsets forms a σ -algebra on X . Moreover, μ is countably-additive (a measure) on X for this σ -algebra.

Proof. Observe that $\emptyset$ is measurable since for $B \subset X$ we have

$$\mu(B) = 0 + \mu(B \cap X) = \mu(\emptyset) + \mu(B \cap X) = \mu(B \cap \emptyset) + \mu(B \cap \emptyset^c).$$

Also A is measurable then A^c also is by symmetry.

We first show that the set of measurable sets is closed under finite intersection. Thus suppose that $A_1, A_2, \dots \subset X$ are measurable and $B \subset X$.

As A_1 is measurable, we obtain that

$$\mu(B \cap A_2^c) = \mu(B \cap A_2^c \cap A_1) + \mu(B \cap A_2^c \cap A_1^c)$$

and

$$\mu(B \cap A_2) = \mu(B \cap A_2 \cap A_1) + \mu(B \cap A_2 \cap A_1^c).$$

Now we add these two equations and use the inequality (which follows from subadditivity and the identity $A_1 \cup A_2 = (A_1^c \cap A_2) \cup (A_1 \cap A_2^c) \cup (A_1 \cap A_2)$)

$$\mu(B \cap A_2^c \cap A_1) + \mu(B \cap A_2 \cap A_1) + \mu(B \cap A_2 \cap A_1^c) \geq \mu(B \cap (A_1 \cup A_2))$$

to get that

$$\mu(B \cap A_2^c) + \mu(B \cap A_2) \geq \mu(B \cap (A_1 \cap A_2)) + \mu(B \cap A_1^c \cap A_2^c).$$

Measurability of A_2 means that the left hand side is $\mu(B)$, thus

$$\mu(B) \geq \mu(B \cap (A_1 \cap A_2)) + \mu(B \cap (A_1 \cup A_2)).$$

The reverse inequality holds from sub-additivity, thus $A_1 \cup A_2$ is measurable.

Thus we have shown that the measurable sets form an algebra. To show that they form a σ -algebra, it thus remains to show that a *disjoint* union of measurable sets is measurable. Thus suppose $A_1, A_2, \dots$ are disjoint measurable sets and $B \subset X$. We claim that

$$\mu\left(B \cap \left(\bigcup_{i=1}^n A_i\right)\right) = \mu(B \cap A_1) + \dots + \mu(B \cap A_n).$$

We prove this by induction as follows. Suppose it holds for n . Then by measurability of A_{n+1} we have that

$$\begin{aligned} \mu\left(B \cap \left(\bigcup_{i=1}^{n+1} A_i\right)\right) &= \mu\left(B \cap \left(\bigcup_{i=1}^{n+1} A_i\right) \cap A_{n+1}\right) + \mu\left(B \cap \left(\bigcup_{i=1}^{n+1} A_i\right) \cap A_{n+1}^c\right) \\ &= \mu(B \cap A_{n+1}) + \mu\left(B \cap \left(\bigcup_{i=1}^n A_i\right)\right) \end{aligned}$$

which completes the induction step (the last equation uses disjointness of the A_i). Now by monotonicity of μ we get

$$\mu\left(B \cap \left(\bigcup_{i=1}^{\infty} A_i\right)\right) \geq \lim_{n \rightarrow \infty} \mu\left(B \cap \left(\bigcup_{i=1}^n A_i\right)\right) = \sum_{i=1}^{\infty} \mu(B \cap A_i).$$

The reverse inequality holds from sub-additivity, thus we have shown that

$$\mu\left(B \cap \left(\bigcup_{i=1}^{\infty} A_i\right)\right) = \sum_{i=1}^{\infty} \mu(B \cap A_i).$$

In particular, for $B = X$ this shows that μ is countably-additive on the measurable sets. It remains to show that this countable union is measurable. Observe that since finite unions are measurable we have

$$\begin{aligned} \mu(B) &= \mu\left(B \cap \left(\bigcup_{i=1}^n A_i\right)\right) + \mu\left(B \cap \left(\bigcup_{i=1}^n A_i\right)^c\right) \\ &\geq \mu\left(B \cap \left(\bigcup_{i=1}^n A_i\right)\right) + \mu\left(B \cap \left(\bigcup_{i=1}^{\infty} A_i\right)^c\right). \end{aligned}$$

Now letting $n \rightarrow \infty$ and using countable sub-additivity, the proof is complete. $\square$

Example 2.3 (Lebesgue outer measure). If $I \subset \mathbb{R}$ is a non-empty bounded interval, we let $\text{len}(I) = \sup(I) - \inf(I)$ and $\text{len}(\emptyset) = 0$. Note I could be open, closed or half open. If $\mathcal{I}$ is a countable set of bounded intervals of $\mathbb{R}$, then we denote $\text{len}(\mathcal{I}) = \sum_{I \in \mathcal{I}} \text{len}(I)$. We say that $\mathcal{I}$ covers B if

$$B \subset \bigcup_{I \in \mathcal{I}} I.$$

Finally, denote the Lebesgue outer-measure of $B \subset \mathbb{R}$ to be

$$\lambda(B) = \inf_{\mathcal{I} \text{ covers } B} \text{len}(\mathcal{I}).$$

Let us now show that any bounded interval J is λ -measurable. To see this, let $B \subset \mathbb{R}$ be any subset and suppose that $\mathcal{I}$ covers B such that $\text{len}(\mathcal{I}) < \lambda(B) + \epsilon$. We can write $J^c = J_1 \cup J_2$ where J_1 and J_2 are disjoint (unbounded) intervals. Let $\mathcal{I}_1 = \{I \cap J \mid I \in \mathcal{I}\}$ and

$$\mathcal{I}_2 = \{I \cap J_m \mid I \in \mathcal{I}, m \in \{1, 2\}\}.$$

Then $\mathcal{I}_1$ covers $B \cap J$ and $\mathcal{I}_2$ covers $B \cap J^c$. Observe that $\text{len}(\mathcal{I}_1) + \text{len}(\mathcal{I}_2) = \text{len}(\mathcal{I})$ since

$$\text{len}(I) = \text{len}(I \cap J_1) + \text{len}(I \cap J) + \text{len}(I \cap J_2).$$

Thus

$$\lambda(B \cap J) + \lambda(B \cap J^c) \leq \text{len}(\mathcal{I}_1) + \text{len}(\mathcal{I}_2) = \text{len}(\mathcal{I}) \leq \lambda(B) + \epsilon.$$

Letting $\epsilon \rightarrow 0$ shows that J is λ -measurable.

3. MONOTONE CLASSES

Definition 3.1. Let X be a set. A monotone class on X is a collection of subsets $\mathcal{A} \subset 2^X$ which satisfies the following properties.

- (1) $\emptyset, X \in \mathcal{A}$.
- (2) If $A_1, A_2, \dots \in \mathcal{A}$ is such that $A_1 \subset A_2 \subset \dots$ then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{A}$.
- (3) If $A_1, A_2, \dots \in \mathcal{A}$ is such that $A_1 \supset A_2 \supset \dots$ then $\bigcap_{i=1}^{\infty} A_i \in \mathcal{A}$.

Recall that an algebra on X is a collection of subsets of X that is closed under finite intersections, finite unions, complements and contains $\emptyset$ and X .

Theorem 3.2 (Monotone Class Lemma). Let X be a set and suppose that $\mathcal{A}$ is an algebra of subsets of X . Then the σ -algebra generated by $\mathcal{A}$ is the smallest monotone class that contains $\mathcal{A}$.

Proof. Let $\mathcal{M}$ denote the smallest monotone class that contains $\mathcal{A}$. We must show that $\mathcal{M}$ is a σ -algebra. It is enough to show that $\mathcal{M}$ is closed under finite unions and complements. For $A \in \mathcal{A}$, let

$$\mathcal{B}_A = \{B \in \mathcal{M} \mid B \cup A \in \mathcal{M}\}.$$

Note that $\mathcal{A} \subset \mathcal{B}_A$ by definition of algebra.

We claim that $\mathcal{B}_A$ is a monotone class. To see this, suppose that $B_1 \subset B_2 \subset \dots$ are all in $\mathcal{B}_A$. Thus $B_i \cup A \in \mathcal{M}$. But $B_i \cup A$ is a nested sequence of sets, and so $\bigcup_i B_i \cup A \in \mathcal{M}$ and so $\bigcup_i B_i \in \mathcal{M}$. Likewise suppose that $B_1 \supset B_2 \supset \dots$ are all in $\mathcal{B}_A$. Thus $B_i \cup A \in \mathcal{M}$. As $B_i \cup A$ monotonically decreases to $A \cup \bigcap_i B_i$ we have that $A \cup \bigcap_i B_i \in \mathcal{M}$ and so $\bigcap_i B_i \in \mathcal{B}_A$.

Thus $\mathcal{B}_A$ is a monotone class containing $\mathcal{A}$, but is a subset of $\mathcal{M}$ by definition. Thus $\mathcal{M} = \mathcal{B}_A$. Now let

$$\mathcal{M}_0 = \{M \in \mathcal{M} \mid M \cup M' \in \mathcal{M} \text{ for all } M' \in \mathcal{M}\}.$$

We see that $\mathcal{A} \subset \mathcal{M}_0$ since $\mathcal{M} = \mathcal{B}_A$. Note also that $\mathcal{M}_0$ is a monotone class. Thus $\mathcal{M}_0 = \mathcal{M}$. Thus we have shown that $\mathcal{M}$ is closed under finite unions, and hence countable unions.

Now let $\mathcal{M}' = \{M \in \mathcal{M} \mid M^c \in \mathcal{M}\}$. Note that $\mathcal{A} \subset \mathcal{M}'$ and $\mathcal{M}'$ is a monotone class. Thus $\mathcal{M}' = \mathcal{M}$ and so $\mathcal{M}$ is closed under complements. Thus $\mathcal{M}$ is indeed a σ -algebra.

□

Example 3.3. Let $\mathcal{A}$ denote the set of all finite unions of half-open intervals $\bigcup_{i=1}^n [a_i, b_i)$ for some $n \geq 0$. Then $\mathcal{A}$ is an algebra, but not a σ -algebra. The monotone class lemma says that the smallest monotone class containing $\mathcal{A}$ is the Borel σ -algebra.

Now let $\mathcal{U}$ be the smallest set containing $\mathcal{A}$ that is closed under countable nested unions. We claim that each element in $\mathcal{U}$ is either empty or has positive Lebesgue measure. Let $\mathcal{U}'$ denote all $U \in \mathcal{U}$ with this property (U is either empty or positive measure). Clearly $\mathcal{A} \subset \mathcal{U}'$. Now let $U_1, U_2, \dots \in \mathcal{U}'$ and $U = \bigcup_i U_i$. If U has zero measure, then each U_i has zero measure and so each U_i empty and so U is empty. Thus $U \in \mathcal{U}'$. Thus $\mathcal{U} = \mathcal{U}'$. This means that $\mathcal{U}$ is not the Borel σ -algebra. This shows that in the definition of monotone class, we really do need closure under countable intersections as well otherwise the monotone class lemma is not true.

4. PRODUCT MEASURES AND FUBINI-TONELLI THEOREM

Let X, Y be measure spaces and let

$$\mathcal{R}(X, Y) = \{U \times V \mid U \subset X \text{ measurable}, V \subset Y \text{ measurable}\}$$

Denote the set of all rectangles with in $X \times Y$ with measurable projections onto the axes. Let μ and ν be two positive measures on X and Y respectively. Then we define the area

$$\text{Area}(U \times V) = \mu(U)\nu(V).$$

If $\mathcal{C} \subset \mathcal{R}$ is a countable collections of rectangles define the

$$\text{Area}(\mathcal{C}) = \sum_{C \in \mathcal{C}} \text{Area}(C).$$

We say that $\mathcal{C}$ covers $A \subset X \times Y$ if

$$A \subset \bigcup_{C \in \mathcal{C}} C.$$

Define the product outer-measure

$$(\mu \times \nu)(A) = \inf_{\mathcal{C} \text{ covers } A} \text{Area}(\mathcal{C}).$$

Proposition 4.1. Each element of $\mathcal{R}$ is $\mu \times \nu$ -measurable.

Proof. Let $R \in \mathcal{R}$ and let $B \subset X \times Y$. Let $\mathcal{C}$ be a countable collection of rectangles that covers B such that

$$\text{Area}(\mathcal{C}) \leq (\mu \times \nu)(B) + \epsilon.$$

Observe that the complement R^c can be written as a disjoint union of finitely many rectangles. More precisely, if $R = U \times V$ then

$$R^c = U^c \times V^c \cup U^c \times V \cup U \times V^c.$$

Call these three rectangles R_1, R_2, R_3 . Observe that any rectangle $C = C_1 \times C_2 \in \mathcal{C}$ we have that

$$\begin{aligned}
& \text{Area}(C \cap R) + \text{Area}(C \cap R_1) + \text{Area}(C \cap R_2) + \text{Area}(C \cap R_3) \\
&= \mu(C_1 \cap U)\nu(C_2 \cap V) + \mu(C_1 \cap U^c)\nu(C_2 \cap V^c) + \mu(C_1 \cap U^c)\nu(C_2 \cap V) + \mu(C_1 \cap U)\nu(C_2 \cap V^c) \\
&= (\mu(C_1 \cap U) + \mu(C_1 \cap U^c))(\nu(C_2 \cap V) + \nu(C_2 \cap V^c)) \\
&= \mu(C_1)\nu(C_2) \\
&= \text{Area}(C).
\end{aligned}$$

Thus if we let

$$\mathcal{C}_2 = \{C \cap R_i \mid i = 1, 2, 3 \text{ and } C \in \mathcal{C}\}$$

and

$$\mathcal{C}_1 = \{C \cap R \mid C \in \mathcal{C}\}$$

then

$$\text{Area}(\mathcal{C}_1) + \text{Area}(\mathcal{C}_2) = \text{Area}(\mathcal{C}).$$

Observe that $\mathcal{C}_1$ covers $B \cap R$ and $\mathcal{C}_2$ covers $B \cap R^c$ and so

$$(\mu \times \nu)(R \cap B) + (\mu \times \nu)(R^c \cap B) \leq \text{Area}(\mathcal{C}_1) + \text{Area}(\mathcal{C}_2) = \text{Area}(\mathcal{C}) \leq (\mu \times \nu)(B) + \epsilon.$$

By letting $\epsilon \rightarrow 0$, the proof is complete. $\square$

We let $\mathcal{B}_{X \times Y}$ denote the smallest σ -algebra on $X \times Y$ containing $\mathcal{R}$. Note that this Proposition and Theorem 2.2 says that $\mu \times \nu$ is a measure (countably sub-additive) on this σ -algebra.

Another convenient definition of $\mu \times \nu$ is as follows. Let $\mathcal{C}$ is a countable collection of rectangles, then define the positive function

$$F_{\mathcal{C}} : X \times Y \rightarrow [0, \infty]$$

to be

$$F_{\mathcal{C}}(a) = \sum_{C \in \mathcal{C}} \mathbf{1}_C(a).$$

Then notice that $\mathcal{C}$ covers A if and only if $\mathbf{1}_A \leq F_{\mathcal{C}}$. Now consider the function $X \rightarrow [0, \infty]$ given by

$$x \mapsto \int_Y F_{\mathcal{C}}(x, y) d\nu(y).$$

Let us show that this function is well defined and measurable. If $F_{\mathcal{C}} = \sum_{i=1}^{\infty} \mathbf{1}_{U_i \times V_i}$ then

$$\int_Y F_{\mathcal{C}}(x, y) d\nu(y) = \int_Y \mathbf{1}_{U_i}(x) \mathbf{1}_{V_i}(y) d\nu(y) = \sum_{i=1}^{\infty} \mathbf{1}_{U_i}(x) \nu(V_i)$$

by the monotone convergence theorem. Thus this function is indeed measurable (by being the countable sum of non-negative measurable functions). By applying the monotone convergence theorem again we have the useful Fubini-type equation

$$\int_X \int_Y F_{\mathcal{C}}(x, y) d\nu(y) d\mu(x) = \sum_{i=1}^{\infty} \mu(U_i) \nu(V_i) = \text{Area}(\mathcal{C}).$$

Proposition 4.2. If $R \in \mathcal{R}$ is a rectangle, then $(\mu \times \nu)(R) = \text{Area}(R)$.

Proof. Suppose that $\mathcal{C}$ covers R . Then we must have that

$$F_{\{R\}} = \mathbf{1}_R \leq F_{\mathcal{C}}.$$

Thus we must have that

$$\text{Area}(R) = \int_X \int_Y F_{\{R\}}(x, y) d\nu(y) d\mu(x) \leq \int_X \int_Y F_{\mathcal{C}}(x, y) d\nu(y) d\mu(x) = \text{Area}(\mathcal{C}).$$

As $\mathcal{C}$ was arbitrary, this shows that

$$(\mu \times \nu)(R) \geq \text{Area}(R).$$

The reverse inequality holds because $\{R\}$ covers R . □

Theorem 4.3 (Fubini-Tonelli theorem for indicator functions). Suppose that μ and ν are σ -finite (on X and Y respectively). Let $B \in \mathcal{B}_{X \times Y}$. Then the mapping

$$x \mapsto \int_Y \mathbf{1}_B(x, y) d\nu(y)$$

is measurable and its integral over X is $(\mu \times \nu)(B)$, that is,

$$\int_X \int_Y \mathbf{1}_B(x, y) d\nu(y) d\mu(x) = (\mu \times \nu)(B) = \int_Y \int_X \mathbf{1}_B(x, y) d\mu(x) d\nu(y).$$

Proof. By a partitioning argument and the monotone convergence theorem, we can reduce to the case where μ and ν are both finite.

Let $\mathcal{M}$ denote the set of all $B \in \mathcal{B}_{X \times Y}$ for which this theorem is true. By the proposition above, $\mathcal{R} \subset \mathcal{M}$. Note that a finite union of rectangles may be written as a finite union of disjoint rectangles, thus the smallest algebra that contains $\mathcal{R}$ is a subset of $\mathcal{M}$. Thus, by the montone class lemma it will be enough to show that $\mathcal{M}$ is a montone class. That $\mathcal{M}$ is closed under countable nested unions follows from the monotone convergence theorem. To show that $\mathcal{M}$ is closed under countable nested intersections, we can use the fact that $\mathcal{M}$ is closed under complements, which uses the finiteness of μ and ν as follows. Indeed if $B \in \mathcal{M}$, then

$$\int_X \int_Y \mathbf{1}_{B^c}(x, y) d\nu(y) d\mu(x) = \int_X \int_Y (1 - \mathbf{1}_B(x, y)) d\nu(y) d\mu(x) = (\mu \times \nu)(X \times Y) - (\mu \times \nu)(B) = (\mu \times \nu)(B^c)$$

and so $B^c \in \mathcal{M}$. □

Example 4.4. The σ -finiteness assumption is necessary. For instance, let $X, Y = \mathbb{R}$ and let μ be the counting measure (not σ -finite) and let ν be the Lebesgue measure and let $B = \{(x, x) \mid x \in \mathbb{R}\}$ be the diagonal. Then

$$\int_X \int_Y \mathbf{1}_B(x, y) d\nu(y) d\mu(x) = \int_X 0 d\mu(x) = 0$$

and yet

$$\int_Y \int_X \mathbf{1}_B(x, y) d\mu(x) d\nu(y) = \int_Y 1 d\nu(y) = \infty.$$

By using the monotone convergence theorem, we get the Fubini-Tonelli theorem for positive measurable functions.

Theorem 4.5 (Tonelli theorem). Suppose that μ and ν are σ -finite (on X and Y respectively) and suppose that $f : X \times Y \rightarrow [0, \infty]$ is $\mathcal{B}_{X \times Y}$ -measurable. Then

$$x \mapsto \int_Y f(x, y) d\nu(y)$$

is measurable and

$$\int_X \int_Y f(x, y) d\nu(y) d\mu(x) = \int_{X \times Y} f(x, y) d(\mu \times \nu)(x, y) = \int_Y \int_X f(x, y) d\mu(x) d\nu(y).$$

Proof. Write $f = \sum_{i=1}^{\infty} c_i \mathbb{1}_{B_i}$ for some $B_i \in \mathcal{B}_{X \times Y}$ and $c_i \in [0, \infty)$. Now apply the previous theorem and the monotone convergence theorem. $\square$

Theorem 4.6. Suppose that μ and ν are σ -finite (on X and Y respectively) and suppose that $f : X \times Y \rightarrow [-\infty, \infty]$ is $\mathcal{B}_{X \times Y}$ -measurable. Suppose that¹

$$\int_{X \times Y} |f(x, y)| d(\mu \times \nu)(x, y) < \infty.$$

Then there exists a measurable set $X' \subset X$ that is μ -conull and such that

$$x \mapsto \int_Y |f(x, y)| d\nu(y) < \infty \quad \text{for all } x \in X'$$

and the map $X' \rightarrow X$ given by

$$x \mapsto \int_Y f(x, y) d\nu(y)$$

is a well defined measurable map. Furthermore²

$$\int_X \int_Y f(x, y) d\nu(y) d\mu(x) = \int_{X \times Y} f(x, y) d(\mu \times \nu)(x, y) = \int_Y \int_X f(x, y) d\mu(x) d\nu(y).$$

Proof. As f is L^1 with respect to $\mu \times \nu$, we can write $f = f^+ - f^-$ where $f^+, f^- : X \times Y \rightarrow [0, \infty]$ and f^+, f^- are both L^1 . Thus by an application of Tonelli's theorem to both of these functions we have that

$$\int_X \int_Y f^{\pm}(x, y) d\nu(y) d\mu(x) < \infty$$

and so there must be a co-null set $X' \subset X$ so that

$$\int_Y f^{\pm}(x, y) d\nu(y) < \infty \quad \text{for all } x \in X'$$

and this quantity is a measurable function on X' . Thus

$$\begin{aligned} \int_{X'} \int_Y f(x, y) d\nu(y) d\mu(x) &= \int_{X'} \int_Y (f^+(x, y) - f^-(x, y)) d\nu(y) d\mu(x) \\ &= \int_{X'} \left(\int_Y f^+(x, y) d\nu(y) - \int_Y f^-(x, y) d\nu(y) \right) d\mu(x) \\ &= \int_{X'} \int_Y f^+(x, y) d\nu(y) d\mu(x) - \int_{X'} \int_Y f^-(x, y) d\nu(y) d\mu(x) \\ &= \int_X \int_Y f^+(x, y) d\nu(y) d\mu(x) - \int_X \int_Y f^-(x, y) d\nu(y) d\mu(x) \\ &= \int_{X \times Y} f^+(x, y) d(\mu \times \nu)(x, y) - \int_{X \times Y} f^-(x, y) d(\mu \times \nu)(x, y) \\ &= \int_{X \times Y} f(x, y) d(\mu \times \nu)(x, y) \end{aligned}$$

$\square$

¹By Tonelli's theorem, we can replace this condition with an iterated integral for $|f|$.

²The integrands are only almost everywhere well defined. For example the first integral is over X' rather than X , but as X' is co-null we can still sensibly integrate over X by extending the domain to whichever function on X we like.

5. JENSEN'S INEQUALITY

Definition 5.1. We say $U \subset \mathbb{R}^d$ is convex if

$$tu + (1 - t)v \in U$$

whenever $u, v \in U$ and $t \in [0, 1]$. We say that $\varphi : U \rightarrow \mathbb{R}$ is convex if

$$\varphi(tu + (1 - t)v) \leq t\varphi(u) + (1 - t)\varphi(v)$$

for all $u, v \in U$ and $t \in [0, 1]$. Note that this is equivalent to saying that the region

$$\{(u, y) \in U \times \mathbb{R} \mid y \geq \varphi(u)\}$$

is convex.

Proposition 5.2. Let $U \subset \mathbb{R}^d$ be an open convex set and suppose that $\varphi : U \rightarrow \mathbb{R}$ is convex. Then φ is continuous.

Remark 5.3. We really need the assumption that U is open. For example, if $U = [0, \infty)$ then the function $\varphi(u) = \mathbb{1}_{\{0\}}(u)$ which vanishes every but equals 1 for $u = 0$ is convex but not continuous.

Proof. We prove this by induction on d . The base case $d = 0$ is trivial (where $U = \{0\} = \mathbb{R}^0$ is a zero-dimensional vector space). Now fix $x_0 \in U$. Let $U' \subset U$ be a cube contained in U such that x_0 is the center of U' . Let B be the boundary of this cube U' . Observe that by the induction hypothesis, φ is continuous on B (as B is made up of finitely many $d - 1$ -dimensional faces) and thus is bounded on B , say $|\varphi(b)| \leq M$ for all $b \in B$. Observe also that any $u \in U'$ may be written as $u = tx_0 + (1 - t)b$ for some $b \in B$ (to see this, note that any ray starting at x_0 that goes through u must intersect B at some point b). Now observe that

$$\|u - x_0\| = \|(t - 1)x_0 + (1 - t)b\| = (1 - t)\|b - x_0\| \geq (1 - t)R$$

where $R > 0$ is half the side-length of the box U' . Now by convexity we have that

$$\begin{aligned} \varphi(u) &= \varphi(tx_0 + (1 - t)b) \leq t\varphi(x_0) + (1 - t)\varphi(b) \\ &\leq t\varphi(x_0) + (1 - t)M \\ &\leq t\varphi(x_0) + MR^{-1}\|u - x_0\| \\ &= \varphi(x_0) + (t - 1)\varphi(x_0) + MR^{-1}\|u - x_0\| \\ &\leq \varphi(x_0) + R^{-1}(|\varphi(x_0)| + M)\|u - x_0\|. \end{aligned}$$

Thus

$$\limsup_{u \rightarrow x_0} \varphi(u) \leq \varphi(x_0).$$

Now suppose for contradiction, that there is some $u_1, u_2, \dots \in U$ such that $u_i \rightarrow x_0$ but $\varphi(u_i) < \varphi(x_0) - \epsilon$ for some fixed $\epsilon > 0$. Now observe that

$$x_0 = t_i u_i + (1 - t_i) b_i$$

for some $t_i \in [0, 1]$ and $b_i \in B$ as we can see by considering a ray that starts at u_i and passes through x_0 and then letting $b_i \in B$ be a point on this ray. We see that $t_i \rightarrow 1$ since

$$\|x_0 - u_i\| = \|(t_i - 1)u_i + (1 - t_i)b_i\| = (1 - t_i)\|b_i - u_i\|$$

and the left hand side converges to zero while $\|b_i - u_i\| \rightarrow \|b_i - x_0\| > 0$. Now observe that

$$\varphi(x_0) \leq t_i \varphi(u_i) + (1 - t_i) \varphi(b_i) \leq t_i (\varphi(x_0) - \epsilon) + (1 - t_i) M.$$

Since $t_i \rightarrow 1$, this upper bound converges to $\varphi(x_0) - \epsilon$ which is a contradiction. $\square$

Proposition 5.4. Let $U \subset \mathbb{R}^d$ be a convex set. Let $t_1, \dots, t_n \in [0, 1]$ such that $t_1 + \dots + t_n = 1$. We have

$$\sum_{i=1}^n t_i u_i \in U$$

for all $u_1, \dots, u_n \in U$. Moreover, if $\varphi : U \rightarrow \mathbb{R}$ is convex then

$$\varphi \left(\sum_{i=1}^n t_i u_i \right) \leq \sum_{i=1}^n t_i \varphi(u_i).$$

Proof. The $n = 2$ case is the definition of convexity. We proceed by induction on n . Write $t = t_1 + \dots + t_{n-1}$. The result is clearly true if $t = 0$, as in that case $t_n = 1$, so we assume $t > 0$. Observe that

$$\sum_{i=1}^n t_i u_i = t \sum_{i=1}^{n-1} t^{-1} t_i u_i + (1 - t) u_n$$

is indeed in U since by the inductive hypothesis,

$$\sum_{i=1}^{n-1} t^{-1} t_i u_i \in U$$

as $\sum_{i=1}^{n-1} t^{-1} t_i = 1$. Likewise by the induction hypothesis we have

$$\begin{aligned} \varphi \left(\sum_{i=1}^n t_i u_i \right) &= \varphi \left(t \sum_{i=1}^{n-1} t^{-1} t_i u_i + (1 - t) u_n \right) \\ &\leq t \varphi \left(\sum_{i=1}^{n-1} t^{-1} t_i u_i \right) + (1 - t) \varphi(u_n) \\ &\leq t \sum_{i=1}^{n-1} t^{-1} t_i \varphi(u_i) + (1 - t) \varphi(u_n) \\ &= \sum_{i=1}^n t_i \varphi(u_i). \end{aligned}$$

$\square$

Note that if $v : X \rightarrow \mathbb{R}^d$ is a vector valued function on a probability space (X, μ) then we can define its integral $\int_X v(x) d\mu(x)$ componentwise, assuming each component is measurable and has a well defined integral (e.g, each component is positive or L^1). If

$$\int_X \|v\| d\mu(x) < \infty$$

then we say that v is L^1 . In this case, each component is L^1 and the integral of v makes sense.

Proposition 5.5 (Compact Jensen's inequality). Let μ be a probability measure supported on a compact subset $K \subset \mathbb{R}^d$ and suppose $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$ is convex. Then

$$\varphi \left(\int_K x d\mu(x) \right) \leq \int_K \varphi(x) d\mu(x).$$

Proof. Note that if μ is supported on a finite set $\{u_1, \dots, u_n\}$ with $\mu(\{u_i\}) = p_i$, then by the inequality above we have that

$$\varphi\left(\int_K x d\mu(x)\right) = \varphi\left(\sum_{i=1}^n p_i u_i\right) \leq \sum_{i=1}^n p_i \varphi(u_i) = \int_K \varphi(x) d\mu(x).$$

Thus the result holds for finite measures. For arbitrary probability measures on K , we use the proposition below and the continuity of ϕ and $x \mapsto x$. $\square$

Proposition 5.6. Let μ be a probability measure on a compact set $[-R, R]^d \subset \mathbb{R}^d$. For each positive integer n , let $\mathcal{C}_n$ be a partition of $[-R, R]^d$ into finitely many cubes, i.e.,

$$[-R, R]^d = \bigcup_{C \in \mathcal{C}_n} C$$

such that each $C \in \mathcal{C}_n$ has diameter at most $1/n$. Let $C_n \subset [-R, R]^d$ be a collection of points consisting of exactly one element from $\mathcal{C}_n$, thus we have a bijection $\iota : \mathcal{C}_n \rightarrow C_n$ where $c \in \iota(C)$. Now define a probability measure on C_n given by $\mu_n(\{c\}) = \mu(\iota(C))$ for $c \in C_n$. Then

$$\int_{\mathbb{R}^d} f d\mu_n \rightarrow \int_{\mathbb{R}^d} f d\mu$$

for all continuous functions $f : [-R, R]^d \rightarrow \mathbb{R}$.

We now obtain immediately Jensen's inequality for integrable random variables. In this section, we fix compact sets $K_1 \subset K_2 \subset \dots$ such that

$$\bigcup_n K_n = \mathbb{R}^d.$$

Theorem 5.7 (Jensen's inequality). Let μ be a probability measure on $\mathbb{R}^d$ such that

$$\int_{\mathbb{R}^d} \|x\| d\mu(x) < \infty$$

and suppose $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$ is convex. Then

$$\varphi\left(\int_{\mathbb{R}^d} x d\mu(x)\right) \leq \liminf_{n \rightarrow \infty} \int_{K_n} \varphi(x) d\mu(x).$$

Proof. For large enough n , we have $\mu(K_n) > 0$. By applying the Compact Jensen's inequality to the probability measure $\mu_n = \mu|_{K_n}$, which is defined by $\mu_n(A) = \mu(K_n)^{-1} \mu(A \cap K_n)$, we have

$$\varphi\left(\mu(K_n)^{-1} \int_{K_n} x d\mu(x)\right) \leq \mu(K_n)^{-1} \int_{K_n} \varphi(x) d\mu(x).$$

Now as $n \rightarrow \infty$ we have $\mu(K_n) \rightarrow 1$ and so the continuity of φ completes the proof. $\square$

Another way of stating this is as follows.

Theorem 5.8. Let ν be a probability measure on a space Ω . Suppose that $f : \Omega \rightarrow \mathbb{R}^d$ is L^1 , i.e.,

$$\int_{\mathbb{R}^d} \|f(\omega)\| d\nu(\omega) < \infty.$$

Suppose that $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$ is convex. Then

$$\varphi\left(\int_{\Omega} f(\omega) d\nu(\omega)\right) \leq \liminf_{N \rightarrow \infty} \int_{f^{-1}(K_N)} \varphi(f(\omega)) d\nu(\omega).$$

In particular, if $\varphi \geq 0$ or $\varphi \circ f \in L^1(\Omega)$ then

$$\varphi \left(\int_{\Omega} f(\omega) d\nu(\omega) \right) \leq \int_{\Omega} \varphi(f(\omega)) d\nu(\omega).$$

6. HAUSDORF MEASURE

6.1. Definition and Borel measurability.

Definition 6.1. Fix $s \geq 0$. Let $\alpha_s > 0$ be chosen such that α_s is the volume of the unit s -ball. For $A \subset \mathbb{R}^d$ (not necessarily Borel) we define

$$\mathcal{H}_{\delta}^s(A) = \inf \left\{ \sum_{i=1}^{\infty} \alpha_s \left(\frac{1}{2} \text{diam}(C_i) \right)^s \mid A \subset \bigcup_{i=1}^{\infty} C_i \text{ for some } C_i \subset \mathbb{R}^d \text{ with } \text{diam}(C_i) \leq \delta \right\}.$$

We define the s -dimensional Hausdorff measure to be

$$\mathcal{H}^s(A) = \lim_{\delta \rightarrow 0} \mathcal{H}_{\delta}^s(A),$$

where this limit exists as the sequence is (non-strictly) increasing.

In other words $\mathcal{H}_{\delta}^s(A)$ measures the optimal way of covering A with sets C_i of diameter at most δ , where the cost of using such a set is $\alpha_s(\text{diam}(C_i)/2)^s$ (i.e., the s -dimensional volume of the unit s -ball of the same diameter). We remark that that C_i can always be taken to be closed (thus Borel) as taking the closure does not change the diameter.

Lemma 6.2. If $B \subset \mathbb{R}^n$, with $n < d$ and let $\hat{B} = B \times \{0\}^{d-n} \subset \mathbb{R}^d$ be the embedding of $\hat{B}$ in $\mathbb{R}^d$. Then $\mathcal{H}^s(\hat{B}) = \mathcal{H}^s(B)$ for all $s \geq 0$.

Proof. This is clear since any cover of $\hat{B}$ can be intersected with the subspace to obtain a cover with sets of diameter no larger. $\square$

Example 6.3. Note that $\mathcal{H}^0$ is the cardinality and $\mathcal{H}^1$ on $\mathbb{R}$ coincides with the Lebesgue measure $\mathcal{L}^1$. Thus as $\mathcal{H}^1$ is rotation invariant we see, by the previous Lemma, that if $L \subset \mathbb{R}^d$ is a line then $\mathcal{H}^1(B)$, for $B \subset L$, is the 1-dimensional Lebesgue measure of $\iota(B)$, where $\iota : L \rightarrow \mathbb{R}$ is an isometric isomorphism. In other words, $\mathcal{H}^1$ corresponds to length.

It is clear the $\mathcal{H}^s$ is an outer measure on $\mathbb{R}^d$.

Definition 6.4. Let μ be an outer measure on a metric space X . We say that μ is a metric outer measure if it has the property that whenever $A_1, A_2 \subset X$ are such that $d(A_1, A_2) > 0$, then $\mu(A_1 \cup A_2) = \mu(A_1) + \mu(A_2)$.

Note that the Hausdorff measure is a metric outer measure as can be seen by the following argument. If the distance between $A_1, A_2 \subset \mathbb{R}^d$ is strictly greater than $\epsilon > 0$, then any set $C \subset \mathbb{R}^d$ of diameter at most ϵ cannot intersect both A_1 and A_2 . Thus for $0 < \delta < \epsilon$ we have that

$$\mathcal{H}_{\delta}^s(A_1 \cup A_2) = \mathcal{H}_{\delta}^s(A_1) + \mathcal{H}_{\delta}^s(A_2).$$

Lemma 6.5. Let μ be a metric outer measure on a metric space X . Suppose that $A \subset X$ is such that $\mu(A) < \infty$ and suppose that $A_1 \subset A_2 \subset \dots$ are such that $A = \bigcup_{n \geq 1} A_n$ and

$$d(A \setminus A_{n+1}, A_n) > 0$$

for all $n \geq 1$. Then

$$\lim_{n \rightarrow \infty} \mu(A_n) = \mu(A).$$

Proof. Let $B_n = A_{n+1} \setminus A_n$. Now observe that $B_1, B_3, \dots, B_{2N-1}$ are all disjoint sets and the distance between any two is non-zero. Thus by definition of metric measure we have that

$$\sum_{i=1}^N \mu(B_{2i-1}) = \mu\left(\bigcup_{i=1}^N B_{2i-1}\right) \leq \mu(A) < \infty.$$

The same argument holds for even indexes. Thus

$$\sum_{n=1}^{\infty} \mu(B_n) < \infty.$$

Now

$$\mu(A \setminus A_n) = \sum_{i=n+1}^{\infty} \mu(B_i) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus from $\mu(A) \leq \mu(A \setminus A_n) + \mu(A_n)$ we see that

$$\mu(A) \leq \lim_{n \rightarrow \infty} \mu(A_n).$$

This completes the proof as $\mu(A_n) \leq \mu(A)$ holds. $\square$

Theorem 6.6. Let μ be a metric outer measure on a metric space X . Then each Borel set is μ -measurable.

Proof. We will show that all closed sets are Borel measurable and this will be enough since the μ -measurable sets form a sigma algebra.

Thus let $C \subset X$ be closed. We must show that for all $A \subset X$ we have that

$$\mu(A) \geq \mu(A \cap C) + \mu(A')$$

where $A' = A \setminus C$. This holds if $\mu(A) = \infty$, so we assume $\mu(A) < \infty$.

Now let

$$A_n = \{a \in A \mid d(a, C) > \frac{1}{n}\}.$$

Now that $A_n \subset A_{n+1}$ and as C is closed, we have that

$$\bigcup_{n=1}^{\infty} A_n = A'.$$

Note also that if $x \in A_n$ and $y \in A' \setminus A_{n+1}$ then we must have

$$d(x, y) \geq \frac{1}{n} - \frac{1}{n+1}$$

which can be seen as follows: We have $d(y, C) \leq 1/(n+1)$ and so there exists $c \in C$ so that

$$d(y, c) < \frac{1}{n+1} + \epsilon.$$

However

$$\frac{1}{n} < d(x, c) \leq d(x, y) + d(y, c) \leq d(x, y) + \frac{1}{n+1} + \epsilon.$$

Thus we have $d(A_n, A' \setminus A_{n+1}) > 0$, so the previous Lemma applies. This means that

$$\mu(A \cap C) + \mu(A') = \mu(A \cap C) + \lim_{n \rightarrow \infty} \mu(A_n) = \lim_{n \rightarrow \infty} \mu((A \cap C) \cup \mu(A_n)) \leq \mu(A),$$

where we use the identity $\mu((A \cap C) \cup \mu(A_n)) = \mu(A \cap C) + \mu(A_n)$ since $d(A \cap C, A_n) \geq 1/n > 0$ and μ is a metric measure. $\square$

Thus the Hausdorff measure forms a Borel measure (i.e., it is σ -additive on Borel sets).

6.2. Equality of Hausdorff and Lebesgue measure when $s = d$.

Proposition 6.7. For (not necessarily Borel) $B \subset \mathbb{R}^d$ we have

$$\mathcal{H}^d(B) = \mathcal{L}_d(B),$$

i.e., the d -dimensional Hausdorff measure is equal to the d -dimensional Lebesgue measure. In fact, $\mathcal{H}^d(B) = \mathcal{H}_\delta^d(B)$ for all $\delta > 0$.

Note that we have already shown the $d = 1$ case in Exampel 6.3. To prove this for $d \geq 1$, we need some results about covering open sets with balls.

Lemma 6.8. Let $\mathcal{C}$ be a collection of (open or closed) balls in $\mathbb{R}^d$ with non-zero radius. Then there exists $\mathcal{C}' \subset \mathcal{C}$ such that $\mathcal{C}'$ is a maximal disjoint subcollection of $\mathcal{C}$ (i.e., the elements of $\mathcal{C}'$ are disjoint and for and $B \in \mathcal{C} \setminus \mathcal{C}'$ we have that $B \cap B' \neq \emptyset$ for some $B' \in \mathcal{C}'$). Moreover, any such $\mathcal{C}'$ must be countable.

Proof. To see that any such $\mathcal{C}'$ is countable, note that each element of $\mathcal{C}'$ contains some rational point (point in $\mathbb{Q}^d$). But there are only finitely many such rational points and no two such balls can share such a point.

To show existence of $\mathcal{C}'$, we can use Zorn's lemma. But let us avoid that. We recursively select a (finite or countable) sequence of balls $B_1, B_2, \dots \in \mathcal{C}$ as follows. Let us order the rational points $q_1, q_2, \dots \in \mathbb{Q}^d$. Let $i(B) = \min\{i \mid q_i \in B\}$. We recursively let B_n be chosen so that

$$i(B_n) = \min\{i(B) \mid B \in \mathcal{C} \setminus \mathcal{B}_{n-1} \text{ and } B \cap B' = \emptyset \text{ for all } B' \in \mathcal{B}_{n-1}\}$$

where $\mathcal{B}_0 = \emptyset$ and $\mathcal{B}_n = \{B_1, \dots, B_n\}$. That is, we keep selecting B_n so that it is disjoint from $B_1, \dots, B_{n-1}$ and $i(B_n)$ is minimal. Note that $i(B_n)$ is strictly increasing sequence. We claim that $\mathcal{C}' = \{B_1, B_2, \dots\}$ is maximal disjoint. This is clear if $\mathcal{C}'$ is finite. Thus suppose $\mathcal{C}'$ is infinite and not maximal disjoint. Hence there exists $B \in \mathcal{C}$ so that $B \cap B' = \emptyset$ for all $B' \in \mathcal{C}'$. Now let q_i be so that $i = i(B)$. As $i(B_k)$ is strictly increasing in k , we have that $i(B_{i+1}) > i$, which violates the choice of B_{i+1} . $\square$

If $B = B(x, r)$ is an open ball, let $D_5 B = B(x, 5r)$ denote the ball with the same centre but 5 times the radius.

Theorem 6.9 (Vitali's covering Lemma). Suppose that $\mathcal{C}$ is a collection of non-empty open balls in $\mathbb{R}^d$ such that

$$D := \sup\{\text{diam}(B) \mid B \in \mathcal{C}\} < \infty$$

and let $U = \bigcup_{B \in \mathcal{C}} B$. Then there exists a **disjoint** subcollection $\mathcal{C}' \subset \mathcal{C}$ so that

$$U \subset \bigcup_{B \in \mathcal{C}'} D_5 B.$$

Moreover, such a $\mathcal{C}'$ must be countable.

Proof. Let

$$\mathcal{F}_j = \{B \in \mathcal{C} \mid 2^{-j}D < \text{diam}(B) \leq 2^{-j+1}D\}$$

for $j = 1, 2, \dots$. Note that the $\mathcal{F}_j$ partition $\mathcal{C}$. Now let $\mathcal{C}'_1$ be a maximal disjoint collection of $\mathcal{F}_1$ and for $j > 1$ we define recursively $\mathcal{C}'_j$ to be a maximal disjoint collection of elements of

$$\{B \in \mathcal{F}_j \mid B \cap B' = \emptyset \text{ for all } B' \in \bigcup_{i=1}^{j-1} \mathcal{C}'_i\}.$$

Now let

$$\mathcal{C}' = \bigcup_{j=1}^{\infty} \mathcal{C}'_j.$$

Clearly the elements of $\mathcal{C}'$ are disjoint. Now let $B \in \mathcal{C} \setminus \mathcal{C}'$. Then $B \in \mathcal{F}_j$ for some j . As $B \notin \mathcal{C}_j$ this means that either B intersects non-trivially some element in $\bigcup_{i=1}^{j-1} \mathcal{C}'_i$ or it intersects non-trivially some element of $\mathcal{C}'_j$. In either case, it intersects non-trivially some element B' of $\bigcup_{i=1}^j \mathcal{C}'_i$. This means that

$$\text{diam}(B') > 2^{-j}D = \frac{1}{2}2^{-j+1}D \geq \frac{1}{2}\text{diam}(B).$$

Thus $B \subset D_5B'$. Hence as B was arbitrary, this shows that

$$U \subset \bigcup_{B \in \mathcal{C}'} D_5B.$$

□

Our next result shows that we can almost write any open set as a countable disjoint union of balls. Here almost means that the union of these balls is a subset of our open set and the remainder has measure zero.

Theorem 6.10. Let $U \subset \mathbb{R}^d$ be an open set and $\delta < 0$. Then there exists a disjoint (hence countable) collection $\mathcal{C}$ of open balls in $\mathbb{R}^d$ such that $\text{diam}(B) < \delta$ and $B \subset U$ for each $B \in \mathcal{C}$ and

$$\mathcal{L}_d\left(U \setminus \bigcup_{B \in \mathcal{C}} B\right) = 0.$$

Proof. We reduce to the case where U is bounded and hence has $\mathcal{L}_d(U) < \infty$ since, after deleting a set of measure 0, we can partition any such U into bounded open sets. For example, after removing the sphere $\{u \in U \mid |u| = n\}$ for each $n \in \mathbb{Z}$.

Let $\mathcal{C}_0$ be the collection of non-empty open balls that are subsets of U and have diameter at most δ . Applying the Vitali covering lemma to this, we can choose a subset $\mathcal{C}_1 \subset \mathcal{C}_0$ of disjoint balls so that

$$V_1 = \bigcup_{B \in \mathcal{C}_1} D_5B \supset \bigcup_{B \in \mathcal{C}_0} B = U.$$

Let

$$W_1 = \bigcup_{B \in \mathcal{C}_1} B.$$

Thus

$$\mathcal{L}_d(U) \leq \mathcal{L}_d(V_1) \leq \sum_{B \in \mathcal{C}_1} 5^d \mathcal{L}_d(B) = 5^d \mathcal{L}_d(W_1).$$

This means that if we let $\alpha = 5^{-d}$ then

$$\mathcal{L}_d(W_1) \geq \alpha \mathcal{L}_d(U).$$

Now fix $0 < \beta < \alpha$. This means that we can choose a finite subset $\mathcal{F}_1 \subset \mathcal{C}_1$ such that $\mathcal{L}_d(F_1) \geq \beta \mathcal{L}_d(U)$ where

$$F_1 = \bigcup_{B \in \mathcal{F}_1} \overline{B}.$$

Now let $U_1 = U \setminus F_1$ and note that U_1 is open (as F_1 , being the union of finitely many closed balls is closed). However

$$\mathcal{L}_d(U_1) = \mathcal{L}_d(U) - \mathcal{L}_d(F_1) \leq (1 - \beta) \mathcal{L}_d(U).$$

Thus we can repeat the same procedure to U_1 in place of U and find a finite set $\mathcal{F}_2$ of open balls in U_1 of diameter at most δ such that if we delete the closure of their union then we get a set $U_2 \subset U_1$ of measure

$$\mathcal{L}_d(U_2) \leq (1 - \beta)\mathcal{L}_d(U_1) \leq (1 - \beta)^2\mathcal{L}_d(U).$$

Repeating this procedure, we can finally let

$$\mathcal{C} = \bigcup_{n=1}^{\infty} \mathcal{F}_n.$$

Note that this is a disjoint collection by construction. The complement of the union of all $\mathcal{C}$ has measure 0 since

$$\mathcal{L}_d(U_n) \leq \beta^n \mathcal{L}_d(U) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$


□

Definition 6.11 (Steiner symmetrization). Let $e_1, \dots, e_d$ be the standard basis for $\mathbb{R}^d$ and let $k \in \{1, \dots, n\}$. Let $\pi_k : \mathbb{R}^d \rightarrow \mathbb{R}^d$ denote the orthogonal projection onto the subspace spanned by $\{e_i \mid i \neq k, 1 \leq i \leq n\}$, thus the image of π_k is the orthogonal complement of $\mathbb{R}e_k$. We define the e_k -Steiner symmetrization of $B \subset \mathbb{R}^d$ to be equal to

$$S_k(B) = \bigcup_{u \in \pi_k(B)} \{u + te_k \mid |t| \leq \frac{1}{2} \mathcal{H}^1(B \cap \pi_k^{-1}(u))\}.$$

In other words, the Steiner-Symmetrization is a union of line segments $\{u + te_k \mid |t| < \frac{1}{2}T_u\}$ parallel to e_k where each line segment has its centre u on the plane $\pi_k(\mathbb{R}^d)$ and the length of each line segment is equal to the Hausdorff measure of the set $B \cap \pi_k^{-1}(u)$, i.e., the set of those points on B that lie on the line $\{u + te_k \mid t \in \mathbb{R}\}$. If B is convex, there is a simpler definition of the Steiner-Symmetrization of B : one simply partitions B into line segments parallel to e_k and then one translates each such line segment in the e_k direction so that each line segment is symmetric about the plane $\pi_k(\mathbb{R}^d)$.

FIGURE 1. The Steiner-Symmetrization of a polygon B . Notice that each vertical cross section of B , given by $\pi_k^{-1}(u) \cap B$, has the same length as the vertical cross section of $S_k(B)$ passing through u .



Proposition 6.12. If $B \subset \mathbb{R}^d$ is bounded and Borel, then so is $S_k(B)$. Moreover,

$$\mathcal{L}_d(B) = \mathcal{L}_d(S_k(B))$$

and

$$\text{diam}(S_k(B)) \leq \text{diam}(B).$$

Proof. We can just focus on the case $d = k$ by symmetry. We can write $\mathcal{L}_d$ as a product measure $\mathcal{L}_{d-1} \times \mathcal{L}_1$ and use Fubini's theorem (and Example 6.3 which says that $\mathcal{H}^1$ corresponds to 1-dimensional Lebesgue measure on lines) to get that the map $F: \mathbb{R}^{d-1} \rightarrow \mathbb{R}$ given by $F(u) = \mathcal{H}^1(B \cap (\{u\} \times \mathbb{R}))$ is Borel and

$$\mathcal{L}_d(B) = \int_{\mathbb{R}^{d-1}} F(u) d\mathcal{L}_{d-1}(u).$$

But

$$S_k(B) = \{(v, y) \in \mathbb{R}^{d-1} \times \mathbb{R} \mid |y| \leq \frac{1}{2}F(v)\}$$

is Borel as F is Borel. Applying Fubini's theorem to $S_k(B)$ we get the desired equality.

Let us now show the inequality involving diameter. Let $v_1 = u_1 + t_1 e_k, v_2 = u_2 + t_2 e_k \in S_k(B)$ where $u_1, u_2 \in \pi_k(B)$ and $|t_i| \leq \frac{1}{2}L_i$ where

$$L_i = \mathcal{H}^1(B \cap \pi_k^{-1}(u_i)).$$

Let m_i and M_i denote the infimum and supremum, respectively, of the set

$$\{y \in \mathbb{R} \mid u_i + y e_k \in B\}.$$

We have that

$$\|v_1 - v_2\|^2 = \|u_1 - u_2\|^2 + |t_1 - t_2|^2.$$

Now assume, WLOG, that

$$\alpha_1 := \frac{1}{2}(M_1 + m_1) \leq \alpha_2 := \frac{1}{2}(M_2 + m_2).$$

This means that $m_1 \leq \alpha_1 \leq \alpha_2 \leq M_2$ and so $M_2 - m_1 \geq 0$. Thus

$$\begin{aligned} M_2 - m_1 &= M_2 - \alpha_2 + \alpha_2 - m_1 \\ &\geq M_2 - \alpha_2 + \alpha_1 - m_1 \\ &= \frac{1}{2}(M_2 - m_2) + \frac{1}{2}(M_1 - m_1) \\ &\geq \frac{1}{2}L_1 + \frac{1}{2}L_2 \\ &\geq |t_1| + |t_2| \\ &\geq |t_1 - t_2| \end{aligned}$$

Now since $u_1 + M_1 e_k \in \bar{B}$ and $u_2 + m_2 e_k \in \bar{B}$ we thus get that

$$\begin{aligned} \text{diam}(B)^2 &= \text{diam}(\bar{B})^2 \\ &\geq \|u_1 + M_1 e_k - (u_2 + m_2 e_k)\|^2 \\ &= \|u_1 - u_2\|^2 + (M_2 - m_1)^2 \\ &\geq \|u_1 - u_2\|^2 + \|t_1 - t_2\|^2 \\ &= \|v_1 - v_2\|^2 \end{aligned}$$

Thus as v_1, v_2 were arbitrary, we have the desired bound $\text{diam}(B)^2 \geq \text{diam}(S_k(B))^2$. $\square$

Let $R_k : \mathbb{R}^d \rightarrow \mathbb{R}^d$ denote the reflection in the orthogonal complement of e_k . That is,

$$R_k(x_1, \dots, x_d) = (x_1, \dots, x_{k-1}, -x_k, x_{k+1}, \dots, x_d).$$

We say that a set $B \subset \mathbb{R}^d$ is e_k -symmetric, if $R_k B = B$.

Lemma 6.13. If $B \subset \mathbb{R}^d$ then $S_k(B)$ is e_k -symmetric. Moreover, if B is e_j -symmetric, for some $j \neq k$, then $S_k(B)$ is also e_j -symmetric.

Proof. That $S_k(B)$ is e_k -symmetric is clear from the definition. We now prove the second claim. Suppose B is e_j -symmetric with $j \neq k$.

Recall the definition,

$$S_k(B) = \bigcup_{u \in \pi_k(B)} \{u + te_k \mid |t| \leq \frac{1}{2} \mathcal{H}^1(B \cap \pi_k^{-1}(u))\}.$$

If $u \in \pi_k(B)$ then there exists $s \in \mathbb{R}$ such that $u + se_k \in B$. But now as B is e_j -symmetric, we have that B also contains $R_j(u + se_k) = R_j u + se_k$. Thus $R_j u \in \pi_k(B)$. Now we use the identity $\pi_k^{-1}(R_j u) = R_j(\pi_k^{-1}(u))$ to get that

$$B \cap \pi_k^{-1}(R_j u) = R_j B \cap R_j(\pi_k^{-1}(u)) = R_j(B \cap \pi_k^{-1}(u))$$

and thus

$$\mathcal{H}^1(B \cap \pi_k^{-1}(R_j u)) = \mathcal{H}^1(B \cap \pi_k^{-1}(u))$$

as $\mathcal{H}^1$ is invariant under R_j (as R_j is an isometry). This shows that if $u + te_k$ is in $S_k(B)$ then so is $R_j(u) + te_k = R_j(u + te_k)$, as required. $\square$

Theorem 6.14 (Isodiametric inequality). If $B \subset \mathbb{R}^d$ is Borel, then

$$\mathcal{L}_d(B) \leq \alpha_d \left(\frac{1}{2} \text{diam}(B) \right)^d.$$

That is, any Borel set has Lebesgue measure at least that equal to a ball of the same diameter.

Remark 6.15. It is not necessarily true that B can be covered by a ball of the same diameter. For example, let $B \subset \mathbb{R}^2$ be an equilateral triangle with sidelengths 1. Then $\text{diam}(B) = 1$, but the smallest disk that covers B has radius strictly greater than $\frac{1}{2}$.

Proof. Let $B' = S_1 S_2 \cdots S_k(B)$, i.e., B' is obtained by successively applying all the Steiner-Symmetrizations. We have that $\text{diam}(B') \leq \text{diam}(B)$ and $\mathcal{L}_d(B') = \mathcal{L}_d(B)$. We also have by the previous Lemma that B' is symmetric about 0. Thus if $x \in B'$ then $-x \in B'$ and $\text{diam}(B') \geq 2\|x\|$. Thus $\text{diam}(B') \geq 2 \sup\{\|x\| \mid x \in B'\}$ and thus B' may be covered by a disk of diameter $\text{diam}(B')$. Thus

$$\mathcal{L}_d(B') \leq \alpha_d \left(\frac{1}{2} \text{diam}(B') \right)^d \leq \alpha_d \left(\frac{1}{2} \text{diam}(B) \right)^d$$

which completes the proof as $\mathcal{L}_d(B') = \mathcal{L}_d(B)$. $\square$

Proof of Proposition 6.7. We first show the inequality

$$\mathcal{L}_d(B) \leq \mathcal{H}^d(B).$$

Let $C_1, C_2, \dots$ be compact (thus Borel) sets that cover B and each have diameter at most δ . We thus have by the Isodiametric inequality that

$$\mathcal{L}_d(B) \leq \sum_{i=1}^{\infty} \mathcal{L}_d(C_i) \leq \sum_{i=1}^{\infty} \alpha_d \left(\frac{1}{2} \text{diam}(C_i) \right)^d.$$

As the C_i are arbitrary, we get that $\mathcal{L}_d(B) \leq \mathcal{H}_\delta^d(B)$ holds for all $\delta > 0$.

Now we prove the reverse inequality. It is sufficient to prove it when $B = U$ is an open set (i.e., if μ is $\mathcal{H}_\delta^d$ or Lebesgue, then there is always an open set V containing B so that $\mu(B) \leq \mu(V) < \mu(B) + \epsilon$, thus $\mu(B)$ is the infimum of $\mu(V)$ over all such V).

For this, we know by the Theorem 6.10 that we can write

$$U = Q \cup \bigcup_{i=1}^{\infty} B_i$$

where each B_i is an open ball of radius at most δ and $\mathcal{L}_d(Q) = 0$. It follows that $\mathcal{H}_\delta^d(Q) = 0$ since Q can be covered with cubes of very small volume and thus balls of volume just a constant multiple of that, hence also arbitrarily small. In particular, this means that if we fix $\epsilon > 0$ then we can cover Q with balls $B'_1, B'_2, \dots$ of radius at most δ such that

$$\sum_{i=1}^{\infty} \mathcal{L}_d(B'_i) < \epsilon.$$

Thus

$$\mathcal{H}_\delta^d(U) \leq \sum_{i=1}^{\infty} \mathcal{L}_d(B_i) + \sum_{i=1}^{\infty} \mathcal{L}_d(B'_i) = \mathcal{L}_d(U) + \sum_{i=1}^{\infty} \mathcal{L}_d(B'_i) \leq \mathcal{L}_d(U) + \epsilon.$$

Taking $\epsilon \rightarrow 0$ completes the proof.

□

7. AREA FORMULA AND CHANGE OF VARIABLES IN INTEGRATION

Suppose $X \subset \mathbb{R}^n$ is open and $f : X \rightarrow \mathbb{R}^m$ with $n \leq m$ is continuously differentiable. We let $Df(x)$ denote the derivative of f at $x \in X$, which is a linear map $Df(x) : \mathbb{R}^n \rightarrow \mathbb{R}^m$ where

$$Df(x)u = \lim_{t \rightarrow 0} t^{-1}(f(x + tu) - f(x)).$$

Recall that to say that f is continuously differentiable is to say that $x \mapsto Df(x)$ is a continuous map in the operator norm on the space of linear maps $\mathbb{R}^n \rightarrow \mathbb{R}^m$. We define the Jacobian to be

$$Jf(x) = \sqrt{\det(Df(x)^* Df(x))}$$

where $Df(x)^* : \mathbb{R}^m \rightarrow \mathbb{R}^n$ denotes the transpose (adjoint) of $Df(x) : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Note that if $n = m$, then $Jf(x) = |\det(Df(x))|$ is the absolute value of the determinant of the derivative.

The goal of this section is to prove the following.

Theorem 7.1 (Area formula). Suppose that $X \subset \mathbb{R}^n$ is open and $n \leq m$ are positive integers. Let $f : X \rightarrow \mathbb{R}^m$ be continuously differentiable function on X that is injective. Then for Borel $B \subset X$ we have that

$$\mathcal{H}^n(f(B)) = \int_B Jf(x) d\mathcal{L}_n(x).$$

TODO: can we remove $Jf > 0$?

Example 7.2 (Arc-length). Suppose $n = 1$. Thus $f(x) = [f_1(x), \dots, f_n(x)]^t$ is a column vector. Thus $Df(x)$ is the column vector of derivatives $[f'_1(x), \dots, f'_n(x)]^t$. Thus

$$Jf(x) = \sqrt{\sum_{i=1}^m (f'_i(x))^2} = \|Df(x)\|$$

is magnitude of the velocity vector (if x denotes time). Thus

$$\mathcal{H}^1(f([a, b])) = \int_a^b \|Df(x)\| dx$$

is the arc-length of the parametrized curve $\{f(x) \mid x \in [a, b]\}$.

We fix for the rest of this section f , X and n, m as in the Area formula.

7.1. Area formula for Linear maps. We first examine the linear case where $f(x) = Lx$ is a linear map. By the Singular Value Decomposition, we may write $L = UDV$ where $U : \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $V : \mathbb{R}^n \rightarrow \mathbb{R}^n$ are isometries and $D : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a diagonal matrix with non-negative entries on the diagonal.

By Proposition 6.7 and the fact that the Hausdorff measure is rotation invariant, we have

$$\mathcal{L}_n(VB) = \mathcal{H}^n(VB) = \mathcal{H}^n(B) = \mathcal{L}_n(B)$$

for any Borel $B \subset \mathbb{R}^n$. Thus $\mathcal{L}_n(DVB) = \det(D)\mathcal{L}_n(VB) = \mathcal{L}_n(B)$ as D is diagonal (i.e., the image of each rectangle is another rectangle where the sidelengths get scaled by the diagonal elements of D).

Lemma 7.3. Let $U : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be an isometry (as above), then for borel $B \subset \mathbb{R}^n$ we have that $\mathcal{H}^n(UB) = \mathcal{L}_n(B)$.

Proof. Note that since $\mathcal{H}^n$ is rotation invariant, we may assume that U is the standard embedding of $\mathbb{R}^n$ into $\mathbb{R}^m$ (i.e., $(x_1, \dots, x_n) \mapsto (x_1, \dots, x_n, 0, \dots, 0)$). It thus suffices to show that $\mathcal{H}^n(B) = \mathcal{H}^n(B \times \{0\}^{m-n})$. This follows from Lemma 6.2. $\square$

We thus get that $\mathcal{H}^n(LB) = \det(D)\mathcal{L}_n(B)$, i.e.,

$$\mathcal{H}^n(f(B)) = \int_B \det(D) d\mathcal{L}_n(x).$$

Thus to complete the proof of the Area formula for linear functions, it remains to show that the Jacobian of $f = L$ is equal to $\det(D)$. Note that $Df = L$ and thus

Now

$$(Jf(x))^2 = \det(L^*L) = \det(V^*D^*U^*UDV) = \det(V^*D^2V) = \det(D)^2.$$

Thus we have shown the area formula when $f(x) = Lx$ is an injective linear function.

7.2. Proof of the Area formula when $Jf(x) > 0$ for all $x \in X$. We now turn to the proof of Theorem 7.1 (the Area formula) in the case that $Jf(x) > 0$ for all $x \in X$, i.e., $Df(x)$ has trivial Kernel for all $x \in X$. We have already shown it for linear maps in the previous section and our strategy will be to use this by approximating a continuously differentiable function by piecewise linear functions.

In this section, we fix the data as in Theorem 7.1, i.e., $f : X \rightarrow \mathbb{R}^m$, where $X \subset \mathbb{R}^n$ is open and $n \leq m$, is continuously differentiable on X and is injective. Moreover, as mentioned above, we assume in this subsection that $Jf(x) > 0$ for all $x \in X$.

Lemma 7.4. Suppose that $B \subset X$ is a bounded convex Borel subset of X . Suppose $\epsilon, \rho > 0$ are such $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear map such that

$$\|Tu\| \geq \rho\|u\| \quad \text{for all } u \in \mathbb{R}^n$$

and

$$\|Df(x) - T\| \leq \epsilon \quad \text{for all } x \in B.$$

Then

$$(1 - \epsilon\rho)^n \mathcal{H}^n(TB) \leq \mathcal{H}^n(f(B)) \leq (1 + \epsilon\rho)^n \mathcal{H}^n(TB).$$

Proof. Note that $h = f - T$ is continuously differentiable with derivative $Dh = Df - T$, thus by the mean value inequality (which we can use as B is convex) we have that

$$\|h(x_1) - h(x_2)\| \leq \sup_{x \in B} \|Dh(x)\| \|x_1 - x_2\|$$

for all $x_1, x_2 \in B$. But $\|Dh(x)\| < \epsilon$ and thus we have that

$$\|f(x_1) - Tx_1 - (f(x_2) - Tx_2)\| \leq \epsilon \|x_1 - x_2\|,$$

we thus have that

$$\|f(x_1) - f(x_2) - (Tx_1 - Tx_2)\| \leq \epsilon \|x_1 - x_2\|.$$

Note that T is injective as $\|Tu\| \geq \rho\|u\| > 0$ for non-zero u and so we can let $g : TB \rightarrow f(B)$ be defined by $g(y) = f(T^{-1}y)$ for $y \in TB$. So for each $y_1, y_2 \in TB$, we have write

$$\|g(y_1) - g(y_2) - (y_1 - y_2)\| \leq \epsilon \|T^{-1}y_1 - T^{-1}y_2\| \leq \epsilon\rho \|y_1 - y_2\|.$$

We thus have

$$(1 - \epsilon\rho)\|y_1 - y_2\| \leq \|g(y_1) - g(y_2)\| \leq (1 + \epsilon\rho)\|y_1 - y_2\|$$

for all $y_1, y_2 \in TB$. As $g : TB \rightarrow f(B)$ is a bijection, we can also write the first inequality as

$$\|g^{-1}(z_1) - g^{-1}(z_2)\| \leq (1 - \epsilon\rho)^{-1} \|z_1 - z_2\|$$

for all $z_1, z_2 \in f(B)$. These inequalities now imply that

$$\mathcal{H}^n(f(B)) = \mathcal{H}^n(g(TB)) \leq (1 + \epsilon\rho)^n \mathcal{H}^n(TB)$$

and

$$\mathcal{H}^n(TB) = \mathcal{H}^n(g^{-1}(f(B))) \leq (1 - \epsilon\rho)^{-n} \mathcal{H}^n(f(B)).$$

□

Lemma 7.5. If $B \subset X$ is Borel, then $f(B)$ is also Borel.

Proof. If B is compact, then $f(B)$ is compact and thus Borel (as f is continuous). Now let $\mathcal{A}$ denote those Borel $B \subset X$ such that $f(B)$ is Borel. We see that $\mathcal{A}$ contains all compact sets. Every open set can be written as a nested union of open sets, thus $\mathcal{A}$ contains all open sets. In particular, this shows that $f(X)$ is open. Thus f is a bijection $X \rightarrow f(X)$ and in particular $\mathcal{A}$ is closed under complements since if $f(B)$ is measurable then $f(X \setminus B) = f(X) \setminus f(B)$, as f is injective, and thus $f(X \setminus B)$ is Borel, thus $X \setminus B$ is Borel. Thus $\mathcal{A}$ is indeed a σ -algebra. □

We are now ready to complete the proof of the Area formula.

Let $K \subset U$ be a compact box (cartesian product of n compact intervals in $\mathbb{R}$). We first prove the Area formula for $B = K$. Now as Df is continuous on K we have that $(x, u) \mapsto \|Df(x)u\|$ is a continuous function on $K \times \mathbb{R}^d$ that never vanishes. Thus $\{Df(x)u \mid x \in K, u \in \mathbb{R}^d \text{ and } \|u\| = 1\}$ is a compact set that does not contain 0, and thus its infimum is $\rho > 0$. Thus there exists $\rho > 0$ such that $\|Df(x)u\| \geq \rho\|u\|$ for all $u \in \mathbb{R}^d$.

Now fix $\epsilon > 0$. As Df is uniformly continuous on K , so we can partition K into finitely many rectangles $B_1, \dots, B_q$ such that

$$\|Df(x) - Df(x')\| \leq \epsilon$$

whenever $x, x' \in B_i$. Choose an $x_i \in B_i$ and let $T_i = Df(x_i)$. Thus $\|T_i u\| \geq \rho \|u\|$ for all $u \in \mathbb{R}^d$ and

$$\|Df(x) - T_i\| \leq \epsilon \quad \text{for all } x \in B_i.$$

Thus we may apply Lemma 7.4 to B_i and T_i and get that

$$(1 - \epsilon\rho)^n \sum_{i=1}^q \mathcal{H}^n(T_i B_i) \leq \mathcal{H}^n(f(B_i)) \leq (1 + \epsilon\rho)^n \sum_{i=1}^q \mathcal{H}^n(T_i B_i).$$

Now apply the area formula to T_i , which we already know is true as T_i is linear and injective, to get that

$$\sum_{i=1}^q \mathcal{H}^n(T_i B_i) = \sum_{i=1}^q (Jf)(x_i) \mathcal{L}_d(B_i).$$

By making the B_i diameters converge to 0, we see that this sum converges to the Riemann integral of Jf . Thus we have shown that

$$(1 - \epsilon\rho)^n \int_B Jf d\mathcal{L}_d \leq \mathcal{H}^n(f(B_i)) \leq (1 + \epsilon\rho)^n \int_B Jf d\mathcal{L}_d.$$

Note that we used here that $f(B_1), \dots, f(B_q)$ are disjoint (as f is injective) and Borel (as shown) and thus were allowed to use the additivity of the Hausdorff measure.

As ϵ was arbitrary, we can let $\epsilon \rightarrow 0$ and this completes the proof (note that ρ was chosen before ϵ and thus is fixed).

Thus we know that the Area formula holds for compact rectangles. It now holds in general since set of all B where it holds is a monotone class.

7.3. Proof of the area formula without assuming $Jf(x) > 0$. We now turn to the general case where of the Area Formula where we drop the assumption that $Jf(x) > 0$ for all $x \in X$. To do this, we simply adjust f slightly so that it does satisfy this hypothesis. To do this, choose Borel $B \subset X$ and write $B = B' \cup B_0$ where $B' = B \cap X'$ where

$$X' = \{x \in X \mid Jf(x) > 0\}$$

and $B_0 = B \cap C_0$ where

$$C_0 = \{x \in X \mid Jf(x) = 0\}.$$

Note that X' is an open subset of $\mathbb{R}^n$ on which Jf never vanishes and so the special case of the area Area Formula we have already proved, applies and thus shows that

$$\int_{B'} Jf(x) d\mathcal{L}_n(x) = \mathcal{H}^n(f(B')).$$

It thus remains to show that

$$0 = \int_{B_0} Jf(x) d\mathcal{L}_n(x) = \mathcal{H}^n(f(B_0))$$

and for this it will suffice to show that

$$\mathcal{H}^n(f(C_0)) = 0.$$

In fact, as U can be written as a countable union of compact sets, it is enough to show that $\mathcal{H}^n(f(C_0 \cap K)) = 0$ for each compact $K \subset X$. We thus let $C = C_0 \cap K$ for some compact $K \subset U$, thus C is a bounded set with $Jf(x) = 0$ for all $x \in C$ and $\{Df(x) \mid x \in C\}$ is a bounded set of matrices (i.e., each entry of this matrix is a bounded function of $x \in C$).

Let $f_\epsilon : \mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ be given by

$$f_\epsilon(x) = (\epsilon x, f(x)).$$

Note that $Df_\epsilon(x)$ has full rank n and thus we can apply the Area Formula to get that

$$\int_C Jf_\epsilon(x) d\mathcal{L}_n(x) = \mathcal{H}^n(f_\epsilon(C)).$$

Let $P : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be the projection. Note that $P(f_\epsilon(C)) = f(C)$ but

$$\mathcal{H}^n(P(f_\epsilon(C))) \leq \mathcal{H}^n(f_\epsilon(C))$$

as $\text{diam}(P(A)) \leq \text{diam}(A)$ for all $A \subset \mathbb{R}^n$. Thus it will be enough to show that as $\epsilon \rightarrow 0$, we have

$$\int_C Jf_\epsilon(x) d\mathcal{L}_n(x) \rightarrow 0.$$

First we observe that since $Df(x)$ is bounded on C we have $Df_\epsilon(x)$ is also bounded on C . Now $Df_\epsilon(x)$, as a matrix, can be decomposed into two matrices: the submatrix formed by the first n rows is ϵI_n , where I_n is the n by n identity matrix while the submatrix formed by the last m rows is $Df(x)$. Thus the transpose is $Df_\epsilon(x)^* = (\epsilon I_n | Df(x))$ where $|$ means concatenate columns. Thus

$$Jf_\epsilon(x) = \sqrt{\det(Df_\epsilon(x)^* Df_\epsilon(x))} = \sqrt{\det(\epsilon^2 I_n + Df(x)^* Df(x))}.$$

This shows that $Jf_\epsilon(x) \rightarrow Jf(x)$ as $\epsilon \rightarrow 0$. Thus the Lebesgue dominated convergence will give us the desired conclusion, as soon as we can show it can be applied. To see that it applies, first observe that C is bounded and thus has finite Lebesgue measure. Also, the set

$$\{Jf_\epsilon(x) \mid x \in C, \epsilon \in [0, 1]\}$$

is bounded since $\{Df(x) \mid x \in C\}$ is bounded as argued above. Thus we have uniform boundedness of the integral by an L^∞ -function, thus an L^1 function as C is bounded (of finite measure). Thus Lebesgue's dominated converge theorem indeed applies.

This completes the full proof of the Area Formula.

7.4. Change of variables. We can now compute surface integrals using the following corollary of the Area formula.

Theorem 7.6. Let $n \leq m$ be integers, $X \subset \mathbb{R}^n$ open, and suppose that $f : X \rightarrow \mathbb{R}^m$ is injective and continuously differentiable. Let $g : \mathbb{R}^m \rightarrow \mathbb{R}$ be a Borel measurable function such that either $g(y) \geq 0$ for all $y \in \mathbb{R}^m$ or g is L^1 w.r.t the Hausdorff measure on $\mathcal{H}^n$ on $f(X)$. Then

$$\int_X g(f(x)) Jf(x) d\mathcal{L}_n(x) = \int_{f(X)} g(y) d\mathcal{H}^n(y).$$

Proof. When $g = 1_B$ for some Borel $B \subset \mathbb{R}^m$ we know this is true by the Area formula applied to $f^{-1}(B)$. The result now follows for $g \geq 0$ by applying the monotone convergence theorem and the fact that g can be written as a monotonic limit of simple functions. g is L^1 w.r.t the Hausdorff measure on $\mathcal{H}^n$ on $f(X)$, then the result follows by writing g as a sum of a positive and a negative function. $\square$

Example 7.7 (Polar co-ordinates). Let $X = \mathbb{R}_{>0} \times (0, 2\pi)$ and $f : X \rightarrow \mathbb{R}^2$ be given by $f(r, \theta) = (r \cos(\theta), r \sin(\theta))$. Note that this is injective thus the Area formula applies. Here

$$Df(r, \theta) = \begin{bmatrix} \cos(\theta) & -r \sin(\theta) \\ \sin(\theta) & r \cos(\theta) \end{bmatrix}$$

and thus $Jf(r, \theta) = |\det(Df(r, \theta))| = r$. Hence we have integration in polar co-ordinates given by

$$\int_{f(X)} g(x, y) dx dy = \int_0^\infty \int_0^{2\pi} g(r \cos(\theta), r \sin(\theta)) r d\theta dr$$

whenever g is integrable or positive. Note that $f(X) = \mathbb{R}^2 \setminus [0, \infty) \times \{0\}$ thus integrating over $f(X)$ is equivalent to integrating over $\mathbb{R}^2$.

In particular, if one sets $g(x, y)$ to be the indicator function of the unit disc, one then gets that the area of the unit disk is given by

$$\int_0^1 \int_0^{2\pi} r d\theta dr = 2\pi \int_0^1 r dr = \pi.$$

We can extend this example to higher dimensions by showing that volume integrals may be computed as an integral of spherical integrals.

Proposition 7.8 (Co-area formula for spheres). Let $n \geq 2$. Let $g : \mathbb{R}^n \rightarrow [0, \infty]$ be any Borel measurable function and let S_r denote the unit sphere in r . Then

$$\int_{\mathbb{R}^n} f d\mathcal{L}_n = \int_0^\infty \int_{S_r} g(u) d\mathcal{H}^n(u) dr,$$

where this integrand is indeed a Borel measurable function of r if g is positive, while if g is merely integrable then it is Borel on a co-null subset of $\mathbb{R}_{>0}$ (thus still makes sense).

Proof. We first parametrize the unit sphere. We have an injective continuously differentiable map $\rho : \mathbb{R}^{n-1} \rightarrow S_1$ whose image is S_1 without a single point, for example we can let ρ be the stereographic projection. Now define $f : \mathbb{R}_{>0} \times \mathbb{R}^{n-1}$ by

$$f(r, x_1, \dots, x_{n-1}) = r\rho(x_1, \dots, x_{n-1}).$$

We thus have that

$$\int_{\mathbb{R}^n} g d\mathcal{L}_n = \int_{\mathbb{R} \times \mathbb{R}^{n-1}} g(f(r, x_1, \dots, x_{n-1})) Jf(r, x_1, \dots, x_{n-1}) dr dx_1 \cdots dx_{n-1}.$$

Note that the image of f is a co-null subset of $\mathbb{R}^n$, thus the integration can be taken over the whole of $\mathbb{R}^n$.

Let us compute this Jacobian. The first column of $Df(r, x_1, \dots, x_{n-1})$ is

$$\frac{\partial}{\partial r} f(r, x_1, \dots, x_{n-1}) = \frac{\partial}{\partial r} r\rho(x_1, \dots, x_{n-1}) = \rho(x_1, \dots, x_{n-1})$$

where we identify vectors with column vectors naturally. The other columns are

$$\frac{\partial}{\partial x_1} r\rho(x_1, \dots, x_{n-1}) = r(\partial_i \rho)(x_1, \dots, x_{n-1}).$$

Observe now that the first column $\rho(x_1, \dots, x_{n-1})$ is a unit vector that is orthogonal to the other columns $r(\partial_i \rho)(x_1, \dots, x_{n-1})$, as these other columns are in the tangent space of the unit sphere at the point $\rho(x_1, \dots, x_{n-1})$. It now follows that $Df^* Df$ has as its first column (and row) the unit vector $(1, 0, \dots, 0)$ while for $1 \leq i, j \leq n$ the (i, j) -entry of $Df^* Df$ is the inner product

$$\langle r(\partial_i \rho)(x_1, \dots, x_{n-1}), r(\partial_j \rho)(x_1, \dots, x_{n-1}) \rangle.$$

It follows that $\det(Df^* Df)$ is the determinant of this submatrix (obtained by deleting first and second column) which we see is

$$r^{2(n-1)} (J\rho(x_1, \dots, x_{n-1}))^2.$$

Thus $Jf(r, x_1, \dots, x_{n-1}) = r^{n-1} J\rho(x_1, \dots, x_{n-1})$.

We thus get that

$$\int_{\mathbb{R}^n} g d\mathcal{L}_n = \int_0^\infty \left(\int_{\mathbb{R}^{n-1}} g(r\rho(x_1, \dots, x_n)) r^{n-1} J\rho(x_1, \dots, x_{n-1}) dx_1 \cdots dx_{n-1} \right) dr$$

and the integrand is indeed Borel measurable (as a function of r) by the Fubini-Tonelli theorem if g is positive, while if g is mere integrable then we have that it is Borel on a co-null subset of $\mathbb{R}_{>0}$.

We now simplify this spherical integral by relating it to an integral over S_r . Note that the mapping $\rho_r = r\rho$ is an injective mapping $\rho_r : \mathbb{R}^{n-1} \rightarrow S_r$ which only omits one point. Thus

$$\int_{S_r} g(u) d\mathcal{H}^{n-1}(u) = \int_{\mathbb{R}^{n-1}} g(\rho_r(x_1, \dots, x_{n-1})) J\rho_r(x_1, \dots, x_{n-1}) dx_1 \cdots dx_{n-1}.$$

But now $D\rho_r(x_1, \dots, x_{n-1})$ has i -th column equal to $r(\partial_i \rho)(x_1, \dots, x_{n-1})$ and thus

$$J\rho_r(x_1, \dots, x_{n-1}) = r^{n-1} J\rho(x_1, \dots, x_{n-1}).$$

Thus

$$\int_{S_r} g(u) d\mathcal{H}^{n-1}(u) = \int_{\mathbb{R}^{n-1}} g(r\rho(x_1, \dots, x_n)) r^{n-1} J\rho(x_1, \dots, x_{n-1}) dx_1 \cdots dx_{n-1},$$

which completes the proof after plugging this expression into the iterated integral above. $\square$

7.5. Volume and surface area of the unit ball. Recall that α_d denotes the volume (n -dimensional Lebesgue measure) of the d -dimensional unit ball. Let

$$\alpha'_d = \mathcal{H}^{d-1}(\{x \in \mathbb{R}^d \mid \|x\| = 1\})$$

denote the *surface area* of the unit sphere. Thus $\alpha'_d r^{d-1}$ is the surface area of the sphere of radius r , while $\alpha_d r^d$ is the volume of the ball of radius r . Thus by Proposition 7.8 (spherical integration) applied to g being the indicator function of the ball of radius R , we see that

$$\alpha_d R^d = \int_0^R \alpha'_d r^{d-1} dr = \frac{1}{d} \alpha'_d R^d,$$

that is,

$$\alpha'_d = d\alpha_d.$$

Thus we get the surface area immediately once we know the volume of the unit ball.