

Assignment_3_QMM

Karthik Badiganti

2022-10-16

Loading Packages

```
library("lpSolveAPI")
library("lpSolve")
library("tinytex")
```

Creating Table as per given data

```
cost <- matrix(c(22,14,30,600,100,
                16,20,24,625,120,
                80,60,70,"-", "-"), ncol=5,byrow=T)

colnames(cost) <- c("Warehouse 1", "Warehouse 2", "Warehouse 3", "Production Cost", "Production Capacity")
rownames(cost) <- c("Plant A", "Plant B", "Demand")

cost <- as.table(cost)
cost
```

##		Warehouse 1	Warehouse 2	Warehouse 3	Production Cost	Production Capacity
## Plant A	22	14	30	600	100	
## Plant B	16	20	24	625	120	
## Demand	80	60	70	-	-	

Formulation of Primal

The Objective Function is to Minimize the Transportation cost

$$Z = 622X_{11} + 614X_{12} + 630X_{13} + 641X_{21} + 645X_{22} + 649X_{23}$$

Subject to:

Supply Constraints:

$$X_{11} + X_{12} + X_{13} + X_{14} \leq 100$$

$$X_{21} + X_{22} + X_{23} + X_{24} \leq 120$$

Demand Constraints

$$X_{11} + X_{21} \geq 80$$

$$X_{12} + X_{22} \geq 60$$

$$X_{13} + X_{23} \geq 70$$

Non – Negativity Constraints

$$X_{ij} \geq 0 \quad \text{Where } i = 1,2 \text{ for Plants A,B and } j = 1,2,3 \text{ for Warehouses}$$

Since demand is not equal to supply we are adding dummy variable for solving the equation

Solving Primal using R

```
#Creating a matrix for the given objective function
```

```
costs <- matrix(c(622,614,630,0,  
                  641,645,649,0), ncol=4, byrow=T)
```

```
#Defining the column names and row names
```

```
colnames(costs) <- c("Warehouse 1", "Warehouse 2", "Warehouse 3", "Dummy")
```

```
rownames(costs) <- c("Plant A", "Plant B")
```

```
costs
```

```
##           Warehouse 1 Warehouse 2 Warehouse 3 Dummy  
## Plant A           622           614           630     0  
## Plant B           641           645           649     0
```

```
#Defining the row signs and row values based on above constraints
```

```
row.signs <- rep("<=",2)
```

```
row.rhs <- c(100,120)
```

```
#Defining the column signs and column values based on above constraints
```

```
col.signs <- rep(">=",4)
```

```
col.rhs <- c(80,60,70,10)
```

```
#Running the lp.transport function
```

```
trans.cost <- lp.transport(costs,"min", row.signs,row.rhs,col.signs,col.rhs)
```

```
#Getting the objective value
```

```
print(paste("The solution of primal is ",trans.cost$objval))
```

```
## [1] "The solution of primal is 132790"
```

```
trans.cost$solution
```

```
##      [,1] [,2] [,3] [,4]  
## [1,]    0   60   40    0  
## [2,]   80    0   30   10
```

From the above results we can say that 132790 is the minimal cost obtained with

$$X_{12} = 60$$

$$X_{13} = 40$$

$$X_{21} = 80$$

$$X_{23} = 30$$

Since the primal was to minimize the transportation cost the dual of it would be to maximize the value added(VA)

$$\text{Maximize VA} = 100P_A + 120P_B - 80W_1 - 60W_2 - 70W_3$$

Subject to the following constraints:

Total Payments Constraints:

$$W_1 - P_A \geq 622$$

$$W_2 - P_A \geq 614$$

$$W_3 - P_A \geq 630$$

$$W_1 - P_B \geq 641$$

$$W_2 - P_2 \geq 645$$

$$W_3 - P_B \geq 649$$

$$W_i > 0, \quad P_j > 0$$

Where i=1,2,3 are Total payments at destination Warehouses

j=A,B are Total payments at Origin plants