
A. Type 2 diabetes
Biobank Japan (East Asian)
IVW (random effects)
Weighted median
MR−Egger
Weighted mode
RAPS
MR−PRESSO (1 outlier removed)
FinnGen R5 (European)
IVW (random effects)
Weighted median
MR−Egger
Weighted mode
RAPS
Sakaue 2021 (European)
IVW (random effects)
Weighted median
MR−Egger
Weighted mode
RAPS
MR−PRESSO (2 outliers removed)
UK Biobank self−report (European)
IVW, linear beta converted to log−odds
1.14 (1.10−1.19)
1.18 (1.11−1.25)
1.27 (1.14−1.42)
1.17 (1.09−1.26)
1.14 (1.09−1.20)
1.15 (1.10−1.19)
1.09 (1.04−1.14)
1.07 (1.00−1.14)
1.04 (0.94−1.16)
1.05 (0.97−1.13)
1.09 (1.04−1.14)
1.11 (1.08−1.14)
1.11 (1.07−1.15)
1.12 (1.05−1.20)
1.12 (1.07−1.17)
1.11 (1.07−1.14)
1.10 (1.07−1.13)
1.07 (1.01−1.15)
0.9
1.0
1.1
1.2
1.3
1.4
OR for cataract per unit log−odds of T2D liability


B. Body mass index
Biobank Japan (East Asian)
IVW (random effects)
Weighted median
MR−Egger
Weighted mode
RAPS
FinnGen R5 (European)
IVW (random effects)
Weighted median
MR−Egger
Weighted mode
RAPS
MR−PRESSO (1 outlier removed)
Sakaue 2021 (European)
IVW (random effects)
Weighted median
MR−Egger
Weighted mode
RAPS
MR−PRESSO (2 outliers removed)
UK Biobank self−report (European)
IVW, linear beta converted to log−odds
1.21 (1.12−1.30)
1.26 (1.12−1.42)
1.40 (1.15−1.70)
1.28 (0.96−1.71)
1.20 (1.11−1.30)
1.13 (1.04−1.22)
1.17 (1.02−1.34)
1.13 (0.92−1.39)
1.19 (0.95−1.48)
1.13 (1.04−1.22)
1.14 (1.05−1.23)
1.20 (1.14−1.26)
1.25 (1.16−1.35)
1.25 (1.10−1.41)
1.27 (1.13−1.43)
1.20 (1.15−1.26)
1.20 (1.14−1.25)
1.10 (0.97−1.24)
0.8
1.0
1.2
1.4
1.6
1.8
OR for cataract per SD of BMI