

1 Erosion of somatic tissue identity with loss of the X-linked

2 intellectual disability factor KDM5C

3

⁴ Katherine M. Bonefas, Ilakkiya Venkatachalam, and Shigeki Iwase.

5 Abstract

It is currently unclear why mutations in numerous chromatin-modifying enzymes cause neurodevelopmental disorders (NDDs). While some chromatin regulators are important for neuron-specific gene expression, loss of repressive chromatin regulators can also lead to the aberrant transcription of tissue-specific genes outside of their intended context. Because very few studies have characterized the these aberrant, tissue-specific transcripts, the mechanisms behind their dysregulation and their functional consequences are largely unknown. Here, we explored this cellular identity crisis in NDDs by investigating the ectopic expression of germline (e.g. sperm and egg) genes in mice lacking lysine demethylase 5c (KDM5C, also known as JARID1C or SMCX), an eraser of histone 3 lysine 4 di and tri-methylation (H3K4me2/3). We found male *Kdm5c* knockout (-KO) mice, which recapitulate key behavioral phenotypes of Claes-Jensen X-linked intellectual disability (MRXSCJ, OMIM: 300534), aberrantly expresses many liver, muscle, ovary, and testis genes within the amygdala and hippocampus. Many genes expressed in the *Kdm5c*-KO brain were typically unique to germ cells, indicating an erosion of the soma-germline boundary. Using germ cell-depleted RNA sequencing datasets, we curated a a list of germline-enriched genes for unbiased characterization of KDM5C in germline gene repression. Germline genes are typically decommissioned in somatic lineages in the post-implantation epiblast, yet *Kdm5c*-KO epiblast-like cells (EpiLCs) aberrantly expressed key regulators of germline identity and meiosis, including *Dazl* and *Stra8*. Germline gene dysregualtion is sexually dimorphic, as female *Kdm5c*-KO EpiLCs were more prone to germline gene dysregulation than knockout males. We found KDM5C acts at a subset of germline gene promoters in EpiLCs to promote the initial placement of DNA methylation at CpG islands. However, germline genes can also become activated in *Kdm5c*-KO cells independent of direct KDM5C regulation. Intriguingly, many of the germline transcripts expressed in the *Kdm5c*-KO brain were late-stage spermatogenesis genes that are not directly bound by KDM5C. This suggests germline developmental programs can progress ectopically in the background of typical *Kdm5c*-KO development. Ultimately, this work characterizes a novel suppressor of germline gene transcription and links impaired soma-vs-germline demarcation to a chromatin-linked neurodevelopmental disorder.

30 Introduction

31 A single genome holds the instructions to generate the myriad of cell types found within the adult
32 organism. This is, in part, accomplished by chromatin regulators that can either promote or impede lineage-
33 specific gene expression through DNA and histone modifications^{1,2}. Many chromatin regulators were initially
34 identified for their roles in shaping cellular and tissue identity^{3–5}. Unexpectedly, next generation sequencing
35 revealed mutations in many chromatin regulators directly cause or are associated with neurodevelopmental
36 disorders (NDDs)⁶. Several studies have suggested this connection between chromatin regulators and
37 neurodevelopment is due to their regulation of brain-specific genes, such as orchestrating transcriptional
38 programs for synaptic maturation⁷ and transitioning between neuronal and glial fates during neural precursor
39 differentiation⁸. However, loss of some chromatin regulators can also result in the ectopic transcription
40 of tissue-specific genes outside of their target environment, such as the misexpression of liver-specific
41 genes within adult neurons⁹. Very few studies have investigated this cellular identity crisis in chromatin-
42 linked NDDs^{9,10} and it is currently unknown if ectopic gene expression contributes to neurodevelopmental
43 impairments.

44 To elucidate the role of tissue identity in chromatin-linked neurodevelopmental disorders, it is essential
45 to first characterize the nature of the dysregulated genes and the molecular mechanisms governing their
46 de-repression. In this study, we characterized the aberrant expression of tissue-enriched genes with
47 loss of lysine demethylase 5C (KDM5C). KDM5C, also known as SMCX or JARID1C, is a chromatin
48 regulator that can repress gene expression through erasure of histone 3 lysine 4 di- and trimethylation
49 (H3K4me2/3)¹¹, marks enriched at active gene promoters. Pathogenic mutations in *KDM5C* cause Intellectual
50 Developmental Disorder, X-linked, Syndromic, Claes-Jensen Type (MRXSCJ, OMIM: 300534). MRXSCJ
51 is more common and often more severe in males and its neurological phenotypes include intellectual
52 disability, seizures, aberrant aggression, and autistic behaviors^{12–14}. Male *Kdm5c* knockout (-KO) mice
53 recapitulate key MRXSCJ phenotypes, including hyperaggression, increased seizure propensity, and learning
54 impairments^{10,15}. Unexpectedly, RNA sequencing (RNA-seq) of the *Kdm5c*-KO hippocampus revealed
55 ectopic expression of testis genes within the brain¹⁰. It is currently unknown what types of testis genes are
56 dysregulated, when in *Kdm5c*-KO development this begins, if this phenomenon is sexually dimorphic, and if
57 other tissue-specific genes are also aberrantly expressed with KDM5C loss.

58 The testis contains both germ cells (cells that pass on their genetic material, e.g. sperm) and somatic
59 cells (cells that perform all other bodily functions). Distinguishing between the germline and the soma is a
60 key feature of multicellularity¹⁶ that occurs during early embryogenesis in many metazoans¹⁷. In mammals,
61 chromatin regulators are crucial for decommissioning germline genes in somatic cells during the transition
62 from naïve to primed pluripotency. Initially, germline gene promoters gain repressive histone H2A lysine 119
63 monoubiquitination (H2AK119ub1)¹⁸ and histone 3 lysine 9 trimethylation (H3K9me3)^{18,19} in embryonic stem
64 cells and are then decorated with DNA CpG methylation in the post-implantation embryo^{19–21}. Currently,

most studies have focused on marker genes important for early germ cell development rather than germline genes as a whole given the lack of a curated list of germline-enriched genes. Thus, it is unclear if the mechanism of repression differs for certain classes of germline genes, e.g. meiotic genes versus spermatid differentiation genes. Furthermore, it is currently unknown if ectopic germline gene expression is influenced by chromosomal sex, as previous studies on germline gene repression have been exclusively in males. Sex is particularly pertinent in the case of sexually dimorphic chromatin regulators like KDM5C, which lies on the X chromosome and partially escapes X chromosome inactivation, resulting in 1.2-2-fold higher in females²²⁻²⁵. Thus, systematically characterizing KDM5C's role in germline gene repression during early embryogenesis, including its interaction with known silencing mechanisms, will unveil key mechanisms underlying the demarcation between soma and germline identity. These results will provide molecular footholds that can then be exploited to test the contribution of ectopic germline genes on neurodevelopment.

To illuminate KDM5C's role in tissue identity, here we characterized the aberrant transcription of tissue-enriched genes within the *Kdm5c*-KO brain and epiblast-like stem cells (EpiLCs), an *in vitro* model of the post-implantation embryo. We observed general dysregulation of tissue-enriched genes in both the *Kdm5c*-KO brain and EpiLCs, including testis, liver, muscle, and ovary-enriched genes. The *Kdm5c*-KO amygdala and hippocampus had significant enrichment of testis genes that are specific to germ cells and not somatic cells of the testis. We found *Kdm5c*-KO EpiLCs aberrantly expressed key drivers of germline identity and meiosis, including *Dazl* and *Stra8*, while the *Kdm5c*-KO brain primarily expressed germline genes important for late spermatogenesis. Female *Kdm5c*-KO EpiLCs expressed over double the number of germline genes than *Kdm5c*-KO males and germline genes dysregulated in both sexes were expressed more highly in females, demonstrating females have increased sensitivity to germline gene dysregulation. KDM5C was bound to only a subset of germline genes expressed in *Kdm5c*-KO EpiLCs, indicating germline-enriched mRNAs can be aberrantly transcribed through indirect mechanisms. Finally, we found KDM5C promotes the long-term silencing of germline genes in somatic cells by aiding the placement of DNA CpG methylation in EpiLCs. Thus, we propose KDM5C plays a fundamental role in the development of tissue identity during early embryogenesis, including the establishment of the soma-germline boundary.

Results

Tissue-enriched genes are aberrantly expressed in the *Kdm5c*-KO brain

Previous RNA sequencing (RNA-seq) in the adult hippocampus revealed ectopic expression of some testis genes within the male *Kdm5c* knockout (-KO) brain¹⁰. It is currently unclear if the testis is the only tissue type misexpressed in the *Kdm5c*-KO brain. We thus characterized dysregulation of *Kdm5c*-KO brain tissue identity by systematically assessing the expression of genes enriched in 17 mouse tissues²⁶, using our published mRNA-seq datasets of the adult amygdala and hippocampus for male mice with constitutive

98 knockout of *Kdm5c*²⁷.

99 We found a large proportion of genes that are significantly upregulated within the *Kdm5c*-KO brain
100 (DESeq2²⁸, log2 fold change > 0.5, q < 0.1) are enriched in non-brain tissues in wild-type mice (Amygdala:
101 35%, Hippocampus: 24%) (Figure 1A-B). The majority of tissue-enriched differentially expressed genes
102 (tissue-enriched DEGs) were testis genes (Figure 1A-C). Even though the testis has the largest total number
103 of tissue-biased genes compared to any other tissue (2,496 genes), testis-biased DEGs were significantly
104 enriched for both brain regions (Amygdala p = 1.83e-05, Odds Ratio = 5.13; Hippocampus p = 4.26e-11,
105 Odds Ratio = 4.45, Fisher's Exact Test). One example of a testis-enriched gene misexpressed in the
106 *Kdm5c*-KO brain is *FK506 binding protein 6* (*Fkbp6*), a known regulator of piRNA expression and meiosis in
107 germ cells^{29,30} (Figure 1C).

108 Interestingly, we also observed significant enrichment of ovary-biased DEGs in both the amygdala and
109 hippocampus (Amygdala p = 0.00574, Odds Ratio = 18.7; Hippocampus p = 0.048, Odds Ratio = 5.88,
110 Fisher's Exact) (Figure 1D). Ovary-enriched DEGs included *Zygotic arrest 1* (*Zar1*), which sequesters mRNAs
111 in oocytes for meiotic maturation and early zygote development³¹ (Figure 1D). Given the *Kdm5c*-KO mice
112 we analyzed are male, this demonstrates ectopic expression of tissue-enriched genes is independent of
113 organismal sex.

114 Although not consistent across brain regions, we also found significant enrichment of DEGs biased
115 towards two non-gonadal tissues - the liver (Amygdala p = 0.0398, Odds Ratio = 6.58, Fisher's Exact Test)
116 and the muscle (Hippocampus p = 0.0104, Odds Ratio = 6.95, Fisher's Exact Test). A liver-biased DEG
117 dysregulated in both the hippocampus and amygdala is *Apolipoprotein C-I* (*Apoc1*), a lipoprotein metabolism
118 and transport gene³² that has been implicated in Alzheimer's disease³³ (Figure 1E). For all tissue-enriched
119 DEGs, aberrantly expressed mRNAs are polyadenylated and spliced into mature transcripts (Figure 1C-E).
120 Of note, we observed little to no dysregulation of brain-enriched genes (Amygdala p = 1; Hippocampus p =
121 0.74, Fisher's Exact), despite the fact these are brain samples and the brain has the second highest total
122 number of tissue-enriched genes (708 genes). Together, these results suggest the aberrant expression of
123 tissue-enriched genes within the brain is a major effect of KDM5C loss.

124 **Germline genes are misexpressed in the *Kdm5c*-KO brain**

125 The testis contains both germ cells (meiotic cells, e.g. spermatogonia) and somatic cells (non-meiotic,
126 e.g. Leydig cells) that support hormone production and germline functions. To determine if *Kdm5c*-KO
127 testis-enriched DEGs are somatic or germline genes, we first evaluated their known functions through gene
128 ontology analysis. We found *Kdm5c*-KO testis-enriched DEGs had high enrichment of germline-relevant
129 ontologies, including spermatid development (GO: 0007286, p.adjust = 6.2e-12) and sperm axoneme
130 assembly (GO: 0007288, p.adjust = 2.45e-14) (Figure 2A).

131 To further validate if these testis DEGs are truly germline genes, we then assessed their expression in
132 somatic versus germ cells within the testis. We first compared their expression in wild-type testes to those

133 without germ cells³⁴, which was accomplished by heterozygous *W* and *Wv* mutations in the enzymatic
134 domain of *c-Kit* (*Kit*^{W/Wv})³⁵. Almost all *Kdm5c*-KO testis-enriched DEGs lost expression with germ cell
135 depletion (Figure 2B). We then assessed testis-enriched DEG expression in a published single cell RNA-seq
136 dataset that identified cell type-specific markers within the testis³⁶. Some *Kdm5c*-KO testis-enriched DEGs
137 were classified as specific markers for different germ cell types (e.g. spermatogonia, spermatocytes, round
138 spermatids, and elongating spermatids), yet none marked somatic cells (Figure 2C). Together, these data
139 demonstrate that the *Kdm5c*-KO brain aberrantly expresses germline genes, but not somatic testicular genes,
140 reflecting an erosion between somatic versus germline identity.

141 We then wanted to characterize germline gene misexpression with *Kdm5c* loss more deeply, but lacked
142 a comprehensive list of mouse germline genes. We therefore generated a list of germline-enriched genes
143 using RNA-seq datasets of *Kit*^{W/Wv} mice that included males and females at embryonic day 12, 14, and 16³⁷
144 and adult male testes³⁴. We defined genes as germline-enriched if their expression met the following criteria:
145 1) their expression is greater than 1 FPKM in wild-type gonads 2) their expression in any non-gonadal tissue
146 of adult wild type mice²⁶ does not exceed 20% of their maximum expression in the wild-type germline, and
147 3) their expression in the germ cell-depleted gonads, for any sex or time point, does not exceed 20% of
148 their maximum expression in the wild-type germline. These criteria yielded 1,288 germline-enriched genes
149 (Figure 2D), which was hereafter used as a resource to globally assess germline gene misexpression with
150 *Kdm5c* loss (Supplementary table 1).

151 ***Kdm5c*-KO epiblast-like cells aberrantly express key regulators of germline identity**

152 Germ cells are typically distinguished from somatic cells soon after the embryo implants into the uterine
153 wall^{38,39}. A subset of epiblast stem cells become the primordial germ cells (PGCs) while the remainder
154 differentiate into the ectoderm, mesoderm, and endoderm to form the somatic tissues⁴⁰. This developmental
155 time point can be modeled *in vitro* through differentiation of embryonic stem cells (ESCs) into epiblast-like
156 stem cells (EpiLCs) (Figure 3A, top)^{41,42}. While some germline-enriched genes are also expressed in
157 naïve embryonic stem cells (ESCs) and in the 2-cell stage^{43–45}, they are silenced as they differentiate into
158 EpiLCs^{19,46}. Therefore, we tested if KDM5C was necessary for silencing germline genes at this developmental
159 stage by evaluating the impact of *Kdm5c* loss in male EpiLCs.

160 We first identified *Kdm5c*-KO EpiLC DEGs in our previously published RNA-seq dataset⁴⁷ (DESeq2, log2
161 fold change > 0.5, q < 0.1) and then assessed tissue-enriched gene expression. Similar to the *Kdm5c*-KO
162 brain, we observed general dysregulation of tissue-enriched genes, with the largest number of genes
163 belonging to the brain and testis, although they were not significantly enriched (Figure 3B). Markers of primed
164 pluripotency, such as *Dnmt3b*, *Fgf5*, *Pou3f1*, and *Otx2*, were unaltered in *Kdm5c*-KO EpiLCs (Figure 3C).
165 Furthermore, *Kdm5c*-KO cell morphology during ESC to EpiLC differentiation (Figure 3D) appeared normal,
166 indicating KDM5C loss does not impair EpiLC formation.

167 To determine if germline DEGs are constitutively dysregulated or if they can change over the course of

development, we next compared EpiLC germline DEGs to those expressed in the *Kdm5c*-KO brain. We found the majority of germline DEGs were unique to either EpiLCs or the brain, with only *Cyct* shared across all tissue/cell types (Figure 3E-F). EpiLCs had particularly high enrichment of meiosis-related gene ontologies (Figure 3G), such as meiotic cell cycle process (GO:1903046, p.adjust = 1.59e-08) and meiotic nuclear division (GO:0140013, p.adjust = 9.76e-09). While there was modest enrichment of meiotic gene ontologies in both brain regions, the *Kdm5c*-KO hippocampus primarily expressed late-stage spermatogenesis genes, such those involved in sperm axoneme assembly (GO:0007288, p.adjust = 0.00621) and sperm motility (GO:0097722, p.adjust = 0.00612).

Notably, DEGs unique to *Kdm5c*-KO EpiLCs included key drivers of germline identity, such as *Stimulated by retinoic acid 8 (Stra8)* and *Deleted in azoospermia like (Dazl)* (Figure 3H). These genes are typically expressed during embryonic germ cell development to commit PGCs to the germline fate, but are also expressed later in life to trigger meiotic gene expression programs^{48–50}. Of note, some germline genes, including *Dazl*, are also expressed in the two-cell embryo^{44,51}. However, we did not see misexpression of two-cell embryo-specific genes, like *Duxf3 (Dux)* ($q = 0.337$) and *Zscan4d* ($q = 0.381$), indicating *Kdm5c*-KO in EpiLCs do not revert back to a 2-cell-like state (Figure 3H). Altogether, these results demonstrate that while the *Kdm5c*-KO brain primarily expresses spermiogenesis genes, *Kdm5c*-KO EpiLCs express key drivers of germline identity and meiosis.

Female epiblast-like cells have increased sensitivity to germline gene misexpression with *Kdm5c* loss

It is currently unknown if the misexpression of germline genes is influenced by chromosomal sex, as previous studies on germline gene repressors have focused on their regulation in males^{18–20,52,53}. We explored the impact of sex upon germline gene misexpression by comparing their dysregulation in male *Kdm5c* hemizygous knockout (XY *Kdm5c*-KO), female homozygous knockout (XX *Kdm5c*-KO) and female heterozygous (XX *Kdm5c*-HET) EpiLCs.⁴⁷ We first identified differentially expressed genes (DEGs) compared to sex-matched wild-type controls (DESeq2²⁸, log2 fold change > 0.5, $q < 0.1$) and then filtered for germline-enriched genes.

Homozygous and heterozygous females expressed over double the number of germline-enriched genes than hemizygous males (Figure 4A). While the majority of germline DEGs in *Kdm5c*-KO males were also dysregulated in females (74%), there were also many male-specific and female-specific germline DEGs (Figure 4B). We the compared the known functions of germline genes dysregulated only in females (XX only - XX *Kdm5c*-KO, XX *Kdm5c*-HET, or both), only in males (XY only), or in all samples (shared). Female-specific germline DEGs were enriched for meiotic (GO:XXX meiotic cell cycle) and flagellar (GO:XXX cilium movement) functions, male-specific DEGs had mitochondrial and cell signaling gene functions (GO:XXX protein localization to mitochondrion). Germline transcripts expressed in both sexes were enriched for meiotic

202 (meiotic nuclear division) and egg-specific functions (female gamete generation).
203 The majority of germline genes expressed in both sexes had a greater log2 fold change from wild-type
204 in females compared to males (Figure 4D). This increased degree of dysregulation in females, along with
205 the increased total number of germline genes, indicates females are more sensitive to the loss of KDM5C-
206 mediated suppression of germline genes. Female sensitivity could be due to improper X chromosome
207 inactivation (XCI) in *Kdm5c* mutants, as many spermatogenesis genes lie on the X chromosome^{54,55}.
208 However, both shared and female-specific germline DEGs were not biased towards the X chromosome, with
209 the majority of genes instead lying on autosomes (Figure 4G). Thus, while female EpiLCs are more prone to
210 germline gene misexpression with KDM5C loss, it is likely independent of potential defects in XCI.

211 **Germline gene misexpression in *Kdm5c* mutants is independent of germ cell sex**

212 While many germline genes act in both the male and female germline, some display sex-biased expression
213 or have functions unique to eggs and sperm. To comprehensively assess whether germline gene sex
214 corresponds with *Kdm5c* mutant sex, we filtered our list of germline-enriched genes for egg (XX) and
215 sperm (XY) biased genes. We defined sex-biased genes as those whose expression in the opposite sex,
216 at any time point, is no greater than 20% of the gene's maximum expression in a given sex. This yielded
217 67 egg-biased, 1,024 sperm-biased, and 197 unbiased germline-enriched genes (Figure 4H), which is
218 consistent with the testis overall having a more unique transcriptome than the ovary²⁶. We found egg,
219 sperm, and unbiased germline genes were dysregulated in all *Kdm5c* mutants (Figure 4I-J). Germline genes
220 dysregulated exclusively in males and females were also not biased towards their corresponding germ
221 cell sex. This indicates differences between male and female germline gene dysregulation is not due to
222 sex-specific activation of sperm or egg-specific transcriptional programs. These results demonstrate that
223 sex influences the degree of germline gene misexpression with KDM5C loss, but not the germ cell sex from
224 which the germline genes originate.

225 **KDM5C binds to a subset of germline gene promoters during early embryogenesis**

226 Previous work suggests KDM5C represses germline genes during early development, as re-expression
227 of KDM5C in knockout neuronal cultures fails to suppress two hippocampal germline DEGs¹⁰. KDM5C binds
228 to several germline gene promoters in embryonic stem cells (ESCs) but is lost in neurons¹⁰. However, the
229 lack of a comprehensive list of germline-enriched genes prohibited systematic characterization of KDM5C
230 binding at germline gene promoters. It is currently unclear if KDM5C is enriched at germline gene promoters,
231 what types of germline genes KDM5C regulates, and if its binding is maintained at any germline genes in
232 neurons.

233 To address these questions, we analyzed KDM5C chromatin immunoprecipitation followed by DNA
234 sequencing (ChIP-seq) datasets in EpiLCs⁴⁷ and primary forebrain neuron cultures (PNCs)¹⁵. EpiLCs had a

235 higher total number of KDM5C peaks than PNCs (EpiLCs: 5,808, PNCs: 1,276, MACS2 q < 0.1 and fold
236 enrichment > 1, removal of *Kdm5c*-KO false positive peaks) and KDM5C was primarily localized to gene
237 promoters in both cell types (EpiLCs: 4,190, PNCs: 745 +/- 500bp from the TSS), although PNCs showed
238 increased localization to non-promoter regions (Figure 5A).

239 The majority of promoters bound by KDM5C in PNCs were also bound in EpiLCs (513 shared promoters),
240 however a large portion of gene promoters were only bound by KDM5C in EpiLCs (3677 EpiLC only
241 promoters) (Figure 5B). Genes bound by KDM5C in both PNCs and EpiLCs were enriched for functions
242 involving nucleic acid turnover, such as deoxyribonucleotide metabolic process (GO:0009262, p.adjust =
243 8.28e-05) (Figure 5C). Germline-specific ontologies were only enriched in promoters unique to EpiLCs, such
244 as meiotic nuclear division (GO: 0007127 p.adjust = 6.77e-16) and meiotic cell cycle process (GO:1903046,
245 p.adjust = 5.05e-16) (Figure 5C). There were no ontologies significantly enriched for genes only bound
246 by KDM5C in PNCs. We next evaluated KDM5C binding around the transcription start site (TSS) of all
247 germline-enriched genes. In EpiLCs, We observed modest KDM5C signal at about half of all germline genes
248 (Figure 5D). Based on our ChIP-seq peak cut-off criteria, KDM5C was strongly bound to about 10% of
249 germline gene promoters in EpiLCs (Figure 5E). In condordance with our gene ontology results, we did not
250 observe KDM5C accumulation at any germline gene promoters in PNCs (Figure 5D). Together, these results
251 demonstrate KDM5C is significantly enriched at a subset of germline gene promoters in EpiLCs, including
252 meiotic genes, but does not directly regulate germline genes in neurons.

253 Many germline-specific genes are suppressed by the transcription factor heterodimers E2F/DP1 and
254 MGA/MAX, which respectively bind E2F and E-box motifs^{20,52,53,56,57}. To elucidate if KDM5C is recruited to
255 germline gene promoters by a similar mechanism, we used HOMER to identify transcription factor motifs
256 enriched at KDM5C-bound or unbound germline gene promoters⁵⁸ (TSS +/- 500 bp, q-value < 0.1). MAX
257 and E2F6 binding sites were significantly enriched at germline genes bound by KDM5C in EpiLCs, but
258 not at germline genes unbound by KDM5C (MAX q-value: 0.0068, E2F6 q-value: 0.0673, E2F q-value:
259 0.0917) (Figure 5F). One third of KDM5C-bound promoters contained the consensus sequence for either
260 E2F6 (E2F, 5'-TCCCGC-3'), MGA (E-box, 5'-CACGTG-3'), or both, but only 17% of KDM5C-unbound genes
261 contained these motifs (Figure 5G). KDM5C-unbound germline genes were instead enriched for multiple
262 RFX transcription factor binding sites (RFX q-value < 0.0001, Rfx2 q-value < 0.0001, Rfx5 q-value < 0.0001)
263 (Figure 5H). RFX transcription factors bind X-box motifs⁵⁹ to promote ciliogenesis^{60,61}. Enriched RFX
264 transcription factors included RFX2, a central regulator of post-meiotic spermatogenesis^{62,63}. Interestingly,
265 RFX2 mRNA is upregulated in *Kdm5c*-KO EpiLCs (Figure 5I), but is not a direct target of KDM5C. Thus,
266 distinct transcription factor programs regulate the expression of KDM5C-bound and unbound germline genes.

267 Finally, we compared KDM5C binding at the promoters of RNA-seq DEGs to determine if the germline
268 genes transcribed in *Kdm5c*-KO cells are direct targets of KDM5C (Figure 5J-L). In EpiLCs, KDM5C was
269 bound to about one third of EpiLC-specific and brain-specific germline DEG promoters (EpiLC only: 36%,
270 Brain only: 33.3%). Again, we did not observe any KDM5C binding at germline gene promoters in PNCs,

even for brain-specific DEGs. Some notable differences in KDM5C binding for EpiLC-specific DEGs included *Dazl* and *Stra8*, two key drivers of germline identity discussed above. Although both mRNAs are expressed in *Kdm5c*-KO EpiLCs, KDM5C is only bound to *Dazl*'s promoter and not *Stra8*'s in EpiLCs (Figure 5K). In contrast to the unique DEGs, 3 out of the 4 genes dysregulated in both the brain and EpiLCs are bound by KDM5C around the TSS. Altogether, this suggests KDM5C decommissions a subset of germline genes in EpiLCs but is no longer recruited to germline gene promoters in neurons. Furthermore, the majority of germline mRNAs expressed in *Kdm5c*-KO cells are dysregulated independent of direct KDM5C recruitment to their promoters.

279 **KDM5C promotes *de novo* DNA methylation at germline genes**

- 280 • if there are differences, say at germline gene CpG islands

281 In the early embryo, germline gene promoters are initially silenced by repressive histone modifications, 282 including histone 2A lysine 119 monoubiquitination (H2AK119ub1) and histone 3 lysine 9 trimethylation 283 (H3K9me3), and are then silenced long-term via DNA CpG methylation (CpGme)^{18,19,46,64}. Our results above 284 suggest KDM5C also acts during this time period, however how KDM5C interacts with other germline gene 285 silencing mechanism is currently unclear. KDM5C is generally thought to suppress transcription through 286 erasure of histone 3 lysine 4 di- and trimethylation (H3K4me2/3)¹¹. However, KDM5C's catalytic activity was 287 recently shown to not be required for suppressing *Dazl* in undifferentiated ESCs⁵¹. Since H3K4me3 impedes 288 *de novo* CpGme placement^{65,66}, KDM5C's catalytic activity may instead be required for CpGme-mediated, 289 long-term silencing of germline genes. In support of this, CpGme is markedly reduced at two germline gene 290 promoters in the *Kdm5c*-KO adult hippocampus¹⁰. Based on these observations, we hypothesized KDM5C 291 erases H3K4me3 to promote the initial placement of CpGme at germline gene promoters in EpiLCs.

292 To test this hypothesis, we first characterized KDM5C's substrates, histone 3 lysine 4 di- and trimethylation 293 (H3K4me2/3) in our previously published ChIP-seq datasets of the wild type and *Kdm5c*-KO amygdala²⁷ 294 and EpiLCs⁴⁷. In congruence with previous work in the *Kdm5c*-KO hippocampus¹⁰, we observed aberrant 295 accumulation of H3K4me3 around the transcription start site (TSS) of germline genes in the *Kdm5c*-KO 296 amygdala (Figure 5A). We additionally found a marked increase in H3K4me2 around the TSS of germline 297 genes in *Kdm5c*-KO EpiLCs (Figure 5B).

298 We then evaluated KDM5C's role in germline gene silencing during ESC to EpiLC differentiation. We first 299 characterized KDM5C's mRNA and protein expression at 0, 24, and 48 hours of differentiation (Figure 5C). 300 While *Kdm5c* mRNA steadily decreased from 0 to 48 hours of differentiation (Figure 5D), KDM5C protein 301 initially increased from 0 to 24 hours but then decreased to near knockout levels by 48 hours (Figure 5E).

- 302 • Germline genes are known to accumulate CpGme at (CGIs).

- 303 – What percentage of germline genes have CGIs

- 304 • Do differential methylation analysis for WT ESCs to WT EpiLCs
305 – What percentage of germline genes significantly gain methylation (at CGI or at promoter)
306 – Out of the ones that gain methylation, which are significantly reduced
- 307 To determine KDM5C's role in the initial placement of DNA methylation at germline gene promoters,
308 we performed whole genome bisulfite sequencing (WGBS) in wild-type and *Kdm5c*-KO ESCs and 96 hour
309 extended EpiLCs (exEpiLCs). Although wild-type cells accumulated high levels of DNA methylation at
310 germline gene promoters over the course of ESCs to exEpiLC differentiation, DNA methylation was markedly
311 reduced in *Kdm5c*-KO exEpiLCs (Figure 5F).
- 312 • discussion
313 • – However, CPG islands greatly determine KDM5C recruitment - KDM5C is known to be enriched at
314 CGIs.)
315 • Altogether, this suggests KDM5C is dynamically regulated during ESC to EpiLC differentiation and
316 promotes germline gene silencing via removal of H3K4me2/3 during early embryogenesis.

317 Discussion

318 In the above study, we demonstrate KDM5C's pivotal role in the development of tissue identity, resulting in
319 substantial misexpression of tissue-enriched genes within the adult *Kdm5c*-KO hippocampus and amygdala.
320 In addition to testis genes identified previously¹⁰, we found significant enrichment of muscle, liver, and even
321 ovary-biased genes aberrantly expressed within the male *Kdm5c*-KO brain. Deeper characterization of
322 testis-enriched DEGs revealed they were not somatic testis genes, but germline genes, thus demonstrating
323 KDM5C's crucial role in establishing the soma-germline boundary. Given the majority of tissue-enriched
324 DEGs are testis and ovary genes with no known brain functions, it is currently unknown if they impair
325 *Kdm5c*-KO brain function and contribute to MRXSCJ-related phenotypes like intellectual disability and
326 aberrant aggression. However, select liver and muscle-biased DEGs do have known roles within the brain,
327 such as the liver-enriched lipid metabolism gene *Apolipoprotein C-I (Apoc1)*³² that is highly expressed in the
328 *Kdm5c*-KO amygdala and hippocampus. Intriguingly, *Apoc1* overexpression in the mouse brain can impair
329 learning and memory⁶⁷ and is implicated in Alzheimer's disease in humans³³.

330 We found a substantial portion of genes upregulated in *Kdm5c*-KO brain are germline genes with no
331 known functions within the brain. Distinguishing the germline from the soma is a key feature of multicellularity
332 and sexual reproduction¹⁶. Previous work characterizing chromatin regulators that silence germ cell-specific
333 transcription has predominantly focused on their repression of key marker genes in embryonic stem cells
334 (ESCs), such as *Dazl* and *Ddx4*^{18,19,53}. To characterize KDM5C's role in germline gene repression at a
335 genome-wide level throughout life, we curated a list of mouse germline-enriched genes using publicly

336 available germ cell-depleted RNA-seq datasets from *Kit^{W/Wv}* mice^{34,37}. This resource enabled us to identify 1)
337 the extent of germline gene dysregulation in *Kdm5c*-KO cells, 2) the types of germline genes misexpressed
338 at different developmental time points, and 3) which groups of germline genes are directly and indirectly
339 regulated by KDM5C. This list of germline-enriched genes serves as a resource for future studies to
340 systematically assess soma-germline dysregulation.

341 RNA-seq of epiblast-like cells (EpiLCs) revealed dysregulation of *Kdm5c*-KO tissue identity begins during
342 early embryogenesis. *In vivo*, germline genes are typically decommissioned in epiblast stem cells and remain
343 silenced as the epiblast differentiates into somatic tissues⁴⁰. However, a small subset of epiblast stem cells
344 will receive signals to reactivate germline gene expression to become the primordial germ cells (PGCs) that
345 will ultimately form the mature germline^{38,39}. This process can be mimicked *in vitro* by differentiating EpiLCs
346 into primordial germ cell-like cells (PGCLCs)⁴¹. Therefore, misexpression of germline genes in EpiLCs might
347 suggest *Kdm5c*-KO ESCs are progressing beyond EpiLC differentiation and instead becoming PGCLCs. Yet,
348 *Kdm5c*-KO EpiLCs properly express primed pluripotency genes, including *Otx2* - an epiblast stem cell marker
349 that is known to repress differentiation into PGCs/PGCLCs⁶⁸. Furthermore, we observed no difference in
350 cellular morphology during *Kdm5c*-KO ESC to EpiLC differentiation. Proper EpiLC differentiation, together
351 with *Kdm5c*-KO mice being viable, suggests germline gene expression is occurring ectopically in conjunction
352 with typical developmental programs, rather than a complete shift to towards germline identity.

- 353 • XX vs XY
- 354 • While many germline genes have roles in both the male and female germline, some display sex-biased
355 expression or have functions unique to eggs and sperm. Many of the germline genes characterized in
356 the male *Kdm5c*-KO brain have sperm-specific functions. It is currently unclear if sperm genes are also
357 dysregulated in female *Kdm5c* mutants, or if they would instead misexpress egg-biased genes.
- 358 • Motif analysis
 - 359 – only 1/3 had max/e2f - Unlikely to be the major recruitment mechanism for KDM5C
 - 360 – KDM5C does not contain motif-specific binding
 - 361 – controls cilia in multiple cell types, initially identified in neurons. Shared axis that could contribute
362 to NDDs
 - 363 * RFX Cilia in NDDs - <https://pubmed.ncbi.nlm.nih.gov/33658631/>
 - 364 – Enriched RFX transcription factors included the spermiogenesis regulator RFX2, whose mRNA is
365 also significantly upregulated in *Kdm5c*-KO EpiLCs (Figure 4).
 - 366 – <https://www.nature.com/articles/srep20435>
 - 367 – <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4498915/>
 - 368 – <https://pubmed.ncbi.nlm.nih.gov/32725242/>

369 We then globally characterized KDM5C binding at germline-enriched gene promoters through KDM5C
370 ChIP-seq in EpiLCs and primary hippocampus and cortex neuron cultures (PNCs).

371 Our analysis of KDM5C ChIP-seq data provided novel insights into direct and indirect mechanisms by
372 which germline genes can be misexpressed in *Kdm5c*-KO cells.

373 While we observed high enrichment of KDM5C peaks at germline genes in EpiLCs, KDM5C was not
374 bound to germline gene promoters in PNCs. This suggests dysregulation of germline genes in the mature
375 *Kdm5c*-KO brain is due to loss of repression during embryogenesis, which is consistent with previous
376 work that found introducing human KDM5C into *Kdm5c*-KO PNCs does not repress two aberrant germline
377 transcripts¹⁰. Although enriched at germline gene promoters in EpiLCs, KDM5C was only bound to about a
378 third of the germline genes aberrantly transcribed in *Kdm5c*-KO EpiLCs. One notable EpiLC DEG not bound
379 by KDM5C is *Stra8*, a transcription factor activated by retinoic acid signaling in germ cells to promote meiotic
380 initiation^{69,70}. However, retinoic acid can only activate *Stra8* expression when DAZL is present. Unlike *Stra8*,
381 *Dazl* is a direct target of KDM5C in EpiLCs and is both transcribed and translated in *Kdm5c*-KO EpiLCs.
382 Expression of indirect DEGs like *Stra8* indicates germline genes can be aberrantly expressed in *Kdm5c*-KO
383 cells through downstream activation by other ectopic germline programs. These ectopic germline programs
384 appear to loosely mimic the trajectory of typical germline development, as germline genes important for early
385 germ cell development and meiosis are expressed in *Kdm5c*-KO EpiLC while late spermatogenesis genes
386 are expressed in the mature *Kdm5c*-KO brain. Altogether, this indicates once a subset of germline genes
387 are activated due to loss of direct KDM5C repression in the early embryo, these ectopic germline programs
388 can continue to progress in the background of *Kdm5c*-KO somatic development.

389 It is important to note that while we focused on KDM5C's regulation of germline genes, some germline-
390 enriched genes are also expressed in naïve ESCs and at the 2-cell stage due to their roles in pluripotency and
391 self-renewal. For example, *Dazl* is primarily known for committing PGCs to the germline fate and regulating
392 the translation of germline-specific RNAs, but is also expressed at the 2-cell stage⁷¹, in naïve ESCs⁴³, and in
393 the inner cell mass⁴³. Very recently, two screens of *Dazl*-repressors identified KDM5C as a direct regulator in
394 ESCs^{51,72}. Interestingly, one study found *Kdm5c*-KO ESCs also aberrantly expressed 2-cell-specific genes,
395 indicating KDM5C negatively regulates totipotency⁵¹. However, out of the four regulators characterized,
396 KDM5C was the only factor whose repression of *Dazl* was independent of the 2-cell specific transcription
397 factor *Dux*⁵¹. Although expressed at the 2-cell stage and in ESCs, *Dazl* is typically silenced when ESCs
398 differentiate into EpiLCs¹⁹. We found KDM5C also represses *Dazl* in EpiLCs, but *Kdm5c*-KO EpiLCs did
399 not express 2-cell specific genes like *Dux* and *Zscan4c*. Together, this suggests the 2-cell-like state in
400 *Kdm5c*-KO ESCs is an indirect consequence of KDM5C's primary role in germline gene repression.

401 *Dazl* and other germline-enriched genes are initially decommissioned in ESCs by repressive histone
402 modifications and then *de novo* DNA CpG methylation (CpGme) in EpiLCs^{19–21,64,73,74}. Unlike the previously
403 characterized germline gene suppressors, KDM5C removes an active histone mark, histone 3 lysine 4 di-
404 and trimethylation (H3K4me2/3)¹¹. In this study, we observed a global increase in H3K4me2/3 in *Kdm5c*-KO

405 EpiLCs and amygdala around the transcription start site (TSS) of germline-enriched genes. However,
406 KDM5C's catalytic activity may not be required for germline gene silencing, as it was recently found to be
407 dispensible for repressing *Dazl* in ESCs⁵¹. Although not necessary in ESCs, KDM5C's catalytic activity be
408 required for CpGme in EpiLCs, as H3K4me3 impedes *de novo* CpGme placement^{65,66}. This is supported
409 by previous work in the *Kdm5C*-KO adult hippocampus, which found CpGme is significantly eroded at at
410 least two germline promoters¹⁰. To elucidate the mechanism behind KDM5C-mediated silencing of germline
411 genes, we characterized KDM5C in ESCs and EpiLCs and found its expression is dynamically regulated
412 during differentiation. Furthermore, KDM5C acts during the transition from naïve to primed pluripotency to
413 promote the initial placement of CpGme at CpG islands for the long-term silencing of germline genes.

414 The above work provides the mechanistic foundation for KDM5C's role in germline gene repression
415 and offers novel insight into how dysregulation in *Kdm5c*-KO tissue identity changes over the course of
416 development. Although their impact upon neurodevelopment is currently unclear, ectopic germline transcripts
417 are also found in models of other related neurodevelopmental disorders⁷⁵, including Immunodeficiency,
418 Centromere instability, Facial anomalies syndrome 1 (ICF1 OMIM: #242860)^{76,77}, Kleefstra syndrome
419 1 (OMIM: #610253)⁹, and MeCP2 duplication syndrome (MDS, OMIM: #300260)⁷⁸. Like KDM5C, the
420 chromatin regulators underlying these conditions, DNA methyltransferase 3b (DNMT3B), H3K9me1/2
421 methyltransferases G9A/GLP, methyl-CpG -binding protein 2 (MECP2), primarily silence gene expression.
422 Thus, KDM5C is among a growing cohort of chromatin-linked neurodevelopmental disorders that may have a
423 similar underlying cause of germline versus soma dysregulation. However, further research is required to
424 determine the ultimate impact of these germline genes and the extent to which this phenomenon occurs in
425 humans.

- 426 • Last paragraph
- 427 – Thus, failure to fine-tune the expression of tissue-enriched, dosage-sensitive genes like *Apoc1*
428 could be one route by which loss of brain tissue identity contributes to *Kdm5c*-KO impairments.

429 Materials and Methods

430 Classifying tissue-enriched and germline-enriched genes

431 Tissue-enriched differentially expresssd genes (DEGs) were determined by their classification in a previ-
432 ously published dataset from 17 male and female mouse tissues²⁶. This study defined tissue expression as
433 greater than 1 Fragments Per Kilobase of transcript per Million mapped read (FPKM) and tissue enrichment
434 as at least 4-fold higher expression than any other tissue.

435 We curated a list of germline-enriched genes using an RNA-seq dataset from wild-type and germline-
436 depleted (*Kit^{W/W^v}*) male and female mouse embryos from embryonic day 12, 14, and 16³⁷, as well as adult
437 male testes³⁴. Germline-enriched genes met the following criteria: 1) their expression is greater than 1

438 FPKM in wild-type germline 2) their expression in any wild-type somatic tissues²⁶ does not exceed 20%
439 of maximum expression in wild-type germline, and 3) their expression in the germ cell-depleted (Kit^{W/Wv})
440 germline, for any sex or time point, does not exceed 20% of maximum expression in wild-type germline.

441 **Cell culture**

442 We utilized our previously established cultures of male wild-type and *Kdm5c* knockout (-KO) embryonic
443 stem cells⁴⁷. Sex was confirmed by genotyping *Uba1/Uba1y* on the X and Y chromosomes with the following
444 primers: 5'-TGGATGGTGTGCCAATG-3', 5'-CACCTGCACGTTGCCCT-3'. Deletion of *Kdm5c* was
445 confirmed through the primers 5'-ATGCCCATATTAAGAGTCCTG-3', 5'-TCTGCCTTGATGGGACTGTT-3',
446 and 5'-GGTTCTCAACACTCACATAGTG-3'.

447 Embryonic stem cells (ESCs) and epiblast-like cells were cultured using previously established
448 methods⁴². Briefly, ESCs were initially cultured in primed ESC (pESC) media consisting of KnockOut
449 DMEM (Gibco#10829–018), fetal bovine serum (Gibco#A5209501), KnockOut serum replacement
450 (Invitrogen#10828–028), Glutamax (Gibco#35050-061), Anti-Anti (Gibco#15240-062), MEM Non-essential
451 amino acids (Gibco#11140-050), and beta-mercaptoethanol (Sigma#M7522). They were then transitioned
452 into ground-state, “naïve” ESCs (nESCs) by culturing for four passages in N2B27 media containing
453 DMEM/F12 (Gibco#11330–032), Neurobasal media (Gibco#21103–049), Gluamax, Anti-Anti, N2 sup-
454 plement (Invitrogen#17502048), and B27 supplement without vitamin A (Invitrogen#12587-010), and
455 beta-mercaptoethanol. Both pESC and nESC media were supplemented with 3 μM GSK3 inhibitor
456 CHIR99021 (Sigma #SML1046-5MG), 1 μM MEK inhibitor PD0325901 (Sigma #PZ0162-5MG), and 1,000
457 units/mL leukemia inhibitory factor (LIF, Millipore#ESG1107).

458 nESCs were differentiated into epiblast-like cells (EpiLCs, 48 hours) and extended EpiLCs (exEpiLCs,
459 96 hours) by culturing in N2B27 media containing DMEM/F12, Neurobasal media, Gluamax, Anti-Anti,
460 N2 supplement, B27 supplement (Invitrogen#17504044), beta-mercaptoethanol, fibroblast growth factor 2
461 (FGF2, R&D Biotechne 233-FB), and activin A (R&D Biotechne 338AC050CF), as previously described⁴².

462 **Immunocytochemistry (ICC)**

463 ICC of DAZL in EpiLCs was performed by first growing cells on fibronectin-coated coverslips. Cells were
464 then washed thrice with phosphobuffered saline (PBS), fixed in 4% paraformaldehyde, washed thrice in PBS,
465 and blocked in PBS containing 0.3% Triton X-100, and 5% fetal bovine serum for 1 hour. They were then
466 washed thrice with PBS and incubating in primary antibody (Rabbit anti DAZL, abcam ab34139, 1:200) in
467 the blocking solution overnight at 4°C with gentle rocking. The next day, cells were rinsed thrice with PBS,
468 and incubated in secondary antibody (Alexafluor 488 Invitrogen #710369, 1:1,000) with DAPI (1:1,000) in
469 blocking buffer for 1 hour at room temperature. Coverslips were then rinsed thrice in PBS, and mounted onto
470 slides using Prolong Gold (Invitrogen #P36930). Images were taken blinded for genotype, chosen based on

471 similar levels of DAPI signal, and quantified via ImageJ before unblinding.

472 RNA sequencing (RNA-seq)

473 After ensuring read quality via FastQC (v0.11.8), reads were then mapped to the mm10 *Mus musculus*
474 genome (Gencode) using STAR (v2.5.3a), during which we removed duplicates and kept only uniquely
475 mapped reads. Count files were generated by FeatureCounts (Subread v1.5.0), and BAM files were
476 converted to bigwigs using deeptools (v3.1.3) and visualized by the UCSC genome browser. RStudio (v3.6.0)
477 was then used to analyze counts files by DESeq2 (v1.26.0)²⁸ to identify differentially expressed genes
478 (DEGs) with a q-value (p-adjusted via FDR/Benjamini–Hochberg correction) less than 0.1 and a log2 fold
479 change greater than 0.5. For all DESeq2 analyses, log2 fold changes were calculated with IfcShrink using
480 the ashr package⁷⁹. MA-plots were generated by ggpubr (v0.6.0), and Eulerr diagrams were generated by
481 eulerr (v6.1.1). Boxplots and scatterplots were generated by ggpubr (v0.6.0) and ggplot2 (v3.3.2). The Upset
482 plot was generated via the package UpSetR (v1.4.0)⁸⁰. Gene ontology (GO) analyses were performed by
483 the R package enrichPlot (v1.16.2) using the biological processes setting and compareCluster.

484 Chromatin immunoprecipitation followed by DNA sequencing (ChIP-seq)

485 ChIP-seq reads were aligned to mm10 using Bowtie1 (v1.1.2) allowing up to two mismatches. Only
486 uniquely mapped reads were used for analysis. Peaks were called using MACS2 software (v2.2.9.1) using
487 input BAM files for normalization, with filters for a q-value < 0.1 and a fold enrichment > 1. We removed
488 “black-listed” genomic regions that often give aberrant signals. Common peak sets were obtained in R via
489 DiffBind (v3.6.5). In the case of KDM5C ChIP-seq, *Kdm5c*-KO peaks were then subtracted from wild-type
490 samples using bedtools (v2.25.0). Peak proximity to genome annotations was determined by ChIPSeeker
491 (v1.32.1). Gene ontology (GO) analyses were performed by the R package enrichPlot (v1.16.2) using the
492 biological processes setting and compareCluster. Enriched motifs were identified using HOMER⁵⁸. Average
493 binding across the genome was visualized using deeptools (v3.1.3). Bigwigs were visualized using the
494 UCSC genome browser.

495 Whole genome bisulfite sequencing (WGBS)

496 Genomic DNA (gDNA) from naïve ESCs and extended EpiLCs was extracted using the Wizard Genomic
497 DNA Purification Kit (Promega A1120), following the instructions for Tissue Culture Cells. gDNA was sent
498 to Novogene for WGBS using the Illumina NovaSeq X Plus platform and sequenced for 150bp paired-end
499 reads (PE150).

500 **Data availability**

501 **Published datasets**

502 All published datasets are available at the Gene Expression Omnibus (GEO) <https://www.ncbi.nlm.nih.gov/geo>. Published RNA-seq datasets analyzed in this study included the male wild-type and *Kdm5c*-KO
503 adult amygdala and hippocampus²⁷ (available at GEO: GSE127722) and male wild-type and *Kdm5c*-KO
504 EpiLCs⁴⁷ (available at GEO: GSE96797).

505 Previously published ChIP-seq experiments included KDM5C in wild-type and *Kdm5c*-KO EpiLCs⁴⁷ (avail-
506 able at GEO: GSE96797) and mouse primary neuron cultures (PNCs) from the cortex and hippocampus¹⁵
507 (available at GEO: GSE61036). ChIP-seq of histone 3 lysine 4 dimethylation in male wild-type and *Kdm5c*-KO
508 EpiLCs⁴⁷ is also available at GEO: GSE96797. ChIP-seq of histone 3 lysine 4 trimethylation in wild-type and
509 *Kdm5c*-KO male amygdala²⁷ are available at GEO: GSE127817.

511 **Data analysis**

512 Scripts used to generate the results, tables, and figures of this study are available via a GitHub repository:
513 XXX

514 **Acknowledgements**

- 515 • Jacob Mueller for providing insight in germline gene regulation.
516 • Sundeep Kalantry for providing reagents and expertise in culturing mouse embryonic stem cells and
517 epiblast-like cells
518 • Funding sources

519 **References**

- 520 1. Strahl, B.D., and Allis, C.D. (2000). The language of covalent histone modifications. *Nature* **403**,
521 41–45. <https://doi.org/10.1038/47412>.
- 522 2. Jenuwein, T., and Allis, C.D. (2001). Translating the Histone Code. *Science* **293**, 1074–1080.
523 <https://doi.org/10.1126/science.1063127>.
- 524 3. Lewis, E.B. (1978). A gene complex controlling segmentation in *Drosophila*. *Nature* **276**, 565–570.
525 <https://doi.org/10.1038/276565a0>.
- 526 4. Kennison, J.A., and Tamkun, J.W. (1988). Dosage-dependent modifiers of polycomb and antennapedia
527 mutations in *Drosophila*. *Proc. Natl. Acad. Sci. U.S.A.* **85**, 8136–8140. <https://doi.org/10.1073/pnas.85.21.8136>.

- 528 5. Kassis, J.A., Kennison, J.A., and Tamkun, J.W. (2017). Polycomb and Trithorax Group Genes in
529 *Drosophila*. *Genetics* 206, 1699–1725. <https://doi.org/10.1534/genetics.115.185116>.
- 530 6. Gabriele, M., Lopez Tobon, A., D'Agostino, G., and Testa, G. (2018). The chromatin basis of
531 neurodevelopmental disorders: Rethinking dysfunction along the molecular and temporal axes. *Prog
Neuropsychopharmacol Biol Psychiatry* 84, 306–327. <https://doi.org/10.1016/j.pnpbp.2017.12.013>.
- 532 7. Zhou, Z., Hong, E.J., Cohen, S., Zhao, W.-N., Ho, H.-Y.H., Schmidt, L., Chen, W.G., Lin, Y., Savner,
E., Griffith, E.C., et al. (2006). Brain-specific phosphorylation of MeCP2 regulates activity-dependent
533 Bdnf transcription, dendritic growth, and spine maturation. *Neuron* 52, 255–269. <https://doi.org/10.1016/j.neuron.2006.09.037>.
- 534 8. Hirabayashi, Y., Suzuki, N., Tsuboi, M., Endo, T.A., Toyoda, T., Shinga, J., Koseki, H., Vidal, M., and
535 Gotoh, Y. (2009). Polycomb Limits the Neurogenic Competence of Neural Precursor Cells to Promote
Astrogenic Fate Transition. *Neuron* 63, 600–613. <https://doi.org/10.1016/j.neuron.2009.08.021>.
- 536 9. Schaefer, A., Sampath, S.C., Intrator, A., Min, A., Gertler, T.S., Surmeier, D.J., Tarakhovsky, A.,
537 and Greengard, P. (2009). Control of cognition and adaptive behavior by the GLP/G9a epigenetic
suppressor complex. *Neuron* 64, 678–691. <https://doi.org/10.1016/j.neuron.2009.11.019>.
- 538 10. Scandaglia, M., Lopez-Atalaya, J.P., Medrano-Fernandez, A., Lopez-Cascales, M.T., Del Blanco, B.,
Lipinski, M., Benito, E., Olivares, R., Iwase, S., Shi, Y., et al. (2017). Loss of Kdm5c Causes Spurious
539 Transcription and Prevents the Fine-Tuning of Activity-Regulated Enhancers in Neurons. *Cell Rep* 21,
47–59. <https://doi.org/10.1016/j.celrep.2017.09.014>.
- 540 11. Iwase, S., Lan, F., Bayliss, P., De La Torre-Ubieta, L., Huarte, M., Qi, H.H., Whetstine, J.R., Bonni, A.,
Roberts, T.M., and Shi, Y. (2007). The X-Linked Mental Retardation Gene SMCX/JARID1C Defines a
Family of Histone H3 Lysine 4 Demethylases. *Cell* 128, 1077–1088. <https://doi.org/10.1016/j.cell.2007.02.017>.
- 541 12. Claes, S., Devriendt, K., Van Goethem, G., Roelen, L., Meireleire, J., Raeymaekers, P., Cassiman,
J.J., and Fryns, J.P. (2000). Novel syndromic form of X-linked complicated spastic paraplegia. *Am J
Med Genet* 94, 1–4.
- 542 13. Jensen, L.R., Amende, M., Gurok, U., Moser, B., Gimmel, V., Tzschach, A., Janecke, A.R., Tariverdian,
G., Chelly, J., Fryns, J.P., et al. (2005). Mutations in the JARID1C gene, which is involved in
transcriptional regulation and chromatin remodeling, cause X-linked mental retardation. *Am J Hum
Genet* 76, 227–236. <https://doi.org/10.1086/427563>.
- 543 14. Carmignac, V., Nambot, S., Lehalle, D., Callier, P., Moortgat, S., Benoit, V., Ghoumid, J., Delobel,
B., Smol, T., Thuillier, C., et al. (2020). Further delineation of the female phenotype with KDM5C
disease causing variants: 19 new individuals and review of the literature. *Clin Genet* 98, 43–55.
544 <https://doi.org/10.1111/cge.13755>.

- 548 15. Iwase, S., Brookes, E., Agarwal, S., Badeaux, A.I., Ito, H., Vallianatos, C.N., Tomassy, G.S., Kasza, T.,
Lin, G., Thompson, A., et al. (2016). A Mouse Model of X-linked Intellectual Disability Associated with
Impaired Removal of Histone Methylation. *Cell Reports* *14*, 1000–1009. <https://doi.org/10.1016/j.celrep.2015.12.091>.
549
- 550 16. Devlin, D.K., Ganley, A.R.D., and Takeuchi, N. (2023). A pan-metazoan view of germline-soma
distinction challenges our understanding of how the metazoan germline evolves. *Current Opinion in
551 Systems Biology* *36*, 100486. <https://doi.org/10.1016/j.coisb.2023.100486>.
- 552 17. Lehmann, R. (2012). Germline Stem Cells: Origin and Destiny. *Cell Stem Cell* *10*, 729–739.
553 <https://doi.org/10.1016/j.stem.2012.05.016>.
- 554 18. Endoh, M., Endo, T.A., Shinga, J., Hayashi, K., Farcas, A., Ma, K.W., Ito, S., Sharif, J., Endoh, T.,
Onaga, N., et al. (2017). PCGF6-PRC1 suppresses premature differentiation of mouse embryonic
555 stem cells by regulating germ cell-related genes. *eLife* *6*. <https://doi.org/10.7554/eLife.21064>.
- 556 19. Mochizuki, K., Sharif, J., Shirane, K., Uranishi, K., Bogutz, A.B., Janssen, S.M., Suzuki, A., Okuda,
A., Koseki, H., and Lorincz, M.C. (2021). Repression of germline genes by PRC1.6 and SETDB1
in the early embryo precedes DNA methylation-mediated silencing. *Nat Commun* *12*, 7020. <https://doi.org/10.1038/s41467-021-27345-x>.
557
- 558 20. Velasco, G., Hubé, F., Rollin, J., Neuillet, D., Philippe, C., Bouzinba-Segard, H., Galvani, A., Viegas-
Péquignot, E., and Francastel, C. (2010). Dnmt3b recruitment through E2F6 transcriptional repressor
mediates germ-line gene silencing in murine somatic tissues. *Proc Natl Acad Sci U S A* *107*, 9281–
559 9286. <https://doi.org/10.1073/pnas.1000473107>.
- 560 21. Hackett, J.A., Reddington, J.P., Nestor, C.E., Dunican, D.S., Branco, M.R., Reichmann, J., Reik,
W., Surani, M.A., Adams, I.R., and Meehan, R.R. (2012). Promoter DNA methylation couples
genome-defence mechanisms to epigenetic reprogramming in the mouse germline. *Development*
561 *139*, 3623–3632. <https://doi.org/10.1242/dev.081661>.
- 562 22. Agulnik, A.I., Mitchell, M.J., Mattei, M.G., Borsani, G., Avner, P.A., Lerner, J.L., and Bishop, C.E.
(1994). A novel X gene with a widely transcribed Y-linked homologue escapes X-inactivation in mouse
563 and human. *Hum Mol Genet* *3*, 879–884. <https://doi.org/10.1093/hmg/3.6.879>.
- 564 23. Carrel, L., Hunt, P.A., and Willard, H.F. (1996). Tissue and lineage-specific variation in inactive
X chromosome expression of the murine Smcx gene. *Hum Mol Genet* *5*, 1361–1366. <https://doi.org/10.1093/hmg/5.9.1361>.
565
- 566 24. Sheardown, S., Norris, D., Fisher, A., and Brockdorff, N. (1996). The mouse Smcx gene exhibits
developmental and tissue specific variation in degree of escape from X inactivation. *Hum Mol Genet*
567 *5*, 1355–1360. <https://doi.org/10.1093/hmg/5.9.1355>.

- 568 25. Xu, J., Deng, X., and Disteche, C.M. (2008). Sex-Specific Expression of the X-Linked Histone
569 Demethylase Gene Jarid1c in Brain. PLoS ONE 3, e2553. <https://doi.org/10.1371/journal.pone.0002553>
- 570 26. Li, B., Qing, T., Zhu, J., Wen, Z., Yu, Y., Fukumura, R., Zheng, Y., Gondo, Y., and Shi, L. (2017). A
571 Comprehensive Mouse Transcriptomic BodyMap across 17 Tissues by RNA-seq. Sci Rep 7, 4200.
<https://doi.org/10.1038/s41598-017-04520-z>.
- 572 27. Vallianatos, C.N., Raines, B., Porter, R.S., Bonefas, K.M., Wu, M.C., Garay, P.M., Collette, K.M.,
573 Seo, Y.A., Dou, Y., Keegan, C.E., et al. (2020). Mutually suppressive roles of KMT2A and KDM5C
in behaviour, neuronal structure, and histone H3K4 methylation. Commun Biol 3, 278. <https://doi.org/10.1038/s42003-020-1001-6>.
- 574 28. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for
575 RNA-seq data with DESeq2. Genome Biol 15, 550. <https://doi.org/10.1186/s13059-014-0550-8>.
- 576 29. Crackower, M.A., Kolas, N.K., Noguchi, J., Sarao, R., Kikuchi, K., Kaneko, H., Kobayashi, E., Kawai, Y.,
577 Kozieradzki, I., Landers, R., et al. (2003). Essential Role of Fkbp6 in Male Fertility and Homologous
Chromosome Pairing in Meiosis. Science 300, 1291–1295. <https://doi.org/10.1126/science.1083022>.
- 578 30. Xiol, J., Cora, E., Koglgruber, R., Chuma, S., Subramanian, S., Hosokawa, M., Reuter, M., Yang, Z.,
Berninger, P., Palencia, A., et al. (2012). A Role for Fkbp6 and the Chaperone Machinery in piRNA
579 Amplification and Transposon Silencing. Molecular Cell 47, 970–979. <https://doi.org/10.1016/j.molcel.2012.07.019>.
- 580 31. Cheng, S., Altmeppen, G., So, C., Welp, L.M., Penir, S., Ruhwedel, T., Menelaou, K., Harasimov, K.,
581 Stützer, A., Blayney, M., et al. (2022). Mammalian oocytes store mRNAs in a mitochondria-associated
membraneless compartment. Science 378, eabq4835. <https://doi.org/10.1126/science.abq4835>.
- 582 32. Rouland, A., Masson, D., Lagrost, L., Vergès, B., Gautier, T., and Bouillet, B. (2022). Role of
583 apolipoprotein C1 in lipoprotein metabolism, atherosclerosis and diabetes: A systematic review.
Cardiovasc Diabetol 21, 272. <https://doi.org/10.1186/s12933-022-01703-5>.
- 584 33. Leduc, V., Jasmin-Bélanger, S., and Poirier, J. (2010). APOE and cholesterol homeostasis in
585 Alzheimer's disease. Trends in Molecular Medicine 16, 469–477. <https://doi.org/10.1016/j.molmed.2010.07.008>.
- 586 34. Mueller, J.L., Skaletsky, H., Brown, L.G., Zaghlul, S., Rock, S., Graves, T., Auger, K., Warren,
587 W.C., Wilson, R.K., and Page, D.C. (2013). Independent specialization of the human and mouse X
chromosomes for the male germ line. Nat Genet 45, 1083–1087. <https://doi.org/10.1038/ng.2705>.
- 588 35. Handel, M.A., and Eppig, J.J. (1979). Sertoli Cell Differentiation in the Testes of Mice Genetically
589 Deficient in Germ Cells. Biology of Reproduction 20, 1031–1038. <https://doi.org/10.1095/biolreprod20.5.1031>.

- 590 36. Green, C.D., Ma, Q., Manske, G.L., Shami, A.N., Zheng, X., Marini, S., Moritz, L., Sultan, C.,
Gurczynski, S.J., Moore, B.B., et al. (2018). A Comprehensive Roadmap of Murine Spermatogenesis
Defined by Single-Cell RNA-Seq. *Dev Cell* *46*, 651–667.e10. <https://doi.org/10.1016/j.devcel.2018.07.025>.
- 591 37. Soh, Y.Q., Junker, J.P., Gill, M.E., Mueller, J.L., van Oudenaarden, A., and Page, D.C. (2015). A
Gene Regulatory Program for Meiotic Prophase in the Fetal Ovary. *PLoS Genet* *11*, e1005531.
<https://doi.org/10.1371/journal.pgen.1005531>.
- 592 38. Magnúsdóttir, E., and Surani, M.A. (2014). How to make a primordial germ cell. *Development* *141*,
245–252. <https://doi.org/10.1242/dev.098269>.
- 593 39. Günesdogan, U., Magnúsdóttir, E., and Surani, M.A. (2014). Primordial germ cell specification: A
context-dependent cellular differentiation event [corrected]. *Philos Trans R Soc Lond B Biol Sci* *369*.
<https://doi.org/10.1098/rstb.2013.0543>.
- 594 40. Bardot, E.S., and Hadjantonakis, A.-K. (2020). Mouse gastrulation: Coordination of tissue patterning,
specification and diversification of cell fate. *Mechanisms of Development* *163*, 103617. <https://doi.org/10.1016/j.mod.2020.103617>.
- 595 41. Hayashi, K., Ohta, H., Kurimoto, K., Aramaki, S., and Saitou, M. (2011). Reconstitution of the
mouse germ cell specification pathway in culture by pluripotent stem cells. *Cell* *146*, 519–532.
<https://doi.org/10.1016/j.cell.2011.06.052>.
- 596 42. Samanta, M., and Kalantry, S. (2020). Generating primed pluripotent epiblast stem cells: A methodol-
ogy chapter. *Curr Top Dev Biol* *138*, 139–174. <https://doi.org/10.1016/bs.ctdb.2020.01.005>.
- 597 43. Welling, M., Chen, H., Muñoz, J., Musheev, M.U., Kester, L., Junker, J.P., Mischerikow, N., Arbab, M.,
Kuijk, E., Silberstein, L., et al. (2015). DAZL regulates Tet1 translation in murine embryonic stem cells.
EMBO Reports *16*, 791–802. <https://doi.org/10.15252/embr.201540538>.
- 598 44. Macfarlan, T.S., Gifford, W.D., Driscoll, S., Lettieri, K., Rowe, H.M., Bonanomi, D., Firth, A., Singer, O.,
Trono, D., and Pfaff, S.L. (2012). Embryonic stem cell potency fluctuates with endogenous retrovirus
activity. *Nature* *487*, 57–63. <https://doi.org/10.1038/nature11244>.
- 599 45. Suzuki, A., Hirasaki, M., Hishida, T., Wu, J., Okamura, D., Ueda, A., Nishimoto, M., Nakachi, Y.,
Mizuno, Y., Okazaki, Y., et al. (2016). Loss of MAX results in meiotic entry in mouse embryonic and
germline stem cells. *Nat Commun* *7*, 11056. <https://doi.org/10.1038/ncomms11056>.
- 600 46. Borgel, J., Guibert, S., Li, Y., Chiba, H., Schübeler, D., Sasaki, H., Forné, T., and Weber, M. (2010).
Targets and dynamics of promoter DNA methylation during early mouse development. *Nat Genet* *42*,
1093–1100. <https://doi.org/10.1038/ng.708>.

- 612 47. Samanta, M.K., Gayen, S., Harris, C., Maclary, E., Murata-Nakamura, Y., Malcore, R.M., Porter, R.S.,
Garay, P.M., Vallianatos, C.N., Samollow, P.B., et al. (2022). Activation of Xist by an evolutionarily
conserved function of KDM5C demethylase. *Nat Commun* *13*, 2602. <https://doi.org/10.1038/s41467-022-30352-1>.
- 613
- 614 48. Koubova, J., Menke, D.B., Zhou, Q., Capel, B., Griswold, M.D., and Page, D.C. (2006). Retinoic
acid regulates sex-specific timing of meiotic initiation in mice. *Proc. Natl. Acad. Sci. U.S.A.* *103*,
615 2474–2479. <https://doi.org/10.1073/pnas.0510813103>.
- 616 49. Lin, Y., Gill, M.E., Koubova, J., and Page, D.C. (2008). Germ Cell-Intrinsic and -Extrinsic Factors
Govern Meiotic Initiation in Mouse Embryos. *Science* *322*, 1685–1687. <https://doi.org/10.1126/science.1166340>.
- 617
- 618 50. Endo, T., Mikedis, M.M., Nicholls, P.K., Page, D.C., and De Rooij, D.G. (2019). Retinoic Acid and Germ
Cell Development in the Ovary and Testis. *Biomolecules* *9*, 775. <https://doi.org/10.3390/biom9120775>.
- 619
- 620 51. Gupta, N., Yakhou, L., Albert, J.R., Azogui, A., Ferry, L., Kirsh, O., Miura, F., Battault, S., Yamaguchi,
K., Laisné, M., et al. (2023). A genome-wide screen reveals new regulators of the 2-cell-like cell state.
621 *Nat Struct Mol Biol*. <https://doi.org/10.1038/s41594-023-01038-z>.
- 622 52. Pohlers, M., Truss, M., Frede, U., Scholz, A., Strehle, M., Kuban, R.-J., Hoffmann, B., Morkel, M.,
Birchmeier, C., and Hagemeier, C. (2005). A Role for E2F6 in the Restriction of Male-Germ-Cell-
623 Specific Gene Expression. *Current Biology* *15*, 1051–1057. <https://doi.org/10.1016/j.cub.2005.04.060>.
- 624 53. Dahlet, T., Truss, M., Frede, U., Al Adhami, H., Bardet, A.F., Dumas, M., Vallet, J., Chicher, J.,
Hammann, P., Kottnik, S., et al. (2021). E2F6 initiates stable epigenetic silencing of germline genes
625 during embryonic development. *Nat Commun* *12*, 3582. <https://doi.org/10.1038/s41467-021-23596-w>.
- 626 54. Wang, P.J., McCarrey, J.R., Yang, F., and Page, D.C. (2001). An abundance of X-linked genes
627 expressed in spermatogonia. *Nat Genet* *27*, 422–426. <https://doi.org/10.1038/86927>.
- 628 55. Khil, P.P., Smirnova, N.A., Romanienko, P.J., and Camerini-Otero, R.D. (2004). The mouse X
chromosome is enriched for sex-biased genes not subject to selection by meiotic sex chromosome
629 inactivation. *Nat Genet* *36*, 642–646. <https://doi.org/10.1038/ng1368>.
- 630 56. Hurlin, P.J. (1999). Mga, a dual-specificity transcription factor that interacts with Max and contains a
T-domain DNA-binding motif. *The EMBO Journal* *18*, 7019–7028. <https://doi.org/10.1093/emboj/18.2.4.7019>.
- 631
- 632 57. Stielow, B., Finkernagel, F., Stiewe, T., Nist, A., and Suske, G. (2018). MGA, L3MBTL2 and E2F6
determine genomic binding of the non-canonical Polycomb repressive complex PRC1.6. *PLoS Genet*
633 *14*, e1007193. <https://doi.org/10.1371/journal.pgen.1007193>.

- 634 58. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H., and Glass, C.K. (2010). Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Molecular Cell* 38, 576–589.
- 635 https://doi.org/10.1016/j.molcel.2010.05.004.
- 636 59. Gajiwala, K.S., Chen, H., Cornille, F., Roques, B.P., Reith, W., Mach, B., and Burley, S.K. (2000). Structure of the winged-helix protein hRFX1 reveals a new mode of DNA binding. *Nature* 403, 916–921. https://doi.org/10.1038/35002634.
- 637
- 638 60. Swoboda, P., Adler, H.T., and Thomas, J.H. (2000). The RFX-Type Transcription Factor DAF-19 Regulates Sensory Neuron Cilium Formation in *C. elegans*. *Molecular Cell* 5, 411–421. https://doi.org/10.1016/S1097-2765(00)80436-0.
- 639
- 640 61. Ashique, A.M., Choe, Y., Karlen, M., May, S.R., Phamluong, K., Solloway, M.J., Ericson, J., and Peterson, A.S. (2009). The Rfx4 Transcription Factor Modulates Shh Signaling by Regional Control of Ciliogenesis. *Sci. Signal.* 2. https://doi.org/10.1126/scisignal.2000602.
- 641
- 642 62. Kistler, W.S., Baas, D., Lemeille, S., Paschaki, M., Seguin-Estevez, Q., Barras, E., Ma, W., Duteyrat, J.-L., Morlé, L., Durand, B., et al. (2015). RFX2 Is a Major Transcriptional Regulator of Spermiogenesis. *PLoS Genet* 11, e1005368. https://doi.org/10.1371/journal.pgen.1005368.
- 643
- 644 63. Wu, Y., Hu, X., Li, Z., Wang, M., Li, S., Wang, X., Lin, X., Liao, S., Zhang, Z., Feng, X., et al. (2016). Transcription Factor RFX2 Is a Key Regulator of Mouse Spermiogenesis. *Sci Rep* 6, 20435. https://doi.org/10.1038/srep20435.
- 645
- 646 64. Liu, M., Zhu, Y., Xing, F., Liu, S., Xia, Y., Jiang, Q., and Qin, J. (2020). The polycomb group protein PCGF6 mediates germline gene silencing by recruiting histone-modifying proteins to target gene promoters. *J Biol Chem* 295, 9712–9724. https://doi.org/10.1074/jbc.RA119.012121.
- 647
- 648 65. Otani, J., Nankumo, T., Arita, K., Inamoto, S., Ariyoshi, M., and Shirakawa, M. (2009). Structural basis for recognition of H3K4 methylation status by the DNA methyltransferase 3A ATRX–DNMT3–DNMT3L domain. *EMBO Reports* 10, 1235–1241. https://doi.org/10.1038/embor.2009.218.
- 649
- 650 66. Guo, X., Wang, L., Li, J., Ding, Z., Xiao, J., Yin, X., He, S., Shi, P., Dong, L., Li, G., et al. (2015). Structural insight into autoinhibition and histone H3-induced activation of DNMT3A. *Nature* 517, 640–644. https://doi.org/10.1038/nature13899.
- 651
- 652 67. Abildayeva, K., Berbée, J.F.P., Blokland, A., Jansen, P.J., Hoek, F.J., Meijer, O., Lütjohann, D., Gautier, T., Pillot, T., De Vente, J., et al. (2008). Human apolipoprotein C-I expression in mice impairs learning and memory functions. *Journal of Lipid Research* 49, 856–869. https://doi.org/10.1194/jlr.M700518JLR200.
- 653
- 654 68. Zhang, J., Zhang, M., Acampora, D., Vojtek, M., Yuan, D., Simeone, A., and Chambers, I. (2018). OTX2 restricts entry to the mouse germline. *Nature* 562, 595–599. https://doi.org/10.1038/s41586-018-0581-5.

- 655
- 656 69. Endo, T., Romer, K.A., Anderson, E.L., Baltus, A.E., De Rooij, D.G., and Page, D.C. (2015). Pe-
657 riodic retinoic acid–STRA8 signaling intersects with periodic germ-cell competencies to regulate
658 spermatogenesis. *Proc. Natl. Acad. Sci. U.S.A.* **112**. <https://doi.org/10.1073/pnas.1505683112>.
- 659 70. Bowles, J., Knight, D., Smith, C., Wilhelm, D., Richman, J., Mamiya, S., Yashiro, K., Chawengsak-
660 sophak, K., Wilson, M.J., Rossant, J., et al. (2006). Retinoid Signaling Determines Germ Cell Fate in
661 Mice. *Science* **312**, 596–600. <https://doi.org/10.1126/science.1125691>.
- 662 71. Yang, M., Yu, H., Yu, X., Liang, S., Hu, Y., Luo, Y., Izsvák, Z., Sun, C., and Wang, J. (2022). Chemical-
663 induced chromatin remodeling reprograms mouse ESCs to totipotent-like stem cells. *Cell Stem Cell* **29**, 400–418.e13. <https://doi.org/10.1016/j.stem.2022.01.010>.
- 664 72. Al Adhami, H., Vallet, J., Schaal, C., Schumacher, P., Bardet, A.F., Dumas, M., Chicher, J., Hammann,
665 P., Daujat, S., and Weber, M. (2023). Systematic identification of factors involved in the silencing
666 of germline genes in mouse embryonic stem cells. *Nucleic Acids Research* **51**, 3130–3149. <https://doi.org/10.1093/nar/gkad071>.
- 667 73. Karimi, M.M., Goyal, P., Maksakova, I.A., Bilenky, M., Leung, D., Tang, J.X., Shinkai, Y., Mager, D.L.,
668 Jones, S., Hirst, M., et al. (2011). DNA Methylation and SETDB1/H3K9me3 Regulate Predominantly
669 Distinct Sets of Genes, Retroelements, and Chimeric Transcripts in mESCs. *Cell Stem Cell* **8**,
670 676–687. <https://doi.org/10.1016/j.stem.2011.04.004>.
- 671 74. Tatsumi, D., Hayashi, Y., Endo, M., Kobayashi, H., Yoshioka, T., Kiso, K., Kanno, S., Nakai, Y., Maeda,
672 I., Mochizuki, K., et al. (2018). DNMTs and SETDB1 function as co-repressors in MAX-mediated
673 repression of germ cell-related genes in mouse embryonic stem cells. *PLoS ONE* **13**, e0205969.
674 <https://doi.org/10.1371/journal.pone.0205969>.
- 675 75. Bonefas, K.M., and Iwase, S. (2021). Soma-to-germline transformation in chromatin-linked neurode-
676 velopmental disorders? *FEBS J.* <https://doi.org/10.1111/febs.16196>.
- 677 76. Velasco, G., Walton, E.L., Sterlin, D., Hédon, S., Nitta, H., Ito, Y., Fouyssac, F., Mégarbané, A.,
678 Sasaki, H., Picard, C., et al. (2014). Germline genes hypomethylation and expression define a
679 molecular signature in peripheral blood of ICF patients: Implications for diagnosis and etiology.
680 *Orphanet J Rare Dis* **9**, 56. <https://doi.org/10.1186/1750-1172-9-56>.
- 681 77. Walton, E.L., Francastel, C., and Velasco, G. (2014). Dnmt3b Prefers Germ Line Genes and Cen-
682 tromeric Regions: Lessons from the ICF Syndrome and Cancer and Implications for Diseases. *Biology*
683 (Basel) **3**, 578–605. <https://doi.org/10.3390/biology3030578>.
- 684 78. Samaco, R.C., Mandel-Brehm, C., McGraw, C.M., Shaw, C.A., McGill, B.E., and Zoghbi, H.Y. (2012).
685 Crh and Oprm1 mediate anxiety-related behavior and social approach in a mouse model of MECP2
686 duplication syndrome. *Nat Genet* **44**, 206–211. <https://doi.org/10.1038/ng.1066>.

- 676 79. Stephens, M. (2016). False discovery rates: A new deal. Biostat, kxw041. <https://doi.org/10.1093/biostatistics/kxw041>.
- 677
- 678 80. Conway, J.R., Lex, A., and Gehlenborg, N. (2017). UpSetR: An R package for the visualization of intersecting sets and their properties. Bioinformatics 33, 2938–2940. <https://doi.org/10.1093/bioinformatics/btx364>.
- 679
- 680 81. Li, H., Liang, Z., Yang, J., Wang, D., Wang, H., Zhu, M., Geng, B., and Xu, E.Y. (2019). DAZL is a master translational regulator of murine spermatogenesis. Natl Sci Rev 6, 455–468. <https://doi.org/10.1093/nsr/nwy163>.
- 681
- 682 82. Mikedis, M.M., Fan, Y., Nicholls, P.K., Endo, T., Jackson, E.K., Cobb, S.A., De Rooij, D.G., and Page, D.C. (2020). DAZL mediates a broad translational program regulating expansion and differentiation of spermatogonial progenitors. eLife 9, e56523. <https://doi.org/10.7554/eLife.56523>.
- 683

684 **Figures and Tables**

- 685 • Supplementary table 1: list of all germline genes.
- 686 – Columns to include:
- 687 * KDM5C bound vs not
- 688 * DEG in EpiLC, brain, both, neither (separate columns?)



Figure 1: Tissue-enriched genes are misexpressed in the *Kdm5c*-KO brain. **A-B.** Expression of tissue-enriched genes (Li et al 2017) in the male *Kdm5c*-KO amygdala (A) and hippocampus (B). Left - MA plot of mRNA-sequencing. Right - Number of tissue-enriched differentially expressed genes (DEGs). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, Fisher's exact test. **C.** Left - Average bigwigs of an example aberrantly expressed testis-enriched DEG, *FK506 binding protein 6* (*Fkbp6*) in the wild-type (WT) and *Kdm5c*-KO (5CKO) amygdala (red) and hippocampus (teal). Right - Expression of *Cyct* in wild-type tissues from NCBI Gene, with testis highlighted in blue and brain tissues highlighted in red. **D.** Left - Average bigwigs of an example ovary-enriched DEG, *Zygotic arrest 1* (*Zar1*). Right - Expression of *Zar1* in wild-type tissues from NCBI Gene, with ovary highlighted in teal and brain tissues highlighted in red. **E.** Left - Average bigwigs of an example liver-enriched DEG, *Apolipoprotein C-I* (*Apoc1*). Right - Expression of *Apoc1* in wild-type tissues from NCBI Gene, with liver highlighted in orange and brain tissues highlighted in red.



Figure 2: Aberrant transcription of germline genes in the *Kdm5c*-KO in the brain. **A.** enrichPlot gene ontology (GO) of *Kdm5c*-KO amygdala and hippocampus testis-enriched DEGs **B.** Expression of testis DEGs in wild-type (WT) testis versus germ cell-depleted (W/Wv) testis (Mueller et al 2013). Expression is in Fragments Per Kilobase of transcript per Million mapped reads (FPKM). **C.** Number of testis DEGs that were classified as cell-type specific markers in a single cell RNA-seq dataset of the testis (Green et al 2018). Germline cell types are highlighted in green, somatic cell types in black. **D.** Sankey diagram of mouse genes filtered for germline enrichment based on their expression in wild-type and germline-depleted mice and in adult mouse non-gonadal tissues (Li et al 2017).




Figure 3: Kdm5c-KO epiblast-like cells express key drivers of germline identity **A.** Top - Diagram of *in vivo* differentiation of embryonic stem cells (ESCs) of the inner cell mass into epiblast stem cells. Bottom - *in vitro* differentiation of ESCs into epiblast-like cells (EpiLCs). **B.** Number of tissue-enriched differentially expressed genes (DEGs) in *Kdm5c*-KO EpiLCs. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, Fisher's exact test. **C.** Average bigwigs of an example germline gene, *Cyct*, that is dysregulated *Kdm5c*-KO EpiLCs. **D.** No significant difference in primed pluripotency marker expression in wild-type versus *Kdm5c*-KO EpiLCs. Welch's t-test, expression in transcripts per million (TPM). **E.** Representative images of wild-type (WT) and *Kdm5c*-KO cells during ESC to EpiLC differentiation. Brightfield images taken at 20X. **F.** Upset plot displaying the overlap of germline DEGs expressed in *Kdm5c*-KO EpiLCs, amygdala (AMY), and hippocampus (HIP) RNA-seq datasets. **G.** enrichPlot comparing enriched biological process gene ontologies for *Kdm5c*-KO EpiLC, amygdala, and hippocampus germline DEGs. **H.** Top left - Example germline identity DEGs unique to EpiLCs. Top right - Example 2-cell genes that are not dysregulated in *Kdm5c*-KO EpiLCs. p-values for Welch's t-test. Bottom - Average bigwigs of *Dazl* and *Stra8* expression in wild-type and *Kdm5c*-KO EpiLCs. **I.** Immunocytochemistry of DAZL in male wild-type (WT) and *Kdm5c*-KO (5CKO) EpiLCs. Percentage of DAZL-positive cells normalized to number of DAPI-positive cells, p-value for Welch's t-test.




Figure 4: Chromosomal sex influences *Kdm5c*-KO germline gene misexpression. **A-B.** Expression of tissue-enriched genes (Li et al 2017) in the male *Kdm5c*-KO amygdala (A) and hippocampus (B). Left - MA plot of mRNA-seq. Right - Number of tissue-enriched differentially expressed genes (DEGs). * $p<0.05$, ** $p<0.01$, *** $p<0.001$, Fisher's exact test. **C.** Left - Average bigwigs of an example aberrantly expressed testis-enriched DEG, *FK506 binding protein 6* (*Fkbp6*) in the wild-type (WT) and *Kdm5c*-KO (5CKO) amygdala (red) and hippocampus (teal). Right - Expression of *Cyct* in wild-type tissues from NCBI Gene, with testis highlighted in blue and brain tissues highlighted in red. **D.** Left - Average bigwigs of an example ovary-enriched DEG, *Zygotic arrest 1* (*Zar1*). Right - Expression of *Zar1* in wild-type tissues from NCBI Gene, with ovary highlighted in teal and brain tissues highlighted in red. **E.** Left - Average bigwigs of an example liver-enriched DEG, *Apolipoprotein C-I* (*Apoc1*). Right - Expression of *Apoc1* in wild-type tissues from NCBI Gene, with liver highlighted in orange and brain tissues highlighted in red.





Figure 5: KDM5C binds to a subset of germline gene promoters during early embryogenesis **A.** ChIPseeker localization of KDM5C peaks at different genomic regions in EpiLCs (top) and hippocampal and cortex primary neuron cultures (PNCs, bottom). **B.** Overlap of genes with KDM5C bound to their promoters in EpiLCs (purple) and PNCs (blue). **C.** Gene ontology (GO) comparison of genes with KDM5C bound to their promoter in EpiLCs and PNCs. Genes were classified as either bound in EpiLCs only (EpiLC only), unique to PNCs (PNC only, no significant ontologies) or bound in both PNCs and EpiLCs (shared). **D.** Average KDM5C binding around the transcription start site (TSS) of all germline-enriched genes in EpiLCs (left) and PNCs (right). **E.** KDM5C binding at the promoters of RNA-seq germline DEGs. Genes were classified as either only dysregulated in EpiLCs (EpiLC only), genes dysregulated in the hippocampus or amygdala but not EpiLCs (brain only), or genes dysregulated in both EpiLCs and the brain (common). (Legend continued on next page.)

Figure 5: KDM5C binds to a subset of germline gene promoters during early embryogenesis. (Legend continued.) **F.** Example KDM5C ChIP-seq bigwigs of DEGs unique to EpiLCs. Although both are expressed in *Kdm5c*-KO EpiLCs, KDM5C is only bound to the *Dazl* promoter and not the *Stra8* promoter in EpiLCs. **G.** Example KDM5C ChIP-seq bigwigs of RNA-seq DEGs common between the brain and EpiLCs. KDM5C is bound to the *D1Pas1* promoter but not the *XXX* promoter in EpiLCs. **H.** Example KDM5C ChIP-seq bigwigs of RNA-seq DEGs unique to the brain. KDM5C is bound to the *XXX* promoter but not the *Meig1* promoter



F

G

Figure 6: KDM5C's catalytic activity promotes long-term silencing of germline genes via DNA methylation. **A.** Left - Bigwigs of representative histone 3 lysine 4 trimethylation (H3K4me3) ChIP-seq peaks at two germline genes in the wild-type (WT) and *Kdm5c*-KO (5CKO) adult amygdala. Right - Average H3K4me3 at the transcripton start site (TSS) of all germline-enriched genes in wild-type (dark red) and *Kdm5c*-KO (light red) amygdala. **B.** Left - Bigwigs of representative histone 3 lysine 4 dimethylation (H3K4me2) ChIP-seq peaks at representative germline genes in wild-type and *Kdm5c*-KO EpiLCs. Right - Average H3K4me2 at the TSS of all germline-enriched genes in wild-type (dark purple) and *Kdm5c*-KO (light purple) EpiLCs. **C.** Diagram of embryonic stem cell (ESC) to epiblast-like cell (EpiLC) differentiation protocol and collection time points for RNA and protein. **D.** Real time quantitative PCR (RT-qPCR) of *Kdm5c* RNA expression, calculated in comparision to TBP expression ($2^{-\Delta\Delta CT}$). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, Welch's t-test. **E.** KDM5C protein expression normalized to DAXX. Quantified intensity using ImageJ (artificial units - au). Right - representative lanes of Western blot for KDM5C and DAXX. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, Welch's t-test **F.** XXX **G.** XXX

689 Notes

690 Things to do

- 691 • Move dazl to new figure if other staining works
- 692 • Use Ddx4 staining (with dazl?). If not should add RNA to diagram because its a key germline gene.
- 693 • Motif analysis
 - 694 – Discussion - talk about motifs

695 Dazl

696 We were particularly interested in the aberrant transcription of *Dazl*, since it is essential for germ cell
697 development and promotes the translation of germline mRNAs⁸¹. A significant portion of germline transcripts
698 misexpressed in *Kdm5c*-KO EpiLCs are known binding targets of DAZL, including *Stra8*⁸² ($p = 1.698e-07$,
699 Fisher's Exact Test). This suggests expression of DAZL protein could promote the translation of other
700 aberrant germline transcripts, influencing their ability to impact *Kdm5c*-KO cellular function. We thus tested
701 DAZL protein expression in *Kdm5c*-KO EpiLCs through immunocytochemistry (Figure 3I). We observed
702 about 25% of *Kdm5c*-KO EpiLCs expressed DAZL protein and it was localized to the cytoplasm ($p = 0.0015$,
703 Welch's t-test), consistent with the pattern of DAZL expression in spermatogonia⁸². Altogether these results
704 suggest tissue-specific genes are misexpressed during *Kdm5c*-KO embryogenesis, including key drivers of
705 germline identity that can be translated into protein.

- 706 • We additionally found *Kdm5c*-KO EpiLCs ectopically express DAZL protein that is localized to the
707 cytoplasm, similar to its morphology in spermatogonia⁸². **note: maybe just put in results.** Could
708 move around depending upon if I get pheno working.

709 Discussion notes

- 710 • For other paper:
 - 711 – for methods: Heatmaps of gene expression were generated using the base R functions scale and
712 hclust and visualized using the R package ComplexHeatmap (v2.12.1).
 - 713 – * Might be good to look at for retinoic acid paper (WT germ expression): <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0005654>
- 715 • Talk about different motifs if we do see differences there and if it explains direct vs indirect dysregulation
 - 716 – Could also mention PRC1.6/KDM5C binds pcgf6 in ESCs
- 717 • Maybe talk about other regulators/if anything is known about them long term? Or save for 2nd paper.

- 718 – another germline repressor: <https://pubmed.ncbi.nlm.nih.gov/35345626/>
- 719 • end with something like: this indicates kdm5c not only modulates pluripotency and self-renewal in
720 ESCs, but also has a role in long-term silencing of germline genes
- 721 – then transition into the long term silencing mechanism paragraph
- 722 • KDM5C is dynamically regulated during the window of embryonic germline gene silencing, our EpiLC
723 ChIP-seq is likely catching the tail end of KDM5C's main involvement.
- 724 • Our results indicate KDM5C is a pivotal repressor of germline programs in somatic cells.
- 725 • Studies on other chromatin regulators have focused on key marker genes to identify germline gene
726 misexpression, such as *Dazl*.
- 727 • Using a germline-depleted mouse model, we curated a list of mouse germline-enriched genes to
728 globally assess germline gene dysregulation.
- 729 • Many of the genes dysregulated in the mature *Kdm5c*-KO brain were important for late stages of
730 spermatogenesis, such as those important for sperm flagellar structure. Contrastingly, *Kdm5c*-KO
731 EpiLCs aberrantly expressed key drivers of germ cell identity, including *Dazl* and *Stra8*.
- 732 • While a significant portion of KDM5C-bound promoters are germline genes, many germline genes
733 expressed during *Kdm5c*-KO embryogenesis are not directly bound by kdm5c.
- 734 • One notable example is *Stra8*, a transcription factor important for germ cell specific transcription and
735 meiotic initiation
- 736 • The including the demarcation between soma and germline fates.
- 737 the *Kdm5c*-KO brain also Previous work identified testis-enriched genes in *Kdm5c* genes
- 738 –
- 739 – However unlike the gonadal-biased DEGs,
- 740 • The demarcation of the germline versus soma is a key feature of multicellularity and sexually dimorphic
741 reproduction
- 742 • Anything known about tissue-biased gene expression in other H3K4me regulators?
- 743 • Our data suggests the germline developmental program is occurring ectopically as *Kdm5c*-KOs pro-
744 gresses through somatic tissue development
- 745 • tissue-biased gene expression:

746 • Altogether, these data indicate that while some germline genes are misexpressed due to direct loss of
747 KDM5C binding during emryogenesis, secondary downstream mechanisms can also promote their
748 aberrant transcription.

749 • Papers to read/reference:

- 750 – Reconstitution of the Mouse Germ Cell Specification Pathway in Culture by Pluripotent Stem Cells:
751 [https://www.cell.com/fulltext/S0092-8674\(11\)00771-9](https://www.cell.com/fulltext/S0092-8674(11)00771-9)
752 – two cell gene list used by Suzuki et al Max paper is based on 2 cell sequencing: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3395470/>

754 **Figure outline:**

755 **Figure 1: Misexpression of tissue-specific genes in the *Kdm5c*-KO brain** * MA-plot and bar graphs of
756 tissue-enriched genes * Example testis-specific genes (NCBI and bigwigs) * An example ovary tissue-specific
757 gene * An example muscle/liver tissue-specific gene (NCBI and bigwigs)

758 **Figure 2: The male *Kdm5c*-KO brain expresses male and female germline-enriched genes** * Gene
759 ontology of testis DEGs in the amygdala and hippocampus - ontologies are germline ontologies * Expression
760 of testis DEGs in germline-depleted testis (this is adult testis data) * scRNASeq of testis - # of testis DEGs that
761 are germline-specific markers * Although far fewer, 5CKO brain also expresses ovary-enriched genes(NCBI
762 and bigwigs of Zar1) * These ovary enriched genes are also germline specific (NCBI/Li tissues are in adult
763 ovary, so it would be best to show they're oocyte-specific in adult ovary. But I don't think there's a published
764 adult female W/Wv dataset. Could try looking at scRNASeq or just do TPM in embryonic W/Wv data since
765 oocytes are developed at this point? Or both?) * Defining what is/isn't a germline gene, and which are
766 male/female biased using embryonic W/Wv data

767 **Figure 3: *Kdm5c*-KO epiblast-like cells express key drivers regulators of germline identity** * A) ESC
768 to EpiLC differentiation Left - Morphology is unchanged, * B) 5CKO EpiLCs express EpiLC differentiation
769 genes similar to WT lvs * C) Male EpiLCs express germline genes (example Cyct again) * Overlap between
770 brain and EpiLC germline genes - show they're mostly unique * GO of Brain and EpiLC germline genes
771 (meiotic enriched) * Bigwigs or TPM of master regulators * Show that while some are also 2-cell genes
772 (Dazl), 2-cell specific genes aren't dysregulated (Zscan4). Important point because published KDM5C dazl
773 paper is saying KDM5C is a 2-cell regulator, but as far as I can tell only genes shared between germline and
774 2-cell are dysregulated.

775 Staining of Dazl (+ Stra8 if I can get it to work)

776 **Figure 4: Loss of KDM5C's catalytic activity impairs DNAme placement and long-term silencing of
777 germline genes** * Increase in H3K4me3 in *Kdm5c*-KO amygdala at germline genes * Increase in H3K4me2
778 in EpiLCs at germline genes * *Kdm5c* binding in EpiLCs vs PNCs to show that germline repression is
779 happening in early embryo * Previous studies only looked at ESCs, unknown if catalytic activity is required

780 for long-term repression, especially since DNA methylation is placed later). KDM5C RNA and protein ESC →
781 EpiLC (increasing then decreasing) * RNA expression of germline genes with catalytic dead rescue (Ilakkia)
782 * DNA methylation in WT and 5CKO EpiLCs (Ilakkia)

783 **Figure 5: Ectopic, germline-like phenotypes in Kdm5c-KO ESCs/EpiLCs** * Sycp3 staining * DDX4
784 staining and repression of retrotransposons * Cilia??

785 Gaps in knowledge addressed: * Are other tissue-enriched genes dysregulated, or only testis, germline
786 genes? * Curating a robust list of male and female germline genes * Should talk about 2-cell genes
787 vs germline genes - way to systematically categorize? * Mechanism behind long-term germline gene
788 misexpression * Recent evidence suggests loss of KDM5C in ESCs express some germline genes * Unclear
789 if catalytic activity is required for long-term silencing * Unclear if their dysregulation lasts throughout life or
790 the same between brain or not * When in development does it begin? - Recent evidence suggests some
791 germline genes expressed in 5CKO ESCs but unclear if their dysregulation lasts throughout differentiation
792 and if the identity of germline genes are different compared to the brain * Are there functional consequences
793 to germline gene misexpression?

794 Introduction: * Chromatin regulators are important for cellular identity * H3K4me1-3 linked to active
795 gene promoters and enhancers * Surprisingly, mutations in Chromatin regulators lead to many NDDs
796 (including many H3K4 regulators) * Recent studies have shown some chromatin regulators are important
797 for regulating neuron-specific gene expression/chromatin stat_compare_means * However, loss of some
798 chromatin regulators can also lead to ectopic expression of tissue-enriched genes * Very few studies have
799 looked at these genes and it's unclear if these genes contribute to NDD impairments. * Necessary to
800 first characterize the mechanism behind their derepression to identify molecular footholds into testing their
801 contribution to neuronal impairments and potential for therapeutic intervention

- 802 • Loss of KDM5C can result in the misexpression of genes typically only found in the testis
- 803 – Misexpression of tissue-enriched genes hasn't been systematically characterized - Unclear if
804 these genes are exceptions or if other tissue-specific genes are dysregulated
- 805 – Interestingly, these genes (Cyct, D1pas1) typically function in the germline
- 806 – Germ cells (meiotic cells) are typically distinguished from somatic cells very early on in embryogen-
807 esis and is a key feature of multicellularity
- 808 – Chromatin regulators are very important for decommissioning germline genes and act successively
809 the embryo implants into the uterine wall
- 810 – Most studies have focused on ESCs, which have a similar transcriptome to germ cells / 2-cells
- 811 * recently, KDM5C was shown to repress DAZL in ESCs, independent of its catalytic activity
- 812 * However, DNA methylation is lost in the mature 5CKO brain, DNA methylation is placed later
813 and it's Unclear if it's required for long-term repression (maybe too specific, just trying to go
814 into the fact that the mechanism is partially understood but unclear)

815 – Systematic characterization of ectopic germline genes hasn't been done
816 * unknown if other germline-enriched genes are dysregulated, including oocyte-specific genes
817 * Crucially, it's unknown if misexpression of the germline program leads to functional conse-
818 quences in 5CKO cells.

819 **Germline gene repression background:**

820 Interestingly, some of the ectopic testis transcripts identified in the *Kdm5c*-KO brain are typically ex-
821 pressed in germ cells¹⁰. Unlike somatic cells, germ cells (e.g. sperm and eggs) undergo meiosis and pass
822 on their genetic material to the next generation. The germline and the soma are typically distinguished during
823 early embryogenesis, when germline genes are silenced in epiblast stem cells soon after implantation and
824 only reactivated in a subset to form the germline. Chromatin regulators play a key role in decommissioning
825 germline genes as the embryo transitions from naïve to primed pluripotency by placing repressive histone
826 H2A lysine 119 monoubiquitination (H2AK119ub1)¹⁸, histone 3 lysine 9 trimethylation (H3K9me3)^{18,19}, and
827 DNA CpG methylation^{19–21} at germline gene promoters. KDM5C may also be involved in this early decom-
828 missioning of germline genes, as re-expression of KDM5C in neurons fails to suppress their dysregulation¹⁰.
829 In support of this, KDM5C was very recently shown to repress *Deleted in azoospermia like (Dazl)*, a key
830 regulator of germline development, in mouse embryonic stem cells (ESCs)^{51,72}. In support of this, two
831 independent screens in mouse embryonic stem cells (ESCs) recently identified KDM5C as a repressor of
832 *Deleted in azoospermia like (Dazl)*, a key regulator of germline development. However, KDM5C's role in
833 embryonic germline gene repression is currently unclear, given that Dazl is also expressed in ESCs and in
834 the 2-cell stage and germline gene misexpression has yet to be globally characterized during *Kdm5c*-KO
835 embryogenesis.