

Computer simulations

The genomes of recombinant inbred lines

Karl Broman

Biostatistics & Medical Informatics, UW–Madison



`kbroman.org`

`github.com/kbroman`

`@kwbroman`


Course web: kbroman.org/AdvData






daviddeen.com


Intercross




QTL mapping




Congenic line/NIL




Advanced intercross lines




Recombinant inbred lines




Recombinant inbred lines




Collaborative Cross




MAGIC




Heterogeneous stock




Collaborative Cross




CC genome




CC genome




CC genome




CC genome



CC genome




Recombination fraction



r is the "recombination fraction"

Simulation results



Haldane & Waddington 1931

INBREEDING AND LINKAGE*

J. B. S. HALDANE AND C. H. WADDINGTON

John Innes Horticultural Institution, London, England

Received August 9, 1930

TABLE OF CONTENTS

	PAGE
Self-fertilization.....	358
Brother-sister mating. Sex-linked genes.....	360
Brother-sister mating. Autosomal genes.....	364
Parent and offspring mating. Sex-linked genes.....	367
Parent and offspring mating. Autosomal genes.....	368
Inbreeding with any initial population.....	370
Double crossing over.....	372
DISCUSSION.....	373
SUMMARY.....	374
LITERATURE CITED.....	374

Result for selfing

Then $c_n + \lambda d_n \equiv c_n + \frac{1}{4}(1 - 2x)d_n + \frac{1}{2}\lambda(1 - 2x)d_n$

$$\therefore \lambda = \frac{1 - 2x}{2 + 4x}.$$

Then since $d_\infty = 0$, and $c_1 = 0$, $d_1 = 2$,

$$c_\infty = c_\infty + \lambda d_\infty = c_1 + \lambda d_1 = \frac{1 - 2x}{1 + 2x}.$$

Put $y = D_\infty$ (the final proportion of crossover zygotes)

$$\therefore C_\infty + D_\infty = 1, C_\infty - D_\infty = c_\infty \therefore y = \frac{1}{2}(1 - c_\infty).$$

$$\therefore y = \frac{2x}{1 + 2x}.$$

(1.3)

Result for sib-mating

Omitting some rather tedious algebra, the solution of these equations is:

$$\zeta = \frac{q}{2-3q}, \quad \theta = \frac{2q}{2-3q}, \quad \kappa = \frac{1}{2-3q},$$
$$\lambda = \frac{1-2q}{2-3q}, \quad \mu = \frac{1-2q}{2-3q}, \quad \nu = \frac{2q}{2-3q}$$

as may easily be verified.

$$\begin{aligned} \therefore c_{\infty} = c_n + 2e_n + \frac{1}{1+6x} & [(1-2x)(d_n + 2f_n + 2j_n + \tfrac{1}{2}k_n) \\ & + 2g_n + 4x(h_n + i_n)] \end{aligned} \quad (3.4)$$

and $y = \frac{1}{2}(1 - c_{\infty})$.

In the case considered, $d_0 = 1$, $\therefore c_{\infty} = \zeta d_0 = 1 - 2x/1 + 6x$. Hence the proportion of crossover zygotes, $y = 4x/1 + 6x$ (3.5).

Result for sib-mating

Omitting some rather tedious algebra, the solution of these equations is:

$$\zeta = \frac{q}{2-3q}, \quad \theta = \frac{2q}{2-3q}, \quad \kappa = \frac{1}{2-3q},$$
$$\lambda = \frac{1-2q}{2-3q}, \quad \mu = \frac{1-2q}{2-3q}, \quad \nu = \frac{2q}{2-3q}$$


as may easily be verified.

$$\begin{aligned} \therefore c_{\infty} = c_n + 2e_n + \frac{1}{1+6x} & [(1-2x)(d_n + 2f_n + 2j_n + \tfrac{1}{2}k_n) \\ & + 2g_n + 4x(h_n + i_n)] \end{aligned} \quad (3.4)$$

and $y = \frac{1}{2}(1 - c_{\infty})$.

In the case considered, $d_0 = 1$, $\therefore c_{\infty} = \zeta d_0 = 1 - 2x/1 + 6x$. Hence the proportion of crossover zygotes, $y = 4x/1 + 6x$ (3.5).

Simulation results



Non-linear regression

```
out <- nls(R ~ a*r/(1 + b*r),  
           data = data.frame(r=r, R=R),  
           start = list(a=4, b=6))  
summary(out)
```

Non-linear regression

```
out <- nls(R ~ a*r/(1 + b*r),  
           data = data.frame(r=r, R=R),  
           start = list(a=4, b=6))  
summary(out)
```


	Estimate	Std. Error
a	7.016	0.011
b	6.023	0.016

Non-linear regression

```
out <- nls(R ~ a*r/(1 + b*r),  
           data = data.frame(r=r, R=R),  
           start = list(a=4, b=6))  
summary(out)
```

				More data	
	Estimate	Std. Error		Estimate	Std. Error
a	7.016	0.011	a	7.003	0.008
b	6.023	0.016	b	6.005	0.012

Simulation results



Markov chain

- ▶ Sequence of random variables $\{X_0, X_1, X_2, \dots\}$ satisfying

$$\Pr(X_{n+1} \mid X_0, X_1, \dots, X_n) = \Pr(X_{n+1} \mid X_n)$$

- ▶ Transition probabilities $P_{ij} = \Pr(X_{n+1} = j \mid X_n = i)$

- ▶ Here, X_n = “parental type” at generation n .

- ▶ We are interested in **absorption probabilities**

$$\pi_j = \Pr(X_n \rightarrow j \mid X_0)$$

Absorption probabilities

Consider the case of **absorption** into the state $\begin{array}{c|c} A & A \\ A & A \end{array}$
(write this $AA|AA$)

Let h_i = probability, starting at i , of being absorbed into $AA|AA$.

Then $h_{AA|AA} = 1$ and $h_{AB|AB} = 0$.

Condition on the first step: $h_i = \sum_k P_{ik} h_k$

For selfing, this gives a system of 3 linear equations.

Equations for selfing

C₂ *AABB* and *aabb*.

 D_n $AAbb$ and $aaBB$.

E, $AA Bb$, $Aa BB$, $Aabb$, and $aa Bb$.

$$F_5 AB.ab.$$
$$G_n \text{ } Ab.aB.$$

We assume $2C_n + 2D_n + 4E_n + F_n + G_n = 2$, so that $C_1 = D_1 = E_1 = G_1 = 0$, and $F_1 = 2$. Clearly $E_\infty = F_\infty = G_\infty = 0$, and D_∞ is the final proportion of crossover zygotes. Then considering the results of selfing each generation, we have:

$$\left. \begin{aligned} C_{n+1} &= C_n + \frac{1}{2}E_n + \frac{1}{4}(1 - \beta - \delta + \beta\delta)F_n + \frac{1}{4}\beta\delta G_n \\ D_{n+1} &= D_n + \frac{1}{2}E_n + \frac{1}{4}\beta\delta F_n + \frac{1}{4}(1 - \beta - \delta + \beta\delta)G_n \\ E_{n+1} &= \frac{1}{2}E_n + \frac{1}{4}(\beta + \delta - 2\beta\delta)(F_n + G_n) \\ F_{n+1} &= \frac{1}{2}(1 - \beta - \delta + \beta\delta)F_n + \frac{1}{2}\beta\delta G_n \\ G_{n+1} &= \frac{1}{2}\beta\delta F_n + \frac{1}{2}(1 - \beta - \delta + \beta\delta)G_n \end{aligned} \right\} \quad (1.1)$$

Put $y = D_{\infty}$ (the final proportion of crossover zygotes)

$$\therefore C_{\infty} + D_{\infty} = 1, C_{\infty} - D_{\infty} = c_{\infty} \therefore y = \frac{1}{2}(1 - c_{\infty}).$$

➡ $\therefore y = \frac{2x}{1+2x} \quad (1.3)$

for C_{n+1} , D_{n+1} , and F_{n+1} , G_{n+1} ,

$$\left. \begin{array}{l} d_n \end{array} \right\} \quad (1.2)$$

for all values of n .

$$-2x) d_n$$

$$= \frac{1 - 2x}{1 + 2x}.$$

Equations for sib-mating

Typical mating	Number of types	
$AABB \times AABB$	2	$C_{n+1} = C_n + H + \frac{1}{4}(\alpha^2 + \gamma^2)L + \frac{1}{4}(\beta^2 + \delta^2)N + \frac{1}{4}Q + \frac{1}{4}R + \frac{1}{4}(\alpha^2 + \gamma^2)U + \frac{1}{4}(\beta^2 + \delta^2)V + \frac{1}{4}\alpha^2\gamma^2W + \frac{1}{4}\beta^2\delta^2X + \frac{1}{4}\alpha^2\beta^2\gamma^2Y$
$AAbb \times AAbb$	2	$D_{n+1} = D + I + \frac{1}{4}(\alpha^2 + \gamma^2)M + \frac{1}{4}(\beta^2 + \delta^2)P + \frac{1}{4}Q + \frac{1}{4}S + \frac{1}{4}(\beta^2 + \delta^2)U + \frac{1}{4}(\alpha^2 + \gamma^2)V + \frac{1}{4}\alpha^2\beta^2\gamma^2W + \frac{1}{4}\beta^2\delta^2X + \frac{1}{4}\alpha^2\beta^2\gamma^2Y$
$AABB \times aabb$	2	$E_{n+1} = \frac{1}{4}\alpha^2\gamma^2W + \frac{1}{4}\beta^2\delta^2X + \frac{1}{4}\alpha^2\beta^2\gamma^2Y$
$AAbb \times aaBB$	2	$F_{n+1} = \frac{1}{4}\beta^2\delta^2W + \frac{1}{4}\alpha^2\delta^2 + \beta^2\gamma^2X + \frac{1}{4}\alpha^2\gamma^2Y$
$AABB \times AAbb$	8	$G_{n+1} = \frac{1}{4}(\alpha\beta + \gamma\delta)(U + V) + \frac{1}{4}\alpha\beta\gamma\delta(W + 2X + Y)$
$AAbb \times AABB$	8	$H_{n+1} = \frac{1}{4}H + \frac{1}{4}(\alpha\delta + \beta\gamma)(U + V) + \frac{1}{4}\alpha\beta\gamma\delta(W + 2X + Y)$
$AAbb \times AABb$	8	$I_{n+1} = \frac{1}{2}I + \frac{1}{4}(\alpha\delta + \beta\gamma)(U + V) + \frac{1}{4}\alpha\beta\gamma\delta(W + 2X + Y)$
$AABB \times Aabb$	8	$J_{n+1} = \frac{1}{4}(\alpha\delta + \beta\gamma)(U + V) + \frac{1}{4}\alpha\beta\gamma\delta(W + 2X + Y)$
$AAbb \times AaBB$	8	$K_{n+1} = \frac{1}{4}(\alpha\delta + \beta\gamma)(U + V) + \frac{1}{4}\alpha\beta\gamma\delta(W + 2X + Y)$
$AABB \times AB.ab$	4	$L_{n+1} = \frac{1}{4}(\alpha^2\gamma^2W + \beta^2\delta^2X)$
$AAbb \times Ab.aB$	4	$M_{n+1} = \frac{1}{4}(\alpha^2\gamma^2W + \beta^2\delta^2X)$
$AABB \times Ab.aB$	4	$N_{n+1} = \frac{1}{4}R + \frac{1}{4}(\alpha\beta + \gamma\delta)(U + V) + \frac{1}{4}\alpha\beta\gamma\delta(W + 2X + Y)$
$AAbb \times AB.ab$	4	$P_{n+1} = \frac{1}{4}S + \frac{1}{4}(\alpha\beta + \gamma\delta)(U + V) + \frac{1}{4}\alpha\beta\gamma\delta(W + 2X + Y)$
$AABb \times AABb$	4	$Q_{n+1} = 2G + \frac{1}{4}(H + I + J + K) + \frac{1}{4}(\alpha^2 + \gamma^2)(L + M) + \frac{1}{4}(\beta^2 + \delta^2)(N + P) + \frac{1}{4}Q + \frac{1}{4}(R + S + T) + \frac{1}{4}(\alpha^2 + \alpha\beta + \beta^2 + \gamma^2 + \gamma\delta + \delta^2)(U + V) + \frac{1}{4}(\alpha\delta + \beta\gamma)^2(W + Y) + \frac{1}{4}(\alpha\gamma + \beta\delta)^2X$
$AABb \times AaBB$	4	$R_{n+1} = \frac{1}{4}(\beta^2 + \delta^2)L + \frac{1}{4}(\alpha^2 + \gamma^2)N + \frac{1}{4}R + \frac{1}{4}(\beta + \delta)U + \frac{1}{4}(\alpha + \gamma)V + \frac{1}{4}(\alpha\delta + \beta\gamma)^2(W + Y) + \frac{1}{4}(\alpha\gamma + \beta\delta)^2X$
$AABb \times Aabb$	4	$S_{n+1} = \frac{1}{4}(\beta^2 + \delta^2)M + \frac{1}{4}(\alpha^2 + \gamma^2)P + \frac{1}{4}S + \frac{1}{4}(\alpha + \gamma)U + \frac{1}{4}(\beta + \delta)V + \frac{1}{4}(\alpha\delta + \beta\gamma)^2(W + Y) + \frac{1}{4}(\alpha\gamma + \beta\delta)^2X$
$AABb \times aaBb$	4	$T_{n+1} = \frac{1}{4}(\alpha\beta + \gamma\delta)(U + V) + \frac{1}{4}(\alpha\delta + \beta\gamma)^2(W + Y) + \frac{1}{4}(\alpha\gamma + \beta\delta)^2X$
$AABb \times AB.ab$	8	$U_{n+1} = \frac{1}{2}J + \frac{1}{4}(\alpha\beta + \gamma\delta)(L + N) + \frac{1}{4}(S + T) + \frac{1}{4}(\alpha + \gamma)U + \frac{1}{4}(\beta + \delta)V + \frac{1}{4}\alpha\gamma(\beta\gamma + \alpha\delta)W + \frac{1}{4}(\alpha\gamma + \beta\delta)(\alpha\delta + \beta\gamma)X + \frac{1}{4}\beta\delta(\beta\gamma + \alpha\delta)Y$
$AABb \times Ab.aB$	8	$V_{n+1} = \frac{1}{2}K + \frac{1}{4}(\alpha\beta + \gamma\delta)(M + P) + \frac{1}{4}(R + T) + \frac{1}{4}(\beta + \delta)U + \frac{1}{4}(\alpha + \gamma)V + \frac{1}{4}\beta\delta(\beta\gamma + \alpha\delta)W + \frac{1}{4}(\alpha\gamma + \beta\delta)(\alpha\delta + \beta\gamma)X + \frac{1}{4}\alpha\gamma(\beta\gamma + \alpha\delta)Y$
$AB.ab \times AB.ab$	1	$W_{n+1} = 2(E + J) + \frac{1}{4}(\alpha^2 + \gamma^2)L + \frac{1}{4}(\beta^2 + \delta^2)N + \frac{1}{4}(S + T) + \frac{1}{4}(\alpha^2 + \gamma^2)U + \frac{1}{4}(\beta^2 + \delta^2)V + \frac{1}{4}\alpha^2\gamma^2W + \frac{1}{4}(\alpha^2\beta^2 + \beta^2\gamma^2)X + \frac{1}{4}\beta^2\delta^2Y$
$AB.ab \times Ab.aB$	2	$X_{n+1} = \frac{1}{4}T + \frac{1}{4}(\alpha\beta + \gamma\delta)(U + V) + \frac{1}{4}\alpha\beta\gamma\delta(W + 2X + Y)$
$Ab.aB \times Ab.aB$	1	$Y_{n+1} = 2(F + K) + \frac{1}{4}(\alpha^2 + \gamma^2)M + \frac{1}{4}(\beta^2 + \delta^2)P + \frac{1}{4}(R + T) + \frac{1}{4}(\beta^2 + \delta^2)U + \frac{1}{4}(\alpha^2 + \gamma^2)V + \frac{1}{4}\beta^2\delta^2W + \frac{1}{4}(\alpha^2\delta^2 + \delta^2\gamma^2)X + \frac{1}{4}\alpha^2\gamma^2Y$

Result for sib-mating

Omitting some rather tedious algebra, the solution of these equations is:

$$\zeta = \frac{q}{2-3q}, \quad \theta = \frac{2q}{2-3q}, \quad \kappa = \frac{1}{2-3q},$$
$$\lambda = \frac{1-2q}{2-3q}, \quad \mu = \frac{1-2q}{2-3q}, \quad \nu = \frac{2q}{2-3q}$$

as may easily be verified.

$$\begin{aligned} \therefore c_{\infty} = c_n + 2e_n + \frac{1}{1+6x} & [(1-2x)(d_n + 2f_n + 2j_n + \tfrac{1}{2}k_n) \\ & + 2g_n + 4x(h_n + i_n)] \end{aligned} \quad (3.4)$$

and $y = \frac{1}{2}(1 - c_{\infty})$.


In the case considered, $d_0 = 1$, $\therefore c_{\infty} = \zeta d_0 = 1 - 2x/1 + 6x$. Hence the proportion of crossover zygotes, $y = 4x/1 + 6x$ (3.5).

3-point coincidence




- ▶ r_{ij} = recombination fraction for interval (i, j)
Assume $r_{12} = r_{23} = r$.
- ▶ **Coincidence** = $c = \Pr(\text{double recombinant})/r^2$
 $= \Pr(\text{rec'n in 23} \mid \text{rec'n in 12})/\Pr(\text{rec'n in 23})$
- ▶ No interference = 1
Positive interference < 1
Negative interference > 1
- ▶ Generally c is a function of r

Coincidence




Coincidence



Coincidence in 8-way RILs

- ▶ The trick that allowed us to get the coincidence for 2-way RILs doesn't work for 8-way RILs.
- ▶ It's sufficient to consider 4-way RILs.
- ▶ Calculations for 3 points in 4-way RILs is still **astoundingly complex**.
 - 2 points in 2-way RILs by sib mating:
55 parental types → **22 states** by symmetry
 - 3 points in 4-way RILs by sib mating:
2,164,240 parental types → **137,488 states** by symmetry
- ▶ Even **counting** the states was difficult.

Coincidence




The formula

$$C = \frac{(1 + 6r)[280 + 1208r - 848r^2 + 5c(7 - 28r - 368r^2 + 344r^3) - 2c^2(49 - 324r + 452r^2)r^2 - 16c^3(1 - 2r)r^4]}{49(1 + 12r - 12cr^2)[5 + 10r - 4(2 + c)r^2 + 8cr^3]}$$


3-point symmetry

$$\Pr(M_2 = x \mid M_1 = A, M_2 \neq A, M_3 = A)$$




Markov property

$$\log_2 \left\{ \frac{\Pr(M_3=A \mid M_2=A, M_1=x)}{\Pr(M_3=A \mid M_2=A)} \right\}$$




Markov property

$$\log_2 \left\{ \frac{\Pr(M_3=A \mid M_2=B, M_1=x)}{\Pr(M_3=A \mid M_2=B)} \right\}$$




Markov property

$$\log_2 \left\{ \frac{\Pr(M_3=A \mid M_2=\text{C}, M_1=x)}{\Pr(M_3=A \mid M_2=\text{C})} \right\}$$



Markov property


$$\log_2 \left\{ \frac{\Pr(M_3=A \mid M_2=E, M_1=x)}{\Pr(M_3=A \mid M_2=E)} \right\}$$




Whole genome simulations

- ▶ 2-way selfing, 2-way sib-mating, 8-way sib-mating
- ▶ Mouse-like genome, 1665 cM
- ▶ Strong positive crossover interference
- ▶ Inbreed to complex fixation
- ▶ 10,000 simulation replicates


No. generations to fixation




No. generations to 99% fixation




Percent genome not fixed




No. breakpoints




Segment lengths




Segment lengths




Probability a segment is inherited intact




Length of smallest segment



No. segments < 1 cM



Collaborative Cross



The PreCC

What happens at G_2F_k ?

$\Pr(g_1 = i)$ as a function of k

$\Pr(g_1 = i, g_2 = j)$ as a function of k and the recombination fraction

Crazy table

Table 4 Two-locus haplotype probabilities at generation F_k in the formation of four-way RIL by sibling mating

Chr.	Individual	Prototype	No. states	Probability of each
A	Random	AA	4	$\frac{1}{4(1+6r)} - \left[\frac{6r^2-7r-3rs}{4(1+6r)s} \right] \left(\frac{1-2r+s}{4} \right)^k + \left[\frac{6r^2-7r+3rs}{4(1+6r)s} \right] \left(\frac{1-2r-s}{4} \right)^k$
		AB	4	$\frac{r}{2(1+6r)} + \left[\frac{10r^2-r-rs}{4(1+6r)s} \right] \left(\frac{1-2r+s}{4} \right)^k - \left[\frac{10r^2-r+rs}{4(1+6r)s} \right] \left(\frac{1-2r-s}{4} \right)^k$
		AC	8	$\frac{r}{2(1+6r)} - \left[\frac{2r^2+3r+rs}{4(1+6r)s} \right] \left(\frac{1-2r+s}{4} \right)^k + \left[\frac{2r^2+3r-rs}{4(1+6r)s} \right] \left(\frac{1-2r-s}{4} \right)^k$
X	Female	AA	2	$\frac{1}{3(1+4r)} + \frac{1}{6(1+r)} \left(-\frac{1}{2} \right)^k - \left[\frac{4r^3-(4r^2+3r)t+3r^2-5r}{4(4r^2+5r+1)t} \right] \left(\frac{1-r+t}{4} \right)^k + \left[\frac{4r^3+(4r^2+3r)t+3r^2-5r}{4(4r^2+5r+1)t} \right] \left(\frac{1-r-t}{4} \right)^k$
		AB	2	$\frac{2r}{3(1+4r)} + \frac{r}{3(1+r)} \left(-\frac{1}{2} \right)^k + \left[\frac{2r^3+6r^2-(2r^2+r)t}{2(4r^2+5r+1)t} \right] \left(\frac{1-r+t}{4} \right)^k - \left[\frac{2r^3+6r^2+(2r^2+r)t}{2(4r^2+5r+1)t} \right] \left(\frac{1-r-t}{4} \right)^k$
		AC	4	$\frac{2r}{3(1+4r)} - \frac{r}{6(1+r)} \left(-\frac{1}{2} \right)^k - \left[\frac{9r^2+5r+rt}{4(4r^2+5r+1)t} \right] \left(\frac{1-r+t}{4} \right)^k + \left[\frac{9r^2+5r-rt}{4(4r^2+5r+1)t} \right] \left(\frac{1-r-t}{4} \right)^k$
		CC	1	$\frac{1}{3(1+4r)} - \frac{1}{3(1+r)} \left(\frac{1}{2} \right)^k + \left[\frac{9r^2+5r+rt}{2(4r^2+5r+1)t} \right] \left(\frac{1-r+t}{4} \right)^k - \left[\frac{9r^2+5r-rt}{2(4r^2+5r+1)t} \right] \left(\frac{1-r-t}{4} \right)^k$
X	Male	AA	2	$\frac{1}{3(1+4r)} - \frac{1}{3(1+r)} \left(-\frac{1}{2} \right)^k + \left[\frac{r^3-(8r^3+r^2-3r)t-10r^2+5r}{2(4r^4-35r^3-29r^2+15r+5)} \right] \left(\frac{1-r+t}{4} \right)^k + \left[\frac{r^3+(8r^3+r^2-3r)t-10r^2+5r}{2(4r^4-35r^3-29r^2+15r+5)} \right] \left(\frac{1-r-t}{4} \right)^k$
		AB	2	$\frac{2r}{3(1+4r)} - \frac{2r}{3(1+r)} \left(-\frac{1}{2} \right)^k + \left[\frac{r^4+(5r^3-r)t-10r^3+5r^2}{4r^4-35r^3-29r^2+15r+5} \right] \left(\frac{1-r+t}{4} \right)^k + \left[\frac{r^4-(5r^3-r)t-10r^3+5r^2}{4r^4-35r^3-29r^2+15r+5} \right] \left(\frac{1-r-t}{4} \right)^k$
		AC	4	$\frac{2r}{3(1+4r)} + \frac{r}{3(1+r)} \left(-\frac{1}{2} \right)^k - \left[\frac{2r^4+(2r^3-r^2+r)t-19r^3+5r}{2(4r^4-35r^3-29r^2+15r+5)} \right] \left(\frac{1-r+t}{4} \right)^k - \left[\frac{2r^4-(2r^3-r^2+r)t-19r^3+5r}{2(4r^4-35r^3-29r^2+15r+5)} \right] \left(\frac{1-r-t}{4} \right)^k$
		CC	1	$\frac{1}{3(1+4r)} + \frac{2}{3(1+r)} \left(-\frac{1}{2} \right)^k + \left[\frac{2r^4+(2r^3-r^2+r)t-19r^3+5r}{4r^4-35r^3-29r^2+15r+5} \right] \left(\frac{1-r+t}{4} \right)^k + \left[\frac{2r^4-(2r^3-r^2+r)t-19r^3+5r}{4r^4-35r^3-29r^2+15r+5} \right] \left(\frac{1-r-t}{4} \right)^k$

$s = \sqrt{4r^2-12r+5}$ and $t = \sqrt{r^2-10r+5}$; the autosomal haplotype probabilities are valid for $r < \frac{1}{2}$.

Lesson

Computer simulations are hugely valuable.

Uses of simulations

- ▶ Study probabilities
- ▶ Estimate power/sample size
- ▶ Evaluate performance of a method
- ▶ Evaluate sensitivity/robustness of a method

Relative advantages?

- ▶ Simulations
- ▶ Numerical calculations
- ▶ Analytic calculations

References

- ▶ Haldane & Waddington (1931) Inbreeding and Linkage. 16:357–374
- ▶ Broman KW (2005) The genomes of recombinant inbred lines. Genetics 169:1133–1146
- ▶ Teuscher & Broman (2007) Haplotype probabilities for multiple-strain recombinant inbred lines. Genetics 175:1267–1274
- ▶ Broman KW (2012) Genotype probabilities at intermediate generations in the construction of recombinant inbred lines. Genetics 190:403–412
- ▶ Broman KW (2012) Haplotype probabilities in advanced intercross populations. G3 2:199–202