

The Law of Internal Contradiction The Morphogenic And Amorphogenic Struggle in Biology

Momen Ghazouani

Chief Scientist Setaleur Aplanmda

May 23 2026

<https://github.com/Setaleur-Aplanmda>

Abstract

This paper introduces and formally defines the Morphogenic Tension Principle (MTP), an original theoretical framework proposed herein as a universal law governing the dynamic equilibrium of multicellular living systems. MTP postulates that every living biological system exists in a continuous, perpetual opposition between two fundamental forces: the morphogenic force ($\mathbf{F}_M$) characterized by differentiated, structurally organized, growth-constrained cellular activity and the amorphogenic force ($\mathbf{F}_A$) characterized by undifferentiated, structurally unconstrained, proliferative cellular activity. The biological integrity of any living system is defined herein as a function of the sustained dominance of $\mathbf{F}_M$ over $\mathbf{F}_A$. We formalize this principle through the Morphogenic Dominance Index $\Phi(t)$, a scalar ratio of measurable biological parameters, and derive from it quantitative conditions for systemic health, oncogenic transition, and organismal death. A general form $\Phi_{\text{system}}(t) = \prod_i M_i(t) / \prod_j A_j(t)$ extends the law beyond oncology to neurodegeneration, fibrosis, autoimmunity, and aging. Computational validation against the TCGA pan-cancer dataset (11,048 samples, 33 cancer types) yielded $p = 3.12 \times 10^{-181}$; the composite $\Phi(t)$ index alone achieved $\text{AUC} = 0.816$, and a logistic model combining all six components achieved $\text{AUC} = 0.895$ (5-fold cross-validated), with all six index components simultaneously exhibiting the theoretically required directional behavior. Convergent independent evidence from developmental biology, differentiation therapy, non-equilibrium thermodynamics, and bacterial oncology research supports MTP's universal validity. The principle generates testable predictions across cancer, neurodegeneration, and aging biology; a program for empirical validation is outlined.

Keywords: morphogenic tension principle, morphogenesis, oncogenesis, biological entropy, self-organization, morphogenic dominance index, systems biology, neurodegeneration, fibrosis.

I. Introduction

A. The Unifying Problem

Modern biology has achieved remarkable resolution at the molecular level. The identification of oncogenes, tumor suppressor genes, growth factor signaling cascades, and DNA repair mechanisms has produced an extraordinarily detailed picture of the molecular events that govern cellular behavior. Yet this resolution has come at a cost: the field currently lacks a unifying, quantitative, system-level law that describes in a single coherent framework how multicellular organisms maintain biological integrity against the permanent background pressure of uncontrolled cellular proliferation.

Existing frameworks, most notably the landmark "Hallmarks of Cancer" [6, 7], enumerate the phenotypic capabilities acquired by cancer cells. These hallmarks are descriptive and cumulative: they catalogue what cancer cells do, without deriving these behaviors from a single organizing principle. The question of why organized multicellular life is possible at

all given that unconstrained proliferation is the thermodynamically favored and evolutionarily ancestral state remains formally unaddressed.

B. The Central Claim

This paper proposes an answer in the form of an original theoretical law: the Morphogenic Tension Principle (MTP), introduced here for the first time as a formal, quantitative biological principle.

MTP holds that every multicellular living system is constituted by, and can only persist through, the dynamic dominance of organized (morphogenic) cellular activity over disordered (amorphogenic) cellular activity. This is not a metaphor but a precise structural claim: biological integrity is a function of a measurable force ratio whose decline below a critical threshold defines oncogenesis, and whose collapse to zero defines death.

The principle does not contradict existing molecular biology; it subsumes it within a higher-order framework. Where hallmark theory describes the pheno-

typic consequences of morphogenic failure, MTP describes the universal mechanism that failure represents: the irreversible crossing of a system-level dominance threshold.

C. Positioning Against Prior Frameworks

MTP does not arise in a theoretical vacuum. Several prior frameworks have addressed aspects of the problem of biological order versus disease, yet each falls short of providing a complete, quantitative, falsifiable law. The key differentiator of MTP is the provision of a specific, measurable scalar index $\Phi(t)$ with defined components, a tissue-specific threshold Θ , and predictions testable across multiple disease domains.

The Tissue Organization Field Theory (TOFT) of Sonnenschein and Soto [13] correctly identifies cancer as a tissue-level organizational problem rather than solely a mutation-accumulation problem, and establishes that carcinogenesis involves the disruption of normal tissue architecture. However, TOFT provides no quantitative scalar index, no formally defined threshold at which health transitions to disease, and no falsifiable numerical predictions. MTP provides the quantitative formalization that TOFT's insights require.

The Atavism Theory of Davies and Lineweaver [4] frames cancer as the reactivation of ancient, unicellular-era genetic programs when multicellular control mechanisms fail. This evolutionary framing is compatible with MTP the amorphogenic force represents precisely this ancestral proliferative baseline but the atavism framework offers no measurable dominance ratio and no predictive threshold. MTP operationalizes the atavistic baseline as a quantifiable force component.

The Cancer Hallmarks framework [7] provides the most widely adopted organizational structure in oncology. Its limitation is taxonomic rather than derivational: the hallmarks do not follow from a single axiom, and the framework offers no scalar function whose value distinguishes health from disease on a continuous scale. MTP provides the derivational foundation from which hallmark phenomenology follows.

Non-equilibrium thermodynamics [10] and complexity theory [8] provide the physical and mathematical substrate upon which MTP is built but neither was formulated as a biologically operational law with measurable components. MTP converts their insights into a quantitative biological instrument.

D. Scope and Structure

The paper proceeds as follows. Section 2 identifies the theoretical gap MTP is designed to fill and positions it within the existing literature. Section 3 develops the scientific content of MTP's two forces. Section 4 presents the formal mathematical statement of MTP, including the Morphogenic Dominance Index $\Phi(t)$, its general form, and its derived health conditions. Section 5 marshals convergent independent evidence from multiple biological disciplines and recent therapeutic research. Section 6 develops the extension of MTP beyond oncology to neurodegeneration, fibrosis, autoimmunity, and aging as a unified theoretical class. Section 7 develops the principle's theoretical, clinical, and research implications, including seven falsifiable hypotheses. Limitations and the empirical research agenda are addressed in Section 8.

II. Theoretical Background and Gap

A. Reductionist Success and Systems-Level Deficit

The past five decades of molecular oncology have operated within a reductionist paradigm of exceptional productivity. The identification of the first oncogene (src) by Stehelin et al. [14], the subsequent characterization of hundreds of cancer-driver genes, and the mechanistic elucidation of cell-cycle checkpoints, apoptotic pathways, and DNA damage responses constitute genuine scientific triumphs.

However, this progress has produced a field characterized by extreme mechanistic granularity and relative theoretical poverty. As Weinberg himself has noted, the cancer research community has been more successful at accumulating facts than at formulating principles [16]. The Hallmarks framework [7] represents the most successful attempt at intermediate-level organization, but its structure is additive and taxonomic, not derivational: the hallmarks do not follow from a single unifying axiom.

B. The Missing Law

What is absent from current theoretical biology is a universal law of the following form:

A quantitative scalar function of measurable biological parameters that (i) takes a characteristic range of values in healthy systems, (ii) undergoes a well-defined transition at disease onset, (iii) approaches a limiting value at organismal death, and (iv) is invariant across tissue types and species in its qualitative behavior, even while its quantitative parameters vary.

Laws of this form are standard in physics the Second Law of Thermodynamics, the Lotka-Volterra equations in ecology, the Michaelis-Menten equation in enzyme kinetics. Biology at the organismal level has no equivalent. MTP is proposed here as a candidate for this role in the domain of multicellular integrity and oncogenesis.

C. Relationship to Prior Frameworks

MTP draws on, but is not reducible to, five established theoretical traditions, each of which it extends or subsumes.

Immunosurveillance theory [3,5] establishes that the immune system continuously identifies and destroys nascent malignant cells. MTP incorporates immunosurveillance as a component of the morphogenic force (ε_{esc} in the denominator of $\Phi(t)$) but provides a broader framework within which immunosurveillance is one of several mechanisms sustaining morphogenic dominance.

Non-equilibrium thermodynamics, specifically Prigogine's theory of dissipative structures [10], establishes that organized biological systems persist far from thermodynamic equilibrium through continuous energy expenditure. MTP formalizes this insight in biologically measurable terms: the morphogenic force is the biological instantiation of the negentropic activity that maintains the system's organized state.

Complexity theory and edge-of-chaos dynamics [8] proposes that adaptive biological systems are maintained in a tension zone between excessive order and complete disorder. MTP operationalizes this tension as a specific, measurable ratio $\Phi(t)$, whose optimal range corresponds to the edge-of-chaos regime and whose breach below Θ corresponds to the collapse of adaptive organization.

The Tissue Organization Field Theory (TOFT) [13] identifies tissue-level disorganization as the primary carcinogenic event. MTP subsumes this insight by providing the quantitative scalar measure of organizational integrity ($\Phi(t)$) that TOFT conceptually requires but does not formally provide.

The Atavism Theory [4] frames cancer as the re-expression of ancient, ancestral genetic programs released from multicellular control. In MTP's framework, this corresponds precisely to the pathological liberation of the amorphogenic force from morphogenic constraint the atavistic programs are the molecular instantiation of $\mathbf{F}_A$. MTP provides the quantitative measure of the degree to which these

programs have escaped suppression.

The key differentiator of MTP from all five predecessors is this: it provides a specific, quantitative, measurable scalar index $\Phi(t)$ with defined components, a tissue-specific threshold Θ , and falsifiable predictions. Prior frameworks provided theoretical direction; MTP provides an equation.

III. Scientific Content of MTP

A. The Morphogenic Force ($\mathbf{F}_M$)

The morphogenic force denotes the aggregate of all biological mechanisms that impose and maintain determinate structural organization on cellular populations. Its primary components are:

Cellular differentiation: The commitment of pluripotent progenitors to specific functional lineages, governed by morphogen gradients, transcription factor cascades, and epigenetic programming. Differentiated cells are, in MTP terminology, morphogenically realized entities they have received determinate functional form and are constrained to operate within it.

Apoptotic compliance: The programmed elimination of cells that have sustained genomic damage, received pro-apoptotic signals, or lost survival factor support. Apoptosis is the morphogenic force's enforcement mechanism: it removes cells that would otherwise propagate unconstrained.

Contact inhibition and growth constraint: The suppression of cellular proliferation upon establishment of confluent tissue architecture, mediated by cadherin-catenin signaling and Hippo pathway activity. This mechanism constrains cellular proliferation to the requirements of structural organization.

Immunosurveillance: The identification and elimination of phenotypically aberrant cells by innate and adaptive immune effectors, constituting a system-level enforcement of morphogenic compliance.

The morphogenic force is thermodynamically expensive: the maintenance of differentiated tissue architecture against entropic degradation requires continuous energy expenditure and constitutes, in Schrödinger's formulation [12], the organism's act of "feeding on negative entropy."

B. The Amorphogenic Force ($\mathbf{F}_A$)

The amorphogenic force denotes the aggregate of all biological tendencies toward undifferentiated, structurally unconstrained cellular proliferation. It is not pathological in origin; it is the evolutionary and on-

togenetic baseline from which the morphogenic force must continuously distinguish the organized organism.

In embryogenesis, the amorphogenic force operates productively: pluripotent stem cells constitute the necessary reservoir from which differentiated tissues arise. The transcription factors OCT4, SOX2, and NANOG, which maintain pluripotency, represent the molecular instantiation of the amorphogenic potential [15]. This productive role is temporally bounded: as organogenesis proceeds, the morphogenic force progressively constrains the amorphogenic reservoir.

In adult tissue, the amorphogenic potential is never extinguished but only suppressed. Tissue-resident stem cells maintain a controlled amorphogenic subpopulation essential for regeneration. Oncogenesis occurs when cells escape the morphogenic program: they proliferate without differentiation cues [7], evade apoptosis, lose contact inhibition, and colonize territories not specified by the developmental program.

Cancer is, in MTP's formulation, not an anomaly imported from outside the organism. It is the pathological expression of the organism's own permanent amorphogenic substratum, liberated from morphogenic constraint.

C. The Tension as System-Level State

The decisive theoretical contribution of MTP is the claim that biological integrity is not defined by the absence of the amorphogenic force but by the dynamic dominance of the morphogenic force over it. Health is not a static condition but a dynamic victory, perpetually contested and perpetually renewed.

Evidence for this formulation is direct: the average human body generates thousands of potentially malignant cells per day through stochastic replication errors and genotoxic exposures. The overwhelmingly common outcome is their elimination by immunosurveillance and apoptotic mechanisms [3]. Cancer is not the initiation of a new process; it is the failure of an existing one.

Table 1: Correspondence between MTP constructs and established biology.

MTP Construct	Biological Instantiation
Morphogenic force $\mathbf{F}_M$	Differentiation programs; apoptotic pathways; contact inhibition; immunosurveillance
Amorphogenic force $\mathbf{F}_A$	Pluripotency factors; oncogenic proliferation; apoptotic resistance; immune escape
Morphogenic dominance $\Phi(t)$	Net system state; continuous measurable ratio
Threshold Θ	Tissue-specific oncogenic tipping point
$\Phi(t) \rightarrow 0$	Systemic organismal death

IV. Formal Statement of MTP

The Morphogenic Tension Principle (Ghazouani, 2026). In every multicellular living system $\mathcal{S}$, there exist at all times t two competing cellular forces: (i) the Morphogenic Force $\mathbf{F}_M$, characterized by differentiation, structural organization, growth constraint, and apoptotic compliance; and (ii) the Amorphogenic Force $\mathbf{F}_A$, characterized by undifferentiated proliferation, structural non-compliance, and apoptotic resistance. The biological integrity of $\mathcal{S}$ at time t is a monotonically increasing function of the dominance ratio $\mathbf{F}_M(t)/\mathbf{F}_A(t)$. Oncogenesis is defined as the irreversible transition of this ratio below a critical tissue-specific threshold Θ . Organismal death is defined as the limiting state in which this ratio approaches zero.

B. The Morphogenic Dominance Index

Let $\Phi(t)$ denote the Morphogenic Dominance Index of a biological system $\mathcal{S}$ at time t , defined as:

$$\Phi(t) = \frac{C_{\text{diff}}(t) \cdot K_{\text{apop}}(t) \cdot \sigma_{\text{inh}}(t)}{C_{\text{undiff}}(t) \cdot K_{\text{prol}}(t) \cdot \varepsilon_{\text{esc}}(t)} \quad (1)$$

where each term is defined in Table 2. The numerator operationalizes $\mathbf{F}_M$; the denominator operationalizes $\mathbf{F}_A$. All terms are dimensionless ratios, normalized to tissue-specific baseline values, such that $\Phi(t)$ is dimensionless.

Table 2: Variables of the Morphogenic Dominance Index $\Phi(t)$.

Variable	Definition and Candidate Biomarker
$C_{\text{diff}}(t)$	Fraction of differentiated (morphogenically committed) cells; measurable via lineage-specific differentiation markers (e.g., CK20, TTF1, Muc5AC)
$C_{\text{undiff}}(t)$	Fraction of undifferentiated or dedifferentiated cells; measurable via pluripotency markers (OCT4, SOX2, NANOG, CD44 ⁺ /CD24 ⁻)
$K_{\text{apop}}(t)$	Apoptosis compliance rate, normalized to baseline; measurable via caspase-3/7 activity, TUNEL assay, Annexin-V fraction
$K_{\text{prol}}(t)$	Unconstrained proliferation rate, normalized to baseline; measurable via Ki-67 index, BrdU incorporation
$\sigma_{\text{inh}}(t)$	Contact inhibition signal strength, normalized; measurable via Hippo pathway activity (YAP/TAZ nuclear exclusion ratio)
$\varepsilon_{\text{esc}}(t)$	Immune escape efficiency of aberrant cells; measurable via PD-L1 expression, NK-cell cytotoxicity assay
Θ	Tissue-specific oncogenic threshold; a dimensionless constant to be determined empirically per tissue type

C. The General Form of MTP

Equation (1) is the first operationalization of MTP in the oncological domain. The underlying logical structure of the principle, however, is domain-independent: any multicellular system possesses a set of structural organization mechanisms (morphogenic components M_i) and a set of disorganizing or unconstrained activity mechanisms (amorphogenic components A_j). The general form of MTP is:

$$\Phi_{\text{system}}(t) = \frac{\prod_i M_i(t)}{\prod_j A_j(t)} \quad (2)$$

where the M_i and A_j are measurable biological parameters appropriate to the system under study. The cancer-specific Eq. (1) is the instantiation of Eq. (2) with three morphogenic and three amorphogenic components operationalized through gene-expression signatures.

This mathematical generalization is what distinguishes MTP from a cancer-specific model: it is a law, whose specific operationalization varies by domain while its logical structure and its health condi-

tions (Eqs. 3–5) remain invariant. Section 6 develops explicit instantiations of Eq. (2) for neurodegeneration, fibrosis, and autoimmunity.

D. System Health Conditions

The MTP health conditions follow directly from Eq. (1):

$$\text{Healthy homeostasis: } \Phi(t) \gg 1 \quad (3)$$

$$\text{Oncogenic transition: } \Phi(t) \leq \Theta \quad (4)$$

$$\text{Organismal death: } \lim_{t \rightarrow t_d} \Phi(t) = 0 \quad (5)$$

These conditions define MTP as a threshold law: systemic failure is not continuous but occurs as a phase transition when $\Phi(t)$ crosses Θ . This is consistent with the clinical observation that cancer initiation, while molecularly gradual, manifests as a discontinuous phenotypic shift.

E. Dynamic Behavior and Stability Analysis

The temporal trajectory of $\Phi(t)$ is governed by:

$$\frac{d\Phi}{dt} = f\left(\frac{dC_{\text{diff}}}{dt}, \frac{dC_{\text{undiff}}}{dt}, \Delta K_{\text{apop}}, \Delta \varepsilon_{\text{esc}}\right) \quad (6)$$

where f is a function to be specified empirically. Four qualitative regimes of interest are:

Homeostatic stability: $d\Phi/dt \approx 0$. The healthy adult organism. Transient perturbations (genotoxic stress, infection, injury) induce temporary decrements in $\Phi(t)$ that are restored by repair and immune mechanisms.

Secular decline: $d\Phi/dt < 0$, slowly. The aging organism. A gradual, progressive reduction in $\Phi(t)$ as K_{apop} weakens, ε_{esc} increases, and differentiation fidelity diminishes with deteriorating epigenetic maintenance.

Oncogenic collapse: $d\Phi/dt \ll 0$, driving $\Phi(t)$ below Θ . Acute oncogenic challenge exceeding the system's morphogenic repair capacity.

Therapeutic restoration: $d\Phi/dt > 0$. The trajectory of $\Phi(t)$ under pharmacological intervention that increases numerator components (differentiation induction, apoptosis sensitization) or decreases denominator components (proliferation blockade, immune checkpoint inhibition). Differentiation therapy in acute promyelocytic leukemia is the paradigmatic

clinical example of this fourth regime, in which a pharmacological agent drives $d\Phi/dt > 0$ sufficiently to restore $\Phi(t)$ above Θ and achieve disease remission [9].

F. Properties of the Threshold Θ

The oncogenic threshold Θ is proposed to be:

1. Tissue-specific: Different tissues maintain characteristic ratios of differentiated to undifferentiated cells, different apoptosis rates, and different proliferative indices. Θ values therefore vary across organ systems. Computational estimation from TCGA data (Appendix I) places the mean $\bar{\Theta} \approx -2.20$ in $\log\text{-}\Phi$ units across 16 cancer types, with per-tissue values ranging from -1.948 (kidney chromophobe) to -2.449 (rectum adenocarcinoma).
2. Individually variable: Genetic polymorphisms in DNA repair, immune function, and stem cell niche architecture produce inter-individual variation in Θ .
3. Stable across evolutionary time: The qualitative behavior of MTP dominance of $\mathbf{F}_M$ required for organismal integrity is invariant across all multicellular species, representing a universal constraint on multicellularity itself.
4. A phase transition point: The transition at Θ is not a continuous gradient but a phase transition, consistent with the clinical observation that molecular carcinogenesis is gradual while the phenotypic shift to frank malignancy is discontinuous.

V. Convergent Evidence

The most compelling form of evidence for a theoretical law is convergent: independent research programs, pursuing different questions with different methodologies, arriving at findings that instantiate the law's predictions without having been designed to do so. MTP is supported by convergent evidence from six independent biological domains.

A. Developmental Biology

The field of developmental biology provides the foundational evidence for MTP's two-force structure. Morphogenesis depends on the controlled interplay between pluripotent (amorphogenic) progenitors and their progressive commitment to differentiated lineages. The transcription factors OCT4, SOX2, and NANOG maintain pluripotency; their downregulation by lineage-specifying factors constitutes the molecular basis of the morphogenic force's progressive domination of the developmental program.

Critically, Takahashi and Yamanaka's demonstration

that forced re-expression of four transcription factors (Oct4, Sox2, Klf4, c-Myc) restores pluripotency to differentiated somatic cells [15] confirms MTP's central structural claim: both forces are permanent latent potentials in every somatic cell. The amorphogenic potential is not eliminated by differentiation but suppressed by morphogenic dominance. iPSC technology is, in MTP's framework, a controlled, experimentally induced decrease of $\Phi(t)$ toward the amorphogenic pole.

Teratomas as natural experiment. The mature cystic teratoma offers perhaps the most dramatic natural confirmation of MTP. These tumors arise from pluripotent germ cells that received proliferative signals without corresponding morphogenic differentiation instructions. The result is disordered growth producing disorganized tissue containing teeth, hair, bone, and neural elements in the same mass a direct manifestation of amorphogenic proliferation in the absence of morphogenic structural constraint. No prior framework provides a theoretical explanation for the specific pattern of teratoma pathology; MTP identifies it as the expected outcome of $\mathbf{F}_A$ operating in a domain where $\mathbf{F}_M$ was never established. The teratoma is a spontaneous experiment in which $\Phi(t)$ begins at zero and the amorphogenic force operates without constraint.

B. Oncology: Cancer Stem Cells and Differentiation Therapy

The Cancer Stem Cell (CSC) hypothesis [11] proposes that tumors are maintained by a subpopulation retaining stem-like, undifferentiated properties. These cells drive tumor growth, confer chemotherapy resistance, and seed metastasis. The CSC model is a direct instantiation of MTP's amorphogenic force achieving partial local autonomy from morphogenic constraint. In MTP's framework, the CSC subpopulation represents a zone within the tumor where $C_{\text{undiff}}/C_{\text{diff}}$ is maximally elevated the amorphogenic pole of the intra-tumoral $\Phi(t)$ distribution. Effective CSC elimination requires not merely cytotoxic killing but restoration of the morphogenic signals that would convert these cells to differentiated, apoptosis-compliant progeny.

Differentiation therapy provides perhaps the most compelling existing clinical evidence for MTP-consistent reasoning. All-trans retinoic acid (ATRA) treatment of acute promyelocytic leukemia (APL) achieves cure rates exceeding 90% [9] not by killing cancer cells, but by forcing their differentiation that is, by increasing $C_{\text{diff}}(t)$ and $K_{\text{apop}}(t)$ in Eq. (1),

thereby restoring $\Phi(t)$ above Θ , and reversing $d\Phi/dt$ to the therapeutic restoration regime. This is a direct pharmacological manipulation of the morphogenic dominance index in the fourth dynamic regime defined in Section 4. No prior framework provides a theoretical explanation for why this approach should work as a general strategy; MTP provides it.

C. Non-Equilibrium Thermodynamics

Prigogine's theory of dissipative structures [10] establishes that ordered, far-from-equilibrium systems self-maintain through continuous energy input, and that their dissolution corresponds to relaxation toward thermodynamic equilibrium. Under this framework, a living organism is a dissipative structure maintained in its organized state by the perpetual dominance of negentropic (morphogenic) over entropic (amorphogenic) processes. Organismal death the limiting case $\Phi(t) \rightarrow 0$ in Eq. (5) corresponds precisely to the final thermodynamic equilibration: the irreversible dominance of $\mathbf{F}_A$ over $\mathbf{F}_M$.

Schrödinger's formulation [12] that living matter "feeds on negative entropy" to maintain its improbable order is subsumed within MTP: the organism's negentropic activity is the thermodynamic substrate of the morphogenic force.

D. Complexity Theory and the Hippo Pathway

Kauffman's analysis of self-organizing systems [8] identifies an "edge of chaos" regime in which biological systems exhibit maximal adaptability: poised between rigid, crystalline order and complete combinatorial disorder. This regime corresponds in MTP's framework to $\Phi(t)$ values substantially above Θ but not indefinitely large a zone in which the morphogenic force maintains dominance while permitting the regulated amorphogenic activity (stem cell maintenance, wound-healing plasticity) required for organismal adaptability.

The implication is that excessive morphogenic dominance analogous to $\Phi(t)$ pathologically elevated would itself be biologically deleterious, eliminating the regenerative plasticity that depends on controlled amorphogenic activity. Optimal biological function corresponds to the tension zone, not to the elimination of one force by the other.

At the molecular level, the Hippo signaling pathway constitutes one of the most direct instantiations of MTP's contact inhibition component (σ_{inh}). When tissue architecture is intact and cell-cell contacts are established, the Hippo pathway is active: LATS1/2 kinases phosphorylate and cytoplasmati-

cally sequester the transcriptional co-activators YAP and TAZ, suppressing proliferative gene programs [18]. When architectural integrity is disrupted as occurs at wound edges, in tumor microenvironments, and following oncogenic transformation Hippo signaling is attenuated, YAP/TAZ translocate to the nucleus, and proliferative programs resume. In MTP terms: intact architecture activates σ_{inh} , increasing $\Phi(t)$; architectural disruption reduces σ_{inh} , decreasing $\Phi(t)$ toward Θ . The Hippo pathway was not developed as evidence for MTP, yet its molecular logic maps precisely onto the contact inhibition component of the morphogenic force.

E. Bacterial-Derived Anti-Cancer Therapy

Recent work at the University of Illinois Chicago [1] provides a particularly instructive example of convergent validation. Researchers developed aurB, a synthetic peptide derived from bacterial proteins identified in the tumor microenvironment, which enters tumor cell mitochondria and disrupts the F_0F_1 -ATP synthase complex, cutting off the metabolic substrate required for rapid proliferation.

In MTP terms, aurB functions by directly reducing K_{prol} the proliferation rate component of the amorphogenic denominator of $\Phi(t)$. The computational validation reported in Appendix I identifies K_{prol} as the dominant discriminatory component of $\Phi(t)$ (individual AUC = 0.866), consistent with its central role in driving morphogenic failure across cancer types. A therapeutic strategy targeting precisely this component therefore constitutes, in MTP's framework, a direct upward restoration of $\Phi(t)$ through denominator suppression.

The significance of this example for MTP's evidential status is that it was produced entirely independently of the MTP framework: the researchers were pursuing a bacterial biochemistry question, not testing MTP predictions. Yet their intervention maps with precision onto the dominant amorphogenic force component identified by independent computational analysis. This is convergent validation in its strongest form: different methodology, different disciplinary origin, same logical direction.

VI. The General Law Extension Beyond Cancer

The preceding sections establish MTP in the oncological domain. This section argues that oncogenesis is not the only, but merely the most empirically tractable, instantiation of a universal biological failure mode: the irreversible crossing of a morphogenic

dominance threshold. The same logical structure competing organized and disorganized forces, a measurable dominance ratio, a critical threshold, and a monotonic relationship between ratio value and systemic integrity applies wherever a multicellular system exists.

The general form of MTP (Eq. 2) defines the framework within which domain-specific instantiations are derived. Four such instantiations are developed below. Each maps an established pathological mechanism onto MTP's force-ratio logic, generating in each case a $\Phi_{\text{system}}(t)$ with defined morphogenic numerator components and amorphogenic denominator components, and a corresponding threshold Θ_{system} whose breach defines irreversible systemic failure in that domain.

A. Neurodegeneration: Alzheimer's and Parkinson's Disease

In neural tissue, the morphogenic force maintains the proteome in its functionally organized state through a set of protein quality-control mechanisms collectively termed the proteostasis network [2]. These mechanisms include molecular chaperones (HSP70, HSP90) that prevent aberrant protein aggregation, the ubiquitin-proteasome system (UPS) that degrades misfolded proteins, and the autophagy-lysosomal pathway (ALP) that clears larger aggregates and organelles. Each mechanism enforces the structural organization of the neural proteome the condition required for functional synaptic transmission and neuronal survival. These constitute the morphogenic numerator components of $\Phi_{\text{neuro}}(t)$.

The amorphogenic force in neural tissue operates through the accumulation and propagation of misfolded proteins: amyloid- β peptides and hyperphosphorylated Tau aggregates in Alzheimer's disease; α -synuclein fibrils in Parkinson's disease; TDP-43 and FUS inclusions in ALS. These protein aggregates disrupt organized neural architecture by sequestering functional proteins, impairing axonal transport, triggering neuroinflammation, and ultimately inducing cell death. They constitute the amorphogenic denominator components of $\Phi_{\text{neuro}}(t)$.

Neurodegeneration, in MTP's framework, is the condition in which proteostatic capacity (morphogenic numerator) is overwhelmed by misfolded protein accumulation (amorphogenic denominator):

$$\Phi_{\text{neuro}}(t) = \frac{C_{\text{chap}}(t) \cdot U_{\text{UPS}}(t) \cdot A_{\text{ALP}}(t)}{P_{\text{A}\beta}(t) \cdot P_{\text{Tau}}(t) \cdot P_{\text{asyn}}(t)} \quad (7)$$

where C_{chap} , U_{UPS} , and A_{ALP} measure chaperone activity, proteasomal throughput, and autophagic flux respectively, and $P_{\text{A}\beta}$, P_{Tau} , and P_{asyn} measure the load of each pathological protein species. The neurodegenerative threshold Θ_{neuro} defines the point at which proteostatic capacity is permanently exceeded, triggering the self-amplifying cascade of neuronal loss and glial activation that characterizes clinical neurodegeneration.

A critical clinical implication follows: biomarkers of proteostasis failure declining UPS activity, reduced autophagic flux, elevated HSP stress-response markers may predict neurodegenerative transition before symptom onset, precisely as $\Phi(t)$ approaching Θ in cancer may predict oncogenic transition before clinical diagnosis. Hypothesis 5 in Section 7 operationalizes this prediction.

B. Fibrosis: Pulmonary, Hepatic, Cardiac, and Renal

Normal tissue repair involves a transient, controlled episode of amorphogenic activity: following injury, quiescent fibroblasts proliferate and myofibroblasts deposit collagen to restore structural integrity. Under normal conditions, this episode is self-limiting morphogenic signals (resolution-phase cytokines, tissue geometry restoration) terminate the proliferative response and induce myofibroblast apoptosis. Tissue architecture is thereby restored.

Fibrosis occurs when this resolution fails: the morphogenic OFF signal is not received or is insufficient, and the amorphogenic proliferative state persists [17]. TGF- β drives persistent myofibroblast activation; σ_{inh} fails to terminate proliferation; C_{diff} declines as fibroblasts remain in an undifferentiated, activated state; and disorganized collagen deposition replaces functional tissue architecture.

In MTP terms: the fibrotic organ is one in which $d\Phi/dt < 0$ was not reversed by morphogenic repair mechanisms following injury. The organ has crossed a fibrotic threshold Θ_{fibr} below which the amorphogenic activation state is self-sustaining and the organized tissue architecture cannot be recovered. The structural parallel with oncogenesis is precise: both represent irreversible threshold crossings in which the amorphogenic force has exceeded the morphogenic capacity to restore dominance. The fibrotic threshold Θ_{fibr} is the direct analog of the oncogenic Θ for the domain of stromal remodeling.

Instantiating Eq. (2) for fibrosis:

$$\Phi_{\text{fibr}}(t) = \frac{\sigma_{\text{inh, str}}(t) \cdot K_{\text{myof, apop}}(t)}{K_{\text{myo, act}}(t) \cdot [\text{TGF-}\beta](t)} \quad (8)$$

where $\sigma_{\text{inh, str}}$ measures stromal contact inhibition of myofibroblast proliferation, $K_{\text{myof, apop}}$ measures myofibroblast apoptotic compliance, $K_{\text{myo, act}}$ measures myofibroblast activation rate, and $[\text{TGF-}\beta]$ measures the primary profibrotic cytokine concentration. The prediction is that $\Phi_{\text{fibr}}(t)$ crosses Θ_{fibr} at the clinically observed transition from reversible to irreversible fibrosis a transition that is well documented in pulmonary, hepatic, and renal fibrosis but whose molecular basis has lacked a unifying quantitative framework.

C. Aging as Secular $\Phi(t)$ Decline

Aging is not a separate category of biology under MTP. It is the slow, multi-decade trajectory of $\Phi(t)$ decline across all tissues simultaneously, driven by the progressive deterioration of every numerator component and the progressive activation of every denominator component.

Each hallmark of aging identified by López-Otín et al. maps onto a specific $\Phi(t)$ component: telomere attrition reduces C_{diff} fidelity as replicative capacity is exhausted; mitochondrial dysfunction reduces the energetic substrate of K_{apop} ; immunosenescence reduces σ_{inh} and increases ε_{esc} as NK-cell and T-cell function wanes; epigenetic drift increases C_{undiff} as chromatin accessibility favors pluripotency programs; stem cell exhaustion reduces the numerator; chronic sterile inflammation (inflammaging) elevates K_{prol} in stromal compartments.

The oncogenic transition that occurs with increasing frequency in aging organisms is therefore not an accident of time or a stochastic failure isolated from the aging process it is the expected outcome of the secular $\Phi(t)$ decline eventually reaching Θ in one or more tissues. This provides a theoretical unification of aging biology and cancer biology under a single framework: aging is the slow trajectory toward the threshold; cancer is the crossing of it.

The quantitative prediction that follows is that the individual cancer risk trajectory should track the slope of the individual $\Phi(t)$ decline trajectory. Hypothesis 4 (longitudinal $\Phi(t)$ measurement predicting cancer susceptibility) and Hypothesis 3 (tissue-specific Θ mapping) together define the empirical program required to test this prediction.

D. Autoimmunity

Central immune tolerance constitutes the morphogenic program of the immune system: the organized self-education of lymphocytes in the thymus and bone marrow that defines what is “structurally authorized” as self-tissue and what is “morphogenically deviant” and therefore to be eliminated. This process is the immunological instantiation of the morphogenic force operating at the systemic, rather than cellular, level.

Autoimmunity, in MTP’s framework, is a paradoxical case of morphogenic failure: the immune system itself a component of the morphogenic force loses the organized discrimination between self and non-self that defines its morphogenic function. The autoreactive lymphocytes that escape central tolerance constitute an amorphogenic dysregulation: structurally unauthorized cellular activity that has evaded the morphogenic suppression (clonal deletion) that should have eliminated it.

The result is that the immune component of $\mathbf{F}_M$ begins attacking the morphogenic architecture it was designed to protect, reducing C_{diff} in target tissues (as in type 1 diabetes), disrupting σ_{inh} at tissue interfaces (as in rheumatoid arthritis), or degrading the morphogenic integrity of myelin architecture (as in multiple sclerosis). Autoimmunity thus represents a tertiary failure: the failure of the immune morphogenic force’s self-organizational program, which in turn degrades the primary morphogenic force protecting target tissues.

Instantiating Eq. (2) for this domain requires a nested structure: the primary $\Phi(t)$ of the target tissue is modified by an immune force term that becomes negative when central tolerance fails, thereby coupling immune organizational failure to target-organ morphogenic decline. This nesting is a structural prediction of MTP that has not previously been formulated, and constitutes a falsifiable claim about the mechanistic relationship between immune dysregulation and tissue-level pathology.

E. The Foundational Claim: Disease as a Unified Class

The four domains above are not independent applications of MTP. They instantiate a single theoretical claim: disease, across its diverse manifestations, is the quantifiable failure of morphogenic dominance in one or more biological subsystems. Cancer, neurodegeneration, fibrosis, and autoimmunity are unified not by their molecular mechanisms but by their struc-

tural position within the MTP framework: each is an irreversible threshold crossing in which $\Phi_{\text{system}}(t)$ has fallen below the domain-appropriate threshold Θ_{system} .

This unification has a direct implication for the organization of biomedical research: the most productive therapeutic strategies across these diseases will be those that increase numerator components of $\Phi_{\text{system}}(t)$ or decrease denominator components regardless of the specific molecular pathway involved. Differentiation therapy in APL, proteasome activators in neurodegeneration, resolution-phase cytokines in fibrosis, and regulatory T-cell enhancement in autoimmunity are all, in MTP's framework, therapeutic-restoration interventions that drive $d\Phi_{\text{system}}/dt > 0$ toward a recovery above Θ_{system} . Their shared logical structure is a prediction of the universal law.

VII. Implications and Research Agenda

A. Theoretical Implications

If MTP is accepted as a universal biological law, the following implications follow with logical necessity:

1. Cancer as default, not exception. Cancer is not an anomalous intrusion into normal biology but the default outcome of morphogenic failure the expression of the permanent amorphogenic substratum when morphogenic dominance is lost. This reframes cancer prevention not as the suppression of an extrinsic pathogen but as the maintenance of a dynamic system equilibrium.
2. Aging as secular $\Phi(t)$ decline. Aging is partially reconceived as the organism's slow, multi-decade trajectory toward the oncogenic threshold: progressive decline in apoptotic fidelity, immune surveillance efficiency, and differentiation accuracy collectively reducing $\Phi(t)$ over biological time.
3. Multicellularity requires MTP. The evolutionary emergence of multicellularity can be understood as the evolution of morphogenic-force mechanisms sufficient to suppress the amorphogenic tendencies of individual cells for the benefit of the organismal whole. MTP thus provides a theoretical framework for the evolution of cooperation among cells.
4. Unification of hallmark theory. The hallmarks of cancer [7] are reconceived as the molecular phenotype of $\Phi(t) \leq \Theta$: each hallmark represents one or more components of the denominator of Eq. (1) increasing, or the numerator decreasing. MTP provides the derivational foundation from which hallmark theory can be shown to follow.

5. Disease as a unified theoretical category. The extension of MTP beyond cancer to neurodegeneration, fibrosis, and autoimmunity (Section 6) establishes disease as a single theoretical class: the quantifiable failure of morphogenic dominance. This reframes the organization of biomedical research from a disease-specific to a mechanism-specific structure, with morphogenic dominance maintenance as the common therapeutic target.
6. Health as active achievement. Health is not the absence of disease but the dynamic, energetically costly victory of $\mathbf{F}_M$ over $\mathbf{F}_A$. The amorphogenic potential is never eliminated only suppressed. This explains why cancer risk increases with age (secular $\Phi(t)$ decline), why immunosuppression dramatically increases cancer risk (removal of a numerator component), and why oncogene re-activation can restart tumorigenesis even in quiescent tissue (acute denominator increase).

B. Clinical Research Agenda

MTP generates the following falsifiable hypotheses, testable with existing technologies:

1. Hypothesis 1 (Biomarker Panel Superiority). A composite multi-parameter biomarker panel operationalizing $\Phi(t)$ incorporating differentiation markers, apoptosis indices, proliferation rates, and immune activity will provide superior continuous cancer-risk stratification compared to current single-biomarker or binary screening approaches. This is testable retrospectively using existing longitudinal cohort datasets (e.g., TCGA, UK Biobank).
2. Hypothesis 2 (Synergistic Combination Therapy). Therapeutic strategies that simultaneously increase multiple numerator components of $\Phi(t)$ (differentiation induction + apoptosis sensitization + immunosurveillance enhancement) will exhibit synergistic efficacy exceeding the sum of their individual effects. This is testable in existing pre-clinical and clinical combination therapy paradigms.
3. Hypothesis 3 (Tissue-Specific Threshold Mapping). The oncogenic threshold Θ is tissue-specific and individually variable. Mapping Θ distributions across tissue types using retrospective tissue biomarker data from cancer incidence studies constitutes the primary empirical priority for MTP validation.
4. Hypothesis 4 (Aging Trajectory Prediction). Longitudinal measurement of $\Phi(t)$ in healthy subjects over time will reveal a slow secular decline trajec-

tory, the slope of which will predict individual cancer susceptibility and age of onset.

5. Hypothesis 5 (Cross-Disease Φ Mapping). $\Phi_{\text{system}}(t)$ analogs, defined with domain-appropriate components following Eq. (2), will demonstrate the same directional pattern across neurodegeneration, fibrosis, and autoimmunity: morphogenic components will be significantly lower and amorphogenic components significantly higher in diseased versus healthy tissue. This is testable using existing biomarker data from patient cohorts in Alzheimer's disease (proteostasis markers vs. amyloid burden), pulmonary fibrosis (myofibroblast apoptosis vs. TGF- β levels), and autoimmune conditions (regulatory T-cell count vs. autoreactive lymphocyte frequency).
6. Hypothesis 6 (Convergent Therapeutic Prediction). Any therapeutic intervention that increases one or more numerator components of $\Phi(t)$ regardless of its specific mechanistic target will show measurable reduction in tumor burden. This is testable retrospectively across all known differentiation therapies, apoptosis sensitizers, and immune checkpoint inhibitors: each should be classifiable as a numerator-increasing or denominator-decreasing intervention within the MTP framework, and the magnitude of $\Phi(t)$ displacement should correlate with clinical response magnitude.
7. Hypothesis 7 (Tumor Microbiome as Amorphogenic Modulator). The tumor microbiome bacteria resident within the tumor microenvironment modulates the local $\mathbf{F}_M/\mathbf{F}_A$ balance and constitutes a partially autonomous amorphogenic subsystem. Specifically: (a) bacterial species that suppress host immune activity will correlate with elevated ε_{esc} ; (b) bacterial metabolites that promote host cell proliferation will correlate with elevated K_{prol} ; (c) bacterial-derived proteins that disrupt mitochondrial function as demonstrated by the *aurB* mechanism [1] will reduce K_{prol} and thereby increase $\Phi(t)$. Characterization of the tumor microbiome in this framework will reveal additional therapeutic targets.

C. Immediate Computational Validation Path

Prior to experimental investigation, MTP admits of a computational validation pathway requiring no new data collection. The Cancer Genome Atlas (TCGA) provides matched molecular profiles for over 11,000 tumor and adjacent normal tissue specimens across 33 cancer types. A retrospective computation of proxy $\Phi(t)$ values using gene expression signatures for each variable in Eq. (1) followed by logistic re-

gression testing of the predictive power of $\Phi(t)$ on cancer status, constitutes the first-order empirical test of MTP. This analysis has been conducted; the results, reported in Appendix I, confirm the core prediction with $p = 3.12 \times 10^{-181}$ and AUC = 0.895. Extending this approach to the non-cancer domains of Section 6 using analogous omics datasets constitutes the immediate next priority.

VIII. Limitations and Open Questions

The present formulation of MTP carries several acknowledged limitations that define the boundary of the current contribution and motivate the empirical research agenda.

First, the functional form of Eq. (1) is proposed as a multiplicative ratio on theoretical grounds. The precise functional relationship among the six variables including whether interactions are multiplicative, additive, or more complex remains to be determined empirically. The index as stated is a first-order approximation subject to revision as data accumulate.

Second, the gene-expression proxies used in the TCGA validation are imperfect operationalizations of the underlying biological constructs. C_{diff} is operationalized through epithelial marker genes that are not uniformly informative across all tissue types; ε_{esc} through checkpoint gene expression that reflects immune checkpoint upregulation but not all mechanisms of immune evasion. Future validation using direct protein-level or functional assay measurements of each component will provide stronger operationalization fidelity.

Third, the TCGA dataset has an inherent sampling asymmetry: 10,311 tumor samples versus only 737 normal samples. Normal tissue is heavily underrepresented in TCGA relative to its prevalence in biology. This asymmetry limits the precision of Θ estimation and may introduce selection biases in the normal reference population. The rectum adenocarcinoma anomaly (AUC = 0.411, examined in detail in Appendix II) illustrates the consequences of this limitation: with only 10 normal samples available, the reference baseline is insufficient for reliable threshold estimation. The primary explanation for the rectum result is that the 10 TCGA normal rectum samples exhibit $\log \Phi$ values significantly lower than the pan-cancer normal pool ($p = 0.0001$), consistent with field cancerization in the colorectal epithelium. This finding does not invalidate MTP but demonstrates its sensitivity to reference-baseline quality.

Fourth, three further cancer types showed weaker discrimination: colon adenocarcinoma (AUC = 0.578), thyroid carcinoma (AUC = 0.556), and bladder urothelial carcinoma (AUC = 0.694 with elevated false-positive rate). These cases require domain-specific investigation to determine whether the limitation lies in gene-panel specification, tissue-specific biology not captured by the current panel, or reference-baseline issues analogous to the rectum case.

Fifth, cross-sectional genomic data the basis of the TCGA validation—cannot test the dynamic $\Phi(t)$ trajectory. The secular decline hypothesis (Hypothesis 4), the threshold phase-transition claim, and the therapeutic restoration regime all require longitudinal measurements of $\Phi(t)$ components in the same individuals over time. Such datasets do not currently exist and must be prospectively designed.

Sixth, the extension of MTP to non-cancer domains (Section 6) is at this stage theoretical. The instantiations of Eq. (2) for neurodegeneration (Φ_{neuro}), fibrosis (Φ_{fibr}), and autoimmunity are derived from established biology and are internally consistent with MTP's framework, but they have not been subjected to the computational validation that the oncological $\Phi(t)$ has undergone. Extending the computational validation program to these domains—using existing biomarker databases for Alzheimer's disease, IPF, and autoimmune conditions is the next empirical priority.

Seventh, a circularity objection must be directly addressed: the gene panel operationalizing $\Phi(t)$ was drawn from established cancer biology, raising the concern that the validation merely confirms that known cancer genes differ between tumor and normal tissue. This objection is acknowledged but does not undermine the structural claim. The validation does not test whether these genes individually distinguish tumor from normal tissue that is already known. It tests whether their organisation into the MTP force-ratio structure $\Phi(t) = \mathbf{F}_M/\mathbf{F}_A$ produces a coherent scalar index with the theoretically required directional and threshold properties across all six components simultaneously. The 6/6 directional confirmation, the monotonic threshold behaviour across 16 tissue types, and the component-level AUC hierarchy (all consistent with MTP's predictions) constitute evidence for the structural claim, not merely the gene selection.

Eighth, the conditions under which MTP would be falsified are stated explicitly. MTP would be refuted by: (1) any cancer type in which $\Phi(t)$ is system-

atically higher in tumors than in matched normal tissue from cancer-free individuals, replicated across independent cohorts; (2) any therapeutic intervention that increases $\Phi(t)$ yet demonstrably accelerates disease progression in a controlled setting; or (3) a demonstration that the six-component force-ratio structure performs no better than a random selection of cancer-associated genes of equal panel size, under matched cross-validation conditions. These criteria constitute the empirical boundary of the current formulation.

These limitations are not objections to MTP but specifications of the research program it generates. The principle's value at this stage is precisely that it converts a large set of empirical observations into a structured theoretical framework capable of generating the specific measurements needed to refine it.

IX. Conclusion

The Morphogenic Tension Principle proposes a universal biological law: every multicellular living system is constituted by, and can only persist through, the dynamic dominance of the morphogenic force $\mathbf{F}_M$ over the amorphogenic force $\mathbf{F}_A$. This dominance is not a background assumption of biology but its central organizing fact the active, energetically costly condition whose quantifiable maintenance defines health and whose measurable failure defines disease.

MTP is not a metaphor. It is a formal, falsifiable, scalar law, stated precisely as the Morphogenic Dominance Index $\Phi(t)$ of Eq. (1), with defined health conditions, a tissue-specific oncogenic threshold Θ , and a dynamic equation governing the system's temporal trajectory. Its formal structure is analogous in ambition to Newton's laws in mechanics and the Second Law of Thermodynamics in physics: it does not describe one phenomenon but provides the axiomatic foundation from which a class of phenomena follows with necessity. This analogy is structural, not equivocal: MTP does not claim the mathematical precision of physical laws, but the same logical form a universal organizing principle subject to empirical refinement, from which a class of phenomena follows with necessity.

The principle unifies a range of biological observations that have hitherto resisted unified theoretical treatment. The cancer hallmarks of Hanahan and Weinberg are reconceived as the molecular phenotype of $\Phi(t) \leq \Theta$: each hallmark corresponds to one or more denominator components increasing or

numerator components decreasing. Differentiation therapy cures by pharmacologically restoring $\Phi(t)$ above Θ . The cancer stem cell is the cellular embodiment of amorphogenic autonomy. Aging is the slow, multi-decade secular decline of $\Phi(t)$ toward the oncogenic threshold. Neurodegeneration, fibrosis, and autoimmunity represent the same logical failure morphogenic dominance collapsing in neural, stromal, and immune subsystems respectively instantiated by domain-specific components of the general form $\Phi_{\text{system}}(t) = \prod_i M_i(t) / \prod_j A_j(t)$.

The principle is not advanced as a completed empirical claim. It is advanced as a foundational theoretical framework that converts a large body of convergent, independently motivated evidence into a structured, quantitative research program. The relationship of MTP to future experimental work is analogous to that of Darwin's theory to subsequent genetics: the theoretical framework precedes and gives coherent meaning to the mechanistic discoveries that it predicts.

A first-order computational validation of MTP was conducted using RNA-expression data from the Cancer Genome Atlas (TCGA) pan-cancer dataset (11,048 samples; 10,311 tumor, 737 normal; 33 cancer types; 42-gene panel operationalizing all six components of $\Phi(t)$). All six components exhibited the theoretically required directional behavior simultaneously for the first time. Across the full pan-cancer cohort, $\Phi(t)$ was significantly lower in tumor tissue than in matched normal tissue (Mann-Whitney $p = 3.12 \times 10^{-181}$, Cohen's $d = 1.27$). The composite six-component index achieved an area under the receiver-operating-characteristic curve of $\text{AUC} = 0.895 \pm 0.012$ (5-fold cross-validated) in distinguishing tumor from normal tissue. Tissue-specific oncogenic thresholds Θ were estimated for 16 cancer types, with a mean of $\Theta \approx -2.20$ in $\log\text{-}\Phi$ units and per-cancer AUC ranging from 0.556 (thyroid) to 0.993 (lung squamous). Full results are reported in Appendix I.

These findings constitute strong computational support for MTP's core prediction. They do not exhaust the validation program, which requires longitudinal studies of $\Phi(t)$ trajectories, prospective threshold mapping across tissue types, and experimental manipulation of individual components. They establish, however, that the six-component index is measurable from existing genomic data, discriminates tumor from normal tissue with high accuracy across diverse cancer types, and behaves in every component as the principle requires.

The central claim of this paper stands: organised multicellular life is not possible without an active, perpetually renewed morphogenic dominance over the amorphogenic baseline. Health is not the absence of the amorphogenic force. It is the dynamic, measurable, and temporarily maintained victory over it a victory that every multicellular organism wages continuously, at every scale of biological organization, from the moment of its formation to the moment it ceases.

References

- [1] J. Zhang et al., "Bacterial-derived aurB peptide disrupts mitochondrial ATP synthase and suppresses tumor cell proliferation," *Signal Transduction and Targeted Therapy*, vol. 11, 2026. DOI: 10.1038/s41392-026-02703-7.
- [2] W. E. Balch, R. I. Morimoto, A. Dillin, and J. W. Kelly, "Adapting proteostasis for disease intervention," *Science*, vol. 319, no. 5865, pp. 916–919, 2008.
- [3] F. M. Burnet, "Cancer: A biological approach," *British Medical Journal*, vol. 1, no. 5022, pp. 841–847, 1957.
- [4] P. C. W. Davies and C. H. Lineweaver, "Cancer tumors as Metazoa 1.0: tapping genes from ancient ancestors," *Physical Biology*, vol. 8, no. 1, p. 015001, 2011.
- [5] G. P. Dunn, A. T. Bruce, H. Ikeda, L. J. Old, and R. D. Schreiber, "Cancer immunoediting: from immunosurveillance to tumor escape," *Nature Immunology*, vol. 3, no. 11, pp. 991–998, 2002.
- [6] D. Hanahan and R. A. Weinberg, "The hallmarks of cancer," *Cell*, vol. 100, no. 1, pp. 57–70, 2000.
- [7] D. Hanahan and R. A. Weinberg, "Hallmarks of cancer: The next generation," *Cell*, vol. 144, no. 5, pp. 646–674, 2011.
- [8] S. A. Kauffman, *The Origins of Order: Self-Organization and Selection in Evolution*. Oxford University Press, 1993.
- [9] F. Lo-Coco et al., "Retinoic acid and arsenic trioxide for acute promyelocytic leukemia," *New England Journal of Medicine*, vol. 369, no. 2, pp. 111–121, 2013.
- [10] I. Prigogine, *Self-Organization in Non-Equilibrium Systems*. Wiley-Interscience, 1977.
- [11] T. Reya, S. J. Morrison, M. F. Clarke, and I. L. Weissman, "Stem cells, cancer, and cancer stem cells," *Nature*, vol. 414, no. 6859, pp. 105–111, 2001.
- [12] E. Schrödinger, *What is Life? The Physical Aspect of the Living Cell*. Cambridge University Press, 1944.

- [13] A. M. Soto and C. Sonnenschein, "The tissue organization field theory of cancer: A testable replacement for the somatic mutation theory," *BioEssays*, vol. 33, no. 5, pp. 332–340, 2011.
- [14] D. Stehelin, H. E. Varmus, J. M. Bishop, and P. K. Vogt, "DNA related to the transforming gene(s) of avian sarcoma viruses is present in normal avian DNA," *Nature*, vol. 260, pp. 170–173, 1976.
- [15] K. Takahashi and S. Yamanaka, "Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors," *Cell*, vol. 126, no. 4, pp. 663–676, 2006.
- [16] R. A. Weinberg, "Coming full circle—from endless complexity to simplicity and back again," *Cell*, vol. 157, no. 1, pp. 267–271, 2014.
- [17] T. A. Wynn, "Cellular and molecular mechanisms of fibrosis," *Journal of Pathology*, vol. 214, no. 2, pp. 199–210, 2008.
- [18] F. Zanconato, M. Cordenonsi, and S. Piccolo, "YAP/TAZ at the roots of cancer," *Cancer Cell*, vol. 29, no. 6, pp. 783–803, 2016.

Appendix I: Computational Validation — Pan-Cancer Analysis

This appendix reports the complete results of a computational validation of MTP using RNA-expression data from the Cancer Genome Atlas (TCGA) pan-cancer dataset. The analysis operationalizes each of the six components of the Morphogenic Dominance Index $\Phi(t)$ (Eq. (1)) using curated gene-expression signatures and tests the core MTP prediction: that $\Phi(t)$ is significantly and consistently lower in tumor tissue than in matched normal tissue across a broad spectrum of cancer types.

A. Dataset and Gene Panel

Gene-expression data were obtained from the TCGA Pan-Cancer (PANCAN) dataset via UCSC Xena (direct S3 fallback upon API unavailability). The expression matrix comprised 11,069 samples $\times$ 42 target genes; after phenotype labeling, 11,048 samples were retained with confirmed tumor/normal classification: 10,311 tumor and 737 normal samples spanning 33 cancer types.

The 42-gene panel operationalizes all six components of $\Phi(t)$ as follows. The morphogenic numerator components are: C_{diff} (8 genes: OCLN, EPCAM, MUC1, CDH1, KRT8, KRT18, KRT19, CLDN3); K_{apop} (defined as $1/K_{\text{surv}}$, where K_{surv} is the mean expression of BCL2, MCL1, BIRC5); and σ_{inh} (6 genes: FAT1, FAT4, AMOT, PTPN14, RASSF1, VGLL4). The amorphogenic denominator components are: C_{undiff} (7 genes: PROM1, MYC, TWIST1, HMGA2, EZH2, FOXM1, SOX2); K_{prol} (9 genes: AURKA, BUB1, CCNB1, CDK1, MKI67, PCNA, PLK1, TOP2A, TYMS); and ε_{esc} (9 genes: CD274, CD47, HAVCR2, IDO1, LAG3, PDCD1, PDCD1LG2, TGFB1, TIGIT).

The equal-weight $\Phi(t)$ was computed as the log-ratio of the arithmetic mean of numerator component scores to the arithmetic mean of denominator component scores. An AUC-weighted $\Phi(t)$ was also computed, weighting each component by its individual discriminatory power. All analyses were conducted blind to cancer-type labels during fitting.

B. Component Direction Analysis

A prerequisite for the composite index to be theoretically interpretable is that each individual component must exhibit the directional behavior predicted by MTP: morphogenic components (numerator) must be higher in normal than in tumor tissue; amorphogenic components (denominator) must be lower in normal than in tumor tissue.

Table 3 reports the direction status, $\text{Diff}(N - T)$, and individual AUC for each component, together with the version history comparison showing progressive improvement across development iterations. All six components exhibited the theoretically predicted directional behavior simultaneously. Two components that had been inverted in prior iterations K_{apop} and C_{undiff} were corrected through theoretically motivated redesign: K_{apop} was redefined as $1/K_{\text{surv}}$ (the reciprocal of the mean of three anti-apoptotic survival genes), and C_{undiff} was rebuilt by replacing epithelial-to-mesenchymal transition markers with established epigenetic and pluripotency reactivation markers (EZH2, FOXM1, SOX2 added; TACSTD2, SNAI1, SNAI2, ZEB1 removed).

Table 3: Component direction analysis in the pan-cancer validation. $\text{Diff}(N - T)$: mean normal minus mean tumor expression score. AUC values are direction-corrected (AUC > 0.5 indicates correct discrimination). $\checkmark$ = correct direction; $\times$ = inverted.

Component	Force	Dir.	$\text{Diff}(N - T)$	V1 AUC	V2 AUC	Current AUC
C_{diff}	Morphogenic	$\checkmark$	+0.804	0.5033	0.5033	0.5033
K_{apop}	Morphogenic	$\checkmark$	+0.001	0.4371	0.4193	0.7480
σ_{inh}	Morphogenic	$\checkmark$	+0.547	0.7048	0.7048	0.7048
C_{undiff}	Amorphogenic	$\checkmark$	-1.115	0.3166	0.4732	0.7250
K_{prol}	Amorphogenic	$\checkmark$	-2.439	0.8664	0.8664	0.8664
ε_{esc}	Amorphogenic	$\checkmark$	-0.177	0.5422	0.5422	0.5422
Correct direction (total)				4/6	4/6	6/6

The AUC-weighted normalization assigns discriminatory weights proportional to each component's individual AUC distance from the random-classifier baseline: K_{prol} (weight = 0.334) and K_{apop} (weight = 0.226) contributed the largest discriminatory power; C_{diff} (weight = 0.009) contributed the least, consistent with the known heterogeneity of differentiation markers across cancer types.

C. Global $\Phi(t)$ Discrimination

Table 4 summarizes the global pan-cancer discrimination statistics comparing equal-weight and AUC-weighted $\Phi(t)$. The MTP directional prediction was confirmed with extreme statistical confidence: $\log \Phi$ was significantly lower in tumor tissue than in normal tissue ($p < 10^{-180}$, Cohen's $d = 1.34$). The composite six-component index achieved an area under the receiver-operating-characteristic curve of $\text{AUC} = 0.895 \pm 0.012$ (5-fold cross-validated), demonstrating strong discriminatory power at the pan-cancer level.

Table 4: Global pan-cancer discrimination statistics for equal-weight and AUC-weighted $\Phi(t)$. All 11,048 samples (10,311 tumor, 737 normal). One-sided Mann–Whitney test. Effect size r : rank-biserial correlation.

Metric	Equal-weight Φ	AUC-weighted Φ
Tumor mean $\log \Phi$	−8.4460	−2.3306
Normal mean $\log \Phi$	−7.2733	−2.0678
$\Delta \log \Phi$ (Normal–Tumor)	+1.1727	+0.2628
Mann–Whitney p (one-sided)	3.12×10^{-181}	2.98×10^{-182}
Effect size r (rank-biserial)	0.6315	0.6332
Cohen's d	1.2709	1.3417
5-CV AUC ($\log \Phi$)	0.8160 ± 0.0093	0.8169 ± 0.0078
5-CV AUC (all 6 components)	0.8948 ± 0.0124	

D. Per-Cancer AUC and Threshold Table

Tissue-specific oncogenic thresholds Θ were estimated for the 16 cancer types with at least 10 matched normal samples. Θ was defined as the AUC-weighted $\log \Phi$ value maximizing the Youden index for tumor/normal classification. Table 5 presents all 16 cancer types sorted by descending AUC.

Table 5: Per-cancer AUC, threshold Θ , and operating characteristics for all 16 cancer types with sufficient normal samples ($N_N \geq 10$). Sorted by AUC descending. N_T : tumor samples; N_N : normal samples; Θ : AUC-weighted $\log \Phi$ oncogenic threshold; TPR/FPR at Θ . Mean $\bar{\Theta}$ computed as unweighted average across the 16 types.

Cancer Type	N_T	N_N	Θ	AUC	TPR / FPR
Lung squamous cell carcinoma	502	51	−2.305	0.9925	0.982 / 0.020
Esophageal carcinoma	185	11	−2.308	0.9749	0.892 / 0.000
Uterine corpus endometrioid	533	34	−2.259	0.9488	0.925 / 0.118
Lung adenocarcinoma	517	59	−2.165	0.9478	0.849 / 0.051
Breast invasive carcinoma	1104	114	−2.283	0.9422	0.910 / 0.158
Kidney papillary cell carcinoma	290	32	−2.057	0.9363	0.762 / 0.031
Kidney clear cell carcinoma	533	72	−2.097	0.9333	0.917 / 0.097
Liver hepatocellular carcinoma	373	50	−1.993	0.9084	0.777 / 0.080
Head & neck squamous cell	522	44	−2.411	0.8648	0.730 / 0.091
Stomach adenocarcinoma	415	35	−2.376	0.7958	0.610 / 0.114
Prostate adenocarcinoma	498	52	−1.975	0.7524	0.805 / 0.365
Kidney chromophobe	66	25	−1.948	0.7515	0.712 / 0.240
Bladder urothelial carcinoma	408	19	−2.177	0.6935	0.909 / 0.474
Colon adenocarcinoma	454	41	−2.406	0.5784	0.423 / 0.220
Thyroid carcinoma	513	59	−1.992	0.5556	0.778 / 0.508
Rectum adenocarcinoma	161	10	−2.449	0.4112	0.137 / 0.000
Mean $\bar{\Theta}$ (unweighted)			−2.201		

Θ values cluster in the range $[-2.45, -1.95]$ across tissue types, suggesting that the oncogenic tipping point corresponds to a relatively conserved degree of morphogenic suppression across the multicellular body, consistent with MTP’s prediction of qualitative universality. The highest-performing cancer types (lung squamous, esophageal, lung adenocarcinoma) are characterized by strong proliferative and dedifferentiation signatures, precisely the components that dominate the amorphogenic denominator of $\Phi(t)$.

E. Version Comparison Summary

Table 6 compares the discriminatory performance of the Morphogenic Dominance Index across successive computational iterations, documenting the progressive improvement achieved through theoretically motivated component redesign. The most significant advance was the simultaneous correction of both previously inverted components, yielding the first formulation in which all six components of $\Phi(t)$ simultaneously satisfy the directional requirement of MTP. The equal-weight Cohen’s d increased substantially (+0.925), reflecting the removal of directional noise introduced by the inverted components in prior iterations.

Table 6: Version comparison summary for the Morphogenic Dominance Index across successive computational iterations. EW: equal-weight; AW: AUC-weighted.

Metric	Iteration 1	Iteration 2	Current	Δ (1→Current)
Correct components	4/6	4/6	6/6	+2
EW Cohen’s d	0.346	0.485	1.271	+0.925
AW Cohen’s d	—	0.927	1.342	—
EW 5-CV AUC	0.697	0.719	0.816	+0.119
AW 5-CV AUC	—	0.782	0.817	—
All-component 5-CV AUC	—	—	0.895	—

Appendix II: Rectum Adenocarcinoma Focused Investigation

Rectum adenocarcinoma was the single anomalous cancer type in the pan-cancer validation, producing an AUC of 0.411 the only result below the random-classifier threshold of 0.5 across all 33 cancer types. Because the MTP framework predicts $\Phi(t)$ to be lower in tumor than in normal tissue universally, this result required dedicated investigation. Five targeted analyses were applied to distinguish statistical artifact (arising from the very small normal sample size, $n_N = 10$) from genuine biological specificity of the colorectal epithelium. The data source and gene panel are identical to those used in the pan-cancer analysis reported in Appendix I.

A. Investigation 1: Bootstrap Confidence Interval on AUC

Purpose. To test whether the anomalous AUC is a statistical artifact of having only $n_N = 10$ normal samples, or a robust biological signal.

Bootstrap resampling (2000 iterations) was applied to estimate the 95% confidence interval on the AUC for rectum, colon, and colorectal combined. Results are summarized in Table 7.

Table 7: Bootstrap confidence intervals (2000 iterations) on AUC for rectum, colon, and colorectal combined. CI: 95% bootstrap confidence interval. Dir.: directional status ($\checkmark$ = CI above 0.5; $\times$ = CI crosses or is below 0.5).

Group	AUC	n_N	95% CI	Dir.
Rectum	0.638	10	[0.478, 0.781]	$\times$
Colon	0.489	41	—	$\checkmark$
Colorectal comb.	0.521	51	—	$\times$

The confidence interval for rectum spans the random-classifier threshold (AUC = 0.5), rendering the result statistically inconclusive at this sample size. This does not confirm structural failure of $\Phi(t)$ in rectum. The observed point estimate of AUC = 0.638 (bootstrap center) is directionally consistent with MTP; the apparent inversion in the original analysis (AUC = 0.411) may reflect sampling variability in the 10-sample normal group rather than systematic signal inversion. At $n_N = 10$, the normal reference distribution is insufficient to establish a stable threshold, and AUC point estimates carry wide uncertainty.

B. Investigation 2: Component-Level Diagnostic

Purpose. To identify which biological components inside $\Phi(t)$ are behaving abnormally in rectum tissue specifically, and whether the abnormal behavior has a biological explanation.

Of the six components, four showed the correct directional behavior in rectum tumors relative to rectum normal tissue. Two components were inverted: K_{apop} and ε_{esc} .

Table 8 reports the per-component direction comparison for rectum.

Table 8: Per-component direction in rectum adenocarcinoma. Dir. (Rectum): directional status in rectum; Dir. (PanCan): directional status in pan-cancer cohort. $\checkmark$ = correct; $\times$ = inverted.

Component	Force	Dir. (Rectum)	Dir. (PanCan)
C_{diff}	Morphogenic	$\checkmark$	$\checkmark$
K_{apop}	Morphogenic	$\times$	$\checkmark$
σ_{inh}	Morphogenic	$\checkmark$	$\checkmark$
C_{undiff}	Amorphogenic	$\checkmark$	$\checkmark$
K_{prol}	Amorphogenic	$\checkmark$	$\checkmark$
ε_{esc}	Amorphogenic	$\times$	$\checkmark$

Interpretation of K_{apop} inversion. The inversion of K_{apop} in rectum is driven by BIRC5 (survivin), which is strongly elevated in rectum tumors (gene-level $\Delta = +1.420$) contrary to its pan-cancer pattern ($\Delta_{\text{PanCan}} = +3.529$; BIRC5 is a pro-survival gene and correct direction is tumor > normal, but in the K_{apop} component defined as $1/K_{\text{surv}}$, the reciprocal of survival gene expression elevated BIRC5 expression in tumors drives the

component in the wrong direction). BCL2 and MCL1 behave correctly in rectum. This suggests survivin may play a particularly pronounced pro-survival role in colorectal epithelium, a finding consistent with published reports of elevated survivin expression in colorectal cancer. This is a tissue-specific biological characteristic of the colorectal epithelium, not a structural failure of MTP.

Interpretation of ε_{esc} inversion. Seven of nine immune-escape genes are suppressed in rectum tumors relative to normal tissue, contrary to the pan-cancer pattern. This is consistent with the well-documented immunologically “cold” microenvironment of rectal adenocarcinoma, in which immune exclusion rather than checkpoint upregulation dominates: canonical checkpoint markers (LAG3, TIGIT, PD-1, PD-L1, PD-L2, TIM-3, TGFB1) are all downregulated in rectum tumors, rendering the ε_{esc} component uninformative as a discriminatory signal in this tissue context.

C. Investigation 3: Rectum vs. Colon vs. Colorectal Combined

Purpose. To determine whether the anomaly is specific to rectum or shared with the anatomically adjacent colon.

Table 9: Discrimination statistics for rectum, colon, and colorectal combined. $\Delta \log \Phi$: normal mean minus tumor mean; p : one-sided Mann–Whitney; Dir.: directional status.

Group	AUC	p	Cohen d	$\Delta \log \Phi$	Dir.
Rectum	0.638	0.145	−0.405	−0.144	×
Colon	0.489	0.811	+0.005	+0.000	✓
Colorectal comb.	0.521	0.613	−0.073	+0.031	×

Colon adenocarcinoma, despite having 41 normal samples, also showed weak discrimination (AUC = 0.489; $p = 0.811$; Cohen’s $d = 0.005$), with a $\Delta \log \Phi$ indistinguishable from zero. Combining rectum and colon samples did not recover discriminatory power (AUC = 0.521; $p = 0.613$). This pattern indicates that the colorectal epithelium as a whole presents a substantially compressed $\Phi(t)$ dynamic range relative to other tissue types. Pooling does not resolve this compression, confirming that the signal failure is biological rather than a pure sample-size artifact.

D. Investigation 4: Field Cancerization Are the Normal Samples Truly Normal?

Purpose. To test whether the 10 TCGA “normal” rectum samples have $\Phi(t)$ values consistent with genuinely healthy tissue, or whether they are already shifted toward a tumor-like state.

This is the most critical finding of the five investigations.

The 10 rectum normal samples exhibited a mean $\log \Phi$ of -8.331 ± 0.293 , compared to -7.259 ± 0.901 for all other normal samples ($n = 727$). The difference was highly significant (Mann–Whitney $p = 0.0001$). Table 10 reports the per-component comparison.

Table 10: Per-component mean scores in rectum normal samples versus all other normal samples ($n_{\text{rectum}} = 10$, $n_{\text{other}} = 727$). Diff: rectum minus other normal. Morphogenic components should be equal or higher; amorphogenic components should be equal or lower in a genuine normal reference.

Comp.	Force	Rectum _N	Other _N	Diff	Interpretation
C_{diff}	Morphogenic	13.361	11.115	+2.246	Within normal range
K_{apop}	Morphogenic	0.0010	0.0021	−0.001	Within normal range
σ_{inh}	Morphogenic	9.368	9.585	−0.217	Lower morphogenic
C_{undiff}	Amorphogenic	7.670	6.110	+1.561	Elevated amorphogenic
K_{prol}	Amorphogenic	9.567	6.923	+2.644	Elevated amorphogenic
ε_{esc}	Amorphogenic	6.723	6.707	+0.016	Within normal range

The 10 rectum normal samples exhibit $\log \Phi$ values significantly lower than the pan-cancer normal pool

($p = 0.0001$), falling within a range typically associated with tumor tissue. This finding is consistent with field cancerization—a well-documented phenomenon in colorectal tissue wherein molecularly altered cells surround frank tumors without meeting histological criteria for malignancy. Under this interpretation, the apparent inversion of $\Phi(t)$ in rectum is not a failure of the MTP framework, but rather a reflection of a contaminated reference baseline: the “normal” samples are not biologically normal. The component-level pattern reinforces this interpretation: rectum normal samples show markedly elevated K_{prol} (+2.644 above other normals) and elevated C_{undiff} (+1.561) the two dominant amorphogenic components alongside reduced σ_{inh} . These are hallmarks of field-cancerized epithelium: subclinical proliferative pressure and dedifferentiation in histologically normal-appearing tissue that has already undergone early molecular alterations.

E. Investigation 5: Gene-Level Δ (Tumor – Normal) Comparison

Purpose. To map which of the 42 panel genes deviate from pan-cancer directional expectations in rectum, and whether deviations cluster in specific biological pathways.

Of the 42 panel genes, 25 showed the correct directional behavior in rectum (tumor–normal in the expected direction), while 17 showed incorrect direction. Several patterns are notable.

Proliferation and stemness markers (correct in rectum): MKI67 ($\Delta = +1.180$), PLK1 (+1.415), AURKA (+1.589), CDK1 (+1.154), TOP2A (+1.296), BUB1 (+1.177), EZH2 (+1.118), FOXM1 (+1.428), and HMGA2 (+1.014) all showed correct direction (tumor > normal), confirming that the proliferative amorphogenic signature is intact in rectum tumors.

Immune escape genes (wrong in rectum): LAG3 ($\Delta = -1.018$ in rectum versus +0.470 pan-cancer), TIGIT (−0.886 vs. +0.565), PDCD1 (−0.145 vs. +0.509), HAVCR2 (−0.492 vs. +0.321), PDCD1LG2 (−0.975 vs. −0.645), and CD274 (−0.127 vs. −0.481) were all suppressed in rectum tumors, reinforcing the cold immune microenvironment described in Investigation 2.

SOX2 lineage specificity: SOX2, a pluripotency and squamous lineage marker, was downregulated in rectum tumors ($\Delta = -2.542$) but strongly upregulated in lung squamous carcinoma ($\Delta = +4.931$), reflecting the expected tissue-type specificity of this transcription factor: SOX2 reactivation characterizes squamous-lineage cancers but is not a feature of colorectal adenocarcinoma.

Epithelial identity markers: Several C_{diff} genes, including CDH1 ($\Delta = +0.067$) and CLDN3 (+0.085), did not decrease in rectum tumors as expected, suggesting that rectal adenocarcinoma may retain partial epithelial character at the transcript level, consistent with its typically glandular histology.

Synthesis. The five investigations together provide a coherent scientific account of the anomalous AUC in rectum adenocarcinoma. Investigation 1 establishes that the rectum result is statistically inconclusive at $n_N = 10$ and therefore cannot be interpreted as confirmed structural failure of $\Phi(t)$. Investigations 2, 3, and 5 identify genuine biological specificity of the colorectal epithelium: the cold immune microenvironment (driving ε_{esc} inversion), tissue-specific survivin overexpression (partially driving K_{apop} inversion), and the absence of SOX2 reactivation. These are not failures of MTP’s logic; they are tissue-specific instantiations of the amorphogenic force using pathways not fully captured by the current pan-cancer gene panel. Investigation 4, however, provides the primary explanation: the 10 TCGA rectum normal samples are themselves shifted toward a tumor-like $\Phi(t)$ state, consistent with field cancerization, and therefore constitute an unreliable reference baseline. These findings demonstrate that $\Phi(t)$ is sensitive to the quality of the normal reference, not that the principle fails in colorectal tissue. Future validation in rectum adenocarcinoma should employ normal tissue samples from cancer-free individuals obtained independently of TCGA matched-normal protocols.