


# Delaunay Triangulations and Voronoi Diagrams

**Motivation:** Spatial Interpolation, Nearest Neighbor Queries

**Algorithmic Technique:** Randomized Incremental Construction


**Data Structures:** Voronoi diagrams, Delaunay triangulations








# Spatial Interpolation



height measurements

$$p = (p_x, p_y, p_z)$$

# Spatial Interpolation



height measurements


$$p = (p_x, p_y, p_z)$$



projection

$$\pi(p) = (p_x, p_y, 0)$$

# Spatial Interpolation



height measurements

$$p = (p_x, p_y, p_z)$$




projection

$$\pi(p) = (p_x, p_y, 0)$$

Question: How do we estimate the height at  $(x, y)$ ?



# Spatial Interpolation



height measurements

$$p = (p_x, p_y, p_z)$$




projection

$$\pi(p) = (p_x, p_y, 0)$$

Interpolation 1: assign height of nearest neighbor

# Spatial Interpolation




height measurements

$$p = (p_x, p_y, p_z)$$



projection

$$\pi(p) = (p_x, p_y, 0)$$



Interpolation 1: assign height of nearest neighbor



# Spatial Interpolation



height measurements

$$p = (p_x, p_y, p_z)$$




projection

$$\pi(p) = (p_x, p_y, 0)$$

Interpolation 1: assign height of nearest neighbor



# Spatial Interpolation



height measurements

$$p = (p_x, p_y, p_z)$$




projection

$$\pi(p) = (p_x, p_y, 0)$$

**Interpolation 2:** triangulate & interpolate within triangles

# Spatial Interpolation




height measurements

$$p = (p_x, p_y, p_z)$$



projection


$$\pi(p) = (p_x, p_y, 0)$$




**Interpolation 2:** triangulate & interpolate within triangles



# Voronoi Diagrams




# Voronoi Diagrams






# Motivation



# Voronoi Diagrams

# Motivation


## Voronoi Diagrams






# Motivation

## Voronoi Diagrams




# Voronoi diagrams



**Voronoi diagram** of  $\{p_1, \dots, p_n\}$ : subdivision of the plane into cells, such that a point  $q$  is in the cell of  $p_i$  if and only if  $\text{dist}(q, p_i) < \text{dist}(q, p_j)$  for all  $i \neq j$ .


# Voronoi diagrams



**Voronoi diagram** of  $\{p_1, \dots, p_n\}$ : subdivision of the plane into cells, such that a point  $q$  is in the cell of  $p_i$  if and only if  $\text{dist}(q, p_i) < \text{dist}(q, p_j)$  for all  $i \neq j$ .




# Voronoi diagrams



**Voronoi diagram** of  $\{p_1, \dots, p_n\}$ : subdivision of the plane into cells, such that a point  $q$  is in the cell of  $p_i$  if and only if  $\text{dist}(q, p_i) < \text{dist}(q, p_j)$  for all  $i \neq j$ .

# Voronoi diagrams



long history

- Descartes 1644
- Dirichlet 1850
- Voronoi 1907

**Voronoi diagram** of  $\{p_1, \dots, p_n\}$ : subdivision of the plane into cells, such that a point  $q$  is in the cell of  $p_i$  if and only if  $\text{dist}(q, p_i) < \text{dist}(q, p_j)$  for all  $i \neq j$ .

# Voronoi diagrams


for 2 points






# Voronoi diagrams

for 2 points




# Voronoi diagrams

for 2 points



# Voronoi diagrams






# Voronoi diagrams


for 2 points

$$b(p, q) = \{x \in \mathbb{R}^2 : |xp| = |xq|\}$$


bisector of  $p$  and  $q$



# Voronoi diagrams



# Voronoi diagrams






# Voronoi diagrams

for  $n$  points

$$P = \{p_1, p_2, \dots, p_n\}$$




# Voronoi diagrams



# Voronoi diagrams

for  $n$  points

$\text{Vor}(P) = \text{Voronoi diagram of } P$



# Quiz

Are Voronoi cells convex?

A: yes

B: no

C: only if they are bounded

# Quiz

Are Voronoi cells convex?

A: yes

B: no

C: only if they are bounded



# Quiz

Are Voronoi cells convex?

A: yes for this lets have a closer look at cells

B: no

C: only if they are bounded


# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .


Voronoi diagram:



# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

Voronoi diagram:




# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

Voronoi diagram:

Voronoi cell

$$\mathcal{V}(\{p\}) =$$






# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

Voronoi diagram:

Voronoi cell

$$\mathcal{V}(\{p\}) = \mathcal{V}(p) =$$




# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

Voronoi diagram:

Voronoi cell

$$\mathcal{V}(\{p\}) = \mathcal{V}(p) = \{x \in \mathbb{R}^2 : |xp| < |xq| \forall q \in P \setminus \{p\}\}$$




# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

Voronoi diagram:

Voronoi cell

$$\begin{aligned}\mathcal{V}(\{p\}) = \mathcal{V}(p) &= \{x \in \mathbb{R}^2 : |xp| < |xq| \forall q \in P \setminus \{p\}\} \\ &= \bigcap_{q \neq p} h(p, q)\end{aligned}$$



# Voronoi diagrams


Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

Voronoi diagram:

Voronoi cell

$$\begin{aligned}\mathcal{V}(\{p\}) = \mathcal{V}(p) &= \{x \in \mathbb{R}^2 : |xp| < |xq| \forall q \in P \setminus \{p\}\} \\ &= \bigcap_{q \neq p} h(p, q)\end{aligned}$$

intersection of convex sets  $\rightarrow$  cells are convex




# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

Voronoi diagram:

Voronoi cell

$$\begin{aligned}\mathcal{V}(\{p\}) = \mathcal{V}(p) &= \{x \in \mathbb{R}^2 : |xp| < |xq| \forall q \in P \setminus \{p\}\} \\ &= \bigcap_{q \neq p} h(p, q)\end{aligned}$$





# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .


Voronoi diagram:

Voronoi cell

$$\begin{aligned}\mathcal{V}(\{p\}) = \mathcal{V}(p) &= \{x \in \mathbb{R}^2 : |xp| < |xq| \forall q \in P \setminus \{p\}\} \\ &= \bigcap_{q \neq p} h(p, q)\end{aligned}$$

Voronoi edge

$$\mathcal{V}(\{p, p'\}) =$$



# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .


Voronoi diagram:

Voronoi cell

$$\begin{aligned}\mathcal{V}(\{p\}) &= \mathcal{V}(p) = \{x \in \mathbb{R}^2 : |xp| < |xq| \ \forall q \in P \setminus \{p\}\} \\ &= \bigcap_{q \neq p} h(p, q)\end{aligned}$$

Voronoi edge

$$\mathcal{V}(\{p, p'\}) = \{x : |xp| = |xp'| \text{ and } |xp| < |xq| \ \forall q \neq p, p'\}$$



# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .


Voronoi diagram:

Voronoi cell

$$\begin{aligned}\mathcal{V}(\{p\}) &= \mathcal{V}(p) = \{x \in \mathbb{R}^2 : |xp| < |xq| \ \forall q \in P \setminus \{p\}\} \\ &= \bigcap_{q \neq p} h(p, q)\end{aligned}$$

Voronoi edge

$$\begin{aligned}\mathcal{V}(\{p, p'\}) &= \{x : |xp| = |xp'| \text{ and } |xp| < |xq| \ \forall q \neq p, p'\} \\ &= \partial\mathcal{V}(p) \cap \partial\mathcal{V}(p')\end{aligned}$$



# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .


Voronoi diagram:

Voronoi cell

$$\begin{aligned}\mathcal{V}(\{p\}) &= \mathcal{V}(p) = \{x \in \mathbb{R}^2 : |xp| < |xq| \ \forall q \in P \setminus \{p\}\} \\ &= \bigcap_{q \neq p} h(p, q)\end{aligned}$$

Voronoi edge

$$\begin{aligned}\mathcal{V}(\{p, p'\}) &= \{x : |xp| = |xp'| \text{ and } |xp| < |xq| \ \forall q \neq p, p'\} \\ &= \text{rel-int}(\partial\mathcal{V}(p) \cap \partial\mathcal{V}(p')), \text{ i.e. without endpoints}\end{aligned}$$



# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .


Voronoi diagram:

Voronoi cell

$$\begin{aligned}\mathcal{V}(\{p\}) = \mathcal{V}(p) &= \{x \in \mathbb{R}^2 : |xp| < |xq| \ \forall q \in P \setminus \{p\}\} \\ &= \bigcap_{q \neq p} h(p, q)\end{aligned}$$

Voronoi edge

$$\begin{aligned}\mathcal{V}(\{p, p'\}) &= \{x : |xp| = |xp'| \text{ and } |xp| < |xq| \ \forall q \neq p, p'\} \\ &= \text{rel-int}(\partial\mathcal{V}(p) \cap \partial\mathcal{V}(p')), \text{ i.e. without endpoints}\end{aligned}$$




# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

Voronoi diagram:

Voronoi cell

$$\begin{aligned}\mathcal{V}(\{p\}) &= \mathcal{V}(p) = \{x \in \mathbb{R}^2 : |xp| < |xq| \ \forall q \in P \setminus \{p\}\} \\ &= \bigcap_{q \neq p} h(p, q)\end{aligned}$$



Voronoi edge

$$\begin{aligned}\mathcal{V}(\{p, p'\}) &= \{x : |xp| = |xp'| \text{ and } |xp| < |xq| \ \forall q \neq p, p'\} \\ &= \text{rel-int}(\partial\mathcal{V}(p) \cap \partial\mathcal{V}(p')), \text{ i.e. without endpoints}\end{aligned}$$

Voronoi vertex

$$\mathcal{V}(\{p, p', p''\}) =$$




# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

Voronoi diagram:

Voronoi cell

$$\begin{aligned}\mathcal{V}(\{p\}) &= \mathcal{V}(p) = \{x \in \mathbb{R}^2 : |xp| < |xq| \ \forall q \in P \setminus \{p\}\} \\ &= \bigcap_{q \neq p} h(p, q)\end{aligned}$$



Voronoi edge

$$\begin{aligned}\mathcal{V}(\{p, p'\}) &= \{x : |xp| = |xp'| \text{ and } |xp| < |xq| \ \forall q \neq p, p'\} \\ &= \text{rel-int}(\partial\mathcal{V}(p) \cap \partial\mathcal{V}(p')), \text{ i.e. without endpoints}\end{aligned}$$

Voronoi vertex

$$\mathcal{V}(\{p, p', p''\}) = \partial\mathcal{V}(p) \cap \partial\mathcal{V}(p') \cap \partial\mathcal{V}(p'')$$

# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .


Voronoi diagram:

Voronoi cell

a subdivision

Voronoi edge

Voronoi vertex



# Voronoi diagrams

Let  $P$  be a set of points in the plane and  $p, p', p'' \in P$ .

Voronoi diagram:


Voronoi cell

Voronoi edge

Voronoi vertex


a subdivision

a plane graph





# Structure

If points in  $P$  are not all collinear, then  $\text{Vor}(P)$  is connected





# Structure

If points in  $P$  are not all collinear, then  $\text{Vor}(P)$  is connected



# Structure


If points in  $P$  are not all collinear, then  $\text{Vor}(P)$  is connected



next: Complexity of  $\text{Vor}(P)$

# Voronoi Diagrams

Complexity



# Quiz

How many sides can a single cell have?

A: 6

B:  $n/2$

C:  $n - 1$



# Quiz

How many sides can a single cell have?

A: 6


B:  $n/2$

C:  $n - 1$

# Complexity of a Cell

How many sides can a single cell have?

A cell may have up to  $n - 1$  sides




# Complexity of a Cell

How many sides can a single cell have?

A cell may have up to  $n - 1$  sides


Can all cells have  $n - 1$  sides?



# Complexity of a Cell

How many sides can a single cell have?

A cell may have up to  $n - 1$  sides




Can all cells have  $n - 1$  sides?

How many cells/edges/vertices may  $\text{Vor}(P)$  have?


# Complexity

Euler's formula for plane connected graphs:  $V - E + F = 2$




# Complexity

Euler's formula for plane connected graphs:  $V - E + F = 2$



# Complexity


Euler's formula for plane connected graphs:  $V - E + F = 2$



$$5 - 7$$

# Complexity

Euler's formula for plane connected graphs:  $V - E + F = 2$




$$5 - 7 + 4 = 2$$



# Complexity


Euler's formula for plane connected graphs:  $V - E + F = 2$



# Complexity

Euler's formula for plane connected graphs:  $V - E + F = 2$


■



# Complexity

Euler's formula for plane connected graphs:  $V - E + F = 2$

$$(n_v + 1) - n_e + n = 2$$




# Complexity

Euler's formula for plane connected graphs:  $V - E + F = 2$

$$(n_v + 1) - n_e + n = 2$$

Handshaking lemma:

$$2n_e = \sum_v \deg(v)$$




# Complexity

Euler's formula for plane connected graphs:  $V - E + F = 2$

$$(n_v + 1) - n_e + n = 2$$

Handshaking lemma:

$$\begin{aligned} 2n_e &= \sum_v \deg(v) \\ &\geq 3(n_v + 1) \\ &\text{(degree at least 3)} \end{aligned}$$



# Quiz

1.  $(n_v + 1) - n_e + n = 2$
2.  $2n_e \geq 3(n_v + 1)$

Which bounds does this imply on  $n_v$  and  $n_e$  (as tight as possible)?

- A:  $n_e \leq n(n + 1)/2, n_v \leq n(n + 1)/3$
- B:  $n_e \leq 3n - 6, n_v \leq 2n - 5$
- C:  $n_e = n_v = n$

# Quiz

1.  $(n_v + 1) - n_e + n = 2$
2.  $2n_e \geq 3(n_v + 1)$

Which bounds does this imply on  $n_v$  and  $n_e$  (as tight as possible)?

A:  $n_e \leq n(n + 1)/2, n_v \leq n(n + 1)/3$

B:  $n_e \leq 3n - 6, n_v \leq 2n - 5$

C:  $n_e = n_v = n$

# Quiz

1.  $(n_v + 1) - n_e + n = 2$
2.  $2n_e \geq 3(n_v + 1)$

Which bounds does this imply on  $n_v$  and  $n_e$  (as tight as possible)?

Plug in 1. into 2.:

$$2n_e \geq 3(2 + n_e - n)$$



# Quiz

1.  $(n_v + 1) - n_e + n = 2$
2.  $2n_e \geq 3(n_v + 1)$

Which bounds does this imply on  $n_v$  and  $n_e$  (as tight as possible)?

Plug in 1. into 2.:

$$2n_e \geq 3(2 + n_e - n)$$

Re-order terms:

3.  $n_e \leq 3n - 6$

# Quiz

1.  $(n_v + 1) - n_e + n = 2$
2.  $2n_e \geq 3(n_v + 1)$

Which bounds does this imply on  $n_v$  and  $n_e$  (as tight as possible)?

Plug in 1. into 2.:

$$2n_e \geq 3(2 + n_e - n)$$

Re-order terms:

$$3. \quad n_e \leq 3n - 6$$

Plug in 3. into 1.:


$$n_v \leq 2n - 5$$

# Voronoi Diagram Complexity


The Voronoi diagram of  $n$  points in the plane has at most

$3n - 6$  Voronoi edges

$2n - 5$  Voronoi vertices




# Delaunay Triangulations



# Delaunay triangulation

Let  $\text{Vor}(P)$  be the Voronoi diagram of  $P$ .




# Delaunay triangulation

Let  $\text{Vor}(P)$  be the Voronoi diagram of  $P$ .

**Definition:** The graph  $\mathcal{G} = (P, E)$  with

$$E = \{(p, q) \mid \mathcal{V}(p) \text{ and } \mathcal{V}(q) \text{ are adjacent}\}$$

is called the **dual graph** of  $\text{Vor}(P)$ .



# Delaunay triangulation


Let  $\text{Vor}(P)$  be the Voronoi diagram of  $P$ .

**Definition:** The graph  $\mathcal{G} = (P, E)$  with

$$E = \{(p, q) \mid \mathcal{V}(p) \text{ and } \mathcal{V}(q) \text{ are adjacent}\}$$


is called the **dual graph** of  $\text{Vor}(P)$ .

**Definition:** The straight-line drawing of  $\mathcal{G}$  is called **Delaunay graph**  $\mathcal{DG}(P)$ .



# Properties


$\mathcal{DG}(P)$  has no crossing edges.





# Properties


$\mathcal{DG}(P)$  has no crossing edges.



**Proof by contradiction.** Suppose there is an intersection. Then it is determined by 4 points.

# Properties


$\mathcal{DG}(P)$  has no crossing edges.



**Proof by contradiction.** Suppose there is an intersection. Then it is determined by 4 points.

# Properties


$\mathcal{DG}(P)$  has no crossing edges.



**Proof by contradiction.** Suppose there is an intersection. Then it is determined by 4 points.

# Properties

$\mathcal{DG}(P)$  has no crossing edges.




**Proof by contradiction.** Suppose there is an intersection. Then it is determined by 4 points.

# Properties

$\mathcal{DG}(P)$  has no crossing edges.


**Delaunay triangulation:** add edges until all faces are triangles



# Properties


$\mathcal{DG}(P)$  has no crossing edges.

**Delaunay triangulation:** add edges until all faces are triangles




# Characterization of Voronoi vertices/edges

**Definition:** Let  $q$  be a point. Define  $C_P(q)$  as the largest disk with center  $q$  containing no points of  $P$  in its interior.



# Characterization of Voronoi vertices/edges


**Definition:** Let  $q$  be a point. Define  $C_P(q)$  as the largest disk with center  $q$  containing no points of  $P$  in its interior.







# Characterization of Voronoi vertices/edges

**Definition:** Let  $q$  be a point. Define  $C_P(q)$  as the largest disk with center  $q$  containing no points of  $P$  in its interior.



**Observation:**



- A point  $q$  is a **Voronoi vertex**  $\Leftrightarrow |C_P(q) \cap P| \geq 3$ ,
- The bisector  $b(p_i, p_j)$  defines a **Voronoi edge**  
 $\Leftrightarrow \exists q \in b(p_i, p_j)$  with  $C_P(q) \cap P = \{p_i, p_j\}$ .



# Empty-circle property

Theorem about Voronoi diagrams:

- a point  $q$  is a Voronoi vertex  
 $\Leftrightarrow |C_P(q) \cap P| \geq 3,$
- the bisector  $b(p_i, p_j)$  defines a Voronoi edge  
 $\Leftrightarrow \exists q \in b(p_i, p_j)$  with  $C_P(q) \cap P = \{p_i, p_j\}.$





# Empty-circle property

Theorem about Voronoi diagrams:

- a point  $q$  is a Voronoi vertex  
 $\Leftrightarrow |C_P(q) \cap P| \geq 3$ ,
- the bisector  $b(p_i, p_j)$  defines a Voronoi edge  
 $\Leftrightarrow \exists q \in b(p_i, p_j)$  with  $C_P(q) \cap P = \{p_i, p_j\}$ .

**Theorem:** Let  $P$  be a set of points.

- points  $p, q, r$  are vertices of the same face in  $\mathcal{DG}(P) \Leftrightarrow$  circle through  $p, q, r$  is empty,
- edge  $pq$  is in  $\mathcal{DG}(P)$   
 $\Leftrightarrow$  there is an empty circle  $C_{p,q}$  through  $p$  and  $q$ .



# Empty-circle property


Theorem about Voronoi diagrams:

- a point  $q$  is a Voronoi vertex  
 $\Leftrightarrow |C_P(q) \cap P| \geq 3$ ,
- the bisector  $b(p_i, p_j)$  defines a Voronoi edge  
 $\Leftrightarrow \exists q \in b(p_i, p_j)$  with  $C_P(q) \cap P = \{p_i, p_j\}$ .

**Theorem:** Let  $P$  be a set of points.

- points  $p, q, r$  are vertices of the same face in  $\mathcal{DG}(P) \Leftrightarrow$  circle through  $p, q, r$  is empty,
- edge  $pq$  is in  $\mathcal{DG}(P)$   
 $\Leftrightarrow$  there is an empty circle  $C_{p,q}$  through  $p$  and  $q$ .

**Corollary:** Let  $P$  be a set of points and  $\mathcal{T}$  a triangulation of  $P$ .  $\mathcal{T}$  is a Delaunay triangulation  $\Leftrightarrow$  circumcircle of every triangle is empty.



# Quiz

How many triangles does a (Delaunay) triangulation of  $n$  points contain at most?

A:  $2n/3 - 1$

B:  $2n - 5$

C:  $3n - 6$

# Quiz

How many triangles does a (Delaunay) triangulation of  $n$  points contain at most?

A:  $2n/3 - 1$

B:  $2n - 5$

C:  $3n - 6$

# Quiz

How many triangles does a (Delaunay) triangulation of  $n$  points contain at most?

A:  $2n/3 - 1$

B:  $2n - 5$  = no. Voronoi vertices (or use Euler's formula)


C:  $3n - 6$

# Angle-optimal Triangulations




# Triangulation of a point set

**Definition:** A **triangulation** of a point set  $P \subset \mathbb{R}^2$  is a maximal planar subdivision with vertex set  $P$ .



# Triangulation of a point set


**Definition:** A **triangulation** of a point set  $P \subset \mathbb{R}^2$  is a maximal planar subdivision with vertex set  $P$ .



**Observations:**

# Triangulation of a point set

**Definition:** A **triangulation** of a point set  $P \subset \mathbb{R}^2$  is a maximal planar subdivision with vertex set  $P$ .




## Observations:

- all inner faces are triangles

# Triangulation of a point set

**Definition:** A **triangulation** of a point set  $P \subset \mathbb{R}^2$  is a maximal planar subdivision with vertex set  $P$ .




**Observations:**

- all inner faces are triangles

# Triangulation of a point set

**Definition:** A **triangulation** of a point set  $P \subset \mathbb{R}^2$  is a maximal planar subdivision with vertex set  $P$ .




## Observations:

- all inner faces are triangles
- outer face is complement of convex hull

# Triangulation of a point set

**Definition:** A **triangulation** of a point set  $P \subset \mathbb{R}^2$  is a maximal planar subdivision with vertex set  $P$ .




## Observations:

- all inner faces are triangles
- outer face is complement of convex hull

# Triangulation of a point set

**Definition:** A **triangulation** of a point set  $P \subset \mathbb{R}^2$  is a maximal planar subdivision with vertex set  $P$ .



## Observations:

- all inner faces are triangles
- outer face is complement of convex hull

**Theorem 1:** Let  $P$  be a set of  $n$  non-collinear points and let  $h$  be the number of vertices of  $CH(P)$ . Then **every** triangulation of  $P$  has  $t(n, h)$  triangles and  $e(n, h)$  edges.

# Triangulation of a point set

**Definition:** A **triangulation** of a point set  $P \subset \mathbb{R}^2$  is a maximal planar subdivision with vertex set  $P$ .

Compute  $t(n, h)$  and  $e(n, h)$ !

## Observations:

- all inner faces are triangles
- outer face is complement of convex hull

**Theorem 1:** Let  $P$  be a set of  $n$  non-collinear points and let  $h$  be the number of vertices of  $CH(P)$ . Then **every** triangulation of  $P$  has  $t(n, h)$  triangles and  $e(n, h)$  edges.



# Triangulation of a point set

**Definition:** A **triangulation** of a point set  $P \subset \mathbb{R}^2$  is a maximal planar subdivision with vertex set  $P$ .

Compute  $t(n, h)$  and  $e(n, h)$ !  
*Euler's formula for connected plane graphs:*  
 $\# \text{ faces} - \# \text{ edges} + \# \text{ vertices} = 2$ ,  
also counting the outer face.


## Observations:

- all inner faces are triangles
- outer face is complement of convex hull

**Theorem 1:** Let  $P$  be a set of  $n$  non-collinear points and let  $h$  be the number of vertices of  $CH(P)$ . Then **every** triangulation of  $P$  has  $t(n, h)$  triangles and  $e(n, h)$  edges.

# Triangulation of a point set

**Definition:** A **triangulation** of a point set  $P \subset \mathbb{R}^2$  is a maximal planar subdivision with vertex set  $P$ .




## Observations:

- all inner faces are triangles
- outer face is complement of convex hull

**Theorem 1:** Let  $P$  be a set of  $n$  non-collinear points and let  $h$  be the number of vertices of  $CH(P)$ . Then **every** triangulation of  $P$  has  $(2n - 2 - h)$  triangles and  $(3n - 3 - h)$  edges.


# Motivation revisited



height measurements

$$p = (p_x, p_y, p_z)$$

# Motivation revisited




height measurements

$$p = (p_x, p_y, p_z)$$




$$\text{projection } \pi(p) = (p_x, p_y, 0)$$



Interpolation 1: assign height of nearest neighbor

→ Voronoi diagrams

# Motivation revisited




height measurements

$$p = (p_x, p_y, p_z)$$



$$\text{projection } \pi(p) = (p_x, p_y, 0)$$




**Interpolation 1:** assign height of nearest neighbor

→ Voronoi diagrams



# Motivation revisited




height measurements

$$p = (p_x, p_y, p_z)$$




$$\text{projection } \pi(p) = (p_x, p_y, 0)$$



Interpolation 2: triangulate & interpolate within triangles

# Motivation revisited




height measurements

$$p = (p_x, p_y, p_z)$$




$$\text{projection } \pi(p) = (p_x, p_y, 0)$$



Interpolation 2: triangulate & interpolate within triangles

# Motivation revisited




height measurements

$$p = (p_x, p_y, p_z)$$



projection  $\pi(p) = (p_x, p_y, 0)$





What is a 'good' triangulation?

**Interpolation 2:** triangulate & interpolate within triangles





# Back to height interpolation

Lets look at the interpolation along an edge:





# Back to height interpolation

Lets look at the interpolation along an edge:





# Back to height interpolation

Lets look at the interpolation along an edge:





# Back to height interpolation

Lets look at the interpolation along an edge:



# Back to height interpolation



Lets look at the interpolation along an edge:



**Intuition:** avoid 'thin' triangles!

# Back to height interpolation

Lets look at the interpolation along an edge:




**Intuition:** avoid 'thin' triangles!

or: maximize the smallest angle within triangles!

# Angle-optimal triangulations

**Definition:** Let  $\mathcal{T}$  be a triangulation of  $P$  with  $m$  triangles and  $3m$  vertices. Its **angle vector** is  $A(\mathcal{T}) = (\alpha_1, \dots, \alpha_{3m})$ , where  $\alpha_1, \dots, \alpha_{3m}$  are the angles of  $\mathcal{T}$  sorted by increasing value.

$$A(\mathcal{T}) = (60^\circ, 60^\circ, 60^\circ, 60^\circ, 60^\circ, 60^\circ)$$





# Angle-optimal triangulations

**Definition:** Let  $\mathcal{T}$  be a triangulation of  $P$  with  $m$  triangles and  $3m$  vertices. Its **angle vector** is  $A(\mathcal{T}) = (\alpha_1, \dots, \alpha_{3m})$ , where  $\alpha_1, \dots, \alpha_{3m}$  are the angles of  $\mathcal{T}$  sorted by increasing value.

- For two triangulations  $\mathcal{T}$  and  $\mathcal{T}'$  of  $P$  define order  $A(\mathcal{T}) > A(\mathcal{T}')$  according to the lexicographical order.

$$A(\mathcal{T}) = (60^\circ, 60^\circ, 60^\circ, 60^\circ, 60^\circ, 60^\circ)$$



$$A(\mathcal{T}') = (30^\circ, 30^\circ, 30^\circ, 30^\circ, 120^\circ, 120^\circ)$$





# Angle-optimal triangulations

**Definition:** Let  $\mathcal{T}$  be a triangulation of  $P$  with  $m$  triangles and  $3m$  vertices. Its **angle vector** is  $A(\mathcal{T}) = (\alpha_1, \dots, \alpha_{3m})$ , where  $\alpha_1, \dots, \alpha_{3m}$  are the angles of  $\mathcal{T}$  sorted by increasing value.

- For two triangulations  $\mathcal{T}$  and  $\mathcal{T}'$  of  $P$  define order  $A(\mathcal{T}) > A(\mathcal{T}')$  according to the lexicographical order.
- $\mathcal{T}$  is **angle optimal**, if  $A(\mathcal{T}) \geq A(\mathcal{T}')$  for all triangulations  $\mathcal{T}'$  of  $P$ .


$$A(\mathcal{T}) = (60^\circ, 60^\circ, 60^\circ, 60^\circ, 60^\circ, 60^\circ)$$



$$A(\mathcal{T}') = (30^\circ, 30^\circ, 30^\circ, 30^\circ, 120^\circ, 120^\circ)$$


# Edge flips

**Definition:** Let  $\mathcal{T}$  be a triangulation. An edge  $e$  of  $\mathcal{T}$  is **illegal**, if the smallest angle of the triangles incident to  $e$  can be increased by **flipping**  $e$ .




# Edge flips

**Definition:** Let  $\mathcal{T}$  be a triangulation. An edge  $e$  of  $\mathcal{T}$  is **illegal**, if the smallest angle of the triangles incident to  $e$  can be increased by **flipping**  $e$ .




# Edge flips

**Definition:** Let  $\mathcal{T}$  be a triangulation. An edge  $e$  of  $\mathcal{T}$  is **illegal**, if the smallest angle of the triangles incident to  $e$  can be increased by **flipping**  $e$ .




# Edge flips

**Definition:** Let  $\mathcal{T}$  be a triangulation. An edge  $e$  of  $\mathcal{T}$  is **illegal**, if the smallest angle of the triangles incident to  $e$  can be increased by **flipping**  $e$ .



# Edge flips


**Definition:** Let  $\mathcal{T}$  be a triangulation. An edge  $e$  of  $\mathcal{T}$  is **illegal**, if the smallest angle of the triangles incident to  $e$  can be increased by **flipping**  $e$ .



# Edge flips


**Definition:** Let  $\mathcal{T}$  be a triangulation. An edge  $e$  of  $\mathcal{T}$  is **illegal**, if the smallest angle of the triangles incident to  $e$  can be increased by **flipping**  $e$ .

**Observation:** Let  $e$  be an illegal edge in  $\mathcal{T}$  and let  $\mathcal{T}' = \text{flip}(\mathcal{T}, e)$ . Then  $A(\mathcal{T}') > A(\mathcal{T})$ .



# Quiz

How many edge flips are needed to remove all illegal edges?



A: 0


B: 2

C: 3



# Quiz

How many edge flips are needed to remove all illegal edges?




A: 0

B: 2

C: 3

# Quiz

How many edge flips are needed to remove all illegal edges?




A: 0

B: 2

C: 3

# Quiz

How many edge flips are needed to remove all illegal edges?





A: 0

B: 2

C: 3




# Legal Triangulations




# Thales theorem

**Theorem:** If  $ab$  is a diameter, then the angle at any third point on the circle  $c$  is  $90^\circ$ .




# Thales theorem

**Theorem:** If  $ab$  is a diameter, then the angle at any third point on the circle  $c$  is  $90^\circ$ .



**Theorem:** Let  $C$  be a circle,  $\ell$  a line intersecting  $C$  in points  $a$  and  $b$ , and  $p, q, r, s$  points lying on the same side of  $\ell$ . Suppose that  $p, q$  lie on  $C$ ,  $r$  lies inside  $C$ , and  $s$  lies outside  $C$ . Then  $\angle arb > \angle apb = \angle aqb > \angle asb$ , where  $\angle abc$  denotes the smaller angle defined by three points  $a, b, c$ .



$$\angle asb < \angle apb = \angle aqb < \angle arb$$


# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.

**Proof sketch:**






# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.

**Proof sketch:**



$$\varphi_{pr} > \theta_{pr}$$

$$\varphi_{ps} > \theta_{ps}$$

$$\varphi_{rq} > \theta_{rq}$$

$$\varphi_{sq} > \theta_{sq}$$


# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.

**Proof sketch:**




$$\begin{aligned} \varphi_{pr} &> \theta_{pr} \\ \varphi_{ps} &> \theta_{ps} \\ \varphi_{rq} &> \theta_{rq} \\ \varphi_{sq} &> \theta_{sq} \\ \varphi_p &= \theta_{rq} + \theta_{sq} \\ \varphi_q &= \theta_{pr} + \theta_{ps} \end{aligned}$$

# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.




# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.




# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.




# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.

**Definition:** A triangulation without illegal edges is called **legal triangulation**.



# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.

**Definition:** A triangulation without illegal edges is called **legal triangulation**.

Are there legal triangulations?

# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.

**Definition:** A triangulation without illegal edges is called **legal triangulation**.

```
1: while  $\mathcal{T}$  has illegal edge  $e$  do  
2:   flip( $\mathcal{T}, e$ )  
3: return  $\mathcal{T}$ 
```



# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.

**Definition:** A triangulation without illegal edges is called **legal triangulation**.

```
1: while  $\mathcal{T}$  has illegal edge  $e$  do  
2:   flip( $\mathcal{T}, e$ )  
3: return  $\mathcal{T}$ 
```

Does the algorithm terminate?

# Legal triangulations

**Lemma 1:** Let  $\Delta prq$  and  $\Delta pqs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta prq$ . Then:

$$\overline{pq} \text{ is illegal} \iff s \in \text{int}(C).$$

If  $p, q, r, s$  form a convex quadrilateral and  $s \notin \partial C$ , then either  $\overline{pq}$  or  $\overline{rs}$  is illegal.

**Definition:** A triangulation without illegal edges is called **legal triangulation**.

```
1: while  $\mathcal{T}$  has illegal edge  $e$  do  
2:   flip( $\mathcal{T}, e$ )  
3: return  $\mathcal{T}$ 
```


Does the algorithm terminate?

yes, since  $A(\mathcal{T})$  increases and #triangulations is finite

# Legal vs angle-optimal


**We know:** Every angle-optimal triangulation is legal.

But is every legal triangulation angle-optimal?



# Legal vs Delaunay

**Theorem:** Let  $P$  be a set of points in  $\mathbb{R}^2$ . A triangulation  $\mathcal{T}$  of  $P$  is legal if and only if  $\mathcal{T}$  is a Delaunay triangulation.




# Legal vs Delaunay

**Theorem:** Let  $P$  be a set of points in  $\mathbb{R}^2$ . A triangulation  $\mathcal{T}$  of  $P$  is legal if and only if  $\mathcal{T}$  is a Delaunay triangulation.

**Proof sketch:**

" $\Leftarrow$ " obvious, use




**Lemma 1:** Let  $\Delta pqr$  and  $\Delta prs$  be two adjacent triangles in  $\mathcal{T}$  and  $C$  the circumcircle of  $\Delta pqr$ . Then:  $\overline{pr}$  is illegal  $\Leftrightarrow s \in \text{int}(C)$ .

# Legal vs Delaunay

**Theorem:** Let  $P$  be a set of points in  $\mathbb{R}^2$ . A triangulation  $\mathcal{T}$  of  $P$  is legal if and only if  $\mathcal{T}$  is a Delaunay triangulation.

**Proof sketch:**

" $\Rightarrow$ "




- suppose  $s$  is in the interior of circumcircle of  $\triangle pqr$
- let  $\overline{pr}$  be the edge maximizing  $\angle psr$

# Legal vs Delaunay

**Theorem:** Let  $P$  be a set of points in  $\mathbb{R}^2$ . A triangulation  $\mathcal{T}$  of  $P$  is legal if and only if  $\mathcal{T}$  is a Delaunay triangulation.

**Proof sketch:**

" $\Rightarrow$ "




- suppose  $s$  is in the interior of circumcircle of  $\triangle pqr$
- let  $\overline{pr}$  be the edge maximizing  $\angle psr$

# Legal vs Delaunay

**Theorem:** Let  $P$  be a set of points in  $\mathbb{R}^2$ . A triangulation  $\mathcal{T}$  of  $P$  is legal if and only if  $\mathcal{T}$  is a Delaunay triangulation.

**Proof sketch:**

" $\Rightarrow$ "



- suppose  $s$  is in the interior of circumcircle of  $\triangle pqr$
- let  $\overline{pr}$  be the edge maximizing  $\angle psr$
- consider  $t$  adjacent to  $p$  and  $r$
- $s$  also lies in circumcircle of  $\triangle prt$




# Legal vs Delaunay

**Theorem:** Let  $P$  be a set of points in  $\mathbb{R}^2$ . A triangulation  $\mathcal{T}$  of  $P$  is legal if and only if  $\mathcal{T}$  is a Delaunay triangulation.

**Proof sketch:**

" $\Rightarrow$ "



- suppose  $s$  is in the interior of circumcircle of  $\triangle pqr$
- let  $\overline{pr}$  be the edge maximizing  $\angle psr$
- consider  $t$  adjacent to  $p$  and  $r$
- $s$  also lies in circumcircle of  $\triangle prt$
- Thales theorem:  $\angle tsr > \angle psr$
- *Contradiction to choice of  $\overline{pr}$*

# Legal vs Delaunay

**Theorem:** Let  $P$  be a set of points in  $\mathbb{R}^2$ . A triangulation  $\mathcal{T}$  of  $P$  is legal if and only if  $\mathcal{T}$  is a Delaunay triangulation.

**Observation:** If  $P$  is in general position, then the Delaunay triangulation of  $P$  is unique.

# Legal vs Delaunay

**Theorem:** Let  $P$  be a set of points in  $\mathbb{R}^2$ . A triangulation  $\mathcal{T}$  of  $P$  is legal if and only if  $\mathcal{T}$  is a Delaunay triangulation.

**Observation:** If  $P$  is in general position, then the Delaunay triangulation of  $P$  is unique.

$\Rightarrow$  legal triangulation unique

# Legal vs Delaunay

**Theorem:** Let  $P$  be a set of points in  $\mathbb{R}^2$ . A triangulation  $\mathcal{T}$  of  $P$  is legal if and only if  $\mathcal{T}$  is a Delaunay triangulation.

**Observation:** If  $P$  is in general position, then the Delaunay triangulation of  $P$  is unique.

$\Rightarrow$  legal triangulation unique

we know:  $\mathcal{T}$  angle optimal  $\Rightarrow \mathcal{T}$  legal

# Legal vs Delaunay

**Theorem:** Let  $P$  be a set of points in  $\mathbb{R}^2$ . A triangulation  $\mathcal{T}$  of  $P$  is legal if and only if  $\mathcal{T}$  is a Delaunay triangulation.

**Observation:** If  $P$  is in general position, then the Delaunay triangulation of  $P$  is unique.

$\Rightarrow$  legal triangulation unique

we know:  $\mathcal{T}$  angle optimal  $\Rightarrow \mathcal{T}$  legal

$\Rightarrow$  angle-optimal triangulation is  $\mathcal{DG}(P)$ !

# Quiz

In general position: legal  $\Leftrightarrow$  Delaunay  $\Leftrightarrow$  angle-optimal

**Question:** If points are not in general position, are Delaunay triangulations still angle optimal?

A: yes, but the proof is more complicated

B: no, but the smallest angle is still maximized

C: no, not even the smallest angle is maximized

# Quiz

In general position: legal  $\Leftrightarrow$  Delaunay  $\Leftrightarrow$  angle-optimal

**Question:** If points are not in general position, are Delaunay triangulations still angle optimal?

A: yes, but the proof is more complicated

B: no, but the smallest angle is still maximized

C: no, not even the smallest angle is maximized

# Quiz

In general position: legal  $\Leftrightarrow$  Delaunay  $\Leftrightarrow$  angle-optimal

**Question:** If points are not in general position, are Delaunay triangulations still angle optimal?


A: yes, but the proof is more complicated

B: no, but the smallest angle is still maximized

C: no, not even the smallest angle is maximized

If  $P$  not in general position, then the smallest angle in every triangulation of the “large” faces in  $\mathcal{DG}(P)$  is the same.  
(proof uses Thales theorem)






# Computing the Delaunay triangulation

Randomized Incremental Construction using Edge Flips


# Randomized incremental construction



# Randomized incremental construction

**Algorithm** DELAUNAYTRIANGULATION( $P$ )


- 1: Initialize  $\mathcal{T}$  as a large triangle  $\triangle p_0 p_{-1} p_{-2}$  containing all points from  $P$
- 2: Compute a random permutation of  $p_1, \dots, p_n$
- 3: **for**  $r \leftarrow 1$  **to**  $n$  **do**
- 4:   INSERT( $p_r, \mathcal{T}$ )
- 5: Discard  $p_0, p_{-1}$  and  $p_{-2}$  with all their incident edges
- 6: **return**  $\mathcal{T}$



# Randomized incremental construction

**Algorithm** INSERT( $p_r, \mathcal{T}$ )



- 1: find triangle  $\in \mathcal{T}$  containing  $p_r$
- 2: **if**  $p_r$  lies in  $\triangle p_i p_j p_k$  **then**
- 3:   add edges from  $p_r$  to  $p_i, p_j, p_k$
- 4:   LEGALIZEEDGE( $p_r, \overline{p_i p_j}, \mathcal{T}$ )
- 5:   LEGALIZEEDGE( $p_r, \overline{p_j p_k}, \mathcal{T}$ )
- 6:   LEGALIZEEDGE( $p_r, \overline{p_k p_i}, \mathcal{T}$ )



# Randomized incremental construction

## Algorithm INSERT( $p_r, \mathcal{T}$ )


- 1: find triangle  $\in \mathcal{T}$  containing  $p_r$
- 2: **if**  $p_r$  lies in  $\triangle p_i p_j p_k$  **then**
- 3:   add edges from  $p_r$  to  $p_i, p_j, p_k$
- 4:   LEGALIZEEDGE( $p_r, \overline{p_i p_j}, \mathcal{T}$ )
- 5:   LEGALIZEEDGE( $p_r, \overline{p_j p_k}, \mathcal{T}$ )
- 6:   LEGALIZEEDGE( $p_r, \overline{p_k p_i}, \mathcal{T}$ )
- 7: **else**                   *//  $p_r$  lies on edge  $\overline{p_i p_j}$*
- 8:   add edges from  $p_r$  to  $p_l, p_k$
- 9:   LEGALIZEEDGE( $p_r, \overline{p_i p_l}, \mathcal{T}$ )
- 10:   LEGALIZEEDGE( $p_r, \overline{p_l p_j}, \mathcal{T}$ )
- 11:   LEGALIZEEDGE( $p_r, \overline{p_j p_k}, \mathcal{T}$ )
- 12:   LEGALIZEEDGE( $p_r, \overline{p_k p_i}, \mathcal{T}$ )



# Randomized incremental construction

Algorithm  $\text{LEGALIZEEDGE}(p_r, \overline{p_i p_j}, \mathcal{T})$

- 1: **if**  $\overline{p_i p_j}$  is illegal **then** // Edge flip
- 2:   let  $\triangle p_i p_j p_h$  be the adjacent triangle
- 3:   replace  $\overline{p_i p_j}$  by  $\overline{p_r p_h}$
- 4:    $\text{LEGALIZEEDGE}(p_r, \overline{p_j p_h}, \mathcal{T})$
- 5:    $\text{LEGALIZEEDGE}(p_r, \overline{p_h p_i}, \mathcal{T})$




# Correctness

1. All edges inserted are legal

# Correctness

1. All edges inserted are legal
  - All edges inserted are adjacent to  $p_r$ .







# Correctness

## 1. All edges inserted are legal

- All edges inserted are adjacent to  $p_r$ .




- All edges inserted are Delaunay edges.



# Correctness

## 1. All edges inserted are legal


- All edges inserted are adjacent to  $p_r$ .



- All edges inserted are Delaunay edges.


## 2. All other edges are legal

- an edge can only be illegal if it is incident to a new triangle.



# Quiz

If  $\deg(p_r) = k$  after inserting  $p_r$  (not on an edge), how many triangles were created during the insertion process?




A: 3

B:  $k$

C:  $2k - 3$

# Quiz


If  $\deg(p_r) = k$  after inserting  $p_r$  (not on an edge), how many triangles were created during the insertion process?



A: 3

B:  $k$

C:  $2k - 3$




# Computing the Delaunay triangulation

Search Structure and Analysis

# Initialization


Choose  $p_0, p_{-1}, p_{-2}$  far enough away from  $P$ , such that they lie in none of the circles of  $P$  and such that  $P$  lies in their triangle.



# Initialization

Better:


Treat  $p_0, p_{-1}, p_{-2}$  *symbolically* by modifying tests/predicates used for point location and testing illegal edges.



# Search structure

Build **search structure** for point location: directed acyclic graph with

- leaves: current triangles
- inner nodes: deleted triangles






# Search structure

Build **search structure** for point location: directed acyclic graph with


- leaves: current triangles
- inner nodes: deleted triangles



# Search structure

Build **search structure** for point location: directed acyclic graph with


- leaves: current triangles
- inner nodes: deleted triangles



# Search structure

Build **search structure** for point location: directed acyclic graph with

- leaves: current triangles
- inner nodes: deleted triangles




# Analysis

**Lemma:** The expected number of triangles created is at most  $9n + 1 = O(n)$ .

# Analysis

**Lemma:** The expected number of triangles created is at most  $9n + 1 = O(n)$ .


**Proof:** How many triangles are created when  $p_r$  is inserted?



# Analysis

**Lemma:** The expected number of triangles created is at most  $9n + 1 = O(n)$ .

**Proof:** How many triangles are created when  $p_r$  is inserted?




**Backwards analysis:**

- every point in  $p_1, \dots, p_r$  has the same probability  $1/r$  to be the last point

# Analysis

**Lemma:** The expected number of triangles created is at most  $9n + 1 = O(n)$ .

**Proof:** How many triangles are created when  $p_r$  is inserted?




**Backwards analysis:**

- every point in  $p_1, \dots, p_r$  has the same probability  $1/r$  to be the last point
- expected degree of  $p_r$  is  $\leq 6$

# Analysis

**Lemma:** The expected number of triangles created is at most  $9n + 1 = O(n)$ .

**Proof:** How many triangles are created when  $p_r$  is inserted?



**Backwards analysis:**


- every point in  $p_1, \dots, p_r$  has the same probability  $1/r$  to be the last point
- expected degree of  $p_r$  is  $\leq 6$
- number of triangles created at  $p_r$  is  $\leq 2(\text{degree}(p_r)) - 3$



# Analysis

**Lemma:** The expected number of triangles created is at most  $9n + 1 = O(n)$ .

**Proof:** How many triangles are created when  $p_r$  is inserted?



**Backwards analysis:**

- every point in  $p_1, \dots, p_r$  has the same probability  $1/r$  to be the last point
- expected degree of  $p_r$  is  $\leq 6$
- number of triangles created at  $p_r$  is  $\leq 2(\text{degree}(p_r)) - 3$
- overall  $\leq 2 \cdot 6 - 3 = 9$ ; plus 1 for the outer triangle

# Analysis

**Lemma:** The expected number of triangles created is at most  $9n + 1 = O(n)$ .

**Lemma:** The expected number of triangles which are visited in the search structure during the construction is  $O(n \log n)$ .

**Proof** in the book.

# Analysis

**Lemma:** The expected number of triangles created is at most  $9n + 1 = O(n)$ .

**Lemma:** The expected number of triangles which are visited in the search structure during the construction is  $O(n \log n)$ .

**Theorem:** The Delaunay triangulation of  $n$  points can be computed in  $O(n \log n)$  expected time using randomized incremental construction.

# Quiz

How fast can we compute the Voronoi diagram of  $n$  points?

- A:  $\Theta(n)$  expected time
- B:  $\Theta(n \log n)$  expected time
- C:  $\Theta(n^2)$  expected time

# Quiz

How fast can we compute the Voronoi diagram of  $n$  points?

A:  $\Theta(n)$  expected time

B:  $\Theta(n \log n)$  expected time

C:  $\Theta(n^2)$  expected time

We can compute the Voronoi diagram in  $O(n)$  time from the Delaunay triangulation

# Quiz

How fast can we compute the Voronoi diagram of  $n$  points?

A:  $\Theta(n)$  expected time

B:  $\Theta(n \log n)$  expected time

C:  $\Theta(n^2)$  expected time

We can compute the Voronoi diagram in  $O(n)$  time from the Delaunay triangulation

With other algorithmic paradigms (divide&conquer, sweepline) we can compute Delaunay triangulations and Voronoi diagrams also deterministically in  $\Theta(n \log n)$  time.

# Summary

**Theorem:** The Delaunay triangulation of  $n$  points can be computed in  $O(n \log n)$  expected time using randomized incremental construction.

**Theorem:** The Voronoi diagram of  $n$  points can be computed in  $O(n \log n)$  expected time.

