


# Linear Programming


## Integer Linear Programming



# Recall Example: Cutting Paper Rolls

What's the fewest number of rolls need to satisfy an order of:

- 97 rolls width 135cm
- 610 rolls width 108cm
- 395 rolls width 93cm
- 211 rolls width 42cm



Possible ways to cut roll with  $<42\text{cm}$  wasted:

P1:  $2 \cdot 135$

P2:  $135 + 108 + 42$

P3:  $135 + 93 + 42$

P4:  $135 + 3 \cdot 42$

P5:  $2 \cdot 108 + 2 \cdot 42$

P6:  $108 + 2 \cdot 93$

P7:  $108 + 93 + 2 \cdot 42$

P8:  $108 + 4 \cdot 42$

P9:  $3 \cdot 93$

P10:  $2 \cdot 93 + 2 \cdot 42$

P11:  $93 + 4 \cdot 42$

P12:  $7 \cdot 42$

# Example: Cutting Paper Rolls

For each possibility  $P_j$ , add a variable  $x_j \geq 0$  representing # rolls cut that way.

$$\text{minimize } \sum_{j=1}^{12} x_j \quad (\text{total \# of rolls cut})$$

$$\text{subject to } 2x_1 + x_2 + x_3 + x_4 \geq 97$$

$$x_2 + 2x_5 + x_6 + x_7 + x_8 \geq 610$$

$$x_3 + 2x_6 + x_7 + 3x_9 + 2x_{10} + x_{11} \geq 395$$

$$x_2 + x_3 + 3x_4 + 2x_5 + 2x_7 + 4x_8 + 2x_{10} + 4x_{11} + 7x_{12} \geq 211$$

# Example: Cutting Paper Rolls

For each possibility  $P_j$ , add a variable  $x_j \geq 0$  representing # rolls cut that way.

$$\text{minimize } \sum_{j=1}^{12} x_j \quad (\text{total \# of rolls cut})$$

$$\text{subject to } 2x_1 + x_2 + x_3 + x_4 \geq 97$$

$$x_2 + 2x_5 + x_6 + x_7 + x_8 \geq 610$$

$$x_3 + 2x_6 + x_7 + 3x_9 + 2x_{10} + x_{11} \geq 395$$

$$x_2 + x_3 + 3x_4 + 2x_5 + 2x_7 + 4x_8 + 2x_{10} + 4x_{11} + 7x_{12} \geq 211$$

Optimal solution:  $x_1 = 48.5$ ,  $x_5 = 206.25$ ,  $x_6 = 197.5$ , all others zero.

# Example: Cutting Paper Rolls

For each possibility  $P_j$ , add a variable  $x_j \geq 0$  representing # rolls cut that way.

$$\begin{aligned} &\text{minimize } \sum_{j=1}^{12} x_j \quad (\text{total \# of rolls cut}) \\ &\text{subject to } 2x_1 + x_2 + x_3 + x_4 \geq 97 \\ &\quad x_2 + 2x_5 + x_6 + x_7 + x_8 \geq 610 \\ &\quad x_3 + 2x_6 + x_7 + 3x_9 + 2x_{10} + x_{11} \geq 395 \\ &\quad x_2 + x_3 + 3x_4 + 2x_5 + 2x_7 + 4x_8 + 2x_{10} + 4x_{11} + 7x_{12} \geq 211 \end{aligned}$$

Optimal solution:  $x_1 = 48.5$ ,  $x_5 = 206.25$ ,  $x_6 = 197.5$ , all others zero.

What if we want an integer solution?

We can round up to 49, 207, 198, i.e. a total of 454.

# Example: Cutting Paper Rolls

For each possibility  $P_j$ , add a variable  $x_j \geq 0$  representing # rolls cut that way.

$$\begin{aligned} &\text{minimize } \sum_{j=1}^{12} x_j \quad (\text{total \# of rolls cut}) \\ &\text{subject to } 2x_1 + x_2 + x_3 + x_4 \geq 97 \\ &\quad x_2 + 2x_5 + x_6 + x_7 + x_8 \geq 610 \\ &\quad x_3 + 2x_6 + x_7 + 3x_9 + 2x_{10} + x_{11} \geq 395 \\ &\quad x_2 + x_3 + 3x_4 + 2x_5 + 2x_7 + 4x_8 + 2x_{10} + 4x_{11} + 7x_{12} \geq 211 \end{aligned}$$

Optimal solution:  $x_1 = 48.5$ ,  $x_5 = 206.25$ ,  $x_6 = 197.5$ , all others zero.

What if we want an integer solution?

We can round up to 49, 207, 198, i.e. a total of 454.

Not bad, since an optimal integral solution is at least 453.

# Example: Cutting Paper Rolls

For each possibility  $P_j$ , add a variable  $x_j \geq 0$  representing # rolls cut that way.

$$\begin{aligned} &\text{minimize } \sum_{j=1}^{12} x_j \quad (\text{total \# of rolls cut}) \\ &\text{subject to } 2x_1 + x_2 + x_3 + x_4 \geq 97 \\ &\quad x_2 + 2x_5 + x_6 + x_7 + x_8 \geq 610 \\ &\quad x_3 + 2x_6 + x_7 + 3x_9 + 2x_{10} + x_{11} \geq 395 \\ &\quad x_2 + x_3 + 3x_4 + 2x_5 + 2x_7 + 4x_8 + 2x_{10} + 4x_{11} + 7x_{12} \geq 211 \end{aligned}$$

Optimal solution:  $x_1 = 48.5$ ,  $x_5 = 206.25$ ,  $x_6 = 197.5$ , all others zero.


What if we want an integer solution?

We can round up to 49, 207, 198, i.e. a total of 454.

Not bad, since an optimal integral solution is at least 453.

Here, 48, 206, 197, 1 (for  $x_9$ ) is an optimal integral solution.

# Integer Linear Programming




maximize  $c^T x$

subject to  $Ax \leq b$

$x \in \mathbb{Z}^n$




# Integer Linear Programming



$$\begin{aligned} &\text{maximize } c^T x \\ &\text{subject to } Ax \leq b \\ &\quad x \in \mathbb{Z}^n \end{aligned}$$

Integer Linear Programming (ILP) is NP-hard.

# Integer Linear Programming



$$\begin{aligned} &\text{maximize } c^T x \\ &\text{subject to } Ax \leq b \\ &\quad x \in \mathbb{Z}^n \end{aligned}$$


Integer Linear Programming (ILP) is **NP-hard**.

**Variants:**

0/1-Linear Programming:  $x \in \{0, 1\}^n$ ,

Mixed Integer Linear Programming: some variables in  $\mathbb{Z}$ , some in  $\mathbb{R}$ .

# Integer Linear Programming



$$\begin{array}{ll} \text{maximize} & c^T x \\ \text{subject to} & Ax \leq b \\ & x \in \mathbb{Z}^n \end{array}$$

Integer Linear Programming (ILP) is **NP-hard**.


**Variants:**

0/1-Linear Programming:  $x \in \{0, 1\}^n$ ,

Mixed Integer Linear Programming: some variables in  $\mathbb{Z}$ , some in  $\mathbb{R}$ .

ILP and variants most wide-spread use of LPs.

# Integer Linear Programming




$$\begin{array}{ll} \text{maximize} & c^T x \\ \text{subject to} & Ax \leq b \\ & x \in \mathbb{Z}^n \end{array}$$

Next: Example “easy, medium, hard” integer programs.

- Maximum weight matching
- Minimum vertex cover
- Maximum independent set


# Maximum weight perfect matching

A company is assigning workers to jobs. In the bipartite graph an edge connects a worker to a job they are willing to take, weighted by their suitability for the job.



# Maximum weight perfect matching

A company is assigning workers to jobs. In the bipartite graph an edge connects a worker to a job they are willing to take, weighted by their suitability for the job.




How to find an optimal assignment?

# Maximum weight perfect matching

## Greedy approach


always assign largest possible score



# Maximum weight perfect matching

## Greedy approach

always assign largest possible score






# Maximum weight perfect matching

## Greedy approach


always assign largest possible score



# Maximum weight perfect matching

## Greedy approach


always assign largest possible score



# Maximum weight perfect matching

## Greedy approach


always assign largest possible score



# Maximum weight perfect matching

## Greedy approach


always assign largest possible score



# Maximum weight perfect matching

## Greedy approach


always assign largest possible score



# Maximum weight perfect matching

## Greedy approach

always assign largest possible score



total score:


$$95 + 87 + 96 + 90 + 0 + 26 + 64 = 458$$

Eleanor is not assigned to any job!

# Maximum weight perfect matching

## Greedy approach

always assign largest possible score




total score:

$$95 + 87 + 96 + 90 + 0 + 26 + 64 = 458$$

Eleanor is not assigned to any job!

## Optimal solution




total score:

$$95 + 87 + 60 + 81 + 75 + 85 + 64 = 547$$

# Maximum weight perfect matching

## Greedy approach

always assign largest possible score




total score:

$$95 + 87 + 96 + 90 + 0 + 26 + 64 = 458$$

Eleanor is not assigned to any job!

## Optimal solution



total score:

$$95 + 87 + 60 + 81 + 75 + 85 + 64 = 547$$

How to model this as an ILP?



# Maximum weight perfect matching

## Integer program

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& x_e \in \{0, 1\} \text{ for each edge } e \in E. \end{aligned}$$

variable  $x_e$  encodes  
whether edge  $e$  is  
contained in the matching

# Maximum weight perfect matching

## Integer program

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& x_e \in \{0, 1\} \text{ for each edge } e \in E. \end{aligned}$$

variable  $x_e$  encodes  
whether edge  $e$  is  
contained in the matching

if we “relax” the requirement that the  $x_e$  are integral we get . . .

# Maximum weight perfect matching

## Integer program

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& x_e \in \{0, 1\} \text{ for each edge } e \in E. \end{aligned}$$

variable  $x_e$  encodes  
whether edge  $e$  is  
contained in the matching

## LP relaxation

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& 0 \leq x_e \leq 1 \text{ for each edge } e \in E. \end{aligned}$$

# Maximum weight perfect matching

## Integer program

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& x_e \in \{0, 1\} \text{ for each edge } e \in E. \end{aligned}$$

What do we learn from solving the LP-relaxation?

## LP relaxation

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& 0 \leq x_e \leq 1 \text{ for each edge } e \in E. \end{aligned}$$

# Maximum weight perfect matching

## Integer program

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& x_e \in \{0, 1\} \text{ for each edge } e \in E. \end{aligned}$$

What do we learn from solving the LP-relaxation?

If the LP is infeasible so is the IP.

## LP relaxation

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& 0 \leq x_e \leq 1 \text{ for each edge } e \in E. \end{aligned}$$

# Maximum weight perfect matching

## Integer program

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& x_e \in \{0, 1\} \text{ for each edge } e \in E. \end{aligned}$$

What do we learn from solving the LP-relaxation?

An optimal solution to the LP relaxation is a bound on an optimal solution to the IP.

## LP relaxation

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& 0 \leq x_e \leq 1 \text{ for each edge } e \in E. \end{aligned}$$

# Maximum weight perfect matching

## Integer program

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& x_e \in \{0, 1\} \text{ for each edge } e \in E. \end{aligned}$$

## LP relaxation

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& 0 \leq x_e \leq 1 \text{ for each edge } e \in E. \end{aligned}$$

What do we learn from solving the LP-relaxation?

An optimal solution to the LP relaxation is a bound on an optimal solution to the IP.

(How) can we get an optimal integral solution from a fractional one?

# Maximum weight perfect matching

## Integer program

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& x_e \in \{0, 1\} \text{ for each edge } e \in E. \end{aligned}$$

## LP relaxation

$$\begin{aligned} &\text{maximize} && \sum_{e \in E} w_e x_e \\ &\text{subject to} && \sum_{e \in E: v \in e} x_e = 1 \text{ for each vertex } v \in V, \text{ and} \\ &&& 0 \leq x_e \leq 1 \text{ for each edge } e \in E. \end{aligned}$$

What do we learn from solving the LP-relaxation?

An optimal solution to the LP relaxation is a bound on an optimal solution to the IP.

(How) can we get an optimal integral solution from a fractional one?


Here this works out nicely.



# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).



# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

**Proof:** Let  $x^*$  be an optimal solution of the LP relaxation.

Let  $w(x^*) = \sum_{e \in E} w_e x_e^*$  its value and  $k(x^*)$  its number of non-integral components.

# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

**Proof:** Let  $x^*$  be an optimal solution of the LP relaxation.

Let  $w(x^*) = \sum_{e \in E} w_e x_e^*$  its value and  $k(x^*)$  its number of non-integral components.

If  $k(x^*) = 0$ , then we are done.

# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

**Proof:** Let  $x^*$  be an optimal solution of the LP relaxation.

Let  $w(x^*) = \sum_{e \in E} w_e x_e^*$  its value and  $k(x^*)$  its number of non-integral components.

If  $k(x^*) > 0$ , we find an optimal solution  $\tilde{x}$  with  $k(x^*) > k(\tilde{x}^*)$ .

# Maximum weight perfect matching

## Theorem 3.2.1


Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

**Proof:** Let  $x^*$  be an optimal solution of the LP relaxation.

Let  $w(x^*) = \sum_{e \in E} w_e x_e^*$  its value and  $k(x^*)$  its number of non-integral components.

If  $k(x^*) > 0$ , we find an optimal solution  $\tilde{x}$  with  $k(x^*) > k(\tilde{x}^*)$ .

Let  $x_{e_1}^*$  be a non-integral component of  $x$  for edge  $e_1 = (a_1, b_1)$ .



# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).


**Proof:** Let  $x^*$  be an optimal solution of the LP relaxation.

Let  $w(x^*) = \sum_{e \in E} w_e x_e^*$  its value and  $k(x^*)$  its number of non-integral components.

If  $k(x^*) > 0$ , we find an optimal solution  $\tilde{x}$  with  $k(x^*) > k(\tilde{x}^*)$ .

Let  $x_{e_1}^*$  be a non-integral component of  $x$  for edge  $e_1 = (a_1, b_1)$ .

Since  $0 < x_{e_1}^* < 1$  and  $\sum_{b_1 \in e} x_e^* = 1$  there exists  $e_2 = (a_2, b_1) \neq e_1$  s.t.  $x_{e_2}^*$  non-integral.



# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

**Proof:** Let  $x^*$  be an optimal solution of the LP relaxation.


Let  $w(x^*) = \sum_{e \in E} w_e x_e^*$  its value and  $k(x^*)$  its number of non-integral components.

If  $k(x^*) > 0$ , we find an optimal solution  $\tilde{x}$  with  $k(x^*) > k(\tilde{x}^*)$ .

Let  $x_{e_1}^*$  be a non-integral component of  $x$  for edge  $e_1 = (a_1, b_1)$ .

Since  $0 < x_{e_1}^* < 1$  and  $\sum_{b_1 \in e} x_e^* = 1$  there exists  $e_2 = (a_2, b_1) \neq e_1$  s.t.  $x_{e_2}^*$  non-integral.

Similarly, there exists edge  $e_3$  and so on, until we find a cycle.



# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

**Proof:** Let  $x^*$  be an optimal solution of the LP relaxation.


Let  $w(x^*) = \sum_{e \in E} w_e x_e^*$  its value and  $k(x^*)$  its number of non-integral components.

If  $k(x^*) > 0$ , we find an optimal solution  $\tilde{x}$  with  $k(x^*) > k(\tilde{x}^*)$ .

Let  $x_{e_1}^*$  be a non-integral component of  $x$  for edge  $e_1 = (a_1, b_1)$ .

Since  $0 < x_{e_1}^* < 1$  and  $\sum_{b_1 \in e} x_e^* = 1$  there exists  $e_2 = (a_2, b_1) \neq e_1$  s.t.  $x_{e_2}^*$  non-integral.

Similarly, there exists edge  $e_3$  and so on, until we find a cycle.





# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

**Proof:** Let  $x^*$  be an optimal solution of the LP relaxation.

Let  $w(x^*) = \sum_{e \in E} w_e x_e^*$  its value and  $k(x^*)$  its number of non-integral components.


If  $k(x^*) > 0$ , we find an optimal solution  $\tilde{x}$  with  $k(x^*) > k(\tilde{x}^*)$ .

Let  $x_{e_1}^*$  be a non-integral component of  $x$  for edge  $e_1 = (a_1, b_1)$ .

Since  $0 < x_{e_1}^* < 1$  and  $\sum_{b_1 \in e} x_e^* = 1$  there exists  $e_2 = (a_2, b_1) \neq e_1$  s.t.  $x_{e_2}^*$  non-integral.

Similarly, there exists edge  $e_3$  and so on, until we find a cycle.

Since  $G$  is bipartite, the cycle – say  $e_1, \dots, e_t$  – has even length  $t$ .



# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

**Proof:** Let  $x^*$  be an optimal solution of the LP relaxation.

Let  $w(x^*) = \sum_{e \in E} w_e x_e^*$  its value and  $k(x^*)$  its number of non-integral components.

If  $k(x^*) > 0$ , we find an optimal solution  $\tilde{x}$  with  $k(x^*) > k(\tilde{x}^*)$ .


Let  $x_{e_1}^*$  be a non-integral component of  $x$  for edge  $e_1 = (a_1, b_1)$ .

Since  $0 < x_{e_1}^* < 1$  and  $\sum_{b_1 \in e} x_e^* = 1$  there exists  $e_2 = (a_2, b_1) \neq e_1$  s.t.  $x_{e_2}^*$  non-integral.

Similarly, there exists edge  $e_3$  and so on, until we find a cycle.

Since  $G$  is bipartite, the cycle – say  $e_1, \dots, e_t$  – has even length  $t$ .

For small  $\varepsilon > 0$  we define

$$\tilde{x}_e = \begin{cases} x_e^* - \varepsilon & \text{for } e \in \{e_1, \dots, e_{t-1}\} \\ x_e^* + \varepsilon & \text{for } e \in \{e_2, \dots, e_t\} \\ x_e^* & \text{else} \end{cases}$$


# Maximum weight perfect matching

## Theorem 3.2.1


Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

### Proof:

It is easy to see that  $\tilde{x}$  still satisfies  $\sum_{e:v \in e} \tilde{x}_e = 1$  for all  $v \in V$ .

---

For small  $\varepsilon > 0$  we define

$$\tilde{x}_e = \begin{cases} x_e^* - \varepsilon & \text{for } e \in \{e_1, \dots, e_{t-1}\} \\ x_e^* + \varepsilon & \text{for } e \in \{e_2, \dots, e_t\} \\ x_e^* & \text{else} \end{cases}$$


# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).


### Proof:

It is easy to see that  $\tilde{x}$  still satisfies  $\sum_{e:v \in e} \tilde{x}_e = 1$  for all  $v \in V$ .

For sufficiently small  $\varepsilon$  also  $0 \leq \tilde{x}_e \leq 1$  for all  $e \in E$ , hence  $\tilde{x}$  is feasible.

---

For small  $\varepsilon > 0$  we define

$$\tilde{x}_e = \begin{cases} x_e^* - \varepsilon & \text{for } e \in \{e_1, \dots, e_{t-1}\} \\ x_e^* + \varepsilon & \text{for } e \in \{e_2, \dots, e_t\} \\ x_e^* & \text{else} \end{cases}$$


# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

### Proof:


It is easy to see that  $\tilde{x}$  still satisfies  $\sum_{e:v \in e} \tilde{x}_e = 1$  for all  $v \in V$ .

For sufficiently small  $\varepsilon$  also  $0 \leq \tilde{x}_e \leq 1$  for all  $e \in E$ , hence  $\tilde{x}$  is feasible.

What about  $w(\tilde{x})$ ?

---

For small  $\varepsilon > 0$  we define

$$\tilde{x}_e = \begin{cases} x_e^* - \varepsilon & \text{for } e \in \{e_1, \dots, e_{t-1}\} \\ x_e^* + \varepsilon & \text{for } e \in \{e_2, \dots, e_t\} \\ x_e^* & \text{else} \end{cases}$$


# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

### Proof:


It is easy to see that  $\tilde{x}$  still satisfies  $\sum_{e:v \in e} \tilde{x}_e = 1$  for all  $v \in V$ .

For sufficiently small  $\varepsilon$  also  $0 \leq \tilde{x}_e \leq 1$  for all  $e \in E$ , hence  $\tilde{x}$  is feasible.

We have  $w(\tilde{x}) = \sum_{e \in E} w_e \tilde{x}_e = w(x^*) + \varepsilon \underbrace{\sum_{i=1}^t (-1)^i w_{e_i}}_{=: \Delta} = w(x^*) + \varepsilon \Delta$

---

For small  $\varepsilon > 0$  we define

$$\tilde{x}_e = \begin{cases} x_e^* - \varepsilon & \text{for } e \in \{e_1, \dots, e_{t-1}\} \\ x_e^* + \varepsilon & \text{for } e \in \{e_2, \dots, e_t\} \\ x_e^* & \text{else} \end{cases}$$


# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

### Proof:


It is easy to see that  $\tilde{x}$  still satisfies  $\sum_{e:v \in e} \tilde{x}_e = 1$  for all  $v \in V$ .

For sufficiently small  $\varepsilon$  also  $0 \leq \tilde{x}_e \leq 1$  for all  $e \in E$ , hence  $\tilde{x}$  is feasible.

We have  $w(\tilde{x}) = \sum_{e \in E} w_e \tilde{x}_e = w(x^*) + \varepsilon \underbrace{\sum_{i=1}^t (-1)^i w_{e_i}}_{=: \Delta = 0, \text{ else optimality of } x^* \text{ violated}} = w(x^*) + \varepsilon \Delta$

---

For small  $\varepsilon > 0$  we define

$$\tilde{x}_e = \begin{cases} x_e^* - \varepsilon & \text{for } e \in \{e_1, \dots, e_{t-1}\} \\ x_e^* + \varepsilon & \text{for } e \in \{e_2, \dots, e_t\} \\ x_e^* & \text{else} \end{cases}$$


# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

### Proof:


It is easy to see that  $\tilde{x}$  still satisfies  $\sum_{e:v \in e} \tilde{x}_e = 1$  for all  $v \in V$ .

For sufficiently small  $\varepsilon$  also  $0 \leq \tilde{x}_e \leq 1$  for all  $e \in E$ , hence  $\tilde{x}$  is feasible.

We have  $w(\tilde{x}) = \sum_{e \in E} w_e \tilde{x}_e = w(x^*) + \varepsilon \underbrace{\sum_{i=1}^t (-1)^i w_{e_i}}_{=: \Delta = 0, \text{ else optimality of } x^* \text{ violated}} = w(x^*) + \varepsilon \Delta = w(x^*)$

---

For small  $\varepsilon > 0$  we define

$$\tilde{x}_e = \begin{cases} x_e^* - \varepsilon & \text{for } e \in \{e_1, \dots, e_{t-1}\} \\ x_e^* + \varepsilon & \text{for } e \in \{e_2, \dots, e_t\} \\ x_e^* & \text{else} \end{cases}$$




# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

### Proof:

It is easy to see that  $\tilde{x}$  still satisfies  $\sum_{e:v \in e} \tilde{x}_e = 1$  for all  $v \in V$ .


For sufficiently small  $\varepsilon$  also  $0 \leq \tilde{x}_e \leq 1$  for all  $e \in E$ , hence  $\tilde{x}$  is feasible.

We have  $w(\tilde{x}) = \sum_{e \in E} w_e \tilde{x}_e = w(x^*) + \varepsilon \sum_{i=1}^t (-1)^i w_{e_i} = w(x^*) + \varepsilon \Delta = w(x^*)$

How do we choose  $\varepsilon$ ?

---

For small  $\varepsilon > 0$  we define

$$\tilde{x}_e = \begin{cases} x_e^* - \varepsilon & \text{for } e \in \{e_1, \dots, e_{t-1}\} \\ x_e^* + \varepsilon & \text{for } e \in \{e_2, \dots, e_t\} \\ x_e^* & \text{else} \end{cases}$$


# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

### Proof:

It is easy to see that  $\tilde{x}$  still satisfies  $\sum_{e:v \in e} \tilde{x}_e = 1$  for all  $v \in V$ .


For sufficiently small  $\varepsilon$  also  $0 \leq \tilde{x}_e \leq 1$  for all  $e \in E$ , hence  $\tilde{x}$  is feasible.

We have  $w(\tilde{x}) = \sum_{e \in E} w_e \tilde{x}_e = w(x^*) + \varepsilon \sum_{i=1}^t (-1)^i w_{e_i} = w(x^*) + \varepsilon \Delta = w(x^*)$

We choose  $\varepsilon$  largest s.t.  $\tilde{x}$  is still feasible. Then  $k(\tilde{x}) < k(x^*)$ .

---

For small  $\varepsilon > 0$  we define

$$\tilde{x}_e = \begin{cases} x_e^* - \varepsilon & \text{for } e \in \{e_1, \dots, e_{t-1}\} \\ x_e^* + \varepsilon & \text{for } e \in \{e_2, \dots, e_t\} \\ x_e^* & \text{else} \end{cases}$$


# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).

### Proof:

It is easy to see that  $\tilde{x}$  still satisfies  $\sum_{e:v \in e} \tilde{x}_e = 1$  for all  $v \in V$ .

For sufficiently small  $\varepsilon$  also  $0 \leq \tilde{x}_e \leq 1$  for all  $e \in E$ , hence  $\tilde{x}$  is feasible.


We have  $w(\tilde{x}) = \sum_{e \in E} w_e \tilde{x}_e = w(x^*) + \varepsilon \sum_{i=1}^t (-1)^i w_{e_i} = w(x^*) + \varepsilon \Delta = w(x^*)$

We choose  $\varepsilon$  largest s.t.  $\tilde{x}$  is still feasible. Then  $k(\tilde{x}) < k(x^*)$ .

We continue this procedure until all components are integral.

---

For small  $\varepsilon > 0$  we define

$$\tilde{x}_e = \begin{cases} x_e^* - \varepsilon & \text{for } e \in \{e_1, \dots, e_{t-1}\} \\ x_e^* + \varepsilon & \text{for } e \in \{e_2, \dots, e_t\} \\ x_e^* & \text{else} \end{cases}$$



# Maximum weight perfect matching

## Theorem 3.2.1

Let  $G$  be an arbitrary weighted bipartite graph. The LP relaxation from before has an integer optimal solution (which also solves the integer program).


**Moral:** Sometimes solving an integer program is no harder than solving a linear program.

# Minimum vertex cover




**Recall:** a minimum **vertex cover**, is a smallest possible subset  $V' \subseteq V$  such that for every edge  $uv \in E$ , it holds that  $u \in V'$  or  $v \in V'$ .

# Minimum vertex cover



**Recall:** a minimum **vertex cover**, is a smallest possible subset  $V' \subseteq V$  such that for every edge  $uv \in E$ , it holds that  $u \in V'$  or  $v \in V'$ .


# Minimum vertex cover



**Recall:** a minimum **vertex cover**, is a smallest possible subset  $V' \subseteq V$  such that for every edge  $uv \in E$ , it holds that  $u \in V'$  or  $v \in V'$ .

## How can we formulate this as an ILP?

# Minimum vertex cover



$$\text{minimize } \sum_{v \in V} x_v$$

subject to  $x_u + x_v \geq 1$  for every edge  $\{u, v\} \in E$


$$x_v \in \{0, 1\} \text{ for all } v \in V.$$

variable  $x_v$  encodes whether vertex  $v$  is contained in the cover



# Minimum vertex cover

$G = (V, E)$



minimize  $\sum_{v \in V} x_v$


subject to  $x_u + x_v \geq 1$  for every edge  $\{u, v\} \in E$

$x_v \in \{0, 1\}$  for all  $v \in V$ .

variable  $x_v$  encodes whether vertex  $v$  is contained in the cover

LP relaxation:  $0 \leq x_v \leq 1$

# Minimum vertex cover



$$\text{minimize } \sum_{v \in V} x_v$$

subject to  $x_u + x_v \geq 1$  for every edge  $\{u, v\} \in E$

$$x_v \in \{0, 1\} \text{ for all } v \in V.$$

variable  $x_v$  encodes whether vertex  $v$  is contained in the cover

LP relaxation:  $0 \leq x_v \leq 1$  How does this help?

# Minimum vertex cover

## Rounding:

Let  $S_{\text{IP}} \subseteq V$  be a vertex cover solving IP.

Let  $S_{\text{LP}} \subseteq V$  be a vertex cover solving LP,  
obtained by  $S_{\text{LP}} = \{v \in V \mid x_v \geq \frac{1}{2}\}$ .

$$\text{minimize } \sum_{v \in V} x_v$$

$$\text{subject to } x_u + x_v \geq 1 \text{ for all } \{u, v\} \in E$$
$$x_v \in [0, 1] \text{ for all } v \in V.$$

# Minimum vertex cover

## Rounding:

Let  $S_{\text{IP}} \subseteq V$  be a vertex cover solving IP.


Let  $S_{\text{LP}} \subseteq V$  be a vertex cover solving LP,  
obtained by  $S_{\text{LP}} = \{v \in V \mid x_v \geq \frac{1}{2}\}$ .

What solution do we get here?

$$\text{minimize } \sum_{v \in V} x_v$$

subject to  $x_u + x_v \geq 1$  for all  $\{u, v\} \in E$

$$x_v \in [0, 1] \text{ for all } v \in V.$$



# Minimum vertex cover


## Rounding:

Let  $S_{\text{IP}} \subseteq V$  be a vertex cover solving IP.  
Let  $S_{\text{LP}} \subseteq V$  be a vertex cover solving LP,  
obtained by  $S_{\text{LP}} = \{v \in V \mid x_v \geq \frac{1}{2}\}$ .

$$\begin{aligned} &\text{minimize } \sum_{v \in V} x_v \\ &\text{subject to } x_u + x_v \geq 1 \text{ for all } \{u, v\} \in E \\ &\quad x_v \in [0, 1] \text{ for all } v \in V. \end{aligned}$$

What solution do we get here?

solution to LP, with cost  $11/2$



# Minimum vertex cover


## Rounding:

Let  $S_{\text{IP}} \subseteq V$  be a vertex cover solving IP.  
Let  $S_{\text{LP}} \subseteq V$  be a vertex cover solving LP,  
obtained by  $S_{\text{LP}} = \{v \in V \mid x_v \geq \frac{1}{2}\}$ .

$$\begin{aligned} &\text{minimize } \sum_{v \in V} x_v \\ &\text{subject to } x_u + x_v \geq 1 \text{ for all } \{u, v\} \in E \\ &\quad x_v \in [0, 1] \text{ for all } v \in V. \end{aligned}$$

What solution do we get here?

solution to LP, with cost  $11/2$



solution to IP, with cost 11

# Minimum vertex cover


## Rounding:

Let  $S_{\text{IP}} \subseteq V$  be a vertex cover solving IP.

Let  $S_{\text{LP}} \subseteq V$  be a vertex cover solving LP,  
obtained by  $S_{\text{LP}} = \{v \in V \mid x_v \geq \frac{1}{2}\}$ .

What solution do we get here?

$$\begin{aligned} &\text{minimize } \sum_{v \in V} x_v \\ &\text{subject to } x_u + x_v \geq 1 \text{ for all } \{u, v\} \in E \\ &\quad x_v \in [0, 1] \text{ for all } v \in V. \end{aligned}$$




# Minimum vertex cover

## Rounding:

Let  $S_{\text{IP}} \subseteq V$  be a vertex cover solving IP.  
Let  $S_{\text{LP}} \subseteq V$  be a vertex cover solving LP,  
obtained by  $S_{\text{LP}} = \{v \in V \mid x_v \geq \frac{1}{2}\}$ .

What solution do we get here?

solution to LP also to IP, with cost 1



$$\text{minimize } \sum_{v \in V} x_v$$

$$\text{subject to } x_u + x_v \geq 1 \text{ for all } \{u, v\} \in E$$
$$x_v \in [0, 1] \text{ for all } v \in V.$$



# Minimum vertex cover

## Rounding:

Let  $S_{\text{IP}} \subseteq V$  be a vertex cover solving IP.

Let  $S_{\text{LP}} \subseteq V$  be a vertex cover solving LP,  
obtained by  $S_{\text{LP}} = \{v \in V \mid x_v \geq \frac{1}{2}\}$ .

**Fact:**  $|S_{\text{IP}}| \leq |S_{\text{LP}}| \leq 2|S_{\text{IP}}|$

$$\text{minimize } \sum_{v \in V} x_v$$

subject to  $x_u + x_v \geq 1$  for all  $\{u, v\} \in E$   
 $x_v \in [0, 1]$  for all  $v \in V$ .

# Minimum vertex cover


## Rounding:


Let  $S_{\text{IP}} \subseteq V$  be a vertex cover solving IP.  
Let  $S_{\text{LP}} \subseteq V$  be a vertex cover solving LP,  
obtained by  $S_{\text{LP}} = \{v \in V \mid x_v \geq \frac{1}{2}\}$ .

**Fact:**  $|S_{\text{IP}}| \leq |S_{\text{LP}}| \leq 2|S_{\text{IP}}|$

## Proof of second inequality

$$|S_{\text{LP}}| \leq 2 \sum_{v \in V} x_v^* \leq 2 \sum_{v \in V} \tilde{x}_v = 2 \cdot |S_{\text{IP}}|$$

 by definition of  $S_{\text{LP}}$

 since any solution to IP is also a feasible solution for the LP relaxation.

$$\begin{aligned} &\text{minimize } \sum_{v \in V} x_v \\ &\text{subject to } x_u + x_v \geq 1 \text{ for all } \{u, v\} \in E \\ &\quad x_v \in [0, 1] \text{ for all } v \in V. \end{aligned}$$

$x_v^*$ : optimal LP solution,  
 $\tilde{x}_v$ : optimal ILP solution

# Minimum vertex cover

## Rounding:

Let  $S_{\text{IP}} \subseteq V$  be a vertex cover solving IP.  
Let  $S_{\text{LP}} \subseteq V$  be a vertex cover solving LP,  
obtained by  $S_{\text{LP}} = \{v \in V \mid x_v \geq \frac{1}{2}\}$ .

**Fact:**  $|S_{\text{IP}}| \leq |S_{\text{LP}}| \leq 2|S_{\text{IP}}|$

## Proof of second inequality

$$|S_{\text{LP}}| \leq 2 \sum_{v \in V} x_v^* \leq 2 \sum_{v \in V} \tilde{x}_v = 2 \cdot |S_{\text{IP}}|$$

by definition of  $S_{\text{LP}}$


since any solution to IP is also a feasible solution for the LP relaxation.

$$\begin{aligned} &\text{minimize } \sum_{v \in V} x_v \\ &\text{subject to } x_u + x_v \geq 1 \text{ for all } \{u, v\} \in E \\ &\quad x_v \in [0, 1] \text{ for all } v \in V. \end{aligned}$$

$x_v^*$ : optimal LP solution,  
 $\tilde{x}_v$ : optimal ILP solution

**Moral:** Sometimes solving an LP relaxation gives an approximate solution to an **NP**-hard integer program.


# Maximum independent set



**Recall:** a maximum **independent**, is a largest possible subset  $V' \subseteq V$  such that for any  $u, v \in V'$ , it holds that  $(u, v) \notin E$ .

How can we formulate this as an ILP?

# Maximum independent set




$$\text{maximize } \sum_{v \in V} x_v$$

subject to  $x_u + x_v \leq 1$  for each edge  $\{u, v\} \in E$

$$x_v \in \{0, 1\} \text{ for all } v \in V.$$

variable  $x_v$  encodes whether  
vertex  $v$  is contained in the set

# Maximum independent set



$$\text{maximize } \sum_{v \in V} x_v$$

subject to  $x_u + x_v \leq 1$  for each edge  $\{u, v\} \in E$

$$x_v \in \{0, 1\} \text{ for all } v \in V.$$

variable  $x_v$  encodes whether vertex  $v$  is contained in the set


LP relaxation:  $0 \leq x_v \leq 1$

# Maximum independent set

In the LP relaxation ( $0 \leq x_v \leq 1$ ) we always have the feasible solution  $x_v = \frac{1}{2}$  for all  $v$ , meaning the optimum is at least  $\frac{1}{2}|V|$ .

# Maximum independent set

In the LP relaxation ( $0 \leq x_v \leq 1$ ) we always have the feasible solution  $x_v = \frac{1}{2}$  for all  $v$ , meaning the optimum is at least  $\frac{1}{2}|V|$ .






# Maximum independent set

In the LP relaxation ( $0 \leq x_v \leq 1$ ) we always have the feasible solution  $x_v = \frac{1}{2}$  for all  $v$ , meaning the optimum is at least  $\frac{1}{2}|V|$ .


In a complete graph, what are the optimum LP and IP solution?



# Maximum independent set

In the LP relaxation ( $0 \leq x_v \leq 1$ ) we always have the feasible solution  $x_v = \frac{1}{2}$  for all  $v$ , meaning the optimum is at least  $\frac{1}{2}|V|$ .

In a complete graph, the LP has optimum  $n/2$ , and the IP optimum 1.



# Maximum independent set

In the LP relaxation ( $0 \leq x_v \leq 1$ ) we always have the feasible solution  $x_v = \frac{1}{2}$  for all  $v$ , meaning the optimum is at least  $\frac{1}{2}|V|$ .

In a complete graph, the LP has optimum  $n/2$ , and the IP optimum 1.

**Moral:** Sometimes an LP relaxation tells us next to nothing about the integer program.

# Maximum independent set

In the LP relaxation ( $0 \leq x_v \leq 1$ ) we always have the feasible solution  $x_v = \frac{1}{2}$  for all  $v$ , meaning the optimum is at least  $\frac{1}{2}|V|$ .

In a complete graph, the LP has optimum  $n/2$ , and the IP optimum 1.

**Moral:** Sometimes an LP relaxation tells us next to nothing about the integer program.

Approximation for this problem is known to be hard:

J. Håstad: Clique is hard to approximate within  $n^{1-\epsilon}$ ,  
Acta Math. 182(1999): 105-142


# Bonus Examples

Shortest Path and TSP

# Shortest Path Problem

Given a directed graph  $G = (V, E)$  with edge weights  $w$ , we are looking for a shortest path from  $s$  to  $t$ .


Can we also model this as ILP?



# Shortest Path Problem

Given a directed graph  $G = (V, E)$  with edge weights  $w$ , we are looking for a shortest path from  $s$  to  $t$ .

Can we also model this as ILP? Yes! How?




# Shortest Path Problem

Given a directed graph  $G = (V, E)$  with edge weights  $w$ , we are looking for a shortest path from  $s$  to  $t$ .

Can we also model this as ILP? Yes! How?


Idea 1: use variable  $x_{uv}$  for whether edge  $(u, v)$  is used.





# Shortest Path Problem

Given a directed graph  $G = (V, E)$  with edge weights  $w$ , we are looking for a shortest path from  $s$  to  $t$ .




Can we also model this as ILP? Yes! How?

**Idea 1:** use variable  $x_{uv}$  for whether edge  $(u, v)$  is used.

$$\begin{aligned} &\text{minimize} && \sum_{(u,v) \in E} w(u,v)x_{uv} \\ &\text{subject to} && \sum_{(u,v) \in E} x_{uv} = \sum_{(v,w) \in E} x_{vw} \text{ for each vertex } v \in V \setminus \{s, t\}, \text{ and} \\ &&& \sum_{(u,t) \in E} x_{ut} = 1. \\ &&& x_{uv} \in \{0, 1\} \text{ for each edge } (u, v) \in E. \end{aligned}$$

# Shortest Path Problem

Given a directed graph  $G = (V, E)$  with edge weights  $w$ , we are looking for a shortest path from  $s$  to  $t$ .



Can we also model this as ILP? Yes! How?

Idea 1: use variable  $x_{uv}$  for whether edge  $(u, v)$  is used.

$$\begin{aligned} &\text{minimize} && \sum_{(u,v) \in E} w(u,v) x_{uv} \\ &\text{subject to} && \sum_{(u,v) \in E} x_{uv} = \sum_{(v,w) \in E} x_{vw} \text{ for each vertex } v \in V \setminus \{s, t\}, \text{ and} \\ &&& \sum_{(u,t) \in E} x_{ut} = 1. \\ &&& x_{uv} \in \{0, 1\} \text{ for each edge } (u, v) \in E. \end{aligned}$$

LP-relaxation has  $\{0, 1\}$ -solution,  
like MinWeight Perfect Matching


# Shortest Path Problem

Given a directed graph  $G = (V, E)$  with edge weights  $w$ , we are looking for a shortest path from  $s$  to  $t$ .

Can we also model this as ILP? Yes! How?


Idea 2: use variable  $d_v$  for distance from  $s$  to  $v$ .

Then we want that  $d_v \leq d_u + w(u, v)$  for all edges  $(u, v)$ .



# Shortest Path Problem

Given a directed graph  $G = (V, E)$  with edge weights  $w$ , we are looking for a shortest path from  $s$  to  $t$ .



Can we also model this as ILP? Yes! How?

**Idea 2:** use variable  $d_v$  for distance from  $s$  to  $v$ .

Then we want that  $d_v \leq d_u + w(u, v)$  for all edges  $(u, v)$ .

maximize  $d_t$

subject to


$$d_v - d_u \leq w(u, v) \quad \forall (u, v) \in E$$

$$d_s = 0$$

Actually an LP!

# Shortest Path Problem

Given a directed graph  $G = (V, E)$  with edge weights  $w$ , we are looking for a shortest path from  $s$  to  $t$ .



Can we also model this as ILP? Yes! How?

**Idea 2:** use variable  $d_v$  for distance from  $s$  to  $v$ .

Then we want that  $d_v \leq d_u + w(u, v)$  for all edges  $(u, v)$ .

maximize  $d_t$

subject to

$$d_v - d_u \leq w(u, v) \quad \forall (u, v) \in E$$


$$d_s = 0$$

Why do we maximize?

Actually an LP!

# Shortest Path Problem

Given a directed graph  $G = (V, E)$  with edge weights  $w$ , we are looking for a shortest path from  $s$  to  $t$ .



Can we also model this as ILP? Yes! How?

**Idea 2:** use variable  $d_v$  for distance from  $s$  to  $v$ .

Then we want that  $d_v \leq d_u + w(u, v)$  for all edges  $(u, v)$ .

maximize  $d_t$

subject to

$$d_v - d_u \leq w(u, v) \quad \forall (u, v) \in E$$

$$d_s = 0$$


Why do we maximize?

otherwise we could  
set all  $d_v$  to zero

Actually an LP!

# Shortest Path Problem

Given a directed graph  $G = (V, E)$  with edge weights  $w$ , we are looking for a shortest path from  $s$  to  $t$ .



Can we also model this as ILP? Yes! How?

**Idea 2:** use variable  $d_v$  for distance from  $s$  to  $v$ .

Then we want that  $d_v \leq d_u + w(u, v)$  for all edges  $(u, v)$ .

maximize  $d_t$

subject to

$$d_v - d_u \leq w(u, v) \quad \forall (u, v) \in E$$

$$d_s = 0$$

Why do we maximize?


otherwise we could  
set all  $d_v$  to zero

simulating a  $\min(\cdot)$   
by  $\leq$  and maximize

Actually an LP!

# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .






# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .


Can we also model this as ILP?



# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .

Can we also model this as ILP? Yes! How?




# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .

Can we also model this as ILP? Yes! How?

Idea: use variable  $x_{ij}$  for edge from city  $i$  to city  $j$ .



# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .

Can we also model this as ILP? Yes! How?

Idea: use variable  $x_{ij}$  for edge from city  $i$  to city  $j$ .


$$\text{minimize } \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq n} x_{ij} c(i, j)$$

subject to

$$\sum_{1 \leq j \leq n} x_{ij} = 1 \text{ for } i = 1, \dots, n$$

$$\sum_{1 \leq i \leq n} x_{ij} = 1 \text{ for } j = 1, \dots, n$$

$$x_{ij} \in \{0, 1\} \quad \text{for } i, j = 1, \dots, n$$



# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .

Can we also model this as ILP? Yes! How?

Idea: use variable  $x_{ij}$  for edge from city  $i$  to city  $j$ .


$$\text{minimize } \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq n} x_{ij} c(i, j)$$

subject to

$$\sum_{1 \leq j \leq n} x_{ij} = 1 \text{ for } i = 1, \dots, n$$

$$\sum_{1 \leq i \leq n} x_{ij} = 1 \text{ for } j = 1, \dots, n$$

$$x_{ij} \in \{0, 1\} \quad \text{for } i, j = 1, \dots, n$$



Does this correctly solve TSP?

# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .

Can we also model this as ILP? Yes! How?

Idea: use variable  $x_{ij}$  for edge from city  $i$  to city  $j$ .


$$\text{minimize } \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq n} x_{ij} c(i, j)$$

subject to

$$\sum_{1 \leq j \leq n} x_{ij} = 1 \text{ for } i = 1, \dots, n$$

$$\sum_{1 \leq i \leq n} x_{ij} = 1 \text{ for } j = 1, \dots, n$$

$$x_{ij} \in \{0, 1\} \quad \text{for } i, j = 1, \dots, n$$



Does this correctly solve TSP? No!

A tour can “decompose” into many smaller tours.

# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .

Can we also model this as ILP? Yes! How?

Idea: use variable  $x_{ij}$  for edge from city  $i$  to city  $j$ .


minimize  $\sum_{1 \leq i \leq n} \sum_{1 \leq j \leq n} x_{ij} c(i, j)$

subject to

$\sum_{1 \leq j \leq n} x_{ij} = 1$  for  $i = 1, \dots, n$

$\sum_{1 \leq i \leq n} x_{ij} = 1$  for  $j = 1, \dots, n$

$x_{ij} \in \{0, 1\}$  for  $i, j = 1, \dots, n$



Does this correctly solve TSP? No!

A tour can “decompose” into many smaller tours.

How to avoid this?

# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .

Can we also model this as ILP? Yes! How?

Idea: use variable  $x_{ij}$  for edge from city  $i$  to city  $j$ .


$$\text{minimize } \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq n} x_{ij} c(i, j)$$

subject to

$$\sum_{1 \leq j \leq n} x_{ij} = 1 \text{ for } i = 1, \dots, n$$

$$\sum_{1 \leq i \leq n} x_{ij} = 1 \text{ for } j = 1, \dots, n$$

$$x_{ij} \in \{0, 1\} \quad \text{for } i, j = 1, \dots, n$$



Does this correctly solve TSP? No!

A tour can “decompose” into many smaller tours.

How to avoid this?

Use variables  $u_j \hat{=}$  position of city  $j$  on tour

Require:  $u_j \geq u_i + 1$  if  $x_{ij} = 1$



# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .

Can we also model this as ILP? Yes! How?

Idea: use variable  $x_{ij}$  for edge from city  $i$  to city  $j$ .


minimize  $\sum_{1 \leq i \leq n} \sum_{1 \leq j \leq n} x_{ij} c(i, j)$

subject to

$\sum_{1 \leq j \leq n} x_{ij} = 1$  for  $i = 1, \dots, n$

$\sum_{1 \leq i \leq n} x_{ij} = 1$  for  $j = 1, \dots, n$

$x_{ij} \in \{0, 1\}$  for  $i, j = 1, \dots, n$



Does this correctly solve TSP? No!

A tour can “decompose” into many smaller tours.

How to avoid this?

Use variables  $u_j \hat{=}$  position of city  $j$  on tour

Require:  $u_j \geq u_i + 1$  if  $x_{ij} = 1$

by  $u_j + (n - 1) \geq u_i + nx_{ij}$

for all  $i \neq j, j > 1$

# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .

Can we also model this as ILP? Yes! How?

Idea: use variable  $x_{ij}$  for edge from city  $i$  to city  $j$ .

$$\text{minimize } \sum_{1 \leq i \leq n} \sum_{1 \leq j \leq n} x_{ij} c(i, j)$$


subject to

$$\sum_{1 \leq j \leq n} x_{ij} = 1 \text{ for } i = 1, \dots, n$$

$$\sum_{1 \leq i \leq n} x_{ij} = 1 \text{ for } j = 1, \dots, n$$

$$u_i - u_j + nx_{ij} \leq n - 1 \text{ for } i \neq j, j > 1$$

$$x_{ij} \in \{0, 1\}, 1 \leq u_i \leq n \text{ for } i, j = 1, \dots, n$$



Miller-Tucker-Zemlin formulation

# Traveling Salesperson Tour

Given an undirected complete graph  $G = (V, E)$  with edge weights  $c: E \rightarrow \mathbb{R}$ , find a shortest Hamilton circuit in  $G$ .

Can we also model this as ILP? Yes! How?

Idea: use variable  $x_{ij}$  for edge from city  $i$  to city  $j$ .

minimize  $\sum_{1 \leq i \leq n} \sum_{1 \leq j \leq n} x_{ij} c(i, j)$


subject to

$$\sum_{1 \leq j \leq n} x_{ij} = 1 \text{ for } i = 1, \dots, n$$

$$\sum_{1 \leq i \leq n} x_{ij} = 1 \text{ for } j = 1, \dots, n$$

$$\sum_{i \neq j : i, j \in Q} x_{ij} \leq |Q| - 1$$

$$\forall Q \subsetneq \{1, \dots, n\}, |Q| \geq 2$$




Miller-Tucker-Zemlin formulation

alternative:

Dantzig-Fulkerson-Johnson formulation  
with subtour-elimination constraint

# Solving Integer Linear Programs


How can we solve an ILP?



# Solving Integer Linear Programs

How can we solve an ILP?

Optimal integer solutions may be arbitrary far from relaxed LP solutions




# Solving Integer Linear Programs

## How can we solve an ILP?

Optimal integer solutions may be arbitrary far from relaxed LP solutions


## Techniques

- Branch-and-Bound
- Cutting Planes
- Branch-and-Cut



# Branch and Bound for Solving ILP

**Idea:** branch: decompose in two subproblems  
bound: discard if possible subproblems




# Branch and Bound for Solving ILP

**Idea:** branch: decompose in two subproblems  
bound: discard if possible subproblems

## Sketch

- solve relaxed problem
- if solution non integer, choose variable  $x_i$  with non-integer value  $\alpha_i$  and split into

problem  $P_1$ :  $P$  with  $x_i \leq \lfloor \alpha_i \rfloor$       problem  $P_2$ :  $P$  with  $x_i \geq \lceil \alpha_i \rceil$






# Branch and Bound for Solving ILP

**Idea:** branch: decompose in two subproblems  
bound: discard if possible subproblems

## Sketch

- solve relaxed problem
- if solution non integer, choose variable  $x_i$  with non-integer value  $\alpha_i$  and split into

problem  $P_1$ :  $P$  with  $x_i \leq \lfloor \alpha_i \rfloor$       problem  $P_2$ :  $P$  with  $x_i \geq \lceil \alpha_i \rceil$




# Branch and Bound for Solving ILP

**Idea:** branch: decompose in two subproblems  
bound: discard if possible subproblems

## Sketch

- solve relaxed problem
- if solution non integer, choose variable  $x_i$  with non-integer value  $\alpha_i$  and split into  
problem  $P_1$ :  $P$  with  $x_i \leq \lfloor \alpha_i \rfloor$       problem  $P_2$ :  $P$  with  $x_i \geq \lceil \alpha_i \rceil$




# Branch and Bound for Solving ILP

**Idea:** branch: decompose in two subproblems  
bound: discard if possible subproblems

## Sketch

- solve relaxed problem
- if solution non integer, choose variable  $x_i$  with non-integer value  $\alpha_i$  and split into  
problem  $P_1$ :  $P$  with  $x_i \leq \lfloor \alpha_i \rfloor$       problem  $P_2$ :  $P$  with  $x_i \geq \lceil \alpha_i \rceil$




three possibilities for subproblems:

- LP infeasible  $\rightarrow$  discard branch
- solution integer  $\rightarrow$  update OPT and discard branch
- solution non integer  $\rightarrow$  stop if solution worse than OPT, else continue branching

# Cutting Planes for Solving ILPs

**Idea:** solve relaxed LP and cut off non-integer solutions found




# Cutting Planes for Solving ILPs

**Idea:** solve relaxed LP and cut off non-integer solutions found

## Sketch

- solve relaxed problem
- if solution integer, return with this
- find a cut, which separates the found solution from all integer solutions
- add the cut as constraint and restart




# Cutting Planes for Solving ILPs

**Idea:** solve relaxed LP and cut off non-integer solutions found

## Sketch

- solve relaxed problem
- if solution integer, return with this
- find a cut, which separates the found solution from all integer solutions
- add the cut as constraint and restart

How to find a cut?



# Cutting Planes for Solving ILPs


**Idea:** solve relaxed LP and cut off non-integer solutions found

## Sketch

- solve relaxed problem
- if solution integer, return with this
- find a cut, which separates the found solution from all integer solutions
- add the cut as constraint and restart

## How to find a cut?

general-purpose cuts, e.g. Gomory Cuts  
problem-specific cuts




# Cutting Planes for Solving ILPs

**Idea:** solve relaxed LP and cut off non-integer solutions found

## Sketch

- solve relaxed problem
- if solution integer, return with this
- find a cut, which separates the found solution from all integer solutions
- add the cut as constraint and restart



## How to find a cut?

general-purpose cuts, e.g. Gomory Cuts  
problem-specific cuts

Even better is to combine both techniques:

## Branch & Cut



# Summary

An integer linear programm (IP) is of the form

Solving IPs in general is NP-hard

We formulated as IP

- Maximum weight matching
- Minimum vertex cover
- Maximum independent set
- Shortest path
- Traveling Salesperson Tour

Techniques: branch&bound, cutting planes, branch&cut

$$\begin{aligned} &\text{maximize } c^T x \\ &\text{subject to } Ax \leq b \\ &\quad x \in \mathbb{Z}^n \end{aligned}$$

# Summary

An integer linear programm (IP) is of the form

Solving IPs in general is NP-hard

We formulated as IP

- Maximum weight matching
- Minimum vertex cover
- Maximum independent set
- Shortest path
- Traveling Salesperson Tour

Techniques: branch&bound, cutting planes, branch&cut

Next steps:

- basics for solving LPs
- simplex algorithm

$$\begin{aligned} &\text{maximize } c^T x \\ &\text{subject to } Ax \leq b \\ &\quad x \in \mathbb{Z}^n \end{aligned}$$