

## Investigating a Unified Model for Feedback in Cool Core Clusters

### Abstract

Cluster mass functions and the evolution of the cluster mass function are useful for measuring cosmological parameters ([1], [2], [3], [4], [5]). Cluster evolution tests the effect of dark matter and dark energy on the evolution of dark matter halos, and therefore provide a complementary and distinct constraint on cosmological parameters to those tests which constrain them geometrically (e.g. supernovae ([6], [7]) and baryon acoustic oscillations ([8])).

However, clusters are a useful cosmological tool only if we can infer cluster masses – the fundamental cluster property inferred from cosmological simulations ([9]) – from observable properties such as X-ray luminosity, X-ray temperature, lensing shear, optical luminosity, and galaxy velocity dispersion. Empirically, the relationship of mass and these observable properties is well-established ([10]). However, if we could identify a “3rd parameter” – possibly reflecting the degree of relaxation in the cluster – we could improve the utility of clusters as cosmological probes.

The general process of galaxy cluster formation through hierarchical merging is well understood, but many details, such as the impact of feedback sources on the cluster environment and radiative cooling in the cluster core are not. My thesis research has focused on studying these details via X-ray properties of the ICM in clusters of galaxies. I have paid particular attention to ICM entropy distribution, the process of virialization, and the role of AGN feedback in shaping large scale cluster properties.

### Statement of Problem

### Background and Relevance to Previous Work

### General Methodology and Procedure

### Explanation of Techniques

### Expected Results, Significance, and Application

## References

- [1] A. E. Evrard. Biased cold dark matter theory - Trouble from rich clusters? *ApJ*, 341:L71–L74, June 1989.
- [2] L. Wang and P. J. Steinhardt. Cluster Abundance Constraints for Cosmological Models with a Time-varying, Spatially Inhomogeneous Energy Component with Negative Pressure. *ApJ*, 508:483–490, December 1998.
- [3] Z. Haiman, J. J. Mohr, and G. P. Holder. Constraints on Cosmological Parameters from Future Galaxy Cluster Surveys. *ApJ*, 553:545–561, June 2001.
- [4] W. Hu. Self-consistency and calibration of cluster number count surveys for dark energy. *Phys. Rev. D*, 67(8):081304–+, April 2003.
- [5] S. Wang, J. Khoury, Z. Haiman, and M. May. Constraining the evolution of dark energy with a combination of galaxy cluster observables. *Phys. Rev. D*, 70(12):123008–+, December 2004.
- [6] A. G. Riess, A. V. Filippenko, P. Challis, A. Clocchiatti, A. Diercks, P. M. Garnavich, R. L. Gilliland, C. J. Hogan, S. Jha, R. P. Kirshner, B. Leibundgut, M. M. Phillips, D. Reiss, B. P. Schmidt, R. A. Schommer, R. C. Smith, J. Spyromilio, C. Stubbs, N. B. Suntzeff, and J. Tonry. Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *AJ*, 116:1009–1038, September 1998.
- [7] A. G. Riess, L.-G. Strolger, S. Casertano, H. C. Ferguson, B. Mobasher, B. Gold, P. J. Challis, A. V. Filippenko, S. Jha, W. Li, J. Tonry, R. Foley, R. P. Kirshner, M. Dickinson, E. MacDonald, D. Eisenstein, M. Livio, J. Younger, C. Xu, T. Dahlé, and D. Stern. New Hubble Space Telescope Discoveries of Type Ia Supernovae at  $z \geq 1$ : Narrowing Constraints on the Early Behavior of Dark Energy. *ApJ*, 659:98–121, April 2007.
- [8] D. J. Eisenstein, I. Zehavi, D. W. Hogg, R. Scoccimarro, M. R. Blanton, R. C. Nichol, R. Scranton, H.-J. Seo, M. Tegmark, Z. Zheng, S. F. Anderson, J. Annis, N. Bahcall, J. Brinkmann, S. Burles, F. J. Castander, A. Connolly, I. Csabai, M. Doi, M. Fukugita, J. A. Frieman, K. Glazebrook, J. E. Gunn, J. S. Hendry, G. Hennessy, Z. Ivezić, S. Kent, G. R. Knapp, H. Lin, Y.-S. Loh, R. H. Lupton, B. Margon, T. A. McKay, A. Meiksin, J. A. Munn, A. Pope, M. W. Richmond, D. Schlegel, D. P. Schneider, K. Shimasaku, C. Stoughton, M. A. Strauss, M. SubbaRao, A. S. Szalay, I. Szapudi, D. L. Tucker, B. Yanny, and D. G. York. Detection of the Baryon Acoustic Peak in the Large-Scale Correlation Function of SDSS Luminous Red Galaxies. *ApJ*, 633:560–574, November 2005.
- [9] A. E. Evrard. Formation and evolution of X-ray clusters - A hydrodynamic simulation of the intracluster medium. *ApJ*, 363:349–366, November 1990.
- [10] G. M. Voit. Tracing cosmic evolution with clusters of galaxies. *Reviews of Modern Physics*, 77:207–258, April 2005.