

LETTER TO THE EDITOR

Quasar feedback revealed by giant molecular outflows

C. Feruglio¹, R. Maiolino², E. Piconcelli², N. Menci², H. Aussel¹, A. Lamastra², and F. Fiore²

¹ Laboratoire AIM, DSM/Irfu/Service d'Astrophysique, CEA Saclay, F 91 191 Gif-sur-Yvette, France
e-mail: chiara.feruglio@cea.fr

² INAF- Osservatorio astronomico di Roma, via Frascati 33, 00040 Monteporzio Catone, Italy
e-mail: roberto.maiolino@oa-roma.inaf.it

Received June 6, 2010; accepted ...

ABSTRACT

In the standard scenario for galaxy evolution young star-forming galaxies transform into red bulge-dominated spheroids, where star formation has been quenched. To explain such a transformation, a strong negative feedback generated by accretion onto a central super-massive black hole is often invoked. The depletion of gas resulting from quasar-driven outflows should eventually stop star-formation across the host galaxy and lead to the black hole "suicide" for starvation. Direct observational evidence for a major quasar feedback onto the host galaxy is still missing, since outflows previously observed in quasars are generally associated with the ionized component of the gas, which only accounts for a minor fraction of the total gas content, and typically occurring in the central regions. We used the IRAM PdB Interferometer to observe the CO(1–0) transition in Mrk 231, the closest quasar known. Thanks to the wide band we detect broad wings of the CO line, with velocities up to 750 km/s and spatially resolved on the kpc scale. Such broad CO wings trace a giant molecular outflow of about 2000 M_☉/year, far larger than the ongoing star-formation rate (~200 M_☉/year) observed in the host galaxy. This wind will totally expel the cold gas reservoir in Markarian 231 in less than 10⁷ yrs, therefore halting the star-formation activity on the same timescale. The inferred kinetic energy in the molecular outflow is $\sim 4 \times 10^{44}$ erg/s, corresponding to 7% of the AGN bolometric luminosity, which is very close to the fraction expected by models ascribing quasar feedback to highly supersonic shocks generated by radiatively accelerated nuclear winds. Instead, the contribution by the SNe associated with the starburst fall short by orders of magnitude to account for the kinetic energy observed in the outflow. The direct observational evidence for quasar feedback reported here provides solid support to the scenarios ascribing the observed properties of local massive galaxies to quasar-induced large scale winds.

Key words. Galaxies: active – Galaxies: individual: Mrk 231 – Galaxies: quasars: general – Galaxies: evolution

1. Introduction

In the standard scenario for galaxy evolution emerging from both observations and models young star-forming galaxies transform into red bulge-dominated spheroids, where star-formation has been quenched (Bell et al. 2004). Three main drivers are proposed to explain such evolution: conditions at the time of galaxy formation (i.e. density at the time of the formation of the progenitors), galaxy interactions and mergers, and AGN activity. The ubiquitous discovery of super-massive black holes (SMBHs) at the centre of bulgy local galaxies and the correlations between their masses and bulge properties like mass, luminosity and velocity dispersion (Ferrarese & Ford 2005 and references therein), suggest tight links between the evolution of AGN and their host galaxies. The three drivers are tightly correlated. Indeed, massive galaxies not only live in reach environments (see e.g. Bolzonella et al. 2010, Drory et al. 2010 and references therein) but are also the remnants of galaxies formed in biased regions of the density field. Furthermore, galaxy mergers, originally proposed to explain the galaxy morphological evolution, are more common in rich environments, and are thought to destabilize cold gas, therefore enhancing star-formation and also funneling gas into the nuclear region. Such gas eventually accretes onto the nuclear SMBH, triggering an Active Galactic Nucleus (AGN hereafter). The gas and dust can intercept the line of sight to the nucleus, and therefore a natural expectation is that the early, powerful AGN phase is also highly obscured. Once SMBHs reach masses

$> 10^{7-8}$ M_☉ the AGN power is high enough to efficiently heat the gas and expel it from the galaxy through powerful winds (Silk & Rees 1998, Fabian et al. 1999). The depletion of cold gas quenches star-formation and makes the QSO nucleus "suicide", forcing galaxies toward the BH-spheroid mass relation observed locally. Such AGN negative feedback on the star formation in the host galaxy is thought to help solving two long-standing problems of galaxy evolution scenarios: the observed small number of massive galaxies relative to the prediction of models (without AGN feedback) and their red colors, indicative of old stellar populations (Granato et al. 2004, Di Matteo et al. 2005, Menci et al. 2006, Menci et al. 2008, Bower et al. 2006).

While observational evidence for feedback on the Intra Cluster Medium has been observed in radio loud AGN (Fabian et al. 2003; McNamanara & Nulsen 2007), direct observational evidence for AGN feedback onto the gas in the host galaxy (out of which stars form) is still missing. In QSOs and nearby Seyfert nuclei prominent outflows are observed, but generally only in the ionized gas component (e.g. Crenshaw et al. 2003; Turnshek 1984; Pounds et al. 2003), which accounts only for a minor fraction of the gas mass in the host galaxy, and are generally confined in the nuclear region on pc scales or in the photoionization cones. High-velocity energetic winds, whose kinetic energy is comparable to the bolometric energy of the quasar, are observed in the galactic nuclei in X-rays (Reeves et al. 2008). The bulk of the gas in QSO hosts, i.e. the molecular phase, appears little affected by the presence of the nuclear AGN. Indeed,




Fig. 1. Continuum subtracted spectrum of the CO(1–0) transition in Markarian 231. The spectrum was extracted from a region twice the beam size (Full Width Half Maximum, FWHM), and the level of the underlying continuum emission was estimated from the region with $v > 800$ km/s and $v < -800$ km/s. Left panel: full flux scale. Right panel: expanded flux scale to highlight the broad wings. The line profile has been fitted with a Gaussian narrow core (black dotted line) and a Gaussian broad component (red long-dashed line). The FWHM of the core component is 180 km/s while the FWHM of the broad component is 870 km/s, and reaching a Full Width Zero Intensity (FWZI) of 1500 km/s.

most studies of the molecular gas in the host galaxies of QSOs and Seyfert galaxies have found narrow CO lines (width of a few 100 km/s), generally tracing regular rotation patterns, with no clear evidence for prominent molecular outflows (Downes & Solomon 1998, Wilson et al. 2008, Scoville et al. 2003), even in the most powerful quasars at high redshift (Solomon & Vanden Bout 2005, Omont 2007). Yet, most of the past CO observations were obtained with relatively narrow bandwidths, which may have prevented the detection of broad wings of the CO lines possibly associated with molecular outflows. Even worse, many CO surveys were performed with single dish, where broad CO wings may have been confused with baseline instabilities and subtracted away along with the continuum.

In this paper we present new CO(1–0) observations of Mrk 231 obtained with the IRAM Plateau de Bure Interferometer (PdBI). Mrk 231 is the nearest example of a quasar object and it is the most luminous Ultra-Luminous Infrared Galaxy (ULIRG) in the local Universe (Sanders et al. 1988) with an infrared luminosity of $3.6 \times 10^{12} L_{\odot}$ (assuming a distance of 186 Mpc). A significant fraction ($\sim 70\%$) of its bolometric luminosity is ascribed to starburst activity (Lonsdale et al. 2003). Radio, millimeter and near-IR observations suggest that the starbursting disk is nearly face-on (Downes & Solomon 1998, Carilli et al. 1998, Taylor et al. 1999). In particular, past CO(1–0) and (2–1) IRAM PdBI observations of Mrk 231 show evidence for a regular rotation pattern and a relatively narrow profile (Downes & Solomon 1998), and a molecular disk (Carilli et al. 1998). The existence of a quasar-like nucleus in Mrk 231 has been unambiguously demonstrated by observations carried out at different wavelengths, that have revealed the presence of a central compact radio core plus pc-scale jets (Ulvestad et al. 1999), broad optical emission lines (Lipari et al. 2009) in the nuclear spectrum, and a hard X-ray (2–10 keV) luminosity of 10^{44} erg/s (Braitto et al. 2004). In addition, both optical and X-ray data have revealed that our line of sight to the active nucleus is heavily obscured, with a measured hydrogen column as high as $N_H = 2 \times 10^{24} \text{ cm}^{-2}$ (Braitto et al. 2004). Mrk 231 displays clear evidence of powerful ionized outflows by the multiple broad absorption lines (BAL) systems seen all over its UV and optical spectrum. In particular, Mrk 231 is classified as a low-ionization BAL quasar, a very rare subclass ($\sim 10\%$ of the entire population) of BAL quasars

characterized by weak [OIII] emission, in which the covering factor of the absorbing outflowing material may be near unity (Boroson & Meyers 1992). Furthermore, giant bubbles and expanding shells on kpc-scale are visible in deep HST imaging (Lipari et al. 2009). Recent observations with the Herschel Space Observatory have revealed a molecular component of the outflow, as traced by H_2O and OH molecular absorption features, but the lack of spatial information has prevented an assessment of the outflow rate (Fischer et al. 2010).

Due to the simultaneous presence of a strong wind, heavy X-ray absorption and of high star-formation rate, Mrk231 has been regarded as one of the promising candidates of a QSO transitioning from the obscured accretion phase, accompanied by vigorous star formation, where AGN feedback onto the host galaxy is in action, to the un-obscured phase (Page et al. 2004, Stevens et al. 2005) in the framework of the AGN-galaxy co-evolutionary sequence.

2. Data

We have exploited the wide bandwidth offered by the PdBI to observe the CO(1–0) transition in Mrk 231. The observations were carried out between June and November 2009 with the PdBI, using five of the 15 m antennas of the array. We observed the CO(1–0) rotational transition, whose rest frequency of 115.271 GHz is redshifted to 110.607 GHz ($z = 0.04217$), by using using both the C and D antenna configurations. The spectral correlator was configured to cover a band width of about 1 GHz in dual polarization. The on-source integration time was ~ 20 hours. The data were reduced, calibrated channel by channel and analyzed by using the CLIC and MAPPING packages of the GILDAS software. The absolute flux was calibrated on MWC349 ($S(3\text{mm}) = 1.27$ Jy) and 1150+497 ($S(3\text{mm}) = 0.50$ Jy). The absolute flux calibration error is of the order $\pm 10\%$. All maps and spectra are continuum subtracted, the continuum emission being estimated in the spectral regions with $v > 800$ km/s and $v < -800$ km/s.

3. Results

Fig. 1 shows the spectrum of the CO(1–0) emission line, dominated by a narrow component (Full Width Half Maximum \sim




Fig. 2. CO(1-0) maps of the broad wings. Maps of the beam (left panel, having a size of $2.74'' \times 3.4''$), of the CO(1-0) blue wing, integrated between -500 and -700 km/s (centre), and of the CO(1-0) red wing, integrated between 500 and 800 km/s (right). Contours are logarithmically equally spaced between the maximum and 10% of the maximum.

200 km/s), already detected in previous observations (Downes & Solomon 1998, Bryant & Scoville 1997). However, our new data reveal, for the first time, the presence of broad wings extending to about ± 750 km/s, which have been missed, or possibly confused with the underlying continuum, in previous narrower bandwidth observations. Both the blue and red CO(1-0) wings are spatially resolved (with a significance of 4.6σ), as illustrated by their maps and by their radial profiles in Fig. 2 and 3. The peak of the blue wing emission is not offset relative to the peak of the red wing, indicating that these wings are not due to the rotation of an inclined disk, leaving outflowing molecular gas as the only viable explanation. A Gaussian fit of the spatial profile of the blue and red wings (by also accounting for the beam broadening), indicates that the out-flowing medium extends over a region of 0.6 kpc ($0.7''$) in radius. However, both the blue and red wings are clearly extended out to a radius of ~ 4 kpc at low surface brightness (Fig. 2).

The determination of the outflow mass rate depends on the wind geometry and on the conversion factor, α , between the CO luminosity and the molecular gas mass. We estimated the CO luminosity by fitting the observed line profile with a narrow plus a broad component (see Fig. 1). The integrated CO luminosity of the broad component is $L(\text{CO})_B = 1.16 \times 10^9$ K km/s pc², about 1/10 the luminosity of the narrow component. We converted the CO luminosity of the broad component into molecular gas mass $M(\text{H}_2)$ by assuming a conservative conversion factor $\alpha = 0.5 M_\odot (\text{K km/s pc}^2)^{-1}$, i.e. 1/10 the Galactic value. This is the lowest conversion factor found in different locations of M 82 (a typical starburst galaxy), including its molecular outflow (Weiss et al. 2001). We derive a mass of the outflowing molecular gas $M(\text{H}_2) = 5.8 \times 10^8 M_\odot$, which is consistent with the lower limit of $10^7 M_\odot$ inferred by Fischer et al. (2010) based on the absorption molecular lines detected by *Herschel*. Assuming that this gas is uniformly distributed in a spherical volume of 0.6 kpc radius, its density is $\sim 25 \text{ cm}^{-3}$. Since the outflow velocity is at least 700 km/s, the inferred mass outflow rate is $dM(\text{H}_2)/dt = 2200 M_\odot/\text{yr}$. An alternative, even more conservative estimate of the outflow rate can be derived by ignoring the Gaussian fit of the broad CO component and by using only the luminosity directly measured from the broad wings, i.e. by integrating their flux at velocities higher than $+400$ km/s and lower than -400 km/s. The luminosity of the wings is $L(\text{CO}) = 3.2 \times 10^8$ K km/s pc² ($\sim 30\%$ of the CO luminosity of the total broad component inferred through the gaussian fitting). In this case we obtain a very conservative lower limit on the outflow rate of $600 M_\odot/\text{year}$. We note that alternative geometries of the outflowing wind, such as shell-like or disk-like configurations, give higher




Fig. 3. Radial profile of the beam (black dashed line), of the CO(1-0) red wing (red solid line) and of the CO(1-0) blue line wing (blue dot-dashed line). Note that the profiles of the line wings are resolved at least up to 3 arcsec (2.5 kpc).

outflow mass rate, both because the inferred outflowing gas density is higher and because the de-projected outflow velocity is higher (e.g. if the molecular outflow occurs on the galaxy disk plane). In any case the inferred outflow rate is much larger than the star-formation rate in the host galaxy of $200 M_\odot/\text{year}$ (Taylor et al. 1999, Davies et al. 2004). The mass loss rate being much larger than the rate at which gas is converted into stars implies a phase of rapid quenching of star formation in the regions reached by the outflow (~ 1 kpc scale).

4. Discussion and Conclusions

The total amount of molecular gas in the galaxy disk as inferred from the integrated emission of the narrow component, by using the CO-to- H_2 conversion factor α appropriate for ULIRGs (Solomon & Vanden Bout 2005), is $M(\text{H}_2) \sim 10^{10} M_\odot$. Assuming the measured outflow rate of $2200 M_\odot/\text{year}$, the total molecular gas mass will be expelled from the host galaxy within 5×10^6 years, implying the suppression of any significant star-formation activity on the same timescale. Even with the more conservative outflow rate of $600 M_\odot/\text{year}$ the timescale required to expel the molecular gas in the host galaxy would still be 10^7 years. Such a timescale is shorter than the Salpeter time ($\sim 5 \times 10^7$ years). We note that the quenching of star-formation has probably already started in the central regions. Indeed, stel-

lar populations younger than $5 - 20 \times 10^6$ years are not observed within the central kpc (Lipari et al. 2009).

The total kinetic power of the out-flowing gas is 3.7×10^{44} ergs/s, corresponding to $f \sim 7\%$ the AGN bolometric luminosity, $L_{\text{Bol}} \sim 5 \times 10^{45}$ ergs/s (Lonsdale et al. 2003) which, for a black hole mass of $\sim 6 \times 10^8 M_{\odot}$ (Tacconi et al. 2002), corresponds to 6% of the Eddington luminosity. Such value of the kinetic energy is very close to that expected in the case of a shock wave produced by radiation pressure onto the interstellar medium (Lapi et al. 2005). The corresponding energy injected into the interstellar medium is $\Delta E \sim f \times L_{\text{Bol}} \times t_{\text{AGN}} \sim 5.5 \times 10^{57}$ erg, at least 4 orders of magnitude larger than the overall contribution of Supernovae inferred from the observed stellar mass and age of Mrk 231 (Davies et al. 2004), and even larger than the energy injected by the radio jets (Lonsdale et al. 2003). This strongly indicates the radiation field of the quasar as the primary engine at the origin of the observed outflow. Indeed, the observed outflow velocity of ~ 750 km/s implies a Mach number $M \sim 30 - 50$, indicating a highly supersonic motion and, therefore, the formation of a shock front expanding in the interstellar medium of the host galaxy. Interestingly, this Mach number value is in agreement with the expectation for the coupling between the quasar radiation field and the gas (Lapi et al. 2005). Indeed, the Mach number can be calculated as $M^2 \sim \Delta E/E + 1$, where $E = K \times T \times M_{\text{gas}}/m_p = 3 \times 10^{54}$ erg is the thermal energy of the interstellar gas (we assumed a temperature $T = 4000$ K and an interstellar gas mass $M_{\text{gas}} \sim 10^{10} M_{\odot}$ within the shock), and $\Delta E = 5.5 \times 10^{57}$ erg is the overall quasar energy dumped into the interstellar medium during the time taken by the shock to reach the observed radius of 0.6 kpc. The above computation yields $M \sim 50$, in good agreement with our estimate based on the measured outflow speed. This suggests a solution to the long standing problem of the transport of energy from the nucleus to the bulk of the galaxy: a highly supersonic shock transports outwards the energy accumulated in the centre by radiatively accelerated nuclear winds.

Although currently based on a single object, these new observations of Markarian 231 provide one of the first direct observational evidences of quasar feedback that is dramatically affecting the evolution of its host galaxy. The quasar-driven giant molecular outflow is expected to expel the disk on short time scale, therefore halting star formation. This discovery, and in particular the inferred fraction of kinetic energy (7%) injected into the ISM relative to the quasar luminosity, confirms the expectations of models predicting a tight connection between the evolution of massive galaxies and the energy released by the accreting black holes harboured in their nuclei. The greatly improved sensitivity and expanded bandwidth of current and forthcoming millimetre interferometers will allow this kind of studies to be extended to larger samples of quasars. Therefore, it will be possible to directly verify whether the quasar feedback onto the host galaxy is really ubiquitous or not.

Acknowledgements. We would like to thank Arancha Castro-Carrizo and the IRAM staff in Grenoble for helping with data reduction and calibration. We are grateful to David Elbaz and Raphael Gobat for interesting discussions. We acknowledge support from grant D-SIGALE ANR-06-BLAN-01 and A-COSMOS-04-08. This work is based on observations carried out with the IRAM Plateau de Bure Interferometer. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain).

References

- Bell, E. F., et al. 2004, *ApJ*, 608, 752
 Bolzonella, M., Kovac, K., Pozzetti, L. et al. 2009, arXiv:0907.0013
 Boroson, T., & Meyers, K. 1992, *ApJ*, 397, 442

- Bower, R. G., et al. 2006, *MNRAS*, 370, 645
 Braito, V., et al. 2004, *A&A*, 420, 79
 Bryant, P. M., & Scoville, N. Z. 1996, *ApJ*, 457, 678
 Carilli, C. L., Wrobel, J. M., & Ulvestad, J. S. 1998, *AJ*, 115, 928
 Crenshaw, D. M., Kraemer, S. B., & George, I. M. 2003, *ARA&A*, 41, 117
 Davies, R. I., Tacconi, L. J., & Genzel, R., 2004, *ApJ*, 613, 781
 Di Matteo, T., Springel, V. & Hernquist, L. 2005, *Nature*, 433, 604
 Downes, D. & Solomon, P. M., *ApJ*, 507, 615
 Drory, N., Bundy, K. Leauthaud, A. et al. 2009, *ApJ*, 707, 1595
 Fabian, A. C., 1999, *MNRAS*, 308, 39
 Fabian, A. C., Sanders, J. S., Allen, S. W., et al. 2003, *MNRAS*, 344, L43
 Ferrarese, L., & Ford, H., 2005, *Space Science Reviews*, 116, 523
 Fischer, et al., 2010, *A&A*, in press (arXiv:1005.2213)
 Granato, G. L., De Zotti, G., Silva, L., et al. 2004, *ApJ*, 600, 580
 Lapi, A. Cavaliere, V. & Menci, N. 2005, *ApJ*, 619, 60
 Lipari, S., et al. 2009, *MNRAS*, 392, 1295
 Lonsdale, C. J., Lonsdale, C. J., Smith, H. E., & Diamond, P. J. 2003, *ApJ*, 592, 804
 McNamara, B. R. & Nulsen, P. E. J. 2007, *ARA&A*, 45, 117
 Menci, N., Fontana, A., Giallongo, E., et al. 2006, *ApJ*, 647, 753
 Menci, N., Fiore, F., Puccetti, S. & Cavaliere, A. 2008, *ApJ*, 686, 219
 Omont, A. 2007, *Rep. Prog. Phys.*, 70, 1099
 Page, M. J., Stevens, J. A., Ivison, R. J., & Carrera, F. J. 2004, *ApJ*, 611, L85
 Pounds, K. A., Reeves, J. N., King, A. R., et al. 2003, *MNRAS*, 345, 705
 Reeves, J., Done, C., Pounds, K. et al. 2008, *MNRAS*, 385, L108
 Sanders, D. B., et al., 1988, *ApJ*, 325, 74
 Scoville, N. Z., Frayer, D. T., Schinnerer, E. & Christopher, M. 2003, *ApJ*, 585, L105
 Silk, J. & Rees, M. J. 1998, *A&A*, 331, L1
 Solomon, P. M., & Vanden Bout, P. A. 2005, *ARA&A*, 43, 677
 Stevens, J. A., et al. 2006, *MNRAS*, 360, 610
 Tacconi, L. J., Genzel, R., Lutz, D. et al. 2002, *ApJ*, 580, 73
 Taylor, G. B., Silver, C. S., Ulvestad, J. S. & Carilli, C. L. 1999, *ApJ*, 519, 185
 Turnshek, D. A. 1984, *ApJ*, 280, 51
 Ulvestad, J. S., Wrobel, J. M., Carilli, C. L. 1999, *ApJ*, 516, 127
 Weiss, A., Neininger, N., Huttemeister, S., & Klein, U. 2001, *A&A*, 365, 571
 Wilson, C. D., et al. 2008, *ApJ*, 178, 189