





Planetary Science with LOFAR *and more*

Philippe Zarka

Observatoire de Paris - CNRS, LESIA, France,
philippe.zarka@obspm.fr










	Jupiter LF	Jupiter HF	Exoplanets	Lightning
What do we know ?				
What don't we know ?				
What can we learn with LOFAR ?				






Jupiter LF : What do we know ?

- Radio emission discovered in 1955
- Most intense planetary emission ($T_b > 10^{15}$ K)
- Studied for 55 years, Ground-based + S/C instruments, ~no angular resolution !
- Various components due to various sources of accelerated electrons in magnetosphere
- Sharp cutoff at 40 MHz





Jupiter LF : What do we know ?

- Above 10 MHz, Io-controlled & Io-independent emissions, L- and S- bursts, correlated with UV main oval & bright spot





Jupiter LF : What do we know ?

- Observations : emission intense, $f \sim f_{ce}$, broadband, elliptical, X-mode, narrow beaming, correlation/UV, /SW, various modulations



SKR Source Localization (from Cassini/RPWS/HFR)

Cecconi, Lamy & Zarka © 2008



Ephemeris


2006268-10:00
2006268-10:05
LT = 05:29
Lat = -29.7 deg
Dist = 7.2 R_s

Color Code

LH	RH
100–200 kHz	■
200–400 kHz	■
400–800 kHz	■
800–1000 kHz	■



Symbol Code

- out of iso-fc
- within iso-fc (2° error contour)




Jupiter LF : What do we know ?

- Theory : coherent CMI (all auroral radio emissions) → strong B , $f_{pe} \ll f_{ce}$, accelerated/unstable (keV) electrons → direct emission, of up to 1-5% of e^- energy



Jupiter LF : What do we know ?




- ExPRES modelling → oblique emission → ≠ Earth



Jupiter LF : What do we know ?

→ sources of acceleration ?



- Main oval: corotation breakdown + SW/MS interaction (latter dominant for other magnetized planets)
- Io: currents / Alfvén waves



Jupiter LF : What do we know ?





→ sources of acceleration ?

- Main oval: corotation breakdown + SW/MS interaction (latter dominant for other magnetized planets)
- Io: currents / Alfvén waves, field-aligned potential drops (DL's)






Jupiter LF : What don't we know ?

- Magnetic field model (before JUNO !) → constrains all modeling efforts
- Physics of satellite-MS interaction (existence/intensity of radio signatures, Alfvén "lead" angle)
- Physics of radio emission process (radio beaming angle)
- Direct confirmation of S-bursts theory → general implications



Jupiter LF : What can we learn with LOFAR ?

- Surface magnetic field mapping
 - Physics of Io(E,G) - Jupiter interaction
 - Radio beaming angle
 - Electron bunches & electric fields along Io flux tube
- +
- Propagation effects through Io's torus (Faraday rotation, diffraction fringes)
 - Multi-wavelength correlations (Radio, UV, IR, X)
- Need for high resolution (1-2'') fast (msec's) imaging below 40 MHz !





Available online at www.sciencedirect.com



Planetary and Space Science 52 (2004) 1455–1467

**Planetary
and
Space Science**

www.elsevier.com/locate/pss

Fast radio imaging of Jupiter's magnetosphere at low-frequencies with LOFAR

P. Zarka*




LESIA, CNRS/Observatoire de Paris, 5 Place J. Janssen, 92195 Meudon, France

Accepted 3 September 2004

+ a dozen papers by Hess et al. 2007-2010


Jupiter LF : commissionning observations

- Intense emission, detectable by single station
- Predicted ephemeris for Io-Jupiter activity
- Early long baselines Nançay-ITS observations $\sim 21\text{-}24\text{ MHz}$ \rightarrow correlation fringes measured, stability $> 1\text{ sec}$ timescale, clock drift




Jupiter LF : commissionning observations

- More recent observations since 2009 → debugging, characterizing LOFAR <30 MHz
- TBD: correlation station-to-station → synthesize coherent TAB, polarization calibration (/Nançay)



Jupiter's radio bursts detected by 3 LOFAR stations (3/6/2010). The correlation of these signals, depending on ionospheric density fluctuations, is a step of the commissionning of the coherent Tied-Array mode of LOFAR



Observations // Nançay decameter array

A&A 471, 1099–1104 (2007)
DOI: 10.1051/0004-6361:20077204
© ESO 2007








**Astronomy
&
Astrophysics**

VLBI observations of Jupiter with the initial test station of LOFAR and the Nançay decametric array

A. Nigl¹, P. Zarka², J. Kuijpers¹, H. Falcke³, L. Bähren³, and L. Denis²



Jupiter HF : What do we know ?

- Weaker continuum emission > 40 MHz
- Incoherent synchrotron, from MeV e- in radiation belts
- Linear polarization, perpendicular to B, beamed in a narrow forward cone (Γ^{-1})
- Slow variations /t (including correlated with SL9 impact)






Jupiter HF : What don't we know ?

- Low energy end of electrons spectrum
- Time variations, especially related to solar wind
- Existence at other planets (Saturn, Mercury ?)



Jupiter HF : What can we learn with LOFAR ?

- High resolution LF imaging at large $\delta f/f$ (low energy e-) \rightarrow origin + transport of energetic e- in Jupiter's inner radiation belts : pitch-angle scattering by PW, coulomb scattering, interaction with dust ?
- Variation / t & solar wind
- Existence at Saturn ? Mercury ?





ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Planetary and Space Science 52 (2004) 1449–1454

Planetary
and
Space Science

www.elsevier.com/locate/pss




LOFAR and Jupiter's radio (synchrotron) emissions

Imke de Pater*

Department of Astronomy, University of California, 601 Campbell Hall, Berkeley, CA 94720, USA

Accepted 3 September 2004

Exoplanets : What do we know ?



Exoplanets : What do we know ?

- Very weak emission if any

Maximum distance for No detection of a source $\zeta \times Jupiter$:

$$d_{\max} = (\zeta S_J A_e / 2NkT)^{1/2} (b\tau)^{1/4} = 5 \times 10^{-8} (A_e \zeta)^{1/2} f^{5/4} (b\tau)^{1/4} \text{ [pc]}$$

$\zeta = 1$	$b\tau = 10^6$ (1 MHz, 1 sec)	$b\tau = 2 \times 10^8$ (3 MHz, 1 min)	$b\tau = 4 \times 10^{10}$ (10 MHz, 1 hour)	
	$f = 10$ MHz	$f = 100$ MHz	$f = 10$ MHz	$f = 100$ MHz
$A_e = 10^4 \text{ m}^2$ (~NDA)	0.003	0.05	0.01	0.2
$A_e = 10^5 \text{ m}^2$ (~UTR-2)	0.01	0.2	0.03	0.6
$A_e = 10^6 \text{ m}^2$ (~LOFAR77)	0.03	0.5	0.1	2.
(distances in parsecs)				


Exoplanets : What do we know ?

- Electron acceleration → emission level not predictable from first principles
- Various flow/obstacle engines
- "Universal" scaling law → observe hot Jupiters

Obstacle	Flow	
Weakly/Not magnetized <i>(Venus, Mars, Io)</i>	Weakly/Not magnetized <i>(Solar wind)</i>	Strongly magnetized <i>(Jovian magnetosphere)</i>
Strongly magnetized <i>(Earth, Jupiter, Saturn, Uranus, Neptune, Ganymede)</i>	No Intense Cyclotron Radio Emission	<u>Unipolar interaction</u> → Io-induced Radio Emission,

$$P_d = \varepsilon P_m$$

$$\varepsilon = 0.1 - 0.2$$









Exoplanets : What do we know ?

- Electron acceleration → emission level not predictable from first principles
- Various flow/engine obstacles
- "Universal" scaling law → observe hot Jupiters


$$P_{\text{Radio}} \propto P_d \sim \eta \times P_m$$

$$\eta \sim 2-10 \times 10^{-3}$$



Exoplanets : What do we know ?

- Electron acceleration → emission level not predictable from first principles
- Various flow/obstacle engines
- "Universal" scaling law → observe hot Jupiters



	$b\tau = 10^6$ (1 MHz, 1 sec)	$b\tau = 2 \times 10^8$ (3 MHz, 1 min)	$b\tau = 4 \times 10^{10}$ (10 MHz, 1 hour)			
	$f = 10$ MHz	$f = 100$ MHz	$f = 10$ MHz	$f = 100$ MHz	$f = 10$ MHz	$f = 100$ MHz
$A_e = 10^4 \text{ m}^2$ (~NDA)	1	16	3	59	13	220
$A_e = 10^5 \text{ m}^2$ (~UTR-2)	3	50	11	190	40	710
$A_e = 10^6 \text{ m}^2$ (~LOFAR77)	9	160	33	600	130	2200

Exoplanets : What do we know ?





- Planetary magnetic fields ? (scaling laws, effect of spin-orbit locking)

$$\omega \downarrow$$

$$m \propto \omega^\alpha$$


$$\frac{1}{2} \leq \alpha \leq 1$$

$\rightarrow m \downarrow$ (B decay) ?



Exoplanets : What do we know ?

- Various detailed effects (role of stellar age, CME, SW estimates via Fx, SW modeling, effect of IS scintillations, M estimates, B^{*}-p topology)
- ExPRES modelling





Exoplanets : What do we know ?

- Observations at VLA @74, 333, 1465 MHz : Tau Boo


Winglee et al. (1986), Bastian et al. (2000), Farrell et al. (2003, 2004), Lazio & Farrell (2007), Lazio et al. (2010)

→ no detection (but low A_{eff} / high frequency)



Exoplanets : What do we know ?

- Observations at GMRT
- @ 153 MHz (Winterhalter et al., 2006; Majid et al., 2006; George and Stevens, 2007, 2008) : Tau Boo, Ups And
→ no detection (strongly polluted data, high frequency)
- @ 244 & 614 MHz (Lecavelier des Etangs et al., 2009) search for anti-transits of 189733
→ no detection in spite of << mJy sensitivity



→ Unlikely to detect synchrotron emission from extrasolar planets: upper estimates
~ a few nJy, vs expected sensitivity \leq microJy

Exoplanets : What do we know ?

- Observations at UTR-2 (12-32 MHz, ~40 targets, Zarka et al.)



Fig. 3. A diagram of the east-west array stations




Fig. 5. Five-beam pattern of the north-south array.




Fig. 3. A diagram of the east-west array stations







FIG. 2. A histogram of the monthly rainfall distribution.



Exoplanets : What don't we know ?

- If radio emission exists, intensity, frequency range, dynamic spectrum, variations ...
everything !
- turn Theory into Reality








Exoplanets : What can we learn with LOFAR ?

Interests:


- Direct detection → (validation of predictive scaling laws, planet-star distinction via polarization & periodicity)
 - Exoplanetary magnetic fields → (internal structure models, scaling laws)
 - Comparative magnetospheric physics (+ star-planet interactions, stellar wind strength / X flux, age, activity, CME ...)
 - Planetary rotation period → (testing spin-orbit coupling)
 - Possible access to orbit inclination
 - Possible discovery of intense radio-exoplanets
- +
- Planetary protection (versus atmospheric escape & cosmic ray bombardment)

Exoplanets : What can we learn with LOFAR ?

- Detection of ONE radio exoplanet would represent a major advance !
 → how to select targets ? ~predictions / census of HJ, magnetic stars, chromospheric activity, COROT targets, + all nearby stars & selected magnetic stars (red dwarfs...)




Tau Boo : 5-10 G (10^{-4} T)
 HD 76151 : ~10 G
 HD 189733 : >50 G
 HD 171488 : 500G



Exoplanets : What can we learn with LOFAR ?

→ Requires deep RFI mitigation





Available online at www.sciencedirect.com



**Planetary
and
Space Science**

www.elsevier.com/locate/pss

Planetary and Space Science 52 (2004) 1469–1478

The radio search for extrasolar planets with LOFAR

W.M. Farrell^{a,*}, T.J.W. Lazio^b, P. Zarka^c, T.J. Bastian^d, M.D. Desch^a, B.P. Ryabov^e



Available online at www.sciencedirect.com



**Planetary
and
Space Science**

www.elsevier.com/locate/pss

Planetary and Space Science 55 (2007) 598–617

Plasma interactions of exoplanets with their parent star and associated radio emissions

Philippe Zarka*

A&A 475, 359–368 (2007)
DOI: 10.1051/0004-6361:20077397
© ESO 2007






**Astronomy
&
Astrophysics**

Predicting low-frequency radio fluxes of known extrasolar planets^{★,★★}

J.-M. Grießmeier¹, P. Zarka¹, and H. Spreeuw²

Lightning : What do we know ?

- Antenna radiation (current discharge), from small-scale charging, large-scale charge separation, E-field build-up
- Spectrum $1/f - 1/f^4$ (depends on Stroke duration, path tortuosity, distance)
- Flash duration & distribution, correlation/optical flashes & storms



Lightning : What do we know ?






- Saturn & Uranus lightning occurrence, intensity, duration
- Meteorology @ Saturn (SED) with Cassini, ionospheric density

SED

- a few events/minute
- 30 to 300 msec
- ≤ 20 kHz to ≥ 40 MHz
- ~ 0.1 to 100 W/Hz
- ~ 0.4 to 400 Jy @ Earth


UED

- spectrum in $\sim f^{-1}$
- ~ 0.3 to 25 Jy @ Earth






Lightning : What do we know ?

- SED detection at UTR-2, monitoring by Cassini until 2017 !




Lightning : What don't we know ?

- Stroke duration at planets other than Earth → energy ?
- Existence (& properties) at Venus ?
- Dust-devils (& properties) @ Mars ?
- Transient ? existence (& properties) at J, N ?



Flux observed about 10 MHz for Saturn's lightning (SED – in yellow) and the two possible interpretations (A) and (B).



Lightning : What can we learn with LOFAR ?


Interests :

- Comparative lightning physics (electrification processes, atmospheric dynamics & cloud structure, geographical and seasonal variations ...)
→ role in the atmospheric chemistry (production of non-equilibrium trace organic constituents, potentially important for biological processes)
- Comparative planetary meteorology (/Earth), correlation with optical / IR surveys

+

- not @Titan or @exoplanets !
- broadband TAB observations
- no need for high resolution imaging
(as 2nd step only)

Planet	\varnothing_{eq} (km)	a (ua)	\varnothing_{eq-max} (")
Venus	12140	0.72	60.5
Mars	6790	1.52	17.9
Jupiter	142600	5.20	46.8
Saturne	120200	9.54	19.4
Titan	5150	9.54	0.8
Uranus	49000	19.18	3.7
Neptune	50200	30.06	2.4





Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Planetary and Space Science 52 (2004) 1435–1447

Planetary
and
Space Science

www.elsevier.com/locate/pss

Study of solar system planetary lightning with LOFAR

P. Zarka^{a,*}, W.M. Farrell^b, M.L. Kaiser^b, E. Blanc^c, W.S. Kurth^d

^a*LESIA, CNRS/Observatoire de Paris, 5 Place J. Janssen, Meudon 92195, France*

^b*Laboratory for Extraterrestrial Physics, Goddard Space Flight Center, Greenbelt, MD 20771, USA*



^c*CEA Bruyères-le-Châtel, France*

^d*Department of Physics and Astronomy, University of Iowa, Iowa-City, IA 52242, USA*

Accepted 3 September 2004


Lightning : commissionning Observations

- Cassini-predicted 3hr observations → SED activity catched



$$\sigma_{\text{sky}} = 2kT_{\text{sky}} / (A(b\tau)^{1/2})$$

$$T_{\text{sky}} \sim 1.15 \times 10^8 / f^{2.5} \quad \& \quad A_{\text{1 Station}} \sim 48\lambda^2 / 3$$



Nb of Stations	f	b	τ	σ_{sky}
1 station	30 MHz	195 kHz	82 μ s	10^4 Jy
10 stations	30 MHz	6 MHz	20ms	37 Jy


SED @ Earth $\sim 10\text{-}1000$ Jy

(1 Jy = 10^{-26} W.m $^{-2}$.Hz $^{-1}$)


Lightning : commissionning Observations

- Ongoing analysis → spectrum & time profile → energy ?


raw data (6 MHz)



clean data (6 MHz)



raw data (195 kHz) ~Off



Correlations with Cassini
in progress ...


Conclusions

	Jupiter LF	Jupiter HF	Exoplanets	Lightning
What do we know ?				
What don't we know ?				
What can we learn with LOFAR ?				


Conclusions

- Planetary and exoplanetary objectives :
 - require challenging observations
 - time demanding
 - very high interest : first radio-exoplanet ? Jupiter (& Io) as archetypes of auroral and flow-obstacle physics ; lightning @ all solar system planets
- Targeted observations (TAB primarily & Jupiter imaging) + piggybacking
- Composition of TKP/PWG :
 - P. Zarka, J.-M. Griessmeier, W. Majid, H. Spreeuw, D. Stam, G. Woan, I. dePater, G. Hallinan, S. Hess (members)
 - J. Girard, T. Bastian, E. Seran, A. Hatzes, J. Nichols [, V. Ryabov, V. Kolyadin] (associate members)

LOFAR super station in Nançay




instantaneous (u,v) coverage




LOFAR super station in Nançay


Antenna selection and tests




Antenna distribution & phasing



NEC simulations



Antenna amplifier concept, simulation, layout



- need for your scientific inputs !

Test receiver

