

Geophysical Research Letters

RESEARCH LETTER

10.1002/2017GL074376

Key Points:

- River networks dampen long-term flow trends by integrating varied climate responses across complex landscapes
- Larger subcatchments integrate greater climate trend diversity, dampening the local extremes of climate change
- Rivers may possess an underappreciated capacity and provide a tool for mitigating the impacts of climate change

Supporting Information:

- Supporting Information S1

Correspondence to:

K. A. Chezik,
kchezik@sfsu.ca

Citation:

Chezik, K. A., S. C. Anderson, and J. W. Moore (2017), River networks dampen long-term hydrological signals of climate change, *Geophys. Res. Lett.*, 44, doi:10.1002/2017GL074376.

Received 2 JUN 2017

Accepted 4 JUL 2017

Accepted article online 6 JUL 2017

River networks dampen long-term hydrological signals of climate change

K. A. Chezik¹ , S. C. Anderson² , and J. W. Moore¹ 

¹Earth to Ocean Research Group, Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia, Canada, ²School of Aquatic and Fishery Sciences, University of Washington, Seattle, Washington, USA

Abstract River networks may dampen local hydrologic signals of climate change through the aggregation of upstream climate portfolio assets. Here we examine this hypothesis using flow and climate trend estimates (1970–2007) at 55 hydrometric gauge stations and across their contributing watersheds' within the Fraser River basin in British Columbia, Canada. Using a null hypothesis framework, we compared our observed attenuation of river flow trends as a function of increasing area and climate trend diversity, with null-simulated estimates to gauge the likelihood and strength of our observations. We found the Fraser River reduced variability in downstream long-term discharge by >91%, with >3.1 times the attenuation than would be expected under null simulation. Although the strength of dampening varied seasonally, our findings indicate that large free-flowing rivers offer a powerful and largely unappreciated process of climate change mitigation. River networks that integrate a diverse climate portfolio can dampen local extremes and offer climate change relief to riverine biota.

1. Introduction

As billions are spent adapting to climate change by such means as building dikes or improving storm water drainage [Narain *et al.*, 2011], certain habitats and processes provide a natural defense system that mitigates climate change impacts [Jones *et al.*, 2012]. For example, coastal habitats such as oyster reefs, mangrove forests, and eelgrass beds shield 67% of the United States coastline and 1.4 million people from sea level rise and storm impacts [Arkema *et al.*, 2013]. Protection and enhancement of these natural systems has become a key component of many climate mitigation strategies [Guerry *et al.*, 2015], as they return multiple cost-effective services in contrast to engineered alternatives [Jones *et al.*, 2012]. For instance, revitalization of the Mississippi delta not only improves storm protection but also increases production and sustainability of fisheries and wildlife resources, protects the water supply, and reduces the impact of wastewater effluent [Louisiana Coastal Wetlands Conservation and Restoration Task Force and the Wetlands Conservation and Restoration Authority, 1998]. The value of climate-buffering habitats stands only to increase as average global temperatures climb. The Paris Agreement of 2016 aims to limit global temperature rise to 1.5°C [Hulme, 2016], a goal that concedes a minimum doubling of observed warming [Hartmann *et al.*, 2013]; therefore, an urgent challenge and opportunity is discovering, maintaining, and restoring systems that naturally buffer against the oncoming impacts of climate change.

In the face of changing global precipitation patterns [Donat *et al.*, 2016], river networks may offer an unappreciated defense against shifting flow regimes under climate change [Hartmann *et al.*, 2013; Palmer *et al.*, 2009]. Earlier snowmelt [Rauscher *et al.*, 2008], reduced snow pack [McCabe and Wolock, 2014], shifts from snow to rain and changes in the annual distribution of precipitation are all impacting stream flow [Hartmann *et al.*, 2013], resulting in flashier flows with increased frequency of flooding [Hirabayashi *et al.*, 2013], longer periods of drought, and critically low flows for wildlife and agriculture [Melillo *et al.*, 2014]. However, we hypothesize that large free-flowing river networks may naturally dampen these signals of climate change by integrating across varied landscapes and different manifestations of climate. Processes that aggregate across asynchronous components tend to dampen the aggregate, a process known as the portfolio effect [Doak *et al.*, 1998]. Climate is expressed differently and asynchronously over the landscape [e.g., Wang *et al.*, 2012], and on account of their branching architecture and directional flow, river networks integrate these varied climate manifestations [Peterson *et al.*, 2013], potentially dampening the local climatic response of subcatchments. Large free-flowing rivers may have diverse climate portfolios, with subcatchments acting as assets in their




Figure 1. Climate trends (1970–2007) in mean annual temperature (MAT) and mean annual precipitation (MAP) within the Fraser River basin overlaid on a digital elevation model within British Columbia, Canada. Flow gauge sites (dots) are scaled by the size of the contributing area. The map is projected in Albers BC UTM to provide an equal area depiction of the region, but labels are expressed in WGS84 latitude and longitude.

portfolio, thereby dampening local climate impacts. Portfolio effects in rivers are already known to dampen short-term variation [Moore *et al.*, 2015; Yeakel *et al.*, 2014], but attenuation of long-term climate driven shifts have yet to be examined.

Here we consider whether river networks attenuate local long-term climate change through analysis of hydro-metric data from one of the largest free-flowing rivers in North America. Located in British Columbia, Canada, the Fraser River drains an area approximately the size of the United Kingdom ($\sim 217,000 \text{ km}^2$) and in a rare combination is fairly well monitored while also having no dams on its main stem [Vörösmarty *et al.*, 2010] (Figure 1). This watershed drains interior high plateau, coastal mountains, and the Canadian Rockies and thus integrates a diverse mosaic of landscapes, weather, and climate. As with other midlatitude rivers [Bindoff *et al.*, 2013], the Fraser River's discharge volume and variability have increased over the last several decades [Déry *et al.*, 2012; Morrison *et al.*, 2002]. Following projected patterns of climate change, the Fraser River basin is increasingly shifting from a snow-dominated to rain-snow hybrid system [Kang *et al.*, 2016]. Using 38 years of flow data collected concurrently at 55 hydrometric gauge stations throughout the watershed, we use a novel analytical approach to examine whether this large river exhibits signals of long-term climate dampening.

2. Materials and Methods

2.1. Discharge Data and Flow Metrics

To assess signals of climate change in hydrology within the Fraser River basin, we downloaded daily flow data from Environment Canada's HYDAT database. Within HYDAT we selected Fraser River basin hydrometric gauge stations that were not observed to be dam influenced and which collected data in each month between 1970 and 2007, where no month was missing more than 5 days data and at least 35 complete years were present. We selected this 38 year range because this period maximized the number of gauge stations in the Fraser River basin ($n = 55$) operating concurrently over a long enough time period in which climate trends may be observable. We linearly interpolated missing data in log-space, but of the 746,544 days of data only 23 days were interpolated. These daily data were smoothed using a 5 day rolling average before summarizing into annual and monthly response variables. This smoothing technique was used to reduce the influence of erroneous and unusually extreme values [e.g., Déry *et al.*, 2009]. Because dams control flow and stabilize

flow trends, we removed sites where daily flow exhibited unusually high autocorrelation and low variability relative to other sites of similar size, and sites that were observed downstream of, and atypically close to, a dam ($n = 3$). The downstream distance of gauge stations from dams was determined using GIS spatial layers in the *BC Data Catalog*, maintained by the BC provincial government.

As climate change shifts precipitation patterns from snow to rain in temperate regions, rivers are generally predicted to have lower low flows and higher high flows, as well as earlier snowmelt [Nijssen *et al.*, 2001]. These manifestations of rising temperatures are impacting species [Xenopoulos and Lodge, 2006] and people [Hirabayashi *et al.*, 2013]. We summarized our daily rolling average discharge data into annual and monthly flow metrics that capture these transitions. We used maximum- and minimum-flow estimates to capture trends in extreme discharge and median-flow estimates to capture general annual and seasonal trends. Because timing of flow is particularly important to the phenology of many organisms, we also calculated the day of year in which half the annual flow was reached.

2.2. River Discharge Trend Analysis

To test for a river network portfolio effect, we need to detect varied and asynchronous, long-term changes in flow throughout the watershed. To do this, we estimated annual and monthly flow trends for each response variable at each gauge station using a generalized least squares (GLS) model with an AR1 autocorrelation function:

$$Q_{s,t} = \alpha_s + b_s Y_{s,t} + \epsilon_{s,t}, \quad \epsilon_{s,t} \sim \mathcal{N}(\phi \epsilon_{s,t-1}, \sigma_Q^2), \quad (1)$$

where $Q_{s,t}$ represents a flow metric (log(maximum), log(minimum), log(median), logit(day of year to half annual flow)) at each site (s) and time point (t), and $Y_{s,t}$ represents the time in years (e.g., 1970, 1971, ..., 2007) with 1988 subtracted to approximately center the predictor. Flow metrics were scaled by site to facilitate cross-site comparisons. Parameters b_s (slope) and α_s (intercept) represent the estimated mean effect of time on $Q_{s,t}$ and the estimate of $Q_{s,t}$ at time zero (i.e., 1988), respectively. Error ($\epsilon_{s,t}$) of the current time step for a given site was allowed to be correlated with that of the previous time step by ϕ and was assumed to be normally distributed with a variance of σ_Q^2 . To ensure our model estimates and simulations remained within the calendar year, we logit transformed our flow-timing response variable (i.e., day of year to half annual flow) after scaling the data between 0 and 1 (i.e., dividing by 365).

2.3. Quantifying Climate Variability

For a river network to act as a portfolio of climate and dampen the effects of climate change, expressions of climate and their trends need to vary throughout the network. In order to quantify climate trend diversity, we developed a climate index for each flow gauge catchment. Using the Western North America Climate tool (ClimateWNA) [Wang *et al.*, 2016], we estimated historic climate values on an evenly spaced 1 km² grid across the Fraser River basin for each year and month in our study. Temperature and precipitation are likely to drive changing hydrology; therefore, we included variables in our climate index that capture changes in extreme and mean temperature, as well as changes in precipitation volume and physical state (snow versus rain). Climate variables included the mean annual temperature (MAT), and precipitation (MAP), extreme minimum (EMT) and maximum (EXT) temperature, and precipitation as snow (PAS). Climate variable trends were calculated at each grid point using a GLS model similar to equation (1):

$$C_{s,t} = \alpha_s + \beta_s Y_{s,t} + \epsilon_{s,t}, \quad \epsilon_{s,t} \sim \mathcal{N}(\phi \epsilon_{s,t-1}, \sigma_C^2), \quad (2)$$

where $C_{s,t}$ represents a climate variable and $Y_{s,t}$ represents the time in years (e.g., 1970, 1971, ..., 2007). Parameters β_s (slope) and α_s (intercept) represent the estimated mean effect of time on $C_{s,t}$ and the estimate of $C_{s,t}$ at time zero, respectively. Autocorrelated errors were incorporated as in equation (1).

Spatial variation in climate change would suggest catchments of different size integrate different amounts of climate variability. In order to calculate a climate index for each gauge station, we delineated flow gauge catchments and summarized climate trends within those catchments. Using Whitebox Geospatial Analysis Tools [Lindsay, 2016], we delineated the Fraser River basin and the catchment areas of each flow gauge station. Digital elevation models were provided by the provincial government of British Columbia, Canada, and state government of Washington, USA, and were preprocessed with a breaching algorithm and stream burned to facilitate proper flow path and accumulation models [Woodrow *et al.*, 2016]. We summarized the climate variability integrated at each flow gauge by calculating the standard deviation in climate trends within their




Figure 2. Climate variability index as a function of catchment area. As watershed area increases the variety of climatic trends within the watershed increases at a diminishing rate.

River sub-catchments have more variable climate portfolios (Figure 2) as well as aggregate more water and carry more weight on downstream dampening. Therefore, to test whether networks dampen climate signals, we regressed site-specific trend estimates (e.g., % change · decade⁻¹) onto each site's catchment area using a GLS model:

$$\hat{b}_s = c + d\sqrt{A_s} + \eta_s, \quad \eta_s \sim \mathcal{N}(0, f(A_s)), \quad (3)$$

where \hat{b}_s represents a flow trend at a given site (s) and A_s represents site s 's watershed area. We adjusted $\sqrt{A_s}$ by subtracting the mean \sqrt{A} , thereby centering our predictor. Fitted d and c parameters represent the mean effect of watershed area (\sqrt{A}), on flow trends and the mean flow trend for an average sized watershed (i.e., ~ 5700 km²), respectively. Captured in our climate portfolio proxy variable (A) is the interaction between climate and land use. Timber harvest, the dominant source of land cover change in the Fraser River basin, scales with subbasin area and contributes to changes in flow. Although exploratory linear models at the site level do suggest an effect of timber harvest on long-term flow trends for some gauge stations (Figure S1 and S2 in the supporting information), when information is shared across sites in a global model, climate effects dominate (Figure S3). Thus, we did not attempt to separate the contributions of land cover change, climate change, and their interaction on long-term flow trends.

To quantify attenuation, we modeled the change in trend variability with the change in catchment area using an exponential variance function [Pinheiro and Bates, 2000, p. 211]:

$$f(A_s) = \sigma_b^2 \exp(2\delta\sqrt{A_s}), \quad (4)$$

where the variance of the estimated error (η_s) exponentially increases with \sqrt{A} as defined by the estimated δ parameter. This variance function allowed us to predict the range of flow trend values that would be expected as watershed area increased. A decreasing range of flow trends with increasing watershed area indicates a dampening network effect.

2.5. River Attenuation Null Model Simulations

In the observed data, small watersheds exhibited greater variability around their trend estimates than large watersheds, likely because of greater short-term variation in small catchments [Moore et al., 2015]. This relationship between watershed size and trend certainty may pull small watershed trend estimates away from zero, thereby creating the appearance of decreasing trend variability among sites as watershed size increases. Using a null hypothesis framework, we simulated time series with no underlying trend from each site's fitted model, using the estimated standard deviation and autocorrelation parameters ($\hat{\sigma}_Q$, $\hat{\phi}$). Using the same GLS model and AR1 correlation structure as applied to the observed data (equation (1)), we estimated trends for 1000 simulated time series at each site for each response variable (e.g., Figure S4). This simulation process, a form of parametric bootstrapping, created distributions of null expectations with which we compared our observed results.

catchments using the *rasterstats* module (<https://github.com/perrygeo/python-rasterstats.git>) in Python v2.7. The standard deviations of these climate variables were scaled between 0 and 1 and summed as a general climate index for each catchment.

2.4. Quantifying the River Network Portfolio Effect

We hypothesize that flow gauge sites with larger climate portfolios will exhibit greater long-term flow trend stability than sites with smaller climate portfolios. We used catchment area as a proxy for the climate portfolio diversity of a given site, given that larger Fraser




Figure 3. Annual flow trend attenuation within the Fraser River basin. Trend estimates ($\hat{\beta}_s$) ± 1 standard error (SE, grey) plotted against watershed area (km^2), colored by climate portfolio strength (green = small; blue = large) (Figures 3a and 3c). Blue lines represent observed attenuation; orange and red represent simulated attenuation that is weaker and stronger than observed. Density plots show null model simulated variance exponents ($\hat{\delta}$) and the proportion on either side of the observed variance exponent (blue) (Figures 3b and 3d). Flow metrics include long-term flow-timing shifts (change per decade in day of year (DOY) to half annual flow), where decreasing trends suggest (a) more annual flow is occurring earlier in the year, and the percent change per decade in (b) minimum, (c) maximum, and (d) median annual flow. Simulated lines ignore variance in the intercept and slope to focus visually on attenuation.

We then applied equation (3) to each of the 1000 basin-wide simulations (e.g., Figure S5), resulting in 1 observed variance exponent parameter ($\hat{\delta}$) and 1000 simulated $\hat{\delta}$ for each flow metric. By comparing our observed attenuation with our basin-wide null model simulations, we addressed the potential that more variable flows in smaller catchments contribute to the observed pattern of flow trends as a function of watershed area (Figure 3). This null model approach asks how likely our observation is due to a sampling effect versus a network portfolio effect.

2.6. Dampening Summary Statistics

We calculated attenuation certainty as the percentage of simulated $\hat{\delta}$ that were less than the observed $\hat{\delta}$. We calculated attenuation strength (AS) as the ratio between the standard deviation at the smallest and largest watersheds as defined by the exponential variance function (i.e., $AS = \sqrt{f(A_{s=1})}/\sqrt{f(A_{s=2})}$). To compare the degree of attenuation observed with the null-simulated attenuation, we compared this ratio with the same ratio calculated from the null-simulated data (i.e., AS/AS_{NULL}). Finally, to estimate how the day of year to half-annual flow was changing, we estimated the steepest point of each site's logistic curve and multiplied this slope value by 365 to revert from our 0–1 scale to our original annual range of 0–365. To get a decadal rate, we multiplied these results by 3.8 (number of decades in 38 years). We estimated the steepest part of the logistic curve using the divide by four rule [Gelman and Hill, 2008] (dividing the coefficient by four equals the first derivative of the logistic curve at its steepest point).




Figure 4. Monthly maximum-flow trend attenuation within the Fraser River basin. (left) Fraser River's basin-wide maximum-flow trend estimates (i.e., intercept = vertical grey lines) by month with density distributions of null model simulations. Observed values falling farther from the center of the density distribution suggest greater evidence for changes in maximum flow and a greater shift in magnitude. (middle) Observed monthly Fraser River maximum-flow variance exponent ($\hat{\delta}$, blue) and associated density distribution of simulated $\hat{\delta}$ estimates. Decimal values represent the percent of simulated data exhibiting weaker attenuation (yellow) than observed. (right) Trend estimates (\hat{b}_s) ± 1 standard error (SE, grey) plotted against watershed area (km^2), colored by climate portfolio strength (green = small; blue = large), for four seasonally representative months. These reflect months in the prior columns and describe the variation in percent change per decade of maximum flow among sites. Simulated lines ignore variance in the intercept and slope to focus visually on attenuation.

3. Results

3.1. Hydrologic Trend Diversity

Long-term changes in flow metrics varied substantially across the watershed. For instance, annual maximum flow decreased in 40 sites but increased in 15 over the last four decades (-26 to 10% change decade $^{-1}$, standard deviation (SD) = 6%). Annual median flow showed similar variability but a greater tendency toward rising flow trends with 49 increasing and only six decreasing (-5 to 25% change · decade $^{-1}$, SD = 5%). Thus, within the Fraser River basin, there are a variety of long-term trends in flow, suggesting a diversity of climate responses and landscape effects.

3.2. Climate Trend Diversity

Our index of climate variability increased with area (Figure 2), asymptoting in catchments larger than 40,000 km². Thus, rivers that drain larger catchments integrate a greater diversity of shifting climates. These findings suggest that larger areas contain greater landscape complexity, which produce a variety of climates locally and respond to global climate shifts differently.

3.3. Climate Portfolios and Flow Trend Dampening

Long-term changes in hydrology were less variable in larger catchments, as predicted, demonstrating between 92 and 96% dampening of flow trends moving from headwater catchments to the network's outlet. The largest catchments were approximately 10 times more stable in their climate response than the smallest catchments. For instance, trends ranged between a 6% reduction and 19% increase in median flow per decade among small watersheds, while large watersheds ranged between 0.8 and 1.6% change per decade. Similar attenuation was seen in flow timing with a 94% reduction in the day-of-year to half-annual-flow trends such that small watersheds were reaching half their annual flow between 8 days later and 25 days earlier while large watersheds have only shifted 2 to 4 days earlier between 1970 and 2007.

Our null model approach illustrates that support for a river network portfolio effect is greater than 90% for all four annual flow trend metrics and as high as 98% for median-annual-flow trends (Figure 3). Trend attenuation was 3.1 (0.6–9.8 this and hereafter are 90% intervals), 4.7 (1.1–14.7), 3.2 (0.9–9.0), and 5.2 (1.4–14.4) times greater than the null model for annual-flow timing and minimum, maximum, and median annual flow, respectively. These statistics provide strong support for the hypothesis that river network portfolio effects contribute to climate dampening in large rivers.

3.4. Seasonally Shifting Flow Trend Dampening

Regional flow trends and attenuation strength varied by season. Analysis of monthly trends revealed that winter flows are getting higher and summer flows are getting lower over time but that these trends have been more variable in smaller catchments (Figures 4, S6, and S7). For example, winter maximum flows in small catchments ranged from nearly no change to dramatic 47% increases per decade with decreases only as large as 9%. Despite these extreme locations, on average, the basin has only experienced moderate winter maximum flow increases of 5–9%. Summer flows exhibited the opposite, with decreases in high flows over time. However, there was weak evidence of attenuation in the spring (Figures 4, S6, and S7). Overall, climate portfolios tended to have greater attenuation during stable periods of the year and decreased during seasonal transitions. In spring, the likelihood of network-driven attenuation of maximum-flow trends was as low as 24% (Figure 4). Deterioration of the network's attenuation strength was coincident with the spring freshet, when snowmelt drives high flows across the basin. Synchronization events such as the freshet may subvert river network dampening mechanisms by homogenizing the region's response to seasonally driven climate shifts.

4. Discussion

Our results suggest that river networks mitigate and dampen long-term hydrologic trends by integrating a diverse climate portfolio over a heterogeneous landscape. By quantifying the diversity of climate and flow trends within many subbasins of the Fraser River watershed (Figures 1 and 2), we demonstrate as much as a tenfold decrease in flow trend variability with increasing watershed area (i.e., climate portfolio) (Figures 3 and 4). Similar to other temperate rivers [Rauscher *et al.*, 2008] and consistent with recent studies [e.g., Kang *et al.*, 2016, 2014; Shrestha *et al.*, 2012], we show the Fraser River basin is exhibiting climate driven shifts with increasing flows over winter transitioning to decreasing flows over the summer (Figure 4). These findings are consistent with a general body of work that suggests climate change is transitioning previously snow driven systems to rain, resulting in a redistribution of water toward winter flows [Bindoff *et al.*, 2013]. As climate change progresses, river hydrology will continue to shift, further stressing riverine ecosystems and subsequently demanding responsive management. However, our work suggests that river networks provide an underappreciated defense against these climate change impacts as they are uniquely organized to leverage locally filtered expressions of climate into stability. A simple product of network form and gravity, stability emerges as river networks integrate landscape heterogeneity, creating a downstream portfolio of climate that smooths local extremes and tempers long-term flow trends. Therefore, larger rivers should have flow regimes that are less sensitive to local climate trends.

The Fraser River basin has seen changes in land cover between 1970 and 2007, which have impacted river discharge [e.g., *Zhang and Wei*, 2014]. Over this period, timber harvest has been the dominant contributor to changes in forest cover (Figure S8), but increasingly, natural events are having a greater impact. For instance, abnormally hot, dry weather in 2003 led to fires that destroyed an unprecedented 2600 km^2 of forest largely within the Fraser River basin [*Filmon*, 2003]. Beginning in the early 2000s, western mountain pine beetle outbreaks began consuming large swaths of forest, impacting 20% of British Columbia by 2013 [*Schnorbus et al.*, 2010]. These natural events are facilitated by climate change and are becoming increasingly common and extreme [*Melillo et al.*, 2014; *Maness et al.*, 2013]. Pine beetle outbreaks are often subsequently heavily logged, and timber removal can occur in anticipation of fires, resulting in harvest practices responding dynamically to climate change. Analyses indicated that while timber harvest rates did have some local impacts on hydrological change, harvest did not have substantial impacts on flows in a global model (Figure S3); thus, we focus on long-term hydrological change and make the simplifying assumption that different forest harvest rates are subsumed by long-term climate change. Indeed, timber harvest was relatively consistent over the last 38 years across the vast majority of our 55 study catchments, with the exception of four relatively small, heavily fire and pine beetle impacted sites (Figure S8). Therefore, our study captures the interwoven direct and indirect impacts of climate change by considering changes in precipitation but also latent affects through climate filtering of altered landscapes due to climate change.

Our study provides insight into the spatial scaling of a river network's dampening capacity. Although we focused on a vast watershed with a remarkably diverse climate portfolio, we would still expect that smaller rivers or watersheds with less topographic complexity will dampen climate variability. For instance, based on our data, we predict that rivers draining a catchment of $60,000 \text{ km}^2$ would have 66% less variability in the rate of change in hydrology than a smaller river draining 5000 km^2 . While this dampening is less than the tenfold decrease observed in the entire $217,000 \text{ km}^2$ Fraser River watershed, smaller watersheds can offer a defense against the impacts of climate change.

5. Conclusion

Climate change is causing economic and conservation challenges for river systems worldwide [*Palmer et al.*, 2009; *Pel et al.*, 2017]. In the Fraser River, sockeye salmon have seen as much as 80% prespawn mortality in years with later runoff and elevated temperatures, leading managers to reduce or close the fishery (e.g., 2013 and 2015). In the absence of climate dampening, the impacts of these climate extremes could have been considerably worse. In contrast to previously studied types of climate mitigation where habitats modify climate drivers (e.g., carbon storage [*Jones et al.*, 2012]) or physically absorb climate change impacts (e.g., mangroves [*Arkema et al.*, 2013]), river network climate dampening smooths out extreme climatic trends thereby reducing the impact of local extremes. But just as the destruction of coastal habitats degrades their natural capacity to mitigate sea level rise or storm events [*Arkema et al.*, 2013], the management of river basins likely impacts the ability for river networks to attenuate climate trends. For example, dams synchronize the flow regimes of rivers [*Poff et al.*, 2007]. These anthropogenic activities may magnify the impacts of climatic shifts such as the transition from snow- to rain-dominated precipitation in some basins. The climate cost of these watershed activities should be considered in environmental decision making. Climate change is a global challenge; here we suggest large rivers dampen local change by leveraging landscape diversity into a portfolio of climate, an important tool in the climate mitigation toolbox.

Acknowledgments

This paper relied on many open source tools including R v3.3.2 (www.r-project.org/), Python v2.7, QGIS v2.18, and WhiteboxGAT [*Lindsay*, 2016]. All R code can be found at <http://github.com/kchezik/River-Network-Flow-Trends.git>. Data processing was facilitated greatly by GNU Parallel [*Tange*, 2011]. This study relied on decades of flow data collected by those at Environment Canada. Raw HYDAT data can be obtained via Environment Canada, or a simplified version can be found on GitHub. K.A. Chezik and J.W. Moore were supported by the Liber Ero Chair of Coastal Science and Management and Simon Fraser University. S.C. Anderson was supported by the David H. Smith Conservation Research Fellowship.

References

- Arkema, K. K., G. Guannel, G. Verutes, S. A. Wood, A. Guerry, M. Ruckelshaus, P. Kareiva, M. Lacayo, and J. M. Silver (2013), Coastal habitats shield people and property from sea-level rise and storms, *Nat. Clim. Change*, 3(10), 913–918, doi:10.1038/nclimate1944.
- Bindoff, N. L., et al. (2013), Detection and attribution of climate change: From global to regional, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker et al., pp. 867–952, Cambridge Univ. Press, Cambridge, U. K., and New York.
- Déry, S. J., K. Stahl, R. Moore, P. H. Whitfield, B. Menounos, and J. E. Burford (2009), Detection of runoff timing changes in pluvial, nival, and glacial rivers of western Canada, *Water Resour. Res.*, 45, W04426, doi:10.1029/2008WR006975.
- Déry, S. J., M. A. Hernández-Henríquez, P. N. Owens, M. W. Parkes, and E. L. Petticrew (2012), A century of hydrological variability and trends in the Fraser River basin, *Environ. Res. Lett.*, 7(2), 024019, doi:10.1088/1748-9326/7/2/024019.
- Doak, D., D. Bigger, E. Harding, M. Marvier, O. RE, and D. Thomson (1998), The statistical inevitability of stability-diversity relationships in community ecology, *Am. Nat.*, 151(3), 264–276, doi:10.1086/286117.
- Donat, M. G., A. L. Lowry, L. V. Alexander, P. A. O'Gorman, and N. Maher (2016), More extreme precipitation in the world's dry and wet regions, *Nat. Clim. Change*, 6(5), 508–513, doi:10.1038/nclimate2941.

- Filmon, G. (2003), *Firestorm 2003: Provincial Review*, Gov. of British Columbia, Victoria, B.C.
- Gelman, A., and J. Hill (2008), *Data Analysis Using Regression and Multilevel Hierarchical Models*, Cambridge Univ. Press, New York.
- Guerry, A. D., et al. (2015), Natural capital and ecosystem services informing decisions: From promise to practice, *Proc. Natl. Acad. Sci. U.S.A.*, 112(24), 7348–7355, doi:10.1073/pnas.1503751112.
- Hartmann, D., et al. (2013), Observations: Atmosphere and surface, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker et al., pp. 159–254, Cambridge Univ. Press, Cambridge, U.K. and New York.
- Hirabayashi, Y., R. Mahendran, S. Koirala, L. Konoshima, D. Yamazaki, S. Watanabe, H. Kim, and S. Kanae (2013), Global flood risk under climate change, *Nat. Clim. Change*, 3(9), 816–821, doi:10.1038/nclimate1911.
- Hulme, M. (2016), 1.5°C and climate research after the Paris Agreement, *Nat. Clim. Change*, 6(3), 222–224, doi:10.1038/nclimate2939.
- Jones, H. P., D. G. Hole, and E. S. Zavaleta (2012), Harnessing nature to help people adapt to climate change, *Nat. Clim. Change*, 2(7), 504–509, doi:10.1038/nclimate1463.
- Kang, D., X. Shi, H. Gao, and S. Déry (2014), On the changing contribution of snow to the hydrology of the Fraser River Basin, Canada, *J. Hydrometeorol.*, 15, 1344–1365.
- Kang, D., H. Gao, X. Shi, S. ul Islam, and S. J. Déry (2016), Impacts of a rapidly declining mountain snowpack on streamflow timing in Canada's Fraser River basin, *Sci. Rep.*, 6, 19299.
- Louisiana Coastal Wetlands Conservation and Restoration Task Force and the Wetlands Conservation and Restoration Authority (1998), *Coast 2050: Toward a Sustainable Coastal Louisiana*, 161 pp., Louisiana Dep. of Nat. Resour., Baton Rouge, La.
- Lindsay, J. (2016), Whitebox GAT: A case study in geomorphometric analysis, *Comput. Geosci.*, 95, 75–84.
- Maness, H., P. J. Kushner, and I. Fung (2013), Summertime climate response to mountain pine beetle disturbance in British Columbia, *Nat. Geosci.*, 6(1), 65–70.
- McCabe, G. J., and D. M. Wolock (2014), Spatial and temporal patterns in conterminous United States streamflow characteristics, *Geophys. Res. Lett.*, 41, 6889–6897, doi:10.1002/2014GL061980.
- Melillo, J., T. Richmond, and G. W. Yohe (2014), *Climate Change Impacts in the United States: The Third National Climate Assessment*, 841 pp., U.S. Gov. Print. Off., Washington, D.C.
- Moore, J. W., et al. (2015), Emergent stability in a large, free-flowing watershed, *Ecology*, 96(2), 340–347, doi:10.1890/14-0326.1.
- Morrison, J., M. Quick, and M. Foreman (2002), Climate change in the Fraser River watershed: Flow and temperature projections, *J. Hydrol.*, 263, 230–244.
- Narain, U., S. Margulis, and T. Essam (2011), Estimating costs of adaptation to climate change, *Clim. Policy*, 11(3), 1001–1019, doi:10.1080/14693062.2011.582387.
- Nijssen, B., G. O'Donnell, A. Hamlet, and D. Lettenmaier (2001), Hydrologic sensitivity of global rivers to climate change, *Clim. Change*, 50, 143–175.
- Palmer, M. A., D. P. Lettenmaier, L. N. Poff, S. L. Postel, B. Richter, and R. Warner (2009), Climate change and river ecosystems: Protection and adaptation options, *Environ. Manage.*, 44(6), 1053–1068, doi:10.1007/s00267-009-9329-1.
- Pecl, G. T., et al. (2017), Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being, *Science*, 355(6332), eaai9214, doi:10.1126/science.aai9214.
- Peterson, E. E., et al. (2013), Modelling dendritic ecological networks in space: An integrated network perspective, *Ecol. Lett.*, 16(5), 707–719, doi:10.1111/ele.12084.
- Pinheiro, J., and D. Bates (2000), *Mixed-Effects Models in S and S-PLUS*, Springer, New York.
- Poff, L. N., J. D. Olden, D. M. Merritt, and D. M. Pepin (2007), Homogenization of regional river dynamics by dams and global biodiversity implications, *Proc. Natl. Acad. Sci. U.S.A.*, 104(14), 5732–5737, doi:10.1073/pnas.0609812104.
- Rauscher, S. A., J. S. Pal, N. S. Diffenbaugh, and M. M. Benedetti (2008), Future changes in snowmelt-driven runoff timing over the western US, *Geophys. Res. Lett.*, 35, L16703, doi:10.1029/2008GL034424.
- Schnorbus, M., A. Werner, and K. Bennett (2010), *Quantifying the Water Resource Impacts of Mountain Pine Beetle and Associated Salvage Harvest Operations Across a Range of Watershed Scales: Hydrologic Modeling of the Fraser River Basin*, p. 423, Pac. For. Cent., Victoria, B.C.
- Shrestha, R. R., M. A. Schnorbus, A. T. Werner, and A. J. Berland (2012), Modelling spatial and temporal variability of hydrologic impacts of climate change in the Fraser River basin, British Columbia, Canada, *Hydro. Processes*, 26(12), 1840–1860, doi:10.1002/hyp.9283.
- Tange, O. (2011), GNU parallel—The command-line power tool, *login: The USENIX Magazine*, 36(1), 42–47, doi:10.5281/zenodo.16303.
- Vörösmarty, C., et al. (2010), Global threats to human water security and river biodiversity, *Nature*, 467, 555–561, doi:10.1038/nature09440.
- Wang, T., A. Hamann, D. L. Spittlehouse, and T. Q. Murdock (2012), ClimateWNA—High-resolution spatial climate data for Western North America, *J. Appl. Meteorol. Climatol.*, 51(1), 16–29, doi:10.1175/JAMC-D-11-0431.
- Wang, T., A. Hamann, D. Spittlehouse, and C. Carroll (2016), Locally downscaled and spatially customizable climate data for historical and future periods for North America, *PLoS One*, 11(6), e0156720, doi:10.1371/journal.pone.0156720.
- Woodrow, K., J. B. Lindsay, A. A. Berg, and J. Hydrol. (2016), Evaluating DEM conditioning techniques, elevation source data, and grid resolution for field-scale hydrological parameter extraction, 540, 1022–1029.
- Xenopoulos, M. A., and D. M. Lodge (2006), Going with the flow: Using species-discharge relationships to forecast losses in fish biodiversity, *Ecology*, 87(8), 1907–1914.
- Yeakel, J., J. Moore, P. Guimarães, and M. Aguiar (2014), Synchronisation and stability in river metapopulation networks, *Ecol. Lett.*, 17(3), 273–283, doi:10.1111/ele.12228.
- Zhang, M., and X. Wei (2014), Alteration of flow regimes caused by large-scale forest disturbance: A case study from a large watershed in the interior of British Columbia, Canada, *Ecohydrology*, 7(2), 544–556, doi:10.1002/eco.1374.