



Field-testing olfactory ability to understand human olfactory ecology

Kara C. Hoover¹ | Denisa Botescu² | Piotr Fedurek³ | Sophie Aarts³ | J. Colette Berbesque³

¹Department of Anthropology, University of Alaska, Fairbanks, Alaska

²Department of Anthropology, University College of London, London, UK

³Centre for Research in Evolutionary, Social and Inter-Disciplinary Anthropology, University of Roehampton, London, UK

Correspondence

Kara C. Hoover, Department of Anthropology, University of Alaska, Fairbanks, AK, United States.
Email: kchoover@alaska.edu

Abstract

Objectives: We know little about human olfactory ability in natural settings because current knowledge derives from lab-based studies using non-representative samples of convenience. The primary objective was to use a validated lab tool, the five-item odor identification test, to assess variation in olfactory ability in different environments.

Methods: Using the five-item test, we conducted two repeated measures experiments that assessed participant ability to correctly identify an odor source in different odor environments. We also examined consistency in odor labelling due to documented potential bias from idiosyncrasies in odor terms.

Results: We found no variation in olfactory ability due to environment, but this may be due to methodological biases. First, subjective bias results from idiosyncratic differences in participant labelling and researcher coding of answer correctness. Second, better ability to learn odors may provide an advantage to women. Third, reducing positive female learning bias by analyzing consistency in response (regardless of correct odor source identification) results in no sex differences but fails to assess the functional aspect of olfactory ability (naming the correct odor source). Fourth, functional olfactory ability is significantly better in women is significantly better in females, especially in food-rich odor environments.

Conclusions: Environment was not a significant factor in olfactory ability in this study but that result may be confounded by methodological biases. We do not recommend odor identification as a field tool. Functional olfactory ability exhibits a sex-based pattern but consistency in recognizing the same odor does not. Food-rich odors may enhance olfactory ability in females. We discuss evolutionary and ecological implications of superior female functional olfactory ability relative to food foraging activity.

1 | INTRODUCTION

The vast majority of our knowledge on human olfaction derives from lab studies conducted in odor-controlled environments using samples of convenience, typically white educated industrialized rich and democratic or

WEIRD populations (Henrich, Heine, & Norenzayan, 2010)—often students (Hanel & Vione, 2016). While ethnographic work in anthropology has examined olfaction, the focus has been on odor symbolism rather than olfactory ability or olfactory biology (for an overview, see Classen, 1993; Classen, Howes, & Synnott, 1994).



Meanwhile, the vast majority of human olfactory experience is in dynamic non-western and nonindustrialized odorscapes. The limited research conducted on olfactory ability in different ecological settings suggests that non-industrial populations may have superior olfactory sensitivity but not discrimination abilities (Sorokowska, Sorokowski, & Hummel, 2014; Sorokowska, Sorokowski, Hummel, & Huanca, 2013). While these studies provide preliminary insights into what natural human variation in olfactory ability might be like, comparing data collected in the field to those in the lab challenges interpretation given the unknown variables in the field influencing test outcomes. Thus, we know little about how humans detect, discriminate, and identify odors in mixtures in natural settings—for humans, natural settings include the built environment, both urban and rural.

Lab data provide insights into smelling odor mixtures. First, while we may detect many to most odors in a complex odor mixture, the whole of that information is not transduced to the brain due to the inhibition of some olfactory receptor cells via receptor competition (Jinks & Laing, 1999). Even if all odors in the mix are subliminally processed in the cerebrum, only some are perceived (Hummel, Olgun, Gerber, Huchel, & Frasnelli, 2013). Second, there may be an upper limit to the number of odors detectable individually from mixes—one experiment suggested that more than 30 equal-intensity odorants are perceived as an “olfactory white” (Weiss et al., 2012). Third, the ability to detect single odors in mixtures may be improved by experience (Sinding et al., 2015) and training (Barkat, Le Berre, Coureaud, Sicard, & Thomas-Danguin, 2012; Morquecho-Campos, Larsson, Boesveldt, & Olofsson, 2019). Fourth, there may be ecological pressures that have acted on the filtering mechanisms of the olfactory system. There is a neuronal pattern that suggests we are particularly tuned to food odors (Saraiva et al., 2019). Thus, the olfactory system is limited by its ability to integrate complex whole mixtures into perception and may be driven by attention to specific compounds. Perhaps environmental variation outside the lab is not a disruptive confounding factor. By widening data collection to field-based settings, a representative sample would be possible and allow us to understand the extent of natural human variation in olfactory ability.

The principal challenge in generating ecologically salient data is the lack of field-based methods (Kern, Wroblewski, Schumm, Pinto, & McClintock, 2014). The problem with modifying a lab-based test for deployment in the field was first noted in the discipline of anthropology during the ethnographic fieldwork in the Torres Straits at the turn of the century (Myers, 1901). The only field-validated method is the Olfactory Function Field Exam (OFEE), an 11-minute olfactory function test that uses a modified and abbreviated version of Sniffin'

Sticks—two odors for threshold detection (Kern et al., 2014) and five odors for identification (Mueller & Renner, 2006). As part of The National Social Life, Health, and Aging Project in the United States, the OFFE was validated on a cohort of 2304 older adults and conducted in participants' homes following a short odor training session. Results were nearly identical to testing conducted in labs on a similarly aged cohort but not using repeated measures on the same subjects (Kern et al., 2014). OFFE validation results suggest that between-subjects environmental variation was not significant. Olfactory adaptation down-regulates receptor stimulation when odors are constant, however, which means that OFFE subjects were not smelling their home environments and the effect from variation would be small (Auffarth, 2013; Dalton, 2000).

Our goal was to assess the impact on olfactory ability from a dynamic environment to which subjects were not adapted. We needed a tool that was highly portable and easy to administer in the field. The ease of administering the two most accurate tests, threshold and discrimination, in the field without a table was prohibitive. That left odor identification. There are three validated short tests of olfactory ability that rely solely on odor identification—a five odor Sniffin' Sticks test (Mueller & Renner, 2006), a three odor Sniffin' Sticks test (Hummel, Pfetzing, & Lötsch, 2010), and a three-odor test using microencapsulation (Jackman & Doty, 2005). We used the five-item test in two repeated measures studies: a field study comparing olfactory ability in a low odor environment to: (a) a polluted environment and (b) a high odor environment; a lab study comparing olfactory ability in a low odor control lab to a high odor experiment condition.

2 | MATERIALS AND METHODS

2.1 | Materials

Our five-item test included four of the original five-item test odors: peppermint, orange, rose, and fish (Mueller & Renner, 2006). Leather was replaced by garlic because it had a lower identification rate in the UK (Neumann et al., 2012). The field study was conducted in August 2018 and the lab study in February 2019. Materials were purchased in July 2018 and stored at the University of Roehampton at a temperature within the recommended manufacturer's range.

2.2 | Screening and testing protocols

Testers wore cotton gloves during testing and wore no scented products. Each pen containing an odor was moved across the nostrils slowly for 3 seconds. Participants were

asked to describe the odor via free response then choose one of 22 terms from a verbal prompt card: five odor terms (one correct per odor), 15 distractor odor terms (none correct for any odor), “undefinable”, and “no odor” (see table 2 of Mueller & Renner, 2006). The field study consisted of three tests per participant and the lab study consisted of two. Between-test gaps were small (25–30 minutes for field, 15–20 minutes for lab).

2.3 | Data coding and statistical analysis

Data are shared at <https://doi.org/10.6084/m9.figshare.11764272.v4> under the CC BY 4.0 license. All data were analyzed and graphics plotted in R for Windows v3.6.1 (Charif & Lobry, 2007) using the R Studio GUI v1.2.1335 (RStudio Team, 2015). All scripts are shared at <https://doi.org/10.6084/m9.figshare.11933193>

Responses were coded as correct, near miss (a related item like lime for orange), far miss (a more general term like citrus for orange), and incorrect (an unrelated item like raspberry for orange) following Cain (1979). Between-coder agreement (JCB, KCH) was tested by Cohen’s kappa (κ) using *irr* (Gamer, Lemon, Fellows, & Singh, 2012) in the *lpsolve* package (Berkelaar, 2015). The total correct provided an ordinal level variable describing olfactory ability (Mueller & Renner, 2006):

- Poor (0): severely impaired or no sense of smell
- Indeterminate (1–3): further testing required
- Normal (4–5 correct): normal or slightly reduced sense of smell

The effect on olfactory ability from differing environments was estimated using ordinal regression (AKA logistic regression and cumulative link models), specifically, *clmm* in the ordinal package (Christensen, 2019). Logistic regression offers a powerful alternative to traditional ordinal tests (eg, Mann-Whitney *U* test, Wilcoxon signed rank sums, Kruskal Wallis) because it allows the estimation of fixed effects on the dependent variable and estimation of repeated measures via a random term (McCullagh, 1980)—an ordinal version of mixed-model linear regression or repeated measures ANOVA. We used the proportional odds model which assesses the probability of being in *x* ranked category or lower vs being in any higher ranked category with olfactory rank as the thresholds between categories. We included additional fixed effects—sex because females perform better in lab studies (Oliveira-Pinto et al., 2014; Sorokowski et al., 2019) and age because olfactory ability declines after age 50 in lab studies (Martinez et al., 2017; Charlotte Sinding, Puschmann, & Hummel, 2014). The field study had two additional effects, day and time of testing,

due to potential temporal variation in test environments. The proportional odds effects plots were generated in the effects package (Fox & Hong, 2009) using output from *polr* in MASS (Venables & Ripley, 2002).

Odor identification is the hardest Sniffin’ Sticks task due to challenges in odor term recall and lack of social consensus in odor naming; for example, the association of cloves with the dentist or lemon and rose with cleaning supplies may result in marking an identification incorrect even though the participant has actually identified the odor but not used the veridical (or true) label (e.g., orange) for orange rather than citrus or fruity (Cornell Kärnekull, Jönsson, Willander, Sikström, & Larsson, 2015; Jönsson & Stevenson, 2016). Thus, we also analyzed the data for internal consistency in odor term (whether correct, near miss, far miss, incorrect) via an ordinal ranked variable with a rank of 0 (inconsistent terms across trials) or 1 (consistent terms across trials). If an individual could not smell the odor in any trial, the data were excluded. An olfactory ability score for each individual was derived from the sum of consistent responses per odor per person, Olfactory ability ranks are the same as above:

- Poor (0): severely impaired or no sense of smell
- Indeterminate (1–3): further testing required
- Normal (4–5 correct): normal or slightly reduced sense of smell

Differences between males and females were analyzed using the Mann-Whitney *U* test via *wilcox.test* in base R statistics.

2.4 | Ethical Considerations

Field research for this project was approved by the University of Roehampton Ethics Committee, approval number LSC 17/213 (JCB, KCH). Written informed consent included no further information beyond a name (signed and printed). Data collection sheets were labeled with a unique numeric ID. The unique numeric ID was not noted on the informed consent which means no personally identifying information was linked to test data. The data collection sheet included basic demographic information (sex, age, ethnicity) and free and verbal prompt responses to each of five odors per location.

3 | RESULTS

3.1 | Coding

The initial coding of descriptive data by two observers (JCB, KCH) was in moderate agreement: Repeat Markets

(free response $\kappa = 0.537$, verbal prompt $\kappa = 0.500$) and Repeat Labs (free response $\kappa = 0.418$, verbal prompt $\kappa = 0.676$). After discussion of discrepancies, we agreed to code only veridical terms as correct (even if given with other terms), unrelated food types and qualitative responses as incorrect rather than far miss, onion as near miss for garlic, and mint as correct for peppermint. After recording, coders were in perfect agreement ($\kappa = 1$).

3.2 | Field study

For the field study ($n = 29$), we used social media (eg, Twitter, meetup.com), blogs (smellofevolution.com), word of mouth, and direct solicitation for participants. Testing was conducted on two separate days with two groups per day (10 AM and 2 PM). We used the area by the Tate Turbine Hall as a low odor control (low pollution, no food vendors, leafy area) for comparison to Southwark Bridge pedestrian area (traffic pollution, urine, cigarettes, and some food odors) and Borough Market (raw and cooked food vendors). We expected pollution to lower scores at Southwark and the odor signal-to-noise ratio from food odors at Borough to lower scores. The logistic regression indicates no significant effect from location (Table 1). A visualization of the change in the proportional odds (eg, probability of being poor vs normal if you are indeterminate) show the opposite trend to what was expected. The odds of being “poor” and “indeterminate” decrease from Tate to the other locations (1 = Tate, 2 = Southwark, 3 = Borough) while the odds of being “normal” increase (Figure 1).

The only significant fixed effect is within-subjects. The difference between a model with and without the within-subjects term is significant (Likelihood Ratio statistic = 16.6, $df = 1$, p -value = 0.000045). Most individuals vary from the sample average predicted response (set to 0) taking into account the estimated impact from fixed effects (Figure 2). While the significant random term in the regression indicates that individual variation impacts the model, it does not indicate if individuals vary in their scores across locations. Since there are no confounding factors from fixed effects, we used a paired ranks test to determine if there were significant changes in olfactory ability by individual (Table 2). There were no significant changes by individual for olfactory ability but there are two findings worth noting. First, the environmental impact on male olfactory ability is variable with some improving and some declining. So, while the summary statistics suggest no change, within-subjects change is taking place. Second, the environmental impact on female olfactory ability is consistent—all females improve at Borough (a dynamic food odor environment).

3.3 | Lab study

For Repeat Labs ($n = 30$), we recruited University of Roehampton students and staff to participate in a lab-based experiment. Testing was conducted in two labs at the same time each day over 3 days. Data for olfactory ability in a control lab (well-ventilated low-odor) were compared to those taken in an experimental condition (food market odors). Odors (garlic, coffee, madras curry

TABLE 1 Ordinal regression results

	AIC	Within subjects var	Between subjects	Est	Std err	z	p	Odds ratio
Field	126.45	6.51	Southwark	0.782	0.447	1.750	0.080	2.187
			Borough	−0.005	0.416	−0.010	0.990	0.995
			Male	−1.007	0.633	−1.590	0.110	0.365
			50+	−0.573	0.770	−0.740	0.460	0.564
			Day2	0.354	0.655	0.540	0.590	1.424
			PM	−0.729	0.652	−1.120	0.260	0.483
Intercepts			Norm:Indet	−0.711	0.702	−1.010		0.491
			Indet:Poor	6.921	1.932	3.580		1013.000
Lab	97.68	3.22	Location	−0.068	0.324	−0.210	0.830	0.934
			Male	−0.400	0.519	−0.770	0.440	0.670
			50+	−0.260	0.678	−0.380	0.700	0.771
			Intercepts	0.604	0.711	0.850		1.830
			Indet:Poor	4.430	1.362	3.250		83.940

FIGURE 1 Proportional odds of changes in olfactory ability due to the environment, field study




FIGURE 2 Individual effect on olfactory ability (conditional modes with 95% confidence intervals based on individual variance across locations), field study




TABLE 2 Sign test results

Subset	Contrast	<i>S</i>	<i>p</i> value	95%CI	
				Lower	Upper
All	Tate:Borough	3	0.100	0.857	0.951
Female	Tate:Borough	0	0.060	0.854	0.961
Male	Tate:Borough	3	1.000	0.854	0.961
All	Tate:Southwark	3	0.500	0.901	0.957
Female	Tate:Southwark	0	0.100	0.922	0.973
Male	Tate:Southwark	3	1.000	0.820	0.961
All	Control:Experimental	6	1.000	0.939	0.976
Female	Control:Experimental	5	0.700	0.939	0.976
Male	Control:Experimental	1	1.000	0.857	0.951

paste, and caramel furanone [3% concentration]) were diffused each day 1 hour prior to testing. We expected olfactory ability to be negatively impacted by the experimental condition but this was not the case (Table 1). The odds of dropping in olfactory ability rank under experimental conditions (Figure 3) is slight and not significant.

Again, the only significant model term is within-subjects. The difference between a model with and without the within-subjects term is significant (Likelihood Ratio statistic = 3.74, $df = 1$, p -value = 0.053). All individuals differ from the average predicted response to varying degrees (Figure 4). The paired ranks test did not return a




FIGURE 3 Proportional odds of changes in olfactory ability due to the environment, lab study

significant result but the direction of change is variable across the sexes—while females tend to improve more than decline, the positive directionality of the field study is not replicated.

3.4 | Field and Lab combined

Despite the lack of significant differences among locations, the overall trend across both studies is that the control lab provided the best environment for olfactory ability (Figure 5). In addition, females consistently outperformed males (Figure 5) with median and modal scores of 3 (normal) compared to male median and modal scores of 2 (indeterminate). Females captured a proportionately larger share of the normosmic diagnosis:




FIGURE 4 Individual effect on olfactory ability (conditional modes with 95% confidence intervals based on individual variance across locations), lab study




FIGURE 5 Proportional odds of changes in olfactory ability due to environment and sex




FIGURE 6 Variation in odor label consistency by sex

38% of results across trials were female and normosmic but only 14% were male and normosmic. While the field settings have a much lower probability of normosmia, females have a higher probability of normosmia and a very low probability of anosmia, compared to males.

3.5 | Consistency

There were no significant differences in olfactory ability when using the consistency scores in either study: Field ($W = 103$, $p\text{-value} = 0.7$, $CI: -0.999\text{--}2.00$) or Lab ($W = 169$, $p\text{-value} = 1$, $CI: -1\text{--}1$). In other words, males and females are not significantly different in olfactory ability scores when responses are rated based on consistency in use of terms (eg, chewing gum for orange across all trials), rather than correct identification of odor using veridical terms (eg, orange for orange). Some odors were more consistently described than others (Figure 6). Orange was the least consistently described odor for both males and females in 'field and lab' studies (but males in the lab had an equally difficult time describing garlic consistently). Fish was the most consistently described odor for females in the Field and peppermint in the Lab.

Garlic was the most consistently described odor for males in the Field (with peppermint a close second) and fish and garlic the most consistently described odor for males in the Lab.

For the Field study and individuals who used a consistent term two of three times, we examined which of the three locations had the most inconsistent terms (eg, consistency at Tate and Southwark but not Borough). Females were most inconsistent terms at Tate ($n = 18$, 66% of the time) compared to Southwark ($n = 4$, 15% of the time) and Borough ($n = 5$, 19% of the time). Males exhibited no pattern: Tate ($n = 0$, 39% of the time), Southwark ($n = 7$, 30.5% of the time), and Borough ($n = 7$, 30.5% of the time).

4 | DISCUSSION

Our main goal was to conduct preliminary data collection on how humans smell in the dynamic odorscapes of everyday living—how well are they able to recognize ecological salient odors (in our study, food) in a variety of settings. The difference between published lab-based olfactory ability data and the limited published field-



based olfactory ability data indicates a large effect size introduced by the non-lab environment on odor sensitivity and discrimination test scores (Sorokowska et al., 2013; Sorokowska et al., 2014; Sorokowska, Sorokowski, & Frackowiak, 2015). Even though we were using odor identification for our study, those data provided a target sample size for our study. Our study differs in one other important methodological issue—we used a repeated measures design and the other study did not. We failed to find an effect from the environment on odor identification ability. While the comparison of results across three settings failed to find a significant effect from environment on olfactory ability, the data visualization suggested it was better in the lab than the experimental setting or the three outdoor locations. Given the previously discussed problems with odor identification, perhaps our sample was too small for this particular test. We would tentatively argue that comparing lab-based data to field-based data is complicated.

Because there is no tool for mobile rapid collection of olfactory ability, our other goal was to test the use of a clinically validated five-item odor identification test—even if only to provide a rough measure of olfactory ability. Previously identified methodological problems with odor identification affected the outcomes. First, we found that participants were learning—they relied on verbal prompt terms after the first test, even when incorrect (choosing raspberry for orange consistently) and ignored the card in subsequent tests (“same as before but stronger,” “same as last time”). Perhaps longer between-test gaps would mitigate the effect but would require greater participant commitment to the process. Second, observer coding of odor terms introduces subjectivity to the process of marking answers as correct, near miss, far miss, or incorrect (Kern et al., 2014). Our first test for inter-observer agreement suggested the two coders were in moderate agreement on marking. Our discussion revealed where the key differences lied and, once we agree on marking terms, we were in agreement. So, while there was negotiation to come to agreement, there was subjectivity that makes replication challenging in any assessment of odor identification (Rabin & Cain, 1984). Data on social consensus for odor terms and usage frequency of those terms would help mitigate marking correct choices as incorrect—such as knowing that a population associates eugenol with the dentist, rose or lemon with cleaning supplies, or peppermint and orange with chewing gum—but those data are not routinely collected. Third, using an alternate approach to code ability based on the consistent use of odor term labels across trials (Rabin & Cain, 1984) resulted in a different outcome than the primary method of assessment using veridical terms—the sex-based differences in olfactory ability were

not found when considering consistency. The difference in results is easily explained by clinical data that suggest females are much better at odor identification tasks (Doty, 2001)—this is true even when males and females are asked to remember odor terms in a foreign language (Dempsey & Stevenson, 2002). Namely, the female advantage in naming is removed when consistency is the primary rubric. In our study, females who are not consistent across three trials but are consistent across two trials tend to be consistent after first exposure to the verbal prompt—they have learned the answer.

Do our results and interpretation suggest the odor identification test is not accurate? No, the test is more likely providing information about a different cognitive process. The cerebellar subregion involved in higher-order cognitive processing for language and memory—key areas for odor labelling (ie, language)—is associated with olfactory identification but not detection or discrimination (the more reliable tests of olfactory ability) (Wabnegger & Schienle, 2019). The differences we found when focusing on correct source identification as compared to consistency in recognizing the same odor (whether correct or not) indicate the importance of the analytical approach relative to the research questions. We are interested in functional olfactory ability, not language or symbolism or consistency in labelling an odor. The difference is simple: being consistent in using an odor label or having an odor rich cosmology does not indicate a functional olfactory ability but naming the source does. For example, consistently labelling mercaptan (the odor added to natural gas) as eggy or smoky three times in a row will not result in a functional behavioral output in the presence of that smell in the environment, namely to take evasive action. The correct identification of mercaptan as the smell of natural gas will lead to functional behavioral output. Thus, analysis of odor identification labels that focus on correct answers (naming the source) assesses ability to identify and respond to environmental information that is functionally important.

Clinical studies on western societies tend to explain sex-based differences in olfactory ability as derived from either gender-based labor or sex-based biological differences (Dempsey & Stevenson, 2002). There certainly are sex-based differences in biological structures. First, females have roughly double the number of olfactory bulb cells, both glial cells and neurons (Oliveira-Pinto et al., 2014). Second, females have a stronger olfactory modality than men (Dalton, Doolittle, & Breslin, 2002; Doty, Applebaum, Zusho, & Settle, 1985; Doty & Cameron, 2009; Doty et al., 1984; Oliveira-Pinto et al., 2014), even when accounting for different cultural settings in industrialized societies (Doty et al., 1985). This does not



appear to be based on hormonal differences because male-to-female transsexuals taking gender-affirmation hormones do not exhibit the same pattern (Kranz, Kaufmann, & Lanzenberger, 2019). There are also differences in experience that may act on female olfactory ability. Women are more interested than men in smell (Seo et al., 2010), are better than men at learning odors (Doty et al., 1984; Ferdenzi et al., 2013; Larsson, 1997; Larsson, Lövdén, & Nilsson, 2003; Lehrner, 1993), and more proficient than men at odor identification tasks (Oberg, Larsson, & Backman, 2002). The most interesting question that emerges is whether there might be an evolutionary explanation for sex-based differences in olfactory ability.

A gendered division of labor describes most of human prehistory and, even if in a modified form, contemporary societies. So, by focusing specifically on differences between men and women in food acquisition in daily tasks, we may gain some insights into either behavioral or biological selective pressures acting on female olfactory ability. Turning to our closest relatives, female chimpanzees forage alone but in spaces that overlap with other females, which generates some competition for space and food resources. High-ranking females tend to have preferential access to resource-rich sites which correlates with greater reproductive success (Pusey & Schroepfer-Walker, 2013). So, reproduction is, in part, determined by foraging ability in chimpanzees. Females also more frequently use tools for food foraging and processing, a trait shared with our closest relatives, chimps and bonobos (Gruber, Clay, & Zuberbühler, 2010; Pruett et al., 2015), which suggests selective pressure from activity predates the *Homo-Pan* split. Studies of hunting ability in hunter-gatherers that have included a sensory component (vision and audition) indicate that neither plays a strong role but strength does, especially upper body strength (Apicella, 2014; Stibbard-Hawkes, Attenborough, & Marlowe, 2018). So, despite a few ethnographic studies that have focused on the rich odor symbolism of some hunter-gatherer societies (for an overview, see Classen et al., 1994) and a study in Malay hunter-gatherers that focused on the increased prevalence of abstract odor terms in hunter-gatherers (Majid & Kruspe, 2018), there is no information on the functional role of olfaction in hunting, let alone other activities. Further the significance of odor terms is complex—despite the increased use of abstract odor terms, smell is the least codable sense in the Malay and, compared to other populations, olfaction is relatively low in their sensory hierarchy (Majid et al., 2018). There is, however, information on the functional role of olfaction in foraging activities. Women outperform men at foraging tasks that involve spatial memory and navigation skills (McBurney, Gaulin, Devineni, & Adams, 1997; New, Krasnow Max,

Truxaw, & Gaulin Steven, 2007) and the spatial encoding of gatherable foods into memory is chronically active in women but only active under explicit motivation in men (Krasnow et al., 2011). An ability to map food resources and navigate a catchment area is adaptively advantageous. Finally, as noted above, females have a greater density of olfactory neurons than males—when coupled with the fact human olfactory sensory neurons are evolutionarily biased towards food-related odors (Saraiva et al., 2019), superior olfactory ability may well be linked to food foraging. The main takeaway is that females appear to have better signal-to-noise ability when detecting, identifying, and discriminating among food odors, a skill that provides an adaptive advantage in successful foraging and group provisioning—such as seen in our data when female olfactory ability was enhanced in the rich food-odor environment of Borough Market.

In conclusion and despite published data indicating a significant environmental effect on olfactory ability (Sorokowska et al., 2013; Sorokowska et al., 2014), we failed to detect any significant differences in two repeated measures studies. We suspect that either within-population repeated measures tests have a more subtle impact from environment on olfactory ability than cross-cultural comparisons or that the comparison of field- to lab-collected data overemphasized differences. We also found that odor identification is a less than ideal tool for assessing olfactory ability despite ease of delivery and speedy test time. We argue that field-based methods are needed to characterize representative variation in human olfactory ability and the importance of ecological context. Lastly, we find that even in the field and despite methodological problems, females exhibit better olfactory ability and that may be an artifact of long-standing evolutionary pressure acting on a mosaic of traits related to food foraging. This last point is important and highlights the need for field-collected data (ie, non-lab settings), which may seem superfluous and lacking utility to those accustomed to lab environments. The reason is straight forward: the clinically-derived dataset is insufficient to capture variation in and the breadth and context of human olfaction. Indeed, compared to the traditional lab environment, a virtual reality lab environment found that behavioral responses to odors is context dependent—in the absence of ecological validity, motivated behavior is not observed (de Groot et al., 2020). Field-based data collection will widen participation beyond current olfactory ability datasets that are comprised of nonrepresentative samples. Why nonrepresentative? First, WEIRD samples are biased by individuals who can afford the costs of travel and time to be tested in a lab. Second, WEIRD populations (or even simply EIRD) are a global minority. Third, many populations are geographically isolated from

lab-testing sites—this describes the majority of indigenous, traditionally living, small-scale societies. Fourth, lab conditions cannot be recreated in the field. Clinical data serve their purposes but field data are needed to address an entirely different question: are meaningful biological differences in olfactory ability explained by ecological context? This study is one of a small number that use contemporary validated methods of assessing olfactory ability in non-lab settings. We know very little about the ecological or evolutionary pressures acting on human olfactory ability and if we are to truly understand the extent of and variation in olfactory ability, we need better field methods and cross-cultural data as well as transdisciplinary efforts.

ACKNOWLEDGEMENTS

Roehampton University staff and students for their support.

CONFLICT OF INTERESTS

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

K.C.H. and J.C.B. conceived and designed all studies. K.C.H., J.C.B., D.B., P.F. conducted field-based data collection. K.C.H., J.C.B., S.A. conducted lab-based data collection. K.C.H. analyzed data and drafted the manuscript. J.C.B. revised the manuscript critically for analytical approaches and intellectual content.

ORCID

Kara C. Hoover  <https://orcid.org/0000-0001-6394-930X>

REFERENCES

- Apicella, C. L. (2014). Upper-body strength predicts hunting reputation and reproductive success in Hadza hunter-gatherers. *Evolution and Human Behavior*, 35(6), 508–518. <https://doi.org/10.1016/j.evolhumbehav.2014.07.001>
- Auffarth, B. (2013). Understanding smell—The olfactory stimulus problem. *Neuroscience and Biobehavioral Reviews*, 37(8), 1667–1679. <https://doi.org/10.1016/j.neubiorev.2013.06.009>
- Barkat, S., Le Berre, E., Coureaud, G., Sicard, G., & Thomas-Danguin, T. (2012). Perceptual blending in odor mixtures depends on the nature of odorants and human olfactory expertise. *Chemical Senses*, 37(2), 159–166. <https://doi.org/10.1093/chemse/bjr086>
- Berkelaar, M. (2015). *lpSolve: Interface to 'Lp_solve' v. 5.5 to Solve Linear/Integer Programs*. R package version 5.6.13. <https://CRAN.R-project.org/package=lpSolve>
- Cain, W. (1979). To know with the nose: Keys to odor identification. *Science*, 203(4379), 467–470. <https://doi.org/10.1126/science.760202>
- Charif, D., & Lobry, J. R. (2007). 1.0-2: A contributed package to the R project for statistical computing devoted to biological sequences retrieval and analysis. In U. Bastolla, M. Porto, H. E. H. E. Roman, & M. Vendruscolo (Eds.), *Structural approaches to sequence evolution: Molecules, networks, populations* (pp. 207–232). New York: Springer Verlag.
- Christensen, R. H. B. (2019). Ordinal – regression models for ordinal data. R Package Version 2019.3–9. <http://www.cran.r-project.org/package=ordinal/>
- Classen, C. (1993). *Worlds of sense: Exploring the senses in history and across cultures*. London: Routledge.
- Classen, C., Howes, D., & Synnott, A. (1994). *Aroma: The cultural history of smell*. London: Routledge.
- Cornell Kärnekull, S., Jönsson, F. U., Willander, J., Sikström, S., & Larsson, M. (2015). Long-term memory for odors: Influences of familiarity and identification across 64 days. *Chemical Senses*, 40(4), 259–267. <https://doi.org/10.1093/chemse/bjv003>
- Dalton, P. (2000). Psychophysical and behavioral characteristics of olfactory adaptation. *Chemical Senses*, 25(4), 487–492.
- Dalton, P., Doolittle, N., & Breslin, P. A. S. (2002). Gender-specific induction of enhanced sensitivity to odors. *Nature Neuroscience*, 5(3), 199–200. doi:http://www.nature.com/neuro/journal/v5/n3/supinfo/n0803_S1.html
- Dempsey, R. A., & Stevenson, R. J. (2002). Gender differences in the retention of Swahili names for unfamiliar odors. *Chemical Senses*, 27(8), 681–689. <https://doi.org/10.1093/chemse/27.8.681>
- Doty, R., Shaman, P., Applebaum, S., Giberson, R., Sikorski, L., & Rosenberg, L. (1984). Smell identification ability: Changes with age. *Science*, 226(4681), 1441–1443. <https://doi.org/10.1126/science.6505700>
- Doty, R. L. (2001). Olfaction. *Annual Review of Psychology*, 52(1), 423–452. <https://doi.org/10.1146/annurev.psych.52.1.423>
- Doty, R. L., Applebaum, S., Zusho, H., & Settle, R. G. (1985). Sex differences in odor identification ability: A cross-cultural analysis. *Neuropsychologia*, 23(5), 667–672. doi:[https://doi.org/10.1016/0028-3932\(85\)90067-3](https://doi.org/10.1016/0028-3932(85)90067-3)
- Doty, R. L., & Cameron, E. L. (2009). Sex differences and reproductive hormone influences on human odor perception. *Physiology and Behavior*, 97(2), 213–228. <https://doi.org/10.1016/j.physbeh.2009.02.032>
- Ferdenzi, C., Roberts, S. C., Schirmer, A., Delplanque, S., Cekic, S., Porcherot, C., ... Grandjean, D. (2013). Variability of affective responses to odors: Culture, gender, and olfactory knowledge. *Chemical Senses*, 38(2), 175–186. <https://doi.org/10.1093/chemse/bjs083>
- Fox, J., & Hong, J. (2009). *Effect displays in R for multinomial and proportional-odds logit models: Extensions to the effects package*. 2009, 32(1), 24. doi:<https://doi.org/10.18637/jss.v032.i01>
- Gamer, M., Lemon, J., Fellows, I., & Singh, P. (2012). *irr: Various coefficients of interrater reliability and agreement. R package version 0.84*. <https://CRAN.R-project.org/package=irr>
- Groot, J. H. B., Beetsma, D. J. V., van Aerts, T. J. A., le Berre, E., Gallagher, D., Shaw, E., ... Smeets, M. A. M. (2020). From sterile labs to rich VR: Immersive multisensory context critical for odors to induce motivated cleaning behavior. *Behav Res*. <https://doi.org/10.3758/s13428-019-01341-y>
- Gruber, T., Clay, Z., & Zuberbühler, K. (2010). A comparison of bonobo and chimpanzee tool use: Evidence for a female bias in the Pan lineage. *Animal Behaviour*, 80(6), 1023–1033. <https://doi.org/10.1016/j.anbehav.2010.09.005>
- Hanel, P. H. P., & Vione, K. C. (2016). Do student samples provide an accurate estimate of the general public? *PLoS One*, 11(12), e0168354. <https://doi.org/10.1371/journal.pone.0168354>

- Henrich, J., Heine, S. J., & Norenzayan, A. (2010). Most people are not WEIRD. *Nature*, 466(7302), 29. <https://doi.org/10.1038/466029a>
- Hummel, T., Olgun, S., Gerber, J., Huchel, U., & Frasnelli, J. (2013). Brain responses to odor mixtures with sub-threshold components. *Frontiers in Psychology*, 4, 786. <https://doi.org/10.3389/fpsyg.2013.00786>
- Hummel, T., Pftzting, U., & Lötsch, J. (2010). A short olfactory test based on the identification of three odors. *Journal of Neurology*, 257(8), 1316–1321. <https://doi.org/10.1007/s00415-010-5516-5>
- Jackman, A. H., & Doty, R. L. (2005). Utility of a three-item smell identification test in detecting olfactory dysfunction. *Laryngoscope*, 115(12), 2209–2212. <https://doi.org/10.1097/01.mlg.0000183194.17484.bb>
- Jinks, A., & Laing, D. G. (1999). A limit in the processing of components in odour mixtures. *Perception*, 28(3), 395–404. <https://doi.org/10.1068/p2898>
- Jönsson, F. U., & Stevenson, R. J. (2016). Odor knowledge, odor naming, and the “tip-of-the-nose” experience. In B. L. Schwartz (Ed.), *Tip-of-the-tongue states and related phenomena* (pp. 305–326). Cambridge UK: Cambridge University Press.
- Kern, D. W., Wroblewski, K. E., Schumm, L. P., Pinto, J. M., & McClintock, M. K. (2014). Field survey measures of olfaction: The olfactory function field exam (OFFE). *Field Methods*, 26(4), 421–434. <https://doi.org/10.1177/1525822x14547499>
- Kranz, G. S., Kaufmann, U., & Lanzenberger, R. (2019). Probing the impact of gender-affirming hormone treatment on odor perception. *Chemical Senses*, 45(1), 37–44. <https://doi.org/10.1093/chemse/bjz069>
- Krasnow, M. M., Truxaw, D., Gaulin, S. J. C., New, J., Ozono, H., Uono, S., ... Minemoto, K. (2011). Cognitive adaptations for gathering-related navigation in humans. *Evolution and Human Behavior: Official Journal of the Human Behavior and Evolution Society*, 32(1), 1–12. <https://doi.org/10.1016/j.evolhumbehav.2010.07.003>
- Larsson, M. (1997). Semantic factors in episodic recognition of common Odors in early and late adulthood: A review. *Chemical Senses*, 22(6), 623–633. <https://doi.org/10.1093/chemse/22.6.623>
- Larsson, M., Lövdén, M., & Nilsson, L.-G. (2003). Sex differences in recollective experience for olfactory and verbal information. *Acta Psychologica*, 112(1), 89–103. [https://doi.org/10.1016/S0001-6918\(02\)00092-6](https://doi.org/10.1016/S0001-6918(02)00092-6)
- Lehrner, J. P. (1993). Gender differences in long-term odor recognition memory: Verbal versus sensory influences and the consistency of label use. *Chemical Senses*, 18(1), 17–26. <https://doi.org/10.1093/chemse/18.1.17>
- Majid, A., & Kruspe, N. (2018). Hunter-gatherer olfaction is special. *Current Biology*, 28(3), 409–413.e402. <https://doi.org/10.1016/j.cub.2017.12.014>
- Majid, A., Roberts, S. G., Cilissen, L., Emmorey, K., Nicodemus, B., O'Grady, L., ... Levinson, S. C. (2018). Differential coding of perception in the world's languages. *Proceedings of the National Academy of Sciences*, 115(45), 11369–11376. <https://doi.org/10.1073/pnas.1720419115>
- Martinez, B., Karunanayaka, P., Wang, J., Tobia, M. J., Vasavada, M., Eslinger, P. J., & Yang, Q. X. (2017). Different patterns of age-related central olfactory decline in men and women as quantified by olfactory fMRI. *Oncotarget*, 8(45), 79212–79222. <https://doi.org/10.18632/oncotarget.16977>
- McBurney, D. H., Gaulin, S. J. C., Devineni, T., & Adams, C. (1997). Superior spatial memory of women: Stronger evidence for the gathering hypothesis. *Evolution and Human Behavior*, 18(3), 165–174. [https://doi.org/10.1016/S1090-5138\(97\)00001-9](https://doi.org/10.1016/S1090-5138(97)00001-9)
- McCullagh, P. (1980). Regression models for ordinal data. *Journal of the Royal Statistical Society. Series B (Methodological)*, 42(2), 109–142.
- Morquecho-Campos, P., Larsson, M., Boesveldt, S., & Olofsson, J. K. (2019). Achieving olfactory expertise: Training for transfer in odor identification. *Chemical Senses*, 44(3), 197–203. <https://doi.org/10.1093/chemse/bjz007>
- Mueller, C., & Renner, B. (2006). A new procedure for the short screening of olfactory function using five items from the “Sniffin’ sticks” identification test kit. *American Journal of Rhinology and Allergy*, 20(1), 113–116.
- Myers, C. S. (1901). Smell. In A. C. Haddon, W. H. R. Rivers, C. G. Seligman, C. S. Myers, W. McDougall, S. H. Ray, & A. Wilkin (Eds.), *Reports of the Cambridge anthropological expedition to torres straits. Vol. II: Physiology and psychology* (pp. 169–185). Cambridge: Cambridge University Press.
- Neumann, C., Tsioulos, K., Merkonidis, C., Salam, M., Clark, A., & Philpott, C. (2012). Validation study of the ‘Sniffin’ sticks’ olfactory test in a British population: A preliminary communication. *Clinical Otolaryngology*, 37(1), 23–27. <https://doi.org/10.1111/j.1749-4486.2012.02431.x>
- New, J., Krasnow Max, M., Truxaw, D., & Gaulin Steven, J. C. (2007). Spatial adaptations for plant foraging: Women excel and calories count. *Proceedings of the Royal Society B: Biological Sciences*, 274(1626), 2679–2684. <https://doi.org/10.1098/rspb.2007.0826>
- Oberg, C., Larsson, M., & Backman, L. (2002). Differential sex effects in olfactory functioning: The role of verbal processing. *Journal of the International Neuropsychological Society*, 8(5), 691–698.
- Oliveira-Pinto, A. V., Santos, R. M., Coutinho, R. A., Oliveira, L. M., Santos, G. B., Alho, A. T. L., ... Lent, R. (2014). Sexual dimorphism in the human olfactory bulb: Females have more neurons and glial cells than males. *PLoS One*, 9(11), e111733. <https://doi.org/10.1371/journal.pone.0111733>
- Pruetz, J. D., Bertolani, P., Ontl, K. B., Lindshield, S., Shelley, M., & Wessling, E. G. (2015). New evidence on the tool-assisted hunting exhibited by chimpanzees (*Pan troglodytes verus*) in a savannah habitat at Fongoli, Sénégal. *Royal Society Open Science*, 2(4), 140507. <https://doi.org/10.1098/rsos.140507>
- Pusey, A. E., & Schroepfer-Walker, K. (2013). Female competition in chimpanzees. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 368(1631), 20130077–20130077. <https://doi.org/10.1098/rstb.2013.0077>
- Rabin, M. D., & Cain, W. S. (1984). Odor recognition: Familiarity, identifiability, and encoding consistency. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 10(2), 316–325. <https://doi.org/10.1037/0278-7393.10.2.316>
- RStudio Team. (2015). *RStudio: Integrated development for R*. Boston, MA: RStudio, Inc.
- Saraiva, L. R., Riveros-McKay, F., Mezzavilla, M., Abou-Moussa, E. H., Arayata, C. J., Makhoulouf, M., ... Logan, D. W. (2019). A transcriptomic atlas of mammalian olfactory mucosae reveals an evolutionary influence on food odor detection in

- humans. *Science Advances*, 5(7), eaax0396. <https://doi.org/10.1126/sciadv.aax0396>
- Seo, H.-S., Guarneros, M., Hudson, R., Distel, H., Min, B.-C., Kang, J.-K., ... Hummel, T. (2010). Attitudes toward olfaction: A cross-regional study. *Chemical Senses*, 36(2), 177–187. <https://doi.org/10.1093/chemse/bjq112>
- Sinding, C., Coureaud, G., Bervialle, B., Martin, C., Schaal, B., & Thomas-Danguin, T. (2015). Experience shapes our odor perception but depends on the initial perceptual processing of the stimulus. *Attention, Perception, & Psychophysics*, 77(5), 1794–1806. <https://doi.org/10.3758/s13414-015-0883-8>
- Sinding, C., Puschmann, L., & Hummel, T. (2014). Is the age-related loss in olfactory sensitivity similar for light and heavy molecules? *Chemical Senses*, 39(5), 383–390. <https://doi.org/10.1093/chemse/bju004>
- Sorokowska, A., Sorokowski, P., & Frackowiak, T. (2015). Determinants of human olfactory performance: A cross-cultural study. *Science of the Total Environment*, 506–507(0), 196–200. <https://doi.org/10.1016/j.scitotenv.2014.11.027>
- Sorokowska, A., Sorokowski, P., & Hummel, T. (2014). Cross-cultural administration of an odor discrimination test. *Chemosensory Perception*, 7(2), 85–90. <https://doi.org/10.1007/s12078-014-9169-0>
- Sorokowska, A., Sorokowski, P., Hummel, T., & Huanca, T. (2013). Olfaction and environment: Tsimane' of Bolivian rainforest have lower threshold of odor detection than industrialized German people. *PLoS One*, 8(7), e69203. <https://doi.org/10.1371/journal.pone.0069203>
- Sorokowski, P., Karwowski, M., Misiak, M., Marczak, M. K., Dziekan, M., Hummel, T., & Sorokowska, A. (2019). Sex Differences in Human Olfaction: A Meta-Analysis. *Front Psychol*, 10 (242). <https://doi.org/10.3389/fpsyg.2019.00242>
- Stibbard-Hawkes, D. N. E., Attenborough, R. D., & Marlowe, F. W. (2018). A noisy signal: To what extent are Hadza hunting reputations predictive of actual hunting skills? *Evolution and Human Behavior*, 39(6), 639–651. <https://doi.org/10.1016/j.evolhumbehav.2018.06.005>
- Venables, W. N., & Ripley, B. D. (2002). *Modern applied statistics with S*. New York: Springer.
- Wabnegger, A., & Schienle, A. (2019). Cerebellar grey matter and olfactory performance. *Chemical Senses*, 44, 507–510. <https://doi.org/10.1093/chemse/bjz045>
- Weiss, T., Snitz, K., Yablonka, A., Khan, R. M., Gafsou, D., Schneidman, E., & Sobel, N. (2012). Perceptual convergence of multi-component mixtures in olfaction implies an olfactory white. *Proceedings of the National Academy of Sciences of the United States of America*, 109(49), 19959–19964. <https://doi.org/10.1073/pnas.1208110109>

How to cite this article: Hoover KC, Botescu D, Fedurek P, Aarts S, Berbesque JC. Field-testing olfactory ability to understand human olfactory ecology. *Am J Hum Biol*. 2020;e23411. <https://doi.org/10.1002/ajhb.23411>