

Propositional Logic (Part 2)

ECE 650
Methods & Tools for Software Engineering (MTSE)
Fall 2023

Presented by
Dr. Albert Wasef

Used by permission from Prof. Arie Gurfinkel




References

Chapter 1 of Logic for Computer Scientists

- https://link.springer.com/chapter/10.1007/978-0-8176-4763-6_2

Library link:

- https://link-springer-com.proxy.lib.uwaterloo.ca/chapter/10.1007/978-0-8176-4763-6_2



Book Example: Secret to long life

"What is the secret of your long life?" a centenarian was asked.

"I strictly follow my diet: If I don't drink beer for dinner, then I always have fish. Any time I have both beer and fish for dinner, then I do without ice cream. If I have ice cream or don't have beer, then I never eat fish."

The questioner found this answer rather confusing.
Can you simplify it?



SINCE 1828

JOIN MWU

GAMES

BROWSE THESAURUS

WORD OF THE DAY

WORDS AT PLAY

normal form

DICTIONARY

THESAURUS

normal form noun

Definition of *normal form*

logic

: a canonical or standard fundamental form of a statement to which others can be reduced

especially : a compound statement in the propositional calculus consisting of nothing but a conjunction of disjunctions whose disjuncts are either elementary statements or negations thereof

<https://www.merriam-webster.com/dictionary/normal%20form>

Negation Normal Form (NNF)

A formula F is in **Negation Normal Form** (NNF) if every occurrence of negation (\neg) in F is applied to an atomic sub-formula of F .

For example,

- $\neg a \vee \neg b \vee c$ is in NNF
- $\neg (a \wedge b \wedge \neg c)$ is NOT in NNF (why?)

Theorem (NNF): For every formula F there is a semantically equivalent form G in NNF. In symbols

- For every F , exists G in NNF, such that $F \equiv G$

Proof of NNF Theorem

1/2

By **structural induction** on the structure of a formula F

(base case): atomic formulas A and $\neg A$ are in NNF

(IH): Assume that the theorem is true for every sub-formula of F : For every sub-formula H of F there exists J in NNF such that $J \equiv H$. Show that there exists G in NNF such that $G \equiv F$

(\neg case): see next slide

(\vee case): Assume $F = H_1 \vee H_2$. Then, $F \equiv J_1 \vee J_2$

(\wedge case): Assume $F = H_1 \wedge H_2$. Then, $F \equiv J_1 \wedge J_2$

Proof of NNF Theorem

2/2

By **structural induction** on the structure of a formula F

(base case): atomic formulas A and $\neg A$ are in NNF

(IH): Assume that the theorem is true for every sub-formula of F : For every sub-formula H of F there exists J in NNF such that $J \equiv H$. Show that there exists G in NNF such that $G \equiv F$

(\neg case):

(case 1) $F = \neg(\neg H_0) = \neg\neg H_0 \equiv H_0 \equiv J_0$

(case 2) $F = \neg(H_0 \wedge H_1) \equiv \neg H_0 \vee \neg H_1 \equiv J_0 \vee J_1$

(case 3) $F = \neg(H_0 \vee H_1) \equiv \neg H_0 \wedge \neg H_1 \equiv J_0 \wedge J_1$

- where J_0 is NNF of $\neg H_0$, and J_1 is NNF of $\neg H_1$ by IH

Normal Form: DNF

A *literal* is either an atomic proposition v or its negation $\neg v$

A *cube* is a conjunction of literals

- e.g., $(v1 \wedge \neg v2 \wedge v3)$

A formula F is in *Disjunctive Normal Form* (DNF) if F is a disjunction of conjunctions of literals

$$\bigvee_{i=1}^n \left(\bigwedge_{j=1}^{m_i} L_{i,j} \right)$$

(Fun) Fact: determining whether a DNF formula F is satisfiable is easy

- easy == linear in the size of the formula

Normal Form: CNF

A *literal* is either an atomic proposition v or its negation $\neg v$

A *clause* is a disjunction of literals

- e.g., $(v1 \vee \neg v2 \vee v3)$

A formula F is in *Conjunctive Normal Form* (CNF) if F is a conjunction of disjunctions of literals

$$\bigwedge_{i=1}^n \left(\bigvee_{j=1}^{m_i} L_{i,j} \right)$$

(Fun) Fact: determining whether a CNF formula F is satisfiable is hard

- hard == NP-complete

Normal Form Theorem

Theorem: For every formula F , there is an equivalent formula F_1 in CNF, and an equivalent formula F_2 in DNF.

That is, CNF and DNF are normal forms:

- Every propositional formula can be converted to CNF and to DNF without affecting its meaning (i.e., semantics)!

Proof: (by induction on the structure of the formula F)

Details are left as an exercise!

Converting a formula to CNF

Given a formula F

1. Substitute in F every occurrence of a sub-formula of the form

$\neg\neg G$ by G

$\neg(G \wedge H)$ by $(\neg G \vee \neg H)$

$\neg(G \vee H)$ by $(\neg G \wedge \neg H)$

The result is a formula in Negation Normal Form (NNF)

2. Substitute in F each occurrence of a sub-formula of the form

$(F \vee (G \wedge H))$ by $((F \vee G) \wedge (F \vee H))$

$((F \wedge G) \vee H)$ by $((F \vee H) \wedge (G \vee H))$

The resulting formula F is in CNF

- the result in CNF might be exponentially bigger than original formula F

Example: From Truth Table to CNF and DNF

DNF

$$\begin{aligned} &(\neg A \wedge \neg B \wedge \neg C) \vee \\ &(A \wedge \neg B \wedge \neg C) \vee \\ &(A \wedge \neg B \wedge C) \end{aligned}$$

CNF

$$\begin{aligned} &(A \vee B \vee \neg C) \wedge \\ &(A \vee \neg B \vee C) \wedge \\ &(A \vee \neg B \vee \neg C) \wedge \\ &(\neg A \vee \neg B \vee C) \wedge \\ &(\neg A \vee \neg B \vee \neg C) \end{aligned}$$

Truth table

| A | B | C | F |
|-----|-----|-----|-----|
| 0 | 0 | 0 | 1 |
| 0 | 0 | 1 | 0 |
| 0 | 1 | 0 | 0 |
| 0 | 1 | 1 | 0 |
| 1 | 0 | 0 | 1 |
| 1 | 0 | 1 | 1 |
| 1 | 1 | 0 | 0 |
| 1 | 1 | 1 | 0 |

From Truth Table to DNF / CNF

From a truth table T to DNF formula F

- For every row i of T with value 1, construct a conjunction F_i that characterizes the row. That is, F_i is a formula that is true **exactly** for the assignment of row i
- Let $F = \vee F_i$, then F is equivalent to T and is in DNF
- Fact: F has as many disjuncts as rows with value 1 in T
- Question: How big can F be? How small can F be?

From a truth table T to CNF formula G

- For every row i of T with value 0, construct a conjunction F_i that characterizes the row
- Let G_i be NNF of $\neg F_i$
- Let $G = \wedge G_i$, then G is equivalent to T and is in CNF
- Fact: F has as many disjuncts as rows with value 0 in T
- Question: How big can F be? How small can F be?

3-CNF Fragment

A formula F is in 3-CNF iff

- F is in CNF
- every clause of F has at most 3 literals

Theorem: Deciding whether a 3-CNF formula F is satisfiable is at least as hard as deciding satisfiability of an arbitrary CNF formula G

Proof: by effective *reduction* from CNF to 3-CNF

Let G be an arbitrary CNF formula. Replaced every clause of the form

$$(\ell_0 \vee \cdots \vee \ell_n)$$

with 3-literal clauses

$$(\ell_0 \vee b_0) \wedge (\neg b_0 \vee \ell_1 \vee b_1) \wedge \cdots \wedge (\neg b_{n-1} \vee \ell_n)$$

where $\{b_i\}$ are fresh atomic propositions not appearing in F

Complexity of 3-CNF Satisfiability

Theorem (Cook-Levin): The Boolean Satisfiability Problem is NP-complete

Consequences

- If a formula F is satisfiable, then there exists a certificate for satisfiability that can be checked in P (polynomial) time.
 - That is, checking solutions is easy
- Any other problem that has polynomial certificates is polynomial reducible to Boolean Satisfiability
 - That is, such problems can be solved by writing a loop-free program, compiling it to a Boolean circuit, and checking whether the circuit ever accepts some input
- **MANY MANY MANY OPTIMIZATION PROBLEMS ARE LIKE THAT**
- Boolean Satisfiability is easy iff $P = NP$
 - i.e., Boolean satisfiability today is a VERY VERY VERY HARD problem!

Boolean Satisfiability (CNF-SAT)

1/2



Let V be a set of variables



A *literal* is either a variable v in V or its negation $\neg v$



A *clause* is a disjunction of literals e.g., $(v_1 \vee \neg v_2 \vee v_3)$



A Boolean formula in *Conjunctive Normal Form* (CNF) is a conjunction of clauses

e.g., $(v_1 \vee \neg v_2) \wedge (v_3 \vee v_2)$

Boolean Satisfiability (CNF-SAT)

2/2



An *assignment* s of Boolean values to variables *satisfies* a clause c if it evaluates at least one literal in c to true



An assignment s *satisfies* a formula C in CNF if it satisfies every clause in C



Boolean Satisfiability Problem (CNF-SAT):

determine whether a given CNF C is satisfiable

Are the following CNFs SAT or UNSAT

Is CNF 1 satisfiable? (3 clauses)

- $\neg b$
- $\neg a \vee \neg b \vee \neg c$
- a
- SAT: $s(a) = \text{True}$; $s(b) = \text{False}$; $s(c) = \text{False}$

Is CNF 2 satisfiable? (4 clauses)

- $\neg b$
- $\neg a \vee b \vee \neg c$
- a
- $\neg a \vee c$
- UNSAT

Algorithms for SAT

SAT is NP-complete

- solution can be checked in polynomial time
- no polynomial algorithms for finding a solution are known

DPLL (Davis-Putnam-Logemann-Loveland, '60)

- smart enumeration of all possible SAT assignments
- worst-case EXPTIME
- alternate between deciding and propagating variable assignments

CDCL (GRASP '96, Chaff '01)

- conflict-driven clause learning
- extends DPLL with
 - smart data structures, backjumping, clause learning, heuristics, restarts...
- scales to millions of variables
- N. Een and N. Sörensson, "An Extensible SAT-solver", in SAT 2013.

(Optional) Background Reading: SAT

← →

http://cacm.acm.org/magazines/2009/8/34498-boolean-satisfiability-from-theoretical-h

Boolean Satisfiability: From ...

Find: currency

Previous Next

Options

TRUSTED INSIGHTS FOR COMPUTING'S LEADING PROFESSIONALS

ACM.org | Join ACM | About Communications | ACM Resources | Alerts & Feeds

SIGN IN

COMMUNICATIONS
OF THE
ACM

HOME | CURRENT ISSUE | NEWS | BLOGS | OPINION | RESEARCH | PRACTICE | CAREERS | MAGAZINE ARCHIVE

Home / Magazine Archive / August 2009 (Vol. 52, No. 8) / Boolean Satisfiability: From Theoretical Hardness... / Full Text

REVIEW ARTICLES

Boolean Satisfiability: From Theoretical Hardness to Practical Success


By Sharad Malik, Lintao Zhang
Communications of the ACM, Vol. 52 No. 8, Pages 76-82
10.1145/1536616.1536637
[Comments](#)

VIEW AS:

DL

DE

SHARE:



There are many practical situations where we need to satisfy several potentially conflicting constraints. Simple examples of this abound in daily life, for example, determining a schedule for a series of games that resolves the availability of players and venues, or finding a seating assignment at dinner consistent with various rules the host would like to impose. This also applies to applications in computing, for example, ensuring that a hardware/software system functions correctly with its overall behavior constrained by the behavior of its components and their composition, or finding a plan for a robot to reach a goal that is

SIGN IN for Full Access

User Name

Password

[» Forgot Password?](#)

[» Create an ACM Web Account](#)

SIGN IN

ARTICLE CONTENTS:


[Introduction](#)

[Boolean Satisfiability](#)

[Theoretical hardness: SAT and NP Completeness](#)

Some Experience with SAT Solving

Speed-up of 2012 solver over other solvers



from M. Vardi, <https://www.cs.rice.edu/~vardi/papers/highlights15.pdf>

SAT - Milestones

Problems impossible 10 years ago are trivial today

| year | Milestone |
|------|----------------------------------|
| 1960 | Davis-Putnam procedure |
| 1962 | Davis-Logeman-Loveland |
| 1984 | Binary Decision Diagrams |
| 1992 | DIMACS SAT challenge |
| 1994 | SATO: clause indexing |
| 1997 | GRASP: conflict clause learning |
| 1998 | Search Restarts |
| 2001 | zChaff: 2-watch literal, VSIDS |
| 2005 | Preprocessing techniques |
| 2007 | Phase caching |
| 2008 | Cache optimized indexing |
| 2009 | In-processing, clause management |
| 2010 | Blocked clause elimination |


Concept



**Millions of
variables from
HW designs**

2002

2010



Courtesy Daniel le Berre

ENCODING PROBLEMS TO SAT

Graph k-Coloring

Given a graph $G = (V, E)$, and a natural number $k > 0$ is it possible to assign colors to vertices of G such that no two adjacent vertices have the same color.


Formally:

- does there exist a function $f : V \rightarrow [0..k)$ such that
- for every edge (u, v) in E , $f(u) \neq f(v)$

Graph coloring for $k > 2$ is NP-complete

Problem: Encode k-coloring of G into CNF

- construct CNF C such that C is SAT iff G is k-colorable



***k*-coloring as CNF**

Let a Boolean variable $f_{v,i}$ denote that vertex v has color i

- if $f_{v,i}$ is true if and only if $f(v) = i$

Every vertex has at least one color

$$\bigvee_{0 \leq i < k} f_{v,i} \quad (v \in V)$$

No vertex is assigned two colors

$$\bigwedge_{0 \leq i < j < k} (\neg f_{v,i} \vee \neg f_{v,j}) \quad (v \in V)$$

No two adjacent vertices have the same color


$$\bigwedge_{0 \leq i < k} (\neg f_{v,i} \vee \neg f_{u,i}) \quad ((v, u) \in E)$$

Vertex Cover

Given a graph $G=(V,E)$. A vertex cover of G is a subset C of vertices in V such that every edge in E is incident to at least one vertex in C

see a4_encoding.pdf for details of reduction to CNF-SAT

- will be given together with assignment 4



USING A SAT SOLVER

DIMACS interface to a SAT Solver

Input:

- a CNF in DIMACS format

Output:

- SAT/UNSAT + satisfying assignment

We will use a SAT solver called MiniSAT

- available at <https://git.uwaterloo.ca/ece650-f23/minisat>
- written in C++
- use as a library in Assignment 4
- use via DIMACS interface today in class
- MiniSat examples:
 - <https://git.uwaterloo.ca/ece650-f23/minisat-example>

DIMACS CNF File Format

Textual format to represent CNF-SAT problems

```
c start with comments
c
c
p cnf 5 3
 1 -5 4 0
-1 5 3 4 0
-3 -4 0
```

Details

- comments start with **c**
- header line: **p cnf nbvar nbclauses**
 - **nbvar** is # of variables, **nbclauses** is # of clauses
- each clause is a sequence of distinct numbers terminating with **0**
 - **positive** numbers are variables, **negative** numbers are negations

MiniSat

MiniSat is one of the most famous modern SAT-solvers

- written in C++
- designed to be easily understandable and customizable
- many new SAT-solvers use MiniSAT as their base

Web page: <http://minisat.se/>

We will use a slightly updated version from:
<https://git.uwaterloo.ca/ece650-f23/minisat>

Good references for understanding SAT solving details

- MiniSat architecture: <http://minisat.se/downloads/MiniSat.pdf>
- Donald Knuth's SAT13 (also based on MiniSat)
 - <http://www-cs-faculty.stanford.edu/~knuth/programs/sat13.w>

<https://git.uwaterloo.ca/ece650-f23/minisat-example>

MINISAT EXAMPLES

PROPOSITIONAL RESOLUTION

From CNF to database of literals

Assume that all propositional formulas are converted to CNF

Each clause is determined by the set of literals

- e.g., $(a \vee b \vee \neg c)$ is same as $\{a, b, \neg c\}$

A CNF is a database (a set) of clauses

- $(a \vee b \vee \neg c) \wedge (c) \wedge (\neg b \wedge d)$ is represented as
- $\{ \{a, b, \neg c\}, \{c\}, \{\neg b, d\} \}$

Propositional Resolution

Let A be a clause of the form $C \vee p$

Let B be a clause of the form $D \vee \neg p$

Propositional Resolution:


A clause $(C \vee D)$ is a resolvent of A and B on pivot p

Propositional Resolution In Symbols

$$\text{Res}(\{C, p\}, \{D, \neg p\}) = \{C, D\}$$

Given two clauses $\{C, p\}$ and $\{D, \neg p\}$ that contain a literal p of different polarity, create a new clause by taking the union of literals in C and D

Notation: Inference Rule



Inference Rules

We express the evaluation rules as inference rules for our judgments.

The rules are also called **evaluation rules**.

An **inference rule**

$$\frac{F_1 \dots F_n}{G} \text{ where } H$$

defines a relation between judgments F_1, \dots, F_n and G .

- The judgments F_1, \dots, F_n are the **premises** of the rule;
- The judgment G is the **conclusion** of the rule;
- The formula H is called the **side condition** of the rule.

If $n=0$ the rule is called an **axiom**. In this case, the line separating premises and conclusion may be omitted.

Propositional Resolution Inference

Pivot

$$\frac{C \vee p \qquad D \vee \neg p}{C \vee D}$$

Resolvent

$$\text{Res}(\{C, p\}, \{D, \neg p\}) = \{C, D\}$$

Given two clauses $\{C, p\}$ and $\{D, \neg p\}$ that contain a literal p of different polarity, create a new clause by taking the union of literals in C and D

Resolution Lemma

Let F be a CNF formula.

Let R be a resolvent with pivot p of two clauses X and Y in F

Then, $F \cup \{ R \}$ is equivalent to F .

- That is, R is implied by F and adding it to F does not change the meaning of F

Proof:

Show that for any assignment M , $M \models F$ if and only if $M \models F \cup \{ R \}$

If $M \models F \cup \{ R \}$ then $M \models F$ is trivial.

Show that if $M \models \{X, Y\}$ then $M \models R$.

Two cases: (case 1) $M \models p$, (case 2) $M \models \neg p$

Resolution Theorem

Let F be a set of clauses

$$Res(F) = F \cup \{R \mid R \text{ is a resolvent of two clauses in } F\}$$

Define Res^n recursively as follows:

$$Res^0(F) = F$$

$$Res^{n+1}(F) = Res(Res^n(F)), \text{ for } n \geq 0$$

$$Res^*(F) = \bigcup_{n \geq 0} Res^n(F)$$

Resolution Theorem:

A CNF F is UNSAT iff $Res^*(F)$ contains an empty clause

Exercise from LCS

For the following set of clauses determine Res^n for $n=0, 1, 2$

$$A \vee \neg B \vee C$$

$$B \vee C$$

$$\neg A \vee C$$

$$B \vee \neg C$$

$$\neg C$$

Proof of the Resolution Theorem

1/3

(*Soundness*) By Resolution Lemma, F is equivalent to $\text{Res}^i(F)$ for any i .

Let n be such that $\text{Res}^{n+1}(F)$ contains an empty clause, but $\text{Res}^n(F)$ does not

- such n must exist because an empty clause was added at some point

Then, $\text{Res}^n(F)$ must contain two unit clauses L and $\neg L$

- because the only way to construct an empty clause is to resolve two unit clauses

Hence, F is UNSAT

- every clause added by resolution is implied (entailed) by F
- hence, $F \rightarrow L$ and $F \rightarrow \neg L$
- Therefore, $F \rightarrow (L \wedge \neg L)$, and $F \rightarrow \text{False}$

Proof of the Resolution Theorem

2/3

(Completeness) By **induction** on the number of different atomic propositions in F .

(base case) if F has 0 atomic propositions and has a clause, then F contains an empty clause

- empty clause is the only clause without any atomic propositions

Proof of the Resolution Theorem

3/3

(inductive case):

Assume F is UNSAT and F has atomic propositions A_1, \dots, A_{n+1}

Let F_0 be the result of replacing atomic proposition A_{n+1} by 0

Let F_1 be the result of replacing atomic proposition A_{n+1} by 1

Since F is UNSAT, so are F_0 and F_1

- e.g., if F_0 is SAT with assignment M , then extend M to $A_{n+1} \rightarrow 0, \dots$

By IH, both F_0 and F_1 derive an empty clause

- Hence, $\text{Res}^*(F)$ contains (A_{n+1}) (or empty clause) and $\text{Res}^*(F)$ contains $(\neg A_{n+1})$ (or empty clause)

Therefore, $\text{Res}^*(F)$ contains an empty clause!

Example for the last step of Pf of Res Theorem

$$F = (a) \wedge (\neg a \vee b) \wedge (\neg b \vee c) \wedge (\neg c)$$

$$F_0 = (a) \wedge (\neg a) \wedge (\neg c)$$

- $\text{Res}^*(F_0)$ contains an empty clause
- By following the same resolution steps in F , we show that $\text{Res}^*(F)$ contains the clause (b)

$$F_1 = (a) \wedge (c) \wedge (\neg c)$$

- $\text{Res}^*(F_1)$ contains an empty clause
- By following the same resolution steps in F , we show that $\text{Res}^*(F)$ contains the clause $(\neg b)$

Therefore, $\text{Res}^*(F)$ contains an empty clause!

END OF LECTURE 8

Proof System

$$P_1, \dots, P_n \vdash C$$

An inference rule is a tuple (P_1, \dots, P_n, C)

- where, P_1, \dots, P_n, C are formulas
- P_i are called **premises** and C is called a **conclusion**
- intuitively, the rule says that the conclusion is true if the premises are

A proof system \mathcal{P} is a collection of inference rules

A proof in a proof system \mathcal{P} is a tree (or a DAG) such that

- nodes are labeled by formulas
- for each node n , the tuple $(\text{parents}(n), n)$ is an inference rule in \mathcal{P}

Propositional Resolution as an Inference Rule


$$\frac{C \vee p \qquad D \vee \neg p}{C \vee D}$$

Propositional resolution is a **sound** inference rule

Proposition resolution **proof system** consists of a **single** propositional resolution **rule**

A Resolution Proof Example

A refutation of $\neg p \vee \neg q \vee r$, $p \vee r$, $q \vee r$, $\neg r$:



Another Resolution Pf Example

Show by resolution that the following CNF is UNSAT

$$\neg b \wedge (\neg a \vee b \vee \neg c) \wedge a \wedge (\neg a \vee c)$$

$$\begin{array}{c} \frac{\neg a \vee b \vee \neg c \quad a}{b \vee \neg c} \quad b \quad \frac{a \quad \neg a \vee c}{c} \\ \hline \neg c \quad c \\ \hline \perp \end{array}$$

Book: Exercise 33

Using resolution show that

$$A \wedge B \wedge C$$

is a consequence of

$$\neg A \vee B$$

$$\neg B \vee C$$

$$A \vee \neg C$$

$$A \vee B \vee C$$

Entailment and Derivation

A set of formulas F **entails** a set of formulas G iff every model of F and is a model of G

$$F \models G$$

A formula G is **derivable** from a formula F by a proof system P if there exists a proof whose leaves are labeled by formulas in F and the root is labeled by G

$$F \vdash_P G$$

Book: Exercise 33

Using resolution show that

$$A \wedge B \wedge C$$

is a consequence of

$$\neg A \vee B$$

$$\neg B \vee C$$

$$A \vee \neg C$$

$$A \vee B \vee C$$

Soundness and Completeness

A proof system **P** is **sound** iff

$$(F \vdash_P G) \implies (F \models G)$$

A proof system **P** is **complete** iff

$$(F \models G) \implies (F \vdash_P G)$$

PR: Soundness and Completeness

Theorem: Propositional resolution is sound and complete for propositional logic

Proof:

Follows immediately from the Resolution Theorem!

Exercise 34

Show using resolution that F is valid

$$F = (\neg B \wedge \neg C \wedge D) \vee (\neg B \wedge \neg D) \vee (C \wedge D) \vee B$$

$$\neg F = (B \vee C \vee \neg D) \wedge (B \vee D) \wedge (\neg C \vee \neg D) \wedge \neg B$$

Compactness Theorem

Theorem:

A (possibly infinite) set M of propositional formulas is satisfiable iff every finite subset of M is satisfiable.

Corollary:

A (possibly infinite) set M of propositional formulas is unsatisfiable iff there exists a finite subset U of M such that U is unsatisfiable

Proof:

- Section 1.4 in Logic for Computer Scientists by Uwe Schöning

Satisfiability and Unsatisfiability

Let F be a propositional formula (large)

Assume that F is satisfiable. What is a short proof / certificate to establish satisfiability without a doubt?

- provide a model. The model is linear in the size of the formula

Now, assume that F is unsatisfiable. What is a short proof / certificate to establish UNSATISFIABILITY without a doubt?

Is the following formula SAT or UNSAT? How do you explain your answer?

$$\neg b \wedge (\neg a \vee b \vee \neg c) \wedge a \wedge (\neg a \vee c)$$