

MATH 341 - LINEAR ALGEBRA

§4.1 - 4.6

Fall 2019

Pure mathematics is, in its way, the poetry of logical ideas.
- Albert Einstein

§1.3 VECTOR EQUATIONS

THEOREM (ALGEBRAIC PROPERTIES OF \mathbb{R}^n)

For all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c, d in \mathbb{R} :

1. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
2. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
3. $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$
4. $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$, where $-\mathbf{u}$ denotes $-1 \cdot \mathbf{u}$
5. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
6. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
7. $c(d\mathbf{u}) = (cd)\mathbf{u}$
8. $1 \cdot \mathbf{u} = \mathbf{u}$.

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*,

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below.

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. For each \mathbf{u} in V , there is a vector $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. For each \mathbf{u} in V , there is a vector $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6. The scalar multiple of \mathbf{u} by c , denoted by $c\mathbf{u}$ is in V

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. For each \mathbf{u} in V , there is a vector $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6. The scalar multiple of \mathbf{u} by c , denoted by $c\mathbf{u}$ is in V
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. For each \mathbf{u} in V , there is a vector $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6. The scalar multiple of \mathbf{u} by c , denoted by $c\mathbf{u}$ is in V
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. For each \mathbf{u} in V , there is a vector $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6. The scalar multiple of \mathbf{u} by c , denoted by $c\mathbf{u}$ is in V
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
9. $c(d\mathbf{u}) = (cd)\mathbf{u}$

§4.1 VECTOR SPACES AND SUBSPACES

A **vector space** is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *addition* and *scalar multiplication*, subject to the ten axioms (or rules) listed below. The axioms must hold for all vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and all scalars c and $d \in \mathbb{R}$.

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. For each \mathbf{u} in V , there is a vector $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6. The scalar multiple of \mathbf{u} by c , denoted by $c\mathbf{u}$ is in V
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
9. $c(d\mathbf{u}) = (cd)\mathbf{u}$
10. $1\mathbf{u} = \mathbf{u}$.

§4.1 VECTOR SPACES AND SUBSPACES

Facts: For each \mathbf{u} in V and scalar c in \mathbb{R} ,

§4.1 VECTOR SPACES AND SUBSPACES

Facts: For each \mathbf{u} in V and scalar c in \mathbb{R} ,

$$0 \cdot \mathbf{u} = \mathbf{0}$$

§4.1 VECTOR SPACES AND SUBSPACES

Facts: For each \mathbf{u} in V and scalar c in \mathbb{R} ,

$$0 \cdot \mathbf{u} = \mathbf{0}$$

$$c \cdot \mathbf{0} = \mathbf{0}$$

§4.1 VECTOR SPACES AND SUBSPACES

Facts: For each \mathbf{u} in V and scalar c in \mathbb{R} ,

$$0 \cdot \mathbf{u} = \mathbf{0}$$

$$c \cdot \mathbf{0} = \mathbf{0}$$

and

$$(-1) \cdot \mathbf{u} = -\mathbf{u}.$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: The space \mathbb{R}^n is a vector space, with the usual vector addition and scalar multiplication.

§4.1 VECTOR SPACES AND SUBSPACES

Example: The space \mathbb{R}^n is a vector space, with the usual vector addition and scalar multiplication.

PROOF.

Check that vector addition and scalar multiplication in \mathbb{R}^n satisfy the 10 vector space axioms. □

§4.1 VECTOR SPACES AND SUBSPACES

Example: The space \mathbb{R}^n is a vector space, with the usual vector addition and scalar multiplication.

PROOF.

Check that vector addition and scalar multiplication in \mathbb{R}^n satisfy the 10 vector space axioms. □

Much of our intuition about \mathbb{R}^n will carry over to other abstract vector spaces (i.e. we can picture general vector spaces much as we visualize \mathbb{R}^n).

§4.1 VECTOR SPACES AND SUBSPACES

For all vectors \mathbf{u}, \mathbf{v} and \mathbf{w} in V and all c, d in \mathbb{R} we have:

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. For each \mathbf{u} in V , there is a vector $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6. The scalar multiple of \mathbf{u} by c , denoted by $c\mathbf{u}$ is in V
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
9. $c(d\mathbf{u}) = (cd)\mathbf{u}$
10. $1\mathbf{u} = \mathbf{u}$.

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $M_{m \times n}$ be the set of all $m \times n$ matrices.

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $M_{m \times n}$ be the set of all $m \times n$ matrices.

$M_{m \times n}$ is a vector space with vector addition and scalar multiplication given by the usual matrix addition and scalar multiplication of matrices.

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $M_{m \times n}$ be the set of all $m \times n$ matrices.

$M_{m \times n}$ is a vector space with vector addition and scalar multiplication given by the usual matrix addition and scalar multiplication of matrices.

Warning! In this vector space each vector is actually an $m \times n$ matrix!

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $M_{m \times n}$ be the set of all $m \times n$ matrices.

$M_{m \times n}$ is a vector space with vector addition and scalar multiplication given by the usual matrix addition and scalar multiplication of matrices.

Warning! In this vector space each vector is actually an $m \times n$ matrix!

It shouldn't be too hard to see, that in some sense $M_{m \times n}$ is essentially the same vector space as $\mathbb{R}^{m \times n}$ (we'll make this precise later).

§4.1 VECTOR SPACES AND SUBSPACES

For all vectors \mathbf{u}, \mathbf{v} and \mathbf{w} in V and all c, d in \mathbb{R} we have:

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. For each \mathbf{u} in V , there is a vector $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6. The scalar multiple of \mathbf{u} by c , denoted by $c\mathbf{u}$ is in V
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
9. $c(d\mathbf{u}) = (cd)\mathbf{u}$
10. $1\mathbf{u} = \mathbf{u}$.

§4.1 VECTOR SPACES AND SUBSPACES

Example: Now consider M_n and for two $n \times n$ matrices A and B , define

$$A \oplus B = AB$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Now consider M_n and for two $n \times n$ matrices A and B , define

$$A \oplus B = AB$$

i.e $A \oplus B$ is just the product of A and B .

§4.1 VECTOR SPACES AND SUBSPACES

Example: Now consider M_n and for two $n \times n$ matrices A and B , define

$$A \oplus B = AB$$

i.e $A \oplus B$ is just the product of A and B .

Is M_n a vector space with addition defined by \oplus , and the usual scalar multiplication?

§4.1 VECTOR SPACES AND SUBSPACES

For all vectors \mathbf{u}, \mathbf{v} and \mathbf{w} in V and all c, d in \mathbb{R} we have:

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. For each \mathbf{u} in V , there is a vector $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6. The scalar multiple of \mathbf{u} by c , denoted by $c\mathbf{u}$ is in V
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
9. $c(d\mathbf{u}) = (cd)\mathbf{u}$
10. $1\mathbf{u} = \mathbf{u}$.

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let \mathbb{P}_n denote the set of all polynomials

$$\mathbf{p} = a_0 + a_1t + \cdots a_nt^n$$

of degree less than or equal to n .

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let \mathbb{P}_n denote the set of all polynomials

$$\mathbf{p} = a_0 + a_1t + \cdots a_nt^n$$

of degree less than or equal to n .

If $\mathbf{q} = b_0 + b_1t + \cdots b_nt^n$ and $c \in \mathbb{R}$, then vector addition and scalar multiplication are given by

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let \mathbb{P}_n denote the set of all polynomials

$$\mathbf{p} = a_0 + a_1t + \cdots a_nt^n$$

of degree less than or equal to n .

If $\mathbf{q} = b_0 + b_1t + \cdots b_nt^n$ and $c \in \mathbb{R}$, then vector addition and scalar multiplication are given by

$$\begin{aligned}\mathbf{p} + \mathbf{q} &= (a_0 + a_1t + \cdots a_nt^n) + (b_0 + b_1t + \cdots b_nt^n) \\ &= (a_0 + b_0) + (a_1 + b_1)t + \cdots (a_n + b_n)t^n\end{aligned}$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let \mathbb{P}_n denote the set of all polynomials

$$\mathbf{p} = a_0 + a_1t + \cdots a_nt^n$$

of degree less than or equal to n .

If $\mathbf{q} = b_0 + b_1t + \cdots b_nt^n$ and $c \in \mathbb{R}$, then vector addition and scalar multiplication are given by

$$\begin{aligned}\mathbf{p} + \mathbf{q} &= (a_0 + a_1t + \cdots a_nt^n) + (b_0 + b_1t + \cdots b_nt^n) \\ &= (a_0 + b_0) + (a_1 + b_1)t + \cdots (a_n + b_n)t^n\end{aligned}$$

and

$$\begin{aligned}c \cdot \mathbf{p} &= c \cdot (a_0 + a_1t + \cdots a_nt^n) \\ &= ca_0 + ca_1t + \cdots ca_nt^n.\end{aligned}$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let \mathbb{P}_n denote the set of all polynomials

$$\mathbf{p} = a_0 + a_1t + \cdots a_nt^n$$

of degree less than or equal to n .

If $\mathbf{q} = b_0 + b_1t + \cdots b_nt^n$ and $c \in \mathbb{R}$, then vector addition and scalar multiplication are given by

$$\begin{aligned}\mathbf{p} + \mathbf{q} &= (a_0 + a_1t + \cdots a_nt^n) + (b_0 + b_1t + \cdots b_nt^n) \\ &= (a_0 + b_0) + (a_1 + b_1)t + \cdots (a_n + b_n)t^n\end{aligned}$$

and

$$\begin{aligned}c \cdot \mathbf{p} &= c \cdot (a_0 + a_1t + \cdots a_nt^n) \\ &= ca_0 + ca_1t + \cdots ca_nt^n.\end{aligned}$$

Verify the axioms.

§4.1 VECTOR SPACES AND SUBSPACES

For all vectors \mathbf{u}, \mathbf{v} and \mathbf{w} in V and all c, d in \mathbb{R} we have:

1. The sum of \mathbf{u} and \mathbf{v} , denoted $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
4. There is a **zero** vector $\mathbf{0} \in V$, such that $\mathbf{u} + \mathbf{0} = \mathbf{u}$
5. For each \mathbf{u} in V , there is a vector $-\mathbf{u}$ such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
6. The scalar multiple of \mathbf{u} by c , denoted by $c\mathbf{u}$ is in V
7. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
8. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
9. $c(d\mathbf{u}) = (cd)\mathbf{u}$
10. $1\mathbf{u} = \mathbf{u}$.

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let \mathbb{S} denote the set of doubly infinite sequences

$$\{y_k\} = (\dots, y_{-2}, y_{-1}, y_0, y_1, y_2, \dots).$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let \mathbb{S} denote the set of doubly infinite sequences

$$\{y_k\} = (\dots, y_{-2}, y_{-1}, y_0, y_1, y_2, \dots).$$

If $\{y_k\}$ and $\{z_k\} \in \mathbb{S}$ and $c \in \mathbb{R}$, then

$$\{y_k\} + \{z_k\} = \{y_k + z_k\} = (\dots, y_{-1} + z_{-1}, y_0 + z_0, y_1 + z_1, \dots)$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let \mathbb{S} denote the set of doubly infinite sequences

$$\{y_k\} = (\dots, y_{-2}, y_{-1}, y_0, y_1, y_2, \dots).$$

If $\{y_k\}$ and $\{z_k\} \in \mathbb{S}$ and $c \in \mathbb{R}$, then

$$\{y_k\} + \{z_k\} = \{y_k + z_k\} = (\dots, y_{-1} + z_{-1}, y_0 + z_0, y_1 + z_1, \dots)$$

and

$$c \cdot \{y_k\} = (\dots, c \cdot y_{-1}, c \cdot y_0, c \cdot y_1, \dots).$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let \mathbb{S} denote the set of doubly infinite sequences

$$\{y_k\} = (\dots, y_{-2}, y_{-1}, y_0, y_1, y_2, \dots).$$

If $\{y_k\}$ and $\{z_k\} \in \mathbb{S}$ and $c \in \mathbb{R}$, then

$$\{y_k\} + \{z_k\} = \{y_k + z_k\} = (\dots, y_{-1} + z_{-1}, y_0 + z_0, y_1 + z_1, \dots)$$

and

$$c \cdot \{y_k\} = (\dots, c \cdot y_{-1}, c \cdot y_0, c \cdot y_1, \dots).$$

The zero vector and $-\{y_k\}$ are given by

$$\mathbf{0} = (\dots, 0, 0, 0, \dots) \quad \text{and} \quad -\{y_k\} = (\dots, -y_{-1}, -y_0, -y_1, \dots)$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $C([0, 1])$ denote the set of continuous functions $f : [0, 1] \rightarrow \mathbb{R}$.

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $C([0, 1])$ denote the set of continuous functions $f : [0, 1] \rightarrow \mathbb{R}$.

If $\mathbf{f}, \mathbf{g} \in C([0, 1])$ and $c \in \mathbb{R}$, then $\mathbf{f} + \mathbf{g}$ and $c \cdot \mathbf{f}$ can be defined by specifying the value these functions take at each point on $[0, 1]$:

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $C([0, 1])$ denote the set of continuous functions $f : [0, 1] \rightarrow \mathbb{R}$.

If $\mathbf{f}, \mathbf{g} \in C([0, 1])$ and $c \in \mathbb{R}$, then $\mathbf{f} + \mathbf{g}$ and $c \cdot \mathbf{f}$ can be defined by specifying the value these functions take at each point on $[0, 1]$:

$$(\mathbf{f} + \mathbf{g})(t) = \mathbf{f}(t) + \mathbf{g}(t) \quad \text{for all } t \in [0, 1],$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $C([0, 1])$ denote the set of continuous functions $f : [0, 1] \rightarrow \mathbb{R}$.

If $\mathbf{f}, \mathbf{g} \in C([0, 1])$ and $c \in \mathbb{R}$, then $\mathbf{f} + \mathbf{g}$ and $c \cdot \mathbf{f}$ can be defined by specifying the value these functions take at each point on $[0, 1]$:

$$(\mathbf{f} + \mathbf{g})(t) = \mathbf{f}(t) + \mathbf{g}(t) \quad \text{for all } t \in [0, 1],$$

and

$$(c \cdot \mathbf{f})(t) = c \cdot \mathbf{f}(t) \quad \text{for all } t \in [0, 1].$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $C([0, 1])$ denote the set of continuous functions $f : [0, 1] \rightarrow \mathbb{R}$.

If $\mathbf{f}, \mathbf{g} \in C([0, 1])$ and $c \in \mathbb{R}$, then $\mathbf{f} + \mathbf{g}$ and $c \cdot \mathbf{f}$ can be defined by specifying the value these functions take at each point on $[0, 1]$:

$$(\mathbf{f} + \mathbf{g})(t) = \mathbf{f}(t) + \mathbf{g}(t) \quad \text{for all } t \in [0, 1],$$

and

$$(c \cdot \mathbf{f})(t) = c \cdot \mathbf{f}(t) \quad \text{for all } t \in [0, 1].$$

The zero vector is given by

$$\mathbf{0}(t) = 0 \quad \text{for all } t \in [0, 1].$$

§4.1 VECTOR SPACES AND SUBSPACES

A **subspace** of a vector space V is a subset W of V which is a vector space itself (with the addition and scalar multiplication from V .)

§4.1 VECTOR SPACES AND SUBSPACES

A **subspace** of a vector space V is a subset W of V which is a vector space itself (with the addition and scalar multiplication from V .)

THEOREM

Suppose W is a subset of the vector space V , such that:

§4.1 VECTOR SPACES AND SUBSPACES

A **subspace** of a vector space V is a subset W of V which is a vector space itself (with the addition and scalar multiplication from V .)

THEOREM

Suppose W is a subset of the vector space V , such that:

1. The zero vector $\mathbf{0}$ is in W .

§4.1 VECTOR SPACES AND SUBSPACES

A **subspace** of a vector space V is a subset W of V which is a vector space itself (with the addition and scalar multiplication from V .)

THEOREM

Suppose W is a subset of the vector space V , such that:

1. The zero vector $\mathbf{0}$ is in W .
2. W is closed under vector addition. That is, for each \mathbf{u} and \mathbf{v} in W , the sum $\mathbf{u} + \mathbf{v}$ is in W .

§4.1 VECTOR SPACES AND SUBSPACES

A **subspace** of a vector space V is a subset W of V which is a vector space itself (with the addition and scalar multiplication from V .)

THEOREM

Suppose W is a subset of the vector space V , such that:

1. The zero vector $\mathbf{0}$ is in W .
2. W is closed under vector addition. That is, for each \mathbf{u} and \mathbf{v} in W , the sum $\mathbf{u} + \mathbf{v}$ is in W .
3. W is closed under multiplication by scalars. That is, for each \mathbf{u} in W and each scalar c , the vector $c \cdot \mathbf{u}$ is in W .

§4.1 VECTOR SPACES AND SUBSPACES

A **subspace** of a vector space V is a subset W of V which is a vector space itself (with the addition and scalar multiplication from V .)

THEOREM

Suppose W is a subset of the vector space V , such that:

1. The zero vector $\mathbf{0}$ is in W .
2. W is closed under vector addition. That is, for each \mathbf{u} and \mathbf{v} in W , the sum $\mathbf{u} + \mathbf{v}$ is in W .
3. W is closed under multiplication by scalars. That is, for each \mathbf{u} in W and each scalar c , the vector $c \cdot \mathbf{u}$ is in W .

Then W is a subspace of V .

§4.1 VECTOR SPACES AND SUBSPACES

A **subspace** of a vector space V is a subset W of V which is a vector space itself (with the addition and scalar multiplication from V .)

THEOREM

Suppose W is a subset of the vector space V , such that:

1. The zero vector $\mathbf{0}$ is in W .
2. W is closed under vector addition. That is, for each \mathbf{u} and \mathbf{v} in W , the sum $\mathbf{u} + \mathbf{v}$ is in W .
3. W is closed under multiplication by scalars. That is, for each \mathbf{u} in W and each scalar c , the vector $c \cdot \mathbf{u}$ is in W .

Then W is a subspace of V .

To tell if a subset of V is a subspace, it is enough to check the above three conditions

§4.1 VECTOR SPACES AND SUBSPACES

A **subspace** of a vector space V is a subset W of V which is a vector space itself (with the addition and scalar multiplication from V .)

THEOREM

Suppose W is a subset of the vector space V , such that:

1. The zero vector $\mathbf{0}$ is in W .
2. W is closed under vector addition. That is, for each \mathbf{u} and \mathbf{v} in W , the sum $\mathbf{u} + \mathbf{v}$ is in W .
3. W is closed under multiplication by scalars. That is, for each \mathbf{u} in W and each scalar c , the vector $c \cdot \mathbf{u}$ is in W .

Then W is a subspace of V .

To tell if a subset of V is a subspace, it is enough to check the above three conditions (all three must hold, if a single property fails then W is *not* a subspace of V).

§4.1 VECTOR SPACES AND SUBSPACES

Example: The subset $H = \{\mathbf{0}\}$ consisting of only the zero vector is always a subspace of V , and is called the *zero subspace*.

§4.1 VECTOR SPACES AND SUBSPACES


Example: The subset $H = \{\mathbf{0}\}$ consisting of only the zero vector is always a subspace of V , and is called the *zero subspace*.

Example: Is the line L given by $y = x + 2$ a subspace of \mathbb{R}^2 ?

§4.1 VECTOR SPACES AND SUBSPACES

Example: The subset $H = \{\mathbf{0}\}$ consisting of only the zero vector is always a subspace of V , and is called the *zero subspace*.


Example: Is the line L given by $y = x + 2$ a subspace of \mathbb{R}^2 ?



§4.1 VECTOR SPACES AND SUBSPACES

Example: The subset $H = \{\mathbf{0}\}$ consisting of only the zero vector is always a subspace of V , and is called the *zero subspace*.


Example: Is the line L given by $y = x + 2$ a subspace of \mathbb{R}^2 ?



§4.1 VECTOR SPACES AND SUBSPACES

Example: The subset $H = \{\mathbf{0}\}$ consisting of only the zero vector is always a subspace of V , and is called the *zero subspace*.


Example: Is the line L given by $y = x + 2$ a subspace of \mathbb{R}^2 ?



§4.1 VECTOR SPACES AND SUBSPACES

Example: The subset $H = \{\mathbf{0}\}$ consisting of only the zero vector is always a subspace of V , and is called the *zero subspace*.

Example: Is the line L given by $y = x + 2$ a subspace of \mathbb{R}^2 ?



Since L doesn't contain $\mathbf{0}$, and isn't closed under scalar multiplication and vector addition it isn't a subspace of \mathbb{R}^2 .

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $G \subset M_n$ be the set of all $n \times n$ matrices A such that $\det A = 0$. Does G form a subspace of M_n ?

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $G \subset M_n$ be the set of all $n \times n$ matrices A such that $\det A = 0$. Does G form a subspace of M_n ?

Example: Let $F \subset \mathbb{P}_3$ be the subset of all polynomials with integer coefficients. Is F a subspace?

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $G \subset M_n$ be the set of all $n \times n$ matrices A such that $\det A = 0$. Does G form a subspace of M_n ?

Example: Let $F \subset \mathbb{P}_3$ be the subset of all polynomials with integer coefficients. Is F a subspace?

Example: Let $J \subset C([0, 1])$ be the subset of all differentiable functions on $[0, 1]$. Is J a subspace?

§4.1 VECTOR SPACES AND SUBSPACES

Example: Let $G \subset M_n$ be the set of all $n \times n$ matrices A such that $\det A = 0$. Does G form a subspace of M_n ?

Example: Let $F \subset \mathbb{P}_3$ be the subset of all polynomials with integer coefficients. Is F a subspace?

Example: Let $J \subset C([0, 1])$ be the subset of all differentiable functions on $[0, 1]$. Is J a subspace?

Example: Let $H \subset C([0, 1])$ be the subset of all differentiable functions f on $[0, 1]$ with $f'(0.5) = 0$. Is H a subspace?

§4.1 VECTOR SPACES AND SUBSPACES

THEOREM

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are in a vector space V , then $\text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a subspace of V .

§4.1 VECTOR SPACES AND SUBSPACES

THEOREM

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are in a vector space V , then $\text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a subspace of V .

PROOF.

We must check the three subspace criteria:

§4.1 VECTOR SPACES AND SUBSPACES

THEOREM

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are in a vector space V , then $\text{span } \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a subspace of V .

PROOF.

We must check the three subspace criteria:

1. $\mathbf{0} = 0 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2 + \cdots + 0 \cdot \mathbf{v}_p \in \text{span } \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.

§4.1 VECTOR SPACES AND SUBSPACES

THEOREM

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are in a vector space V , then $\text{span } \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a subspace of V .

PROOF.

We must check the three subspace criteria:

1. $\mathbf{0} = 0 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2 + \dots + 0 \cdot \mathbf{v}_p \in \text{span } \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.
2. Let \mathbf{u} and $\mathbf{w} \in \text{span } \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$. Then

§4.1 VECTOR SPACES AND SUBSPACES

THEOREM

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are in a vector space V , then $\text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a subspace of V .

PROOF.

We must check the three subspace criteria:

1. $\mathbf{0} = 0 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2 + \cdots + 0 \cdot \mathbf{v}_p \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.
2. Let \mathbf{u} and $\mathbf{w} \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$. Then

$$\begin{aligned}\mathbf{u} + \mathbf{w} &= (u_1 \mathbf{v}_1 + \cdots + u_p \mathbf{v}_p) + (w_1 \mathbf{v}_1 + \cdots + w_p \mathbf{v}_p) \\ &= (u_1 + w_1) \mathbf{v}_1 + \cdots + (u_p + w_p) \mathbf{v}_p \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}.\end{aligned}$$

§4.1 VECTOR SPACES AND SUBSPACES

THEOREM

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are in a vector space V , then $\text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a subspace of V .

PROOF.

We must check the three subspace criteria:

1. $\mathbf{0} = 0 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2 + \dots + 0 \cdot \mathbf{v}_p \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.
2. Let \mathbf{u} and $\mathbf{w} \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$. Then

$$\begin{aligned}\mathbf{u} + \mathbf{w} &= (u_1 \mathbf{v}_1 + \dots + u_p \mathbf{v}_p) + (w_1 \mathbf{v}_1 + \dots + w_p \mathbf{v}_p) \\ &= (u_1 + w_1) \mathbf{v}_1 + \dots + (u_p + w_p) \mathbf{v}_p \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}.\end{aligned}$$

3. Let $\mathbf{u} \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ and $c \in \mathbb{R}$. Then

§4.1 VECTOR SPACES AND SUBSPACES

THEOREM

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are in a vector space V , then $\text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a subspace of V .

PROOF.

We must check the three subspace criteria:

1. $\mathbf{0} = 0 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2 + \cdots + 0 \cdot \mathbf{v}_p \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.
2. Let \mathbf{u} and $\mathbf{w} \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$. Then

$$\begin{aligned}\mathbf{u} + \mathbf{w} &= (u_1 \mathbf{v}_1 + \cdots + u_p \mathbf{v}_p) + (w_1 \mathbf{v}_1 + \cdots + w_p \mathbf{v}_p) \\ &= (u_1 + w_1) \mathbf{v}_1 + \cdots + (u_p + w_p) \mathbf{v}_p \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}.\end{aligned}$$

3. Let $\mathbf{u} \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ and $c \in \mathbb{R}$. Then

$$\begin{aligned}c \cdot \mathbf{u} &= c(u_1 \mathbf{v}_1 + \cdots + u_p \mathbf{v}_p) \\ &= cu_1 \mathbf{v}_1 + \cdots + cu_p \mathbf{v}_p \in \text{span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}.\end{aligned}$$



§4.1 VECTOR SPACES AND SUBSPACES

Example: Does the set H form a subspace?

$$H = \left\{ \begin{bmatrix} t + 2r \\ 3t - r \\ 5r \end{bmatrix} : \text{for } t, r \in \mathbb{R} \right\}$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Does the set H form a subspace?

$$H = \left\{ \begin{bmatrix} t + 2r \\ 3t - r \\ 5r \end{bmatrix} : \text{for } t, r \in \mathbb{R} \right\}$$

Since any vector $\mathbf{x} \in H$ splits as

$$\mathbf{x} = \begin{bmatrix} t + 2r \\ 3t - r \\ 5r \end{bmatrix} = t \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix} + r \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix},$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Does the set H form a subspace?

$$H = \left\{ \begin{bmatrix} t + 2r \\ 3t - r \\ 5r \end{bmatrix} : \text{for } t, r \in \mathbb{R} \right\}$$

Since any vector $\mathbf{x} \in H$ splits as

$$\mathbf{x} = \begin{bmatrix} t + 2r \\ 3t - r \\ 5r \end{bmatrix} = t \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix} + r \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix},$$

$$\implies H = \text{span} \left\{ \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix} \right\},$$

§4.1 VECTOR SPACES AND SUBSPACES

Example: Does the set H form a subspace?

$$H = \left\{ \begin{bmatrix} t + 2r \\ 3t - r \\ 5r \end{bmatrix} : \text{for } t, r \in \mathbb{R} \right\}$$

Since any vector $\mathbf{x} \in H$ splits as

$$\mathbf{x} = \begin{bmatrix} t + 2r \\ 3t - r \\ 5r \end{bmatrix} = t \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix} + r \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix},$$

$$\implies H = \text{span} \left\{ \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix} \right\},$$

and thus by the theorem, H is a subspace.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

The **null space** of an $m \times n$ matrix A , written as $\text{Nul } A$, is the set of all solutions of the homogeneous equation $A\mathbf{x} = \mathbf{0}$.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

The **null space** of an $m \times n$ matrix A , written as $\text{Nul } A$, is the set of all solutions of the homogeneous equation $A\mathbf{x} = \mathbf{0}$. In set notation,

$$\text{Nul } A = \{\mathbf{x} : \mathbf{x} \text{ is in } \mathbb{R}^n \text{ and } A\mathbf{x} = \mathbf{0}\}.$$

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

The **null space** of an $m \times n$ matrix A , written as $\text{Nul } A$, is the set of all solutions of the homogeneous equation $A\mathbf{x} = \mathbf{0}$. In set notation,

$$\text{Nul } A = \{\mathbf{x} : \mathbf{x} \text{ is in } \mathbb{R}^n \text{ and } A\mathbf{x} = \mathbf{0}\}.$$

THEOREM

The null space of an $m \times n$ matrix A is a subspace of \mathbb{R}^n .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

The **null space** of an $m \times n$ matrix A , written as $\text{Nul } A$, is the set of all solutions of the homogeneous equation $A\mathbf{x} = \mathbf{0}$. In set notation,

$$\text{Nul } A = \{\mathbf{x} : \mathbf{x} \text{ is in } \mathbb{R}^n \text{ and } A\mathbf{x} = \mathbf{0}\}.$$

THEOREM

The null space of an $m \times n$ matrix A is a subspace of \mathbb{R}^n .

In other words, the set of all solutions to a system $A\mathbf{x} = \mathbf{0}$ of m homogeneous linear equations in n unknowns is a subspace of \mathbb{R}^n .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Finding an explicit description of $\text{Nul } A$:

We can find an explicit description of $\text{Nul } A$ by solving the homogenous system $A\mathbf{x} = \mathbf{0}$ and expressing the solution in parametric form.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Finding an explicit description of $\text{Nul } A$:

We can find an explicit description of $\text{Nul } A$ by solving the homogenous system $A\mathbf{x} = \mathbf{0}$ and expressing the solution in parametric form.

Example: Find $\text{Nul } A$, where $A = \begin{bmatrix} -3 & 6 & -1 & 1 & -7 \\ 1 & -2 & 2 & 3 & -1 \\ 2 & -4 & 5 & 8 & -4 \end{bmatrix}$.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Finding an explicit description of $\text{Nul } A$:

We can find an explicit description of $\text{Nul } A$ by solving the homogenous system $A\mathbf{x} = \mathbf{0}$ and expressing the solution in parametric form.

Example: Find $\text{Nul } A$, where $A = \begin{bmatrix} -3 & 6 & -1 & 1 & -7 \\ 1 & -2 & 2 & 3 & -1 \\ 2 & -4 & 5 & 8 & -4 \end{bmatrix}$.

- The spanning vectors will be linearly independent (because of the free variables).

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Finding an explicit description of $\text{Nul } A$:

We can find an explicit description of $\text{Nul } A$ by solving the homogenous system $A\mathbf{x} = \mathbf{0}$ and expressing the solution in parametric form.

Example: Find $\text{Nul } A$, where $A = \begin{bmatrix} -3 & 6 & -1 & 1 & -7 \\ 1 & -2 & 2 & 3 & -1 \\ 2 & -4 & 5 & 8 & -4 \end{bmatrix}$.

- ▶ The spanning vectors will be linearly independent (because of the free variables).
- ▶ The number of vectors in the spanning set for $\text{Nul } A$ equals the number of free variables in the equation $A\mathbf{x} = \mathbf{0}$.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

The **column space** of an $m \times n$ matrix A , written as $\text{Col } A$ is the set of all linear combinations of the columns of A . If

$A = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \dots & \mathbf{a}_n \end{bmatrix}$, then

$$\text{Col } A = \text{Span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\}.$$

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

The **column space** of an $m \times n$ matrix A , written as $\text{Col } A$ is the set of all linear combinations of the columns of A . If

$A = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \dots & \mathbf{a}_n \end{bmatrix}$, then

$$\text{Col } A = \text{Span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\}.$$

THEOREM

The column space of an $m \times n$ matrix A is a subspace of \mathbb{R}^m .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Another way of describing $\text{Col } A$ is:

$$\text{Col } A = \{\mathbf{b} : \mathbf{b} = A\mathbf{x} \text{ for some } \mathbf{x} \text{ in } \mathbb{R}^n\}.$$

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Another way of describing $\text{Col } A$ is:

$$\text{Col } A = \{\mathbf{b} : \mathbf{b} = A\mathbf{x} \text{ for some } \mathbf{x} \text{ in } \mathbb{R}^n\}.$$

In other words, $\text{Col } A$ is the set of all vectors \mathbf{b} such that the equation $A\mathbf{x} = \mathbf{b}$ is consistent.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Another way of describing $\text{Col } A$ is:

$$\text{Col } A = \{\mathbf{b} : \mathbf{b} = A\mathbf{x} \text{ for some } \mathbf{x} \text{ in } \mathbb{R}^n\}.$$

In other words, $\text{Col } A$ is the set of all vectors \mathbf{b} such that the equation $A\mathbf{x} = \mathbf{b}$ is consistent.

In other other words, $\text{Col } A$ is the *range* of the linear transformation $T(\mathbf{x}) = A\mathbf{x}$.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Another way of describing $\text{Col } A$ is:

$$\text{Col } A = \{\mathbf{b} : \mathbf{b} = A\mathbf{x} \text{ for some } \mathbf{x} \text{ in } \mathbb{R}^n\}.$$

In other words, $\text{Col } A$ is the set of all vectors \mathbf{b} such that the equation $A\mathbf{x} = \mathbf{b}$ is consistent.

In other other words, $\text{Col } A$ is the *range* of the linear transformation $T(\mathbf{x}) = A\mathbf{x}$.

Warning! If A is an $m \times n$ matrix, then $\text{Col } A$ is a subspace of \mathbb{R}^m , while $\text{Nul } A$ is a subspace of \mathbb{R}^n .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Example: Let

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 1 & 2 & 5 & 0 \\ 2 & 4 & 9 & 1 \end{bmatrix}.$$

Find Nul A and Col A .

EXAMPLE (CONT.)

EXAMPLE (CONT.)

Let

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 1 & 2 & 5 & 0 \\ 2 & 4 & 9 & 2 \end{bmatrix}$$

1. Determine if $\begin{bmatrix} 1 \\ 2 \\ 2 \\ 1 \end{bmatrix}$ is in Nul A . Could it also be in Col A ?
2. Determine if $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$ is in Col A . Could it also be in Nul A ?

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Nul A	Col A
---------	---------

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Nul A	Col A
1. Nul A is a subspace of \mathbb{R}^n .	1. Col A is a subspace of \mathbb{R}^m .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Nul A	Col A
1. Nul A is a subspace of \mathbb{R}^n .	1. Col A is a subspace of \mathbb{R}^m .
2. Nul A is <i>implicitly</i> defined.	2. Col A is <i>explicitly</i> defined.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Nul A	Col A
1. Nul A is a subspace of \mathbb{R}^n .	1. Col A is a subspace of \mathbb{R}^m .
2. Nul A is <i>implicitly</i> defined.	2. Col A is <i>explicitly</i> defined.
3. It takes time to find vectors in Nul A .	3. It is easy to find vectors in Col A .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Nul A	Col A
1. Nul A is a subspace of \mathbb{R}^n .	1. Col A is a subspace of \mathbb{R}^m .
2. Nul A is <i>implicitly</i> defined.	2. Col A is <i>explicitly</i> defined.
3. It takes time to find vectors in Nul A .	3. It is easy to find vectors in Col A .
4. No clear relation between Nul A and entries of A .	4. Obvious relation between Col A and entries of A .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Nul A	Col A
1. Nul A is a subspace of \mathbb{R}^n .	1. Col A is a subspace of \mathbb{R}^m .
2. Nul A is <i>implicitly</i> defined.	2. Col A is <i>explicitly</i> defined.
3. It takes time to find vectors in Nul A .	3. It is easy to find vectors in Col A .
4. No clear relation between Nul A and entries of A .	4. Obvious relation between Col A and entries of A .
5. If \mathbf{v} in Nul A , then $A\mathbf{v} = \mathbf{0}$.	5. If \mathbf{v} in Col A , then $A\mathbf{x} = \mathbf{v}$ is consistent.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Nul A	Col A
---------	---------

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Nul A	Col A
6. Given \mathbf{v} simply compute $A\mathbf{v}$ to see if \mathbf{v} in Nul A .	6. Takes time to see if \mathbf{v} is in Col A .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Nul A	Col A
6. Given \mathbf{v} simply compute $A\mathbf{v}$ to see if \mathbf{v} in Nul A .	6. Takes time to see if \mathbf{v} is in Col A .
7. Nul $A = \{\mathbf{0}\}$ if and only if $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.	7. Col $A = \mathbb{R}^m$ if and only if $A\mathbf{x} = \mathbf{b}$ has a solution for every \mathbf{b} .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

Nul A	Col A
6. Given \mathbf{v} simply compute $A\mathbf{v}$ to see if \mathbf{v} in Nul A .	6. Takes time to see if \mathbf{v} is in Col A .
7. Nul $A = \{\mathbf{0}\}$ if and only if $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.	7. Col $A = \mathbb{R}^m$ if and only if $A\mathbf{x} = \mathbf{b}$ has a solution for every \mathbf{b} .
8. Nul $A = \{\mathbf{0}\}$ if and only if $\mathbf{x} \mapsto A\mathbf{x}$ is one-to-one.	8. Col $A = \mathbb{R}^m$ if and only if $\mathbf{x} \mapsto A\mathbf{x}$ is onto \mathbb{R}^m .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

A **linear transformation** T from a vector space V into a vector space W is a rule that assigns to each vector \mathbf{x} in V a unique vector $T(\mathbf{x})$ in W , such that

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

A **linear transformation** T from a vector space V into a vector space W is a rule that assigns to each vector \mathbf{x} in V a unique vector $T(\mathbf{x})$ in W , such that

1. $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ for all \mathbf{u}, \mathbf{v} in V , and

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

A **linear transformation** T from a vector space V into a vector space W is a rule that assigns to each vector \mathbf{x} in V a unique vector $T(\mathbf{x})$ in W , such that

1. $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ for all \mathbf{u}, \mathbf{v} in V , and
2. $T(c\mathbf{u}) = cT(\mathbf{u})$ for an \mathbf{u} in V and all scalars c .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

A **linear transformation** T from a vector space V into a vector space W is a rule that assigns to each vector \mathbf{x} in V a unique vector $T(\mathbf{x})$ in W , such that

1. $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ for all \mathbf{u}, \mathbf{v} in V , and
2. $T(c\mathbf{u}) = cT(\mathbf{u})$ for an \mathbf{u} in V and all scalars c .

The **kernel** (or **null space**) of such a T is the set of all \mathbf{u} in V such that $T(\mathbf{u}) = \mathbf{0}$ (the zero vector in V).

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

A **linear transformation** T from a vector space V into a vector space W is a rule that assigns to each vector \mathbf{x} in V a unique vector $T(\mathbf{x})$ in W , such that

1. $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ for all \mathbf{u}, \mathbf{v} in V , and
2. $T(c\mathbf{u}) = cT(\mathbf{u})$ for an \mathbf{u} in V and all scalars c .

The **kernel** (or **null space**) of such a T is the set of all \mathbf{u} in V such that $T(\mathbf{u}) = \mathbf{0}$ (the zero vector in V).

The **range** of T is the set of all vectors in W of the form $T(\mathbf{x})$ for some \mathbf{x} in V .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

A **linear transformation** T from a vector space V into a vector space W is a rule that assigns to each vector \mathbf{x} in V a unique vector $T(\mathbf{x})$ in W , such that

1. $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ for all \mathbf{u}, \mathbf{v} in V , and
2. $T(c\mathbf{u}) = cT(\mathbf{u})$ for an \mathbf{u} in V and all scalars c .

The **kernel** (or **null space**) of such a T is the set of all \mathbf{u} in V such that $T(\mathbf{u}) = \mathbf{0}$ (the zero vector in V).

The **range** of T is the set of all vectors in W of the form $T(\mathbf{x})$ for some \mathbf{x} in V .

Warning! Unlike linear transformations $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, linear transformation on arbitrary vector spaces may *not* be able to be expressed in terms of a standard matrix.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

THEOREM

If $T : V \rightarrow W$ is a linear transformation, then the kernel of T is a subspace of V , and the range of T is a subspace of W .

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

THEOREM

If $T : V \rightarrow W$ is a linear transformation, then the kernel of T is a subspace of V , and the range of T is a subspace of W .

Example: Let $W = C(\mathbb{R})$ (the space of all continuous functions on the real line), and V be the subspace of W consisting of functions with continuous derivative.

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

THEOREM

If $T : V \rightarrow W$ is a linear transformation, then the kernel of T is a subspace of V , and the range of T is a subspace of W .

Example: Let $W = C(\mathbb{R})$ (the space of all continuous functions on the real line), and V be the subspace of W consisting of functions with continuous derivative.

Define $D : V \rightarrow W$ by

$$D(f) = \frac{df}{dx}.$$

§4.2 NULL SPACES, COLUMN SPACES & LIN. TRANS.

THEOREM

If $T : V \rightarrow W$ is a linear transformation, then the kernel of T is a subspace of V , and the range of T is a subspace of W .

Example: Let $W = C(\mathbb{R})$ (the space of all continuous functions on the real line), and V be the subspace of W consisting of functions with continuous derivative.

Define $D : V \rightarrow W$ by

$$D(f) = \frac{df}{dx}.$$

Find the kernel (or null space) and range of D .

§4.3 LINEARLY INDEPENDENT SETS; BASES

An indexed set of vectors $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ in V is called **linearly independent** if the vector equation

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}$$

has *only* the trivial solution, $c_1 = 0, \dots, c_p = 0$.

§4.3 LINEARLY INDEPENDENT SETS; BASES

An indexed set of vectors $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ in V is called **linearly independent** if the vector equation

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}$$

has *only* the trivial solution, $c_1 = 0, \dots, c_p = 0$.

The set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is said to be **linearly dependent** if there is a nontrivial solution to the equation above. (This means there must be some weights, c_1, \dots, c_p not all zero, which make the equation hold).

§4.3 LINEARLY INDEPENDENT SETS; BASES

An indexed set of vectors $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ in V is called **linearly independent** if the vector equation

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}$$

has *only* the trivial solution, $c_1 = 0, \dots, c_p = 0$.

The set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is said to be **linearly dependent** if there is a nontrivial solution to the equation above. (This means there must be some weights, c_1, \dots, c_p not all zero, which make the equation hold).

In such a case, the equation above is called a **linear dependence relation** among $\mathbf{v}_1, \dots, \mathbf{v}_p$.

§4.3 LINEARLY INDEPENDENT SETS; BASES

The set $\{\mathbf{v}\}$ is **linearly independent** if and only if $\mathbf{v} \neq \mathbf{0}$.

§4.3 LINEARLY INDEPENDENT SETS; BASES

The set $\{\mathbf{v}\}$ is **linearly independent** if and only if $\mathbf{v} \neq \mathbf{0}$.

The set $\{\mathbf{v}_1, \mathbf{v}_2\}$ is **linearly dependent** if and only if either $\mathbf{v}_2 = c\mathbf{v}_1$ or $\mathbf{v}_1 = c\mathbf{v}_2$.

§4.3 LINEARLY INDEPENDENT SETS; BASES

The set $\{\mathbf{v}\}$ is **linearly independent** if and only if $\mathbf{v} \neq \mathbf{0}$.

The set $\{\mathbf{v}_1, \mathbf{v}_2\}$ is **linearly dependent** if and only if either $\mathbf{v}_2 = c\mathbf{v}_1$ or $\mathbf{v}_1 = c\mathbf{v}_2$.

Any set containing the zero vector $\mathbf{0}$ is **linearly dependent**.

§4.3 LINEARLY INDEPENDENT SETS; BASES

The set $\{\mathbf{v}\}$ is **linearly independent** if and only if $\mathbf{v} \neq \mathbf{0}$.

The set $\{\mathbf{v}_1, \mathbf{v}_2\}$ is **linearly dependent** if and only if either $\mathbf{v}_2 = c\mathbf{v}_1$ or $\mathbf{v}_1 = c\mathbf{v}_2$.

Any set containing the zero vector $\mathbf{0}$ is **linearly dependent**.

THEOREM

An indexed set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ of two or more vectors, with $\mathbf{v}_1 \neq \mathbf{0}$, is linearly dependent if and only if some \mathbf{v}_j (with $j > 0$) is a linear combination of the preceding vectors, v_1, \dots, v_{j-1} .

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: Let $\mathbf{p}_1 = 1 - t$, $\mathbf{p}_2 = t + t^2$, and $\mathbf{p}_3 = 2 - t + t^2$ be vectors in \mathbb{P}_3 (the set of all polynomials of degree ≤ 3).

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: Let $\mathbf{p}_1 = 1 - t$, $\mathbf{p}_2 = t + t^2$, and $\mathbf{p}_3 = 2 - t + t^2$ be vectors in \mathbb{P}_3 (the set of all polynomials of degree ≤ 3). Is the set $\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$ linearly independent?

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: Let $\mathbf{p}_1 = 1 - t$, $\mathbf{p}_2 = t + t^2$, and $\mathbf{p}_3 = 2 - t + t^2$ be vectors in \mathbb{P}_3 (the set of all polynomials of degree ≤ 3). Is the set $\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$ linearly independent?

Example: Let $\mathbf{v}_1 = \sin x$, $\mathbf{v}_2 = x^2 - x$, and $\mathbf{v}_3 = e^x$ be vectors in $C([0, 1])$.

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: Let $\mathbf{p}_1 = 1 - t$, $\mathbf{p}_2 = t + t^2$, and $\mathbf{p}_3 = 2 - t + t^2$ be vectors in \mathbb{P}_3 (the set of all polynomials of degree ≤ 3). Is the set $\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$ linearly independent?

Example: Let $\mathbf{v}_1 = \sin x$, $\mathbf{v}_2 = x^2 - x$, and $\mathbf{v}_3 = e^x$ be vectors in $C([0, 1])$. Is the set $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ linearly independent?

§4.3 LINEARLY INDEPENDENT SETS; BASES

DEFINITION (BASIS OF A VECTOR SPACE)

Let H be a subspace of a vector space V . An indexed set of vectors $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_p\}$ in V is a **basis** for H if

§4.3 LINEARLY INDEPENDENT SETS; BASES

DEFINITION (BASIS OF A VECTOR SPACE)

Let H be a subspace of a vector space V . An indexed set of vectors $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_p\}$ in V is a **basis** for H if

1. \mathcal{B} is a linearly independent set, and

§4.3 LINEARLY INDEPENDENT SETS; BASES

DEFINITION (BASIS OF A VECTOR SPACE)

Let H be a subspace of a vector space V . An indexed set of vectors $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_p\}$ in V is a **basis** for H if

1. \mathcal{B} is a linearly independent set, and
2. the vectors in \mathcal{B} span H ; that is

$$H = \text{Span}\{\mathbf{b}_1, \dots, \mathbf{b}_p\}.$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

DEFINITION (BASIS OF A VECTOR SPACE)

Let H be a subspace of a vector space V . An indexed set of vectors $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_p\}$ in V is a **basis** for H if

1. \mathcal{B} is a linearly independent set, and
2. the vectors in \mathcal{B} span H ; that is

$$H = \text{Span}\{\mathbf{b}_1, \dots, \mathbf{b}_p\}.$$

This definition also applies to the whole space V ;

§4.3 LINEARLY INDEPENDENT SETS; BASES

DEFINITION (BASIS OF A VECTOR SPACE)

Let H be a subspace of a vector space V . An indexed set of vectors $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_p\}$ in V is a **basis** for H if

1. \mathcal{B} is a linearly independent set, and
2. the vectors in \mathcal{B} span H ; that is

$$H = \text{Span}\{\mathbf{b}_1, \dots, \mathbf{b}_p\}.$$

This definition also applies to the whole space V ; a basis of V is a linearly independent subset of V which spans V .

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: The columns of an invertible $n \times n$ matrix

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: The columns of an invertible $n \times n$ matrix are both linearly independent, and span \mathbb{R}^n (by the Invertible Matrix theorem)

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: The columns of an invertible $n \times n$ matrix are both linearly independent, and span \mathbb{R}^n (by the Invertible Matrix theorem) \implies they form a basis for \mathbb{R}^n .

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: The columns of an invertible $n \times n$ matrix are both linearly independent, and span \mathbb{R}^n (by the Invertible Matrix theorem) \implies they form a basis for \mathbb{R}^n .

Example: The vectors

$$\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, \mathbf{e}_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

are called the **standard basis** for \mathbb{R}^n .

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: The columns of an invertible $n \times n$ matrix are both linearly independent, and span \mathbb{R}^n (by the Invertible Matrix theorem) \implies they form a basis for \mathbb{R}^n .

Example: The vectors

$$\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, \mathbf{e}_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

are called the **standard basis** for \mathbb{R}^n .

Example: The set $\{1, t, t^2, \dots, t^n\}$ forms a basis of \mathbb{P}_n , and is called the **standard basis** for \mathbb{P}_n .

§4.3 LINEARLY INDEPENDENT SETS; BASES

THEOREM (SPANNING SET THEOREM)

Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ be a set in V , and $H = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.

§4.3 LINEARLY INDEPENDENT SETS; BASES

THEOREM (SPANNING SET THEOREM)

Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ be a set in V , and $H = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.

1. If one of the vectors in S – say, \mathbf{v}_k – is a linear combination of the remaining vectors in S , then the set formed from S by removing \mathbf{v}_k still spans H .

§4.3 LINEARLY INDEPENDENT SETS; BASES

THEOREM (SPANNING SET THEOREM)

Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ be a set in V , and $H = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.

1. If one of the vectors in S – say, \mathbf{v}_k – is a linear combination of the remaining vectors in S , then the set formed from S by removing \mathbf{v}_k still spans H .
2. If $H \neq \{\mathbf{0}\}$, some subset of S is a basis for H .

§4.3 LINEARLY INDEPENDENT SETS; BASES

If A is an $n \times m$ matrix, we find a basis for $\text{Nul } A$ by

§4.3 LINEARLY INDEPENDENT SETS; BASES

If A is an $n \times m$ matrix, we find a basis for $\text{Nul } A$ by expressing the solution of $A\mathbf{x} = \mathbf{0}$ in parametric form.

§4.3 LINEARLY INDEPENDENT SETS; BASES

If A is an $n \times m$ matrix, we find a basis for $\text{Nul } A$ by expressing the solution of $A\mathbf{x} = \mathbf{0}$ in parametric form.

Example: Find a basis for $\text{Col } B$, where

$$B = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

If A is an $n \times m$ matrix, we find a basis for $\text{Nul } A$ by expressing the solution of $A\mathbf{x} = \mathbf{0}$ in parametric form.

Example: Find a basis for $\text{Col } B$, where

$$B = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

THEOREM

The pivot columns of a matrix A form a basis for $\text{Col } A$.

§4.3 LINEARLY INDEPENDENT SETS; BASES

If A is an $n \times m$ matrix, we find a basis for $\text{Nul } A$ by expressing the solution of $A\mathbf{x} = \mathbf{0}$ in parametric form.

Example: Find a basis for $\text{Col } B$, where

$$B = \begin{bmatrix} 1 & 4 & 0 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

THEOREM

The pivot columns of a matrix A form a basis for $\text{Col } A$.

Warning! You have to row reduce the matrix to find the pivot columns, but the basis for $\text{Col } A$ are the original pivot columns in A , *not* the row reduced matrix.

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: Find a basis for $\text{Col } A$, where

$$A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: Find a basis for $\text{Col } A$, where

$$A = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}.$$

Hint: A is row-equivalent to the matrix B from the previous slide.

§4.3 LINEARLY INDEPENDENT SETS; BASES

Examples:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \right\}$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

Examples:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \right\} \quad \text{Linearly independent,}$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

Examples:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \right\} \quad \text{Linearly independent, does not span } \mathbb{R}^3$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

Examples:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \right\} \quad \text{Linearly independent, does not span } \mathbb{R}^3$$

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \right\}$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

Examples:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \right\} \quad \text{Linearly independent, does not span } \mathbb{R}^3$$

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \right\} \quad \text{A basis for } \mathbb{R}^3$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

Examples:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \right\} \quad \text{Linearly independent, does not span } \mathbb{R}^3$$

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \right\} \quad \text{A basis for } \mathbb{R}^3$$

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} \right\}$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

Examples:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \right\} \quad \text{Linearly independent, does not span } \mathbb{R}^3$$

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \right\} \quad \text{A basis for } \mathbb{R}^3$$

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} \right\} \quad \text{Spans } \mathbb{R}^3,$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

Examples:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} \right\} \quad \text{Linearly independent, does not span } \mathbb{R}^3$$

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \right\} \quad \text{A basis for } \mathbb{R}^3$$

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}, \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} \right\} \quad \text{Spans } \mathbb{R}^3, \text{ but is linearly dep.}$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: Find a basis for $\text{Nul } A$ and $\text{Col } A$, where

$$A = \begin{bmatrix} 1 & 4 & 0 & 2 \\ 3 & 12 & 1 & 5 \\ 2 & 8 & 1 & 3 \\ 5 & 20 & 2 & 8 \end{bmatrix}$$

§4.3 LINEARLY INDEPENDENT SETS; BASES

Example: Find a basis for $\text{Nul } A$ and $\text{Col } A$, where

$$A = \begin{bmatrix} 1 & 4 & 0 & 2 \\ 3 & 12 & 1 & 5 \\ 2 & 8 & 1 & 3 \\ 5 & 20 & 2 & 8 \end{bmatrix} \longrightarrow \begin{bmatrix} 1 & 4 & 0 & 2 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

§4.4 COORDINATE SYSTEMS

THEOREM (UNIQUE REPRESENTATION THEOREM)

Let $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ be a basis for a vector space V . Then for each \mathbf{x} in V , there exists a unique set of scalars c_1, \dots, c_n such that

$$\mathbf{x} = c_1 \mathbf{b}_1 + \dots + c_n \mathbf{b}_n.$$

§4.4 COORDINATE SYSTEMS

THEOREM (UNIQUE REPRESENTATION THEOREM)

Let $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ be a basis for a vector space V . Then for each \mathbf{x} in V , there exists a unique set of scalars c_1, \dots, c_n such that

$$\mathbf{x} = c_1 \mathbf{b}_1 + \dots + c_n \mathbf{b}_n.$$

We call the weights c_1, \dots, c_n the **coordinates of \mathbf{x} relative to the basis \mathcal{B}** (or the **\mathcal{B} -coordinates of \mathbf{x}**).

§4.4 COORDINATE SYSTEMS

If c_1, \dots, c_n are the \mathcal{B} -coordinates of \mathbf{x} , then the vector in \mathbb{R}^n

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$$

is called the **coordinate vector of \mathbf{x} (relative to \mathcal{B})**, or the **\mathcal{B} -coordinate vector of \mathbf{x}** .

§4.4 COORDINATE SYSTEMS

If c_1, \dots, c_n are the \mathcal{B} -coordinates of \mathbf{x} , then the vector in \mathbb{R}^n


$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$$

is called the **coordinate vector of \mathbf{x} (relative to \mathcal{B})**, or the **\mathcal{B} -coordinate vector of \mathbf{x}** .



The mapping $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ is the **coordinate mapping (determined by \mathcal{B})**.

§4.4 COORDINATE SYSTEMS



§4.4 COORDINATE SYSTEMS





§4.4 COORDINATE SYSTEMS





§4.4 COORDINATE SYSTEMS





§4.4 COORDINATE SYSTEMS




§4.4 COORDINATE SYSTEMS




§4.4 COORDINATE SYSTEMS




§4.4 COORDINATE SYSTEMS



$$\mathbf{v} = 1 \cdot \mathbf{e}_1 + 2 \cdot \mathbf{e}_2$$




§4.4 COORDINATE SYSTEMS




$$\mathbf{v} = 1 \cdot \mathbf{e}_1 + 2 \cdot \mathbf{e}_2$$

$$\Rightarrow [\mathbf{v}]_{\mathcal{E}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$




§4.4 COORDINATE SYSTEMS




$$\mathbf{v} = 1 \cdot \mathbf{e}_1 + 2 \cdot \mathbf{e}_2$$

$$\Rightarrow [\mathbf{v}]_{\mathcal{E}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$




§4.4 COORDINATE SYSTEMS




$$\mathbf{v} = 1 \cdot \mathbf{e}_1 + 2 \cdot \mathbf{e}_2$$

$$\Rightarrow [\mathbf{v}]_{\mathcal{E}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$




$$\mathbf{v} = -1 \cdot \mathbf{b}_1 + 2 \cdot \mathbf{b}_2$$

§4.4 COORDINATE SYSTEMS



$$\mathbf{v} = 1 \cdot \mathbf{e}_1 + 2 \cdot \mathbf{e}_2$$

$$\Rightarrow [\mathbf{v}]_{\mathcal{E}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$



$$\mathbf{v} = -1 \cdot \mathbf{b}_1 + 2 \cdot \mathbf{b}_2$$

$$\Rightarrow [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

§4.4 COORDINATE SYSTEMS

Example: Let $\mathcal{E} = \{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ denote the standard basis of \mathbb{R}^n .

§4.4 COORDINATE SYSTEMS

Example: Let $\mathcal{E} = \{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ denote the standard basis of \mathbb{R}^n .

For the vector

$$\mathbf{x} = \begin{bmatrix} -1 \\ 3 \\ 5 \\ 0 \end{bmatrix}$$

find the coordinate vector of \mathbf{x} relative to \mathcal{E} .

§4.4 COORDINATE SYSTEMS

Example: Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ be a the basis of \mathbb{R}^3 given by

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{b}_3 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}.$$

§4.4 COORDINATE SYSTEMS

Example: Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ be a the basis of \mathbb{R}^3 given by

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{b}_3 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}.$$

$$\text{Let } [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}.$$

§4.4 COORDINATE SYSTEMS

Example: Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ be a the basis of \mathbb{R}^3 given by

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{b}_3 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}.$$

Let $[\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$. Then find \mathbf{v} .

§4.4 COORDINATE SYSTEMS

Example: Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ be a the basis of \mathbb{R}^3 given by

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{b}_3 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}.$$

Let $[\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$. Then find \mathbf{v} .

For the vector

$$\mathbf{x} = \begin{bmatrix} 2 \\ 0 \\ -3 \end{bmatrix}$$

find the coordinate vector of \mathbf{x} relative to \mathcal{B} .

§4.4 COORDINATE SYSTEMS

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ be a basis, and suppose

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix},$$

for some $\mathbf{x} \in \mathbb{R}^n$.

§4.4 COORDINATE SYSTEMS

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ be a basis, and suppose

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix},$$

for some $\mathbf{x} \in \mathbb{R}^n$. Also, let $P_{\mathcal{B}} = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_n]$.

§4.4 COORDINATE SYSTEMS

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ be a basis, and suppose

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix},$$

for some $\mathbf{x} \in \mathbb{R}^n$. Also, let $P_{\mathcal{B}} = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_n]$.

So,

$$\mathbf{x} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n$$

§4.4 COORDINATE SYSTEMS

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ be a basis, and suppose

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix},$$

for some $\mathbf{x} \in \mathbb{R}^n$. Also, let $P_{\mathcal{B}} = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_n]$.

So,

$$\mathbf{x} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n = P_{\mathcal{B}} [\mathbf{x}]_{\mathcal{B}}$$

§4.4 COORDINATE SYSTEMS

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ be a basis, and suppose

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix},$$

for some $\mathbf{x} \in \mathbb{R}^n$. Also, let $P_{\mathcal{B}} = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_n]$.

So,

$$\mathbf{x} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n = P_{\mathcal{B}} [\mathbf{x}]_{\mathcal{B}}$$

and since $P_{\mathcal{B}}$ is invertible (why?)

§4.4 COORDINATE SYSTEMS

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ be a basis, and suppose

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix},$$

for some $\mathbf{x} \in \mathbb{R}^n$. Also, let $P_{\mathcal{B}} = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_n]$.

So,

$$\mathbf{x} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n = P_{\mathcal{B}} [\mathbf{x}]_{\mathcal{B}}$$

and since $P_{\mathcal{B}}$ is invertible (why?), we have

$$[\mathbf{x}]_{\mathcal{B}} = P_{\mathcal{B}}^{-1} \mathbf{x}$$

§4.4 COORDINATE SYSTEMS

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ be a basis, and suppose

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix},$$

for some $\mathbf{x} \in \mathbb{R}^n$. Also, let $P_{\mathcal{B}} = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_n]$.

So,

$$\mathbf{x} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n = P_{\mathcal{B}} [\mathbf{x}]_{\mathcal{B}}$$

and since $P_{\mathcal{B}}$ is invertible (why?), we have

$$[\mathbf{x}]_{\mathcal{B}} = P_{\mathcal{B}}^{-1} \mathbf{x}$$

We call $P_{\mathcal{B}}$ the **change-of-coordinates matrix** from \mathcal{B} to the standard basis in \mathbb{R}^n .

§4.4 COORDINATE SYSTEMS

Example: Let $\mathbf{b}_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ and $\mathbf{b}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$.

§4.4 COORDINATE SYSTEMS

Example: Let $\mathbf{b}_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ and $\mathbf{b}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$.

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$, and let $\mathbf{x} = \begin{bmatrix} 3 \\ 7 \end{bmatrix}$.

§4.4 COORDINATE SYSTEMS

Example: Let $\mathbf{b}_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ and $\mathbf{b}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$.

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$, and let $\mathbf{x} = \begin{bmatrix} 3 \\ 7 \end{bmatrix}$.

Find the change of coordinate matrix $\mathcal{P}_{\mathcal{B}}$, and use it to find the coordinate vector of \mathbf{x} relative to the basis \mathcal{B} .

§4.4 COORDINATE SYSTEMS

THEOREM

Let $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ be a basis for a vector space V . Then the coordinate mapping $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ is a **one-to-one** linear transformation from V **onto** \mathbb{R}^n .

§4.4 COORDINATE SYSTEMS

THEOREM

Let $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ be a basis for a vector space V . Then the coordinate mapping $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ is a **one-to-one** linear transformation from V **onto** \mathbb{R}^n .

A **one-to-one** linear transformation from V **onto** W is called an **isomorphism**.

§4.4 COORDINATE SYSTEMS

THEOREM

Let $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ be a basis for a vector space V . Then the coordinate mapping $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ is a **one-to-one** linear transformation from V **onto** \mathbb{R}^n .

A **one-to-one** linear transformation from V **onto** W is called an **isomorphism**.

In this case, even though the vector spaces V and W may look different, they are essentially the same.

§4.4 COORDINATE SYSTEMS

THEOREM

Let $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ be a basis for a vector space V . Then the coordinate mapping $\mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ is a **one-to-one** linear transformation from V **onto** \mathbb{R}^n .

A **one-to-one** linear transformation from V **onto** W is called an **isomorphism**.

In this case, even though the vector spaces V and W may look different, they are essentially the same.

Thus if a vector space V has a basis $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ then it is essentially “the same” as \mathbb{R}^n .

§4.4 COORDINATE SYSTEMS

PROOF.

§4.4 COORDINATE SYSTEMS

PROOF.

Let $T_{\mathcal{B}} : V \rightarrow \mathbb{R}^n$ be the coord. map which sends $\mathbf{v} = v_1 \mathbf{b}_1 + \cdots + v_n \mathbf{b}_n$ to

$$T_{\mathcal{B}}(\mathbf{v}) = [\mathbf{v}]_{\mathcal{B}}$$

§4.4 COORDINATE SYSTEMS

PROOF.

Let $T_{\mathcal{B}} : V \rightarrow \mathbb{R}^n$ be the coord. map which sends $\mathbf{v} = v_1\mathbf{b}_1 + \cdots + v_n\mathbf{b}_n$ to

$$T_{\mathcal{B}}(\mathbf{v}) = [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}.$$

§4.4 COORDINATE SYSTEMS

PROOF.

Let $T_{\mathcal{B}} : V \rightarrow \mathbb{R}^n$ be the coord. map which sends $\mathbf{v} = v_1\mathbf{b}_1 + \cdots + v_n\mathbf{b}_n$ to

$$T_{\mathcal{B}}(\mathbf{v}) = [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}.$$

Let $\mathbf{w} = w_1\mathbf{b}_1 + \cdots w_n\mathbf{b}_n$ and $c \in \mathbb{R}$.

§4.4 COORDINATE SYSTEMS

PROOF.

Let $T_{\mathcal{B}} : V \rightarrow \mathbb{R}^n$ be the coord. map which sends $\mathbf{v} = v_1\mathbf{b}_1 + \cdots + v_n\mathbf{b}_n$ to

$$T_{\mathcal{B}}(\mathbf{v}) = [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}.$$

Let $\mathbf{w} = w_1\mathbf{b}_1 + \cdots w_n\mathbf{b}_n$ and $c \in \mathbb{R}$. Then

$$\mathbf{v} + \mathbf{w} = (v_1 + w_1)\mathbf{b}_1 + \cdots + (v_n + w_n)\mathbf{b}_n$$

§4.4 COORDINATE SYSTEMS

PROOF.

Let $T_{\mathcal{B}} : V \rightarrow \mathbb{R}^n$ be the coord. map which sends $\mathbf{v} = v_1 \mathbf{b}_1 + \cdots + v_n \mathbf{b}_n$ to

$$T_{\mathcal{B}}(\mathbf{v}) = [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}.$$

Let $\mathbf{w} = w_1 \mathbf{b}_1 + \cdots + w_n \mathbf{b}_n$ and $c \in \mathbb{R}$. Then

$$\mathbf{v} + \mathbf{w} = (v_1 + w_1) \mathbf{b}_1 + \cdots + (v_n + w_n) \mathbf{b}_n$$

and

$$T_{\mathcal{B}}(\mathbf{v} + \mathbf{w})$$

§4.4 COORDINATE SYSTEMS

PROOF.

Let $T_{\mathcal{B}} : V \rightarrow \mathbb{R}^n$ be the coord. map which sends $\mathbf{v} = v_1\mathbf{b}_1 + \cdots + v_n\mathbf{b}_n$ to

$$T_{\mathcal{B}}(\mathbf{v}) = [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}.$$

Let $\mathbf{w} = w_1\mathbf{b}_1 + \cdots + w_n\mathbf{b}_n$ and $c \in \mathbb{R}$. Then

$$\mathbf{v} + \mathbf{w} = (v_1 + w_1)\mathbf{b}_1 + \cdots + (v_n + w_n)\mathbf{b}_n$$

and

$$T_{\mathcal{B}}(\mathbf{v} + \mathbf{w}) = \begin{bmatrix} v_1 + w_1 \\ \vdots \\ v_n + w_n \end{bmatrix}$$

§4.4 COORDINATE SYSTEMS

PROOF.

Let $T_{\mathcal{B}} : V \rightarrow \mathbb{R}^n$ be the coord. map which sends $\mathbf{v} = v_1\mathbf{b}_1 + \cdots + v_n\mathbf{b}_n$ to

$$T_{\mathcal{B}}(\mathbf{v}) = [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}.$$

Let $\mathbf{w} = w_1\mathbf{b}_1 + \cdots + w_n\mathbf{b}_n$ and $c \in \mathbb{R}$. Then

$$\mathbf{v} + \mathbf{w} = (v_1 + w_1)\mathbf{b}_1 + \cdots + (v_n + w_n)\mathbf{b}_n$$

and

$$T_{\mathcal{B}}(\mathbf{v} + \mathbf{w}) = \begin{bmatrix} v_1 + w_1 \\ \vdots \\ v_n + w_n \end{bmatrix} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix} + \begin{bmatrix} w_1 \\ \vdots \\ w_n \end{bmatrix}$$

§4.4 COORDINATE SYSTEMS

PROOF.

Let $T_{\mathcal{B}} : V \rightarrow \mathbb{R}^n$ be the coord. map which sends $\mathbf{v} = v_1\mathbf{b}_1 + \cdots + v_n\mathbf{b}_n$ to

$$T_{\mathcal{B}}(\mathbf{v}) = [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}.$$

Let $\mathbf{w} = w_1\mathbf{b}_1 + \cdots + w_n\mathbf{b}_n$ and $c \in \mathbb{R}$. Then

$$\mathbf{v} + \mathbf{w} = (v_1 + w_1)\mathbf{b}_1 + \cdots + (v_n + w_n)\mathbf{b}_n$$

and

$$T_{\mathcal{B}}(\mathbf{v} + \mathbf{w}) = \begin{bmatrix} v_1 + w_1 \\ \vdots \\ v_n + w_n \end{bmatrix} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix} + \begin{bmatrix} w_1 \\ \vdots \\ w_n \end{bmatrix} = T_{\mathcal{B}}(\mathbf{v}) + T_{\mathcal{B}}(\mathbf{w}).$$

§4.4 COORDINATE SYSTEMS

Also,

$$c \cdot \mathbf{v} = cv_1 \mathbf{b}_1 + \cdots + cv_n \mathbf{b}_n$$

§4.4 COORDINATE SYSTEMS

Also,

$$c \cdot \mathbf{v} = cv_1 \mathbf{b}_1 + \cdots + cv_n \mathbf{b}_n$$

and

$$T_{\mathcal{B}}(c \cdot \mathbf{v}) = \begin{bmatrix} cv_1 \\ \vdots \\ cv_n \end{bmatrix}$$

§4.4 COORDINATE SYSTEMS

Also,

$$c \cdot \mathbf{v} = cv_1 \mathbf{b}_1 + \cdots + cv_n \mathbf{b}_n$$

and

$$T_{\mathcal{B}}(c \cdot \mathbf{v}) = \begin{bmatrix} cv_1 \\ \vdots \\ cv_n \end{bmatrix} = c \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}$$

§4.4 COORDINATE SYSTEMS

Also,

$$c \cdot \mathbf{v} = cv_1 \mathbf{b}_1 + \cdots + cv_n \mathbf{b}_n$$

and

$$T_{\mathcal{B}}(c \cdot \mathbf{v}) = \begin{bmatrix} cv_1 \\ \vdots \\ cv_n \end{bmatrix} = c \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix} = c \cdot T_{\mathcal{B}}(\mathbf{v}).$$

§4.4 COORDINATE SYSTEMS

Also,

$$c \cdot \mathbf{v} = cv_1 \mathbf{b}_1 + \cdots + cv_n \mathbf{b}_n$$

and

$$T_{\mathcal{B}}(c \cdot \mathbf{v}) = \begin{bmatrix} cv_1 \\ \vdots \\ cv_n \end{bmatrix} = c \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix} = c \cdot T_{\mathcal{B}}(\mathbf{v}).$$

Thus $T_{\mathcal{B}}$ is linear.

§4.4 COORDINATE SYSTEMS

One-to-one:

§4.4 COORDINATE SYSTEMS

One-to-one:

If $T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w})$

§4.4 COORDINATE SYSTEMS

One-to-one:

$$\text{If } T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w}) = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix},$$

§4.4 COORDINATE SYSTEMS

One-to-one:

If $T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w}) = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$, then we must have

$$\mathbf{v} =$$

§4.4 COORDINATE SYSTEMS

One-to-one:

If $T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w}) = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$, then we must have

$$\mathbf{v} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n$$

§4.4 COORDINATE SYSTEMS

One-to-one:

If $T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w}) = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$, then we must have

$$\mathbf{v} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n \quad \text{and} \quad \mathbf{w} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n$$

§4.4 COORDINATE SYSTEMS

One-to-one:

If $T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w}) = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$, then we must have

$$\mathbf{v} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n \quad \text{and} \quad \mathbf{w} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n$$

hence $\mathbf{v} = \mathbf{w}$,

§4.4 COORDINATE SYSTEMS

One-to-one:

If $T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w}) = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$, then we must have

$$\mathbf{v} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n \quad \text{and} \quad \mathbf{w} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n$$

hence $\mathbf{v} = \mathbf{w}$, and $T_{\mathcal{B}}$ is one-to-one.

§4.4 COORDINATE SYSTEMS

One-to-one:

If $T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w}) = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$, then we must have

$$\mathbf{v} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n \quad \text{and} \quad \mathbf{w} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n$$

hence $\mathbf{v} = \mathbf{w}$, and $T_{\mathcal{B}}$ is one-to-one.

Onto:

§4.4 COORDINATE SYSTEMS

One-to-one:

If $T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w}) = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$, then we must have

$$\mathbf{v} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n \quad \text{and} \quad \mathbf{w} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n$$

hence $\mathbf{v} = \mathbf{w}$, and $T_{\mathcal{B}}$ is one-to-one.

Onto:

For any vector $\mathbf{b} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} \in \mathbb{R}^n$,

§4.4 COORDINATE SYSTEMS

One-to-one:

If $T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w}) = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$, then we must have

$$\mathbf{v} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n \quad \text{and} \quad \mathbf{w} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n$$

hence $\mathbf{v} = \mathbf{w}$, and $T_{\mathcal{B}}$ is one-to-one.

Onto:

For any vector $\mathbf{b} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} \in \mathbb{R}^n$, the vector $\mathbf{x} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n \in V$

satisfies

$$T_{\mathcal{B}}(\mathbf{x}) = \mathbf{b},$$

§4.4 COORDINATE SYSTEMS

One-to-one:

If $T_{\mathcal{B}}(\mathbf{v}) = T_{\mathcal{B}}(\mathbf{w}) = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$, then we must have

$$\mathbf{v} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n \quad \text{and} \quad \mathbf{w} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n$$

hence $\mathbf{v} = \mathbf{w}$, and $T_{\mathcal{B}}$ is one-to-one.

Onto:

For any vector $\mathbf{b} = \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} \in \mathbb{R}^n$, the vector $\mathbf{x} = c_1 \mathbf{b}_1 + \cdots + c_n \mathbf{b}_n \in V$ satisfies

$$T_{\mathcal{B}}(\mathbf{x}) = \mathbf{b},$$

and hence $T_{\mathcal{B}}$ is onto.



§4.4 COORDINATE SYSTEMS

Example: The set $\mathcal{P} = \{1, t, t^2, \dots, t^n\}$ is a basis for \mathbb{P}_n .

§4.4 COORDINATE SYSTEMS

Example: The set $\mathcal{P} = \{1, t, t^2, \dots, t^n\}$ is a basis for \mathbb{P}_n . For any polynomial $\mathbf{p} = a_0 + a_1t + \dots + a_nt^n$ we have

§4.4 COORDINATE SYSTEMS

Example: The set $\mathcal{P} = \{1, t, t^2, \dots, t^n\}$ is a basis for \mathbb{P}_n . For any polynomial $\mathbf{p} = a_0 + a_1t + \dots + a_nt^n$ we have

$$[\mathbf{p}]_{\mathcal{P}} = \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_n \end{bmatrix}.$$

§4.4 COORDINATE SYSTEMS

Example: The set $\mathcal{P} = \{1, t, t^2, \dots, t^n\}$ is a basis for \mathbb{P}_n . For any polynomial $\mathbf{p} = a_0 + a_1t + \dots + a_nt^n$ we have

$$[\mathbf{p}]_{\mathcal{P}} = \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_n \end{bmatrix}.$$

We can thus do any operations in the vector space \mathbb{P}_n as if we were working in \mathbb{R}^{n+1} .

§4.4 COORDINATE SYSTEMS

Example: The set $\mathcal{P} = \{1, t, t^2, \dots, t^n\}$ is a basis for \mathbb{P}_n . For any polynomial $\mathbf{p} = a_0 + a_1t + \dots + a_nt^n$ we have

$$[\mathbf{p}]_{\mathcal{P}} = \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_n \end{bmatrix}.$$

We can thus do any operations in the vector space \mathbb{P}_n as if we were working in \mathbb{R}^{n+1} .

Caution: Not every vector space has a basis of the form $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$. For example, the space $C(\mathbb{R})$ has no such basis.

§4.4 COORDINATE SYSTEMS

Example: The set $\mathcal{P} = \{1, t, t^2, \dots, t^n\}$ is a basis for \mathbb{P}_n . For any polynomial $\mathbf{p} = a_0 + a_1t + \dots + a_nt^n$ we have

$$[\mathbf{p}]_{\mathcal{P}} = \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_n \end{bmatrix}.$$

We can thus do any operations in the vector space \mathbb{P}_n as if we were working in \mathbb{R}^{n+1} .

Caution: Not every vector space has a basis of the form $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$. For example, the space $C(\mathbb{R})$ has no such basis. It does have a basis with infinitely many basis vectors though.

§4.4 COORDINATE SYSTEMS

Example: Using the coordinate vectors of the polynomials $\mathbf{p}_1 = t^2 + t + 1$, $\mathbf{p}_2 = t^2 + 2t$, and $\mathbf{p}_3 = t^2 - 2$, determine whether they are linearly independent.

§4.5 THE DIMENSION OF A VECTOR SPACE

THEOREM

If a vector space V has a basis $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$, then any subset of V containing more than n vectors must be linearly dependent.

§4.5 THE DIMENSION OF A VECTOR SPACE

THEOREM

If a vector space V has a basis $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$, then any subset of V containing more than n vectors must be linearly dependent.

THEOREM

If a vector space V has a basis of n vectors, then every basis of V must consist of exactly n vectors.

§4.5 THE DIMENSION OF A VECTOR SPACE

DEFINITION

If V is spanned by a finite set, then V is said to be **finite-dimensional**, and the **dimension** of V , written as $\dim V$, is the number of vectors in a basis for V .

§4.5 THE DIMENSION OF A VECTOR SPACE

DEFINITION

If V is spanned by a finite set, then V is said to be **finite-dimensional**, and the **dimension** of V , written as $\dim V$, is the number of vectors in a basis for V .

The dimension of the zero vector space $\{\mathbf{0}\}$ is defined to be zero.

§4.5 THE DIMENSION OF A VECTOR SPACE

DEFINITION

If V is spanned by a finite set, then V is said to be **finite-dimensional**, and the **dimension** of V , written as $\dim V$, is the number of vectors in a basis for V .

The dimension of the zero vector space $\{\mathbf{0}\}$ is defined to be zero.

If V is not spanned by a finite set, then V is said to be **infinite dimensional**.

§4.5 THE DIMENSION OF A VECTOR SPACE

Example: The standard basis $\mathcal{E} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ has n vectors, and hence $\dim \mathbb{R}^n = n$.

§4.5 THE DIMENSION OF A VECTOR SPACE

Example: The standard basis $\mathcal{E} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ has n vectors, and hence $\dim \mathbb{R}^n = n$.

Example: Find the dimension of \mathbb{P}_n (i.e. the space of all polynomials of degree $\leq n$).

§4.5 THE DIMENSION OF A VECTOR SPACE

Example: The standard basis $\mathcal{E} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ has n vectors, and hence $\dim \mathbb{R}^n = n$.

Example: Find the dimension of \mathbb{P}_n (i.e. the space of all polynomials of degree $\leq n$).

Example: Find the dimension of $C(\mathbb{R})$.

§4.5 THE DIMENSION OF A VECTOR SPACE

Example: The standard basis $\mathcal{E} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ has n vectors, and hence $\dim \mathbb{R}^n = n$.

Example: Find the dimension of \mathbb{P}_n (i.e. the space of all polynomials of degree $\leq n$).

Example: Find the dimension of $C(\mathbb{R})$.

Example: Find the dimension of the subspace

$$H = \left\{ \begin{bmatrix} a - 3b + 6c \\ 5a + 4d \\ b - 2c - d \\ 5d \end{bmatrix} : a, b, c, d \in \mathbb{R} \right\}.$$

§4.5 THE DIMENSION OF A VECTOR SPACE

THEOREM

Let H be a subspace of a finite-dimensional vector space V . Any linearly independent set in H can be expanded, if necessary, to a basis for H . Also H is finite-dimensional and

$$\dim H \leq \dim V.$$

§4.5 THE DIMENSION OF A VECTOR SPACE

THEOREM

Let H be a subspace of a finite-dimensional vector space V . Any linearly independent set in H can be expanded, if necessary, to a basis for H . Also H is finite-dimensional and

$$\dim H \leq \dim V.$$

THEOREM (THE BASIS THEOREM)

Let V be a p -dimensional vector space, for $p \geq 1$.

§4.5 THE DIMENSION OF A VECTOR SPACE

THEOREM

Let H be a subspace of a finite-dimensional vector space V . Any linearly independent set in H can be expanded, if necessary, to a basis for H . Also H is finite-dimensional and

$$\dim H \leq \dim V.$$

THEOREM (THE BASIS THEOREM)

Let V be a p -dimensional vector space, for $p \geq 1$. Any linearly independent set of exactly p elements in V is automatically a basis for V .

§4.5 THE DIMENSION OF A VECTOR SPACE

THEOREM

Let H be a subspace of a finite-dimensional vector space V . Any linearly independent set in H can be expanded, if necessary, to a basis for H . Also H is finite-dimensional and

$$\dim H \leq \dim V.$$

THEOREM (THE BASIS THEOREM)

Let V be a p -dimensional vector space, for $p \geq 1$. Any linearly independent set of exactly p elements in V is automatically a basis for V . Any set of exactly p elements that spans V is automatically a basis for V .

§4.5 THE DIMENSION OF A VECTOR SPACE

Fact: The dimension of $\text{Nul } A$ is the number of free variables in the equation $A\mathbf{x} = \mathbf{0}$,

§4.5 THE DIMENSION OF A VECTOR SPACE

Fact: The dimension of $\text{Nul } A$ is the number of free variables in the equation $A\mathbf{x} = \mathbf{0}$, and the dimension of $\text{Col } A$ is the number of pivot columns in A .

§4.5 THE DIMENSION OF A VECTOR SPACE

Fact: The dimension of $\text{Nul } A$ is the number of free variables in the equation $A\mathbf{x} = \mathbf{0}$, and the dimension of $\text{Col } A$ is the number of pivot columns in A .

Example: Find the dimension of $\text{Nul } A$ and $\text{Col } A$, where

$$A = \begin{bmatrix} 2 & -1 & 4 & 1 \\ 1 & 3 & 3 & 1 \\ 2 & 1 & 1 & 3 \end{bmatrix}.$$

§4.6 RANK

If A is an $m \times n$ matrix, each row of A has n entries, and thus can be identified with a vector in \mathbb{R}^n .

§4.6 RANK

If A is an $m \times n$ matrix, each row of A has n entries, and thus can be identified with a vector in \mathbb{R}^n .

The set of all linear combinations of the row vectors is called the **row space** of A and is denoted by $\text{Row } A$.

§4.6 RANK

If A is an $m \times n$ matrix, each row of A has n entries, and thus can be identified with a vector in \mathbb{R}^n .

The set of all linear combinations of the row vectors is called the **row space** of A and is denoted by $\text{Row } A$.

i.e. $\text{Row } A$ is the span of all the rows of A .

§4.6 RANK

If A is an $m \times n$ matrix, each row of A has n entries, and thus can be identified with a vector in \mathbb{R}^n .

The set of all linear combinations of the row vectors is called the **row space** of A and is denoted by $\text{Row } A$.

i.e. $\text{Row } A$ is the span of all the rows of A .

THEOREM

If two matrices A and B are row equivalent, then their row spaces are the same. If B is in echelon form, the nonzero rows of B form a basis for the row space of A as well as for that of B .

§4.6 RANK

Example: Find a basis for Row A , Col A , and Nul A , where

$$A = \begin{bmatrix} -2 & -5 & 8 & 0 \\ 1 & 3 & -5 & 1 \\ 3 & 11 & -19 & 7 \end{bmatrix}.$$

§4.6 RANK

DEFINITION (RANK)

The **rank** of a matrix A is the dimension of the column space of A .

§4.6 RANK

DEFINITION (RANK)

The **rank** of a matrix A is the dimension of the column space of A .

THEOREM

The dimensions of the column space and the row space of an $m \times n$ matrix A are equal.

§4.6 RANK

DEFINITION (RANK)

The **rank** of a matrix A is the dimension of the column space of A .

THEOREM

The dimensions of the column space and the row space of an $m \times n$ matrix A are equal. This common dimension, the rank of A , also equals the number of pivot positions in A and satisfies the equation,

§4.6 RANK

DEFINITION (RANK)

The **rank** of a matrix A is the dimension of the column space of A .

THEOREM

The dimensions of the column space and the row space of an $m \times n$ matrix A are equal. This common dimension, the rank of A , also equals the number of pivot positions in A and satisfies the equation,

$$\text{rank } A + \dim \text{Nul } A = n.$$

§4.6 RANK

Example: If a 12×15 matrix has a 4 dimensional null space, what is rank A ?

§4.6 RANK

Example: If a 12×15 matrix has a 4 dimensional null space, what is rank A ?

§4.6 RANK

Example: If a 12×15 matrix has a 4 dimensional null space, what is rank A ?

Example: Could a 12×15 matrix have a 2 dimensional null space?

§4.6 RANK

Example: If a 12×15 matrix has a 4 dimensional null space, what is rank A ?

Example: Could a 12×15 matrix have a 2 dimensional null space?

Example: A scientist has found two solutions to a homogeneous system of linear 40 equations in 42 variables. The two solutions are not multiples, and all other solutions can be constructed by adding together appropriate multiples of these two solutions. Can the scientist be *certain* that an associated non homogeneous system (with the same coefficients) has a solution?

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

47. The columns of A form a basis for \mathbb{R}^n .

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- 47. The columns of A form a basis for \mathbb{R}^n .
- 48. $\text{Col } A =$

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- 47. The columns of A form a basis for \mathbb{R}^n .
- 48. $\text{Col } A = \mathbb{R}^n$

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- 47. The columns of A form a basis for \mathbb{R}^n .
- 48. $\text{Col } A = \mathbb{R}^n$
- 49. $\dim \text{Col } A =$

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- 47. The columns of A form a basis for \mathbb{R}^n .
- 48. $\text{Col } A = \mathbb{R}^n$
- 49. $\dim \text{Col } A = n$

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- 47. The columns of A form a basis for \mathbb{R}^n .
- 48. $\text{Col } A = \mathbb{R}^n$
- 49. $\dim \text{Col } A = n$
- 50. $\text{rank } A =$

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- 47. The columns of A form a basis for \mathbb{R}^n .
- 48. $\text{Col } A = \mathbb{R}^n$
- 49. $\dim \text{Col } A = n$
- 50. $\text{rank } A = n$

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

47. The columns of A form a basis for \mathbb{R}^n .

48. $\text{Col } A = \mathbb{R}^n$

49. $\dim \text{Col } A = n$

50. $\text{rank } A = n$

51. $\text{Nul } A =$

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- 47. The columns of A form a basis for \mathbb{R}^n .
- 48. $\text{Col } A = \mathbb{R}^n$
- 49. $\dim \text{Col } A = n$
- 50. $\text{rank } A = n$
- 51. $\text{Nul } A = \{\mathbf{0}\}$

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- 47. The columns of A form a basis for \mathbb{R}^n .
- 48. $\text{Col } A = \mathbb{R}^n$
- 49. $\dim \text{Col } A = n$
- 50. $\text{rank } A = n$
- 51. $\text{Nul } A = \{\mathbf{0}\}$
- 52. $\dim \text{Nul } A =$

§4.6 RANK

THEOREM (THE INVERTIBLE MATRIX THEOREM (CONTINUED))

Let A be an $n \times n$ matrix. Then the following statements are each equivalent to the statement that A is an invertible matrix.

- 47. The columns of A form a basis for \mathbb{R}^n .
- 48. $\text{Col } A = \mathbb{R}^n$
- 49. $\dim \text{Col } A = n$
- 50. $\text{rank } A = n$
- 51. $\text{Nul } A = \{\mathbf{0}\}$
- 52. $\dim \text{Nul } A = 0$.