

MATH 341 - LINEAR ALGEBRA

§1.1 - 1.5

Fall 2019

Success isn't the absence of failure,
but going from failure to failure without any loss of
enthusiasm.

CONTACT INFORMATION

Instructor: Brother John Sinkovic

Email: sinkovicj@byui.edu

Office: 232Q RCKS

Phone: 208-496-7562

Office Hours: TTh 1-3pm or by appt through Calendly

WHY LINEAR ALGEBRA?

Linear algebra is used in

WHY LINEAR ALGEBRA?

Linear algebra is used in

- ▶ physics

WHY LINEAR ALGEBRA?

Linear algebra is used in

- ▶ physics
- ▶ engineering

WHY LINEAR ALGEBRA?

Linear algebra is used in

- ▶ physics
- ▶ engineering
- ▶ computer science

WHY LINEAR ALGEBRA?

Linear algebra is used in

- ▶ physics
- ▶ engineering
- ▶ computer science
- ▶ finance

WHY LINEAR ALGEBRA?

Linear algebra is used in

- ▶ physics
- ▶ engineering
- ▶ computer science
- ▶ finance
- ▶ statistics

WHY LINEAR ALGEBRA?

Linear algebra is used in

- ▶ physics
- ▶ engineering
- ▶ computer science
- ▶ finance
- ▶ statistics
- ▶ medical imaging

WHY LINEAR ALGEBRA?

Linear algebra is used in

- ▶ physics
- ▶ engineering
- ▶ computer science
- ▶ finance
- ▶ statistics
- ▶ medical imaging
- ▶ facial recognition

WHY LINEAR ALGEBRA?

Linear algebra is used in

- ▶ physics
- ▶ engineering
- ▶ computer science
- ▶ finance
- ▶ statistics
- ▶ medical imaging
- ▶ facial recognition
- ▶ petrochemical exploration

WHY LINEAR ALGEBRA?

Linear algebra is used in

- ▶ physics
- ▶ engineering
- ▶ computer science
- ▶ finance
- ▶ statistics
- ▶ medical imaging
- ▶ facial recognition
- ▶ petrochemical exploration
- ▶ quantum information theory

§1.1 SYSTEMS OF LINEAR EQUATIONS

DEFINITION

A **linear equation** in the unknowns x_1, \dots, x_n is an equation of the form

§1.1 SYSTEMS OF LINEAR EQUATIONS

DEFINITION

A **linear equation** in the unknowns x_1, \dots, x_n is an equation of the form

$$a_1x_1 + \cdots + a_nx_n = b,$$

where the coefficients a_1, \dots, a_n and the value b are real numbers.

§1.1 SYSTEMS OF LINEAR EQUATIONS

DEFINITION

A **linear equation** in the unknowns x_1, \dots, x_n is an equation of the form

$$a_1x_1 + \cdots + a_nx_n = b,$$

where the coefficients a_1, \dots, a_n and the value b are real numbers.

Many real world applications will involve equations with hundreds or thousands of unknowns.

§1.1 SYSTEMS OF LINEAR EQUATIONS

Examples:

$$x_1 + 3x_2 - 2x_3 = 15$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

Examples:

$$x_1 + 3x_2 - 2x_3 = 15$$

$$x_1 - (\pi + \sqrt{2})x_2 = x_3 - \sin(5)x_4 + 3^2$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

Examples:

$$x_1 + 3x_2 - 2x_3 = 15$$

$$x_1 - (\pi + \sqrt{2})x_2 = x_3 - \sin(5)x_4 + 3^2$$

Non-Examples:

$$x_1 + x_2^2 - x_2 = 1$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

Examples:

$$x_1 + 3x_2 - 2x_3 = 15$$

$$x_1 - (\pi + \sqrt{2})x_2 = x_3 - \sin(5)x_4 + 3^2$$

Non-Examples:

$$x_1 + x_2^2 - x_2 = 1$$

$$\sin(x_1) + \sqrt{x_2} - x_1 \cdot x_4 = 5.$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

A **system of linear equations** is a collection of equations which we try to solve simultaneously.

§1.1 SYSTEMS OF LINEAR EQUATIONS

A **system of linear equations** is a collection of equations which we try to solve simultaneously.

Example: The following equations

$$x_1 + x_2 - 2x_3 = 1$$

$$3x_1 - x_2 + 3x_3 = 4$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

A **system of linear equations** is a collection of equations which we try to solve simultaneously.

Example: The following equations

$$x_1 + x_2 - 2x_3 = 1$$

$$3x_1 - x_2 + 3x_3 = 4$$

have solution $x_1 = 1, x_2 = 2, x_3 = 1$.

§1.1 SYSTEMS OF LINEAR EQUATIONS

A **system of linear equations** is a collection of equations which we try to solve simultaneously.

Example: The following equations

$$x_1 + x_2 - 2x_3 = 1$$

$$3x_1 - x_2 + 3x_3 = 4$$

have solution $x_1 = 1, x_2 = 2, x_3 = 1$.

Two systems are called **equivalent** if they have the same **solution set**.

§1.1 SYSTEMS OF LINEAR EQUATIONS

A **system of linear equations** is a collection of equations which we try to solve simultaneously.

Example: The following equations

$$x_1 + x_2 - 2x_3 = 1$$

$$3x_1 - x_2 + 3x_3 = 4$$

have solution $x_1 = 1, x_2 = 2, x_3 = 1$.

Two systems are called **equivalent** if they have the same **solution set**.

Example: The following system is equivalent to the above example. Notice that it also has solution $x_1 = 1, x_2 = 2, x_3 = 1$:

§1.1 SYSTEMS OF LINEAR EQUATIONS

A **system of linear equations** is a collection of equations which we try to solve simultaneously.

Example: The following equations

$$\begin{aligned}x_1 + x_2 - 2x_3 &= 1 \\ 3x_1 - x_2 + 3x_3 &= 4\end{aligned}$$

have solution $x_1 = 1, x_2 = 2, x_3 = 1$.

Two systems are called **equivalent** if they have the same **solution set**.

Example: The following system is equivalent to the above example. Notice that it also has solution $x_1 = 1, x_2 = 2, x_3 = 1$:

$$\begin{aligned}2x_1 + 2x_2 - 4x_3 &= 2 \\ 2x_1 - 2x_2 + 5x_3 &= 3.\end{aligned}$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:

§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:

The solution set correspond to intersection points between the lines determined by the equations.

§1.1 SYSTEMS OF LINEAR EQUATIONS


2 LINEAR EQUATIONS IN 2 VARIABLES:

The solution set correspond to intersection points between the lines determined by the equations. Three cases:

§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:


The solution set correspond to intersection points between the lines determined by the equations. Three cases:



§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:

The solution set correspond to intersection points between the lines determined by the equations. Three cases:




► One solution

§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:

The solution set correspond to intersection points between the lines determined by the equations. Three cases:



- ▶ One solution
- ▶ Example:



$$y - x = 2$$

$$y + x = 1$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:

The solution set correspond to intersection points between the lines determined by the equations. Three cases:



► One solution

► Example:


$$y - x = 2$$

$$y + x = 1$$


§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:

The solution set correspond to intersection points between the lines determined by the equations. Three cases:



► One solution



► No solution

► Example:


$$y - x = 2$$

$$y + x = 1$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:

The solution set correspond to intersection points between the lines determined by the equations. Three cases:




► One solution

► Example:

$$y - x = 2$$

$$y + x = 1$$



► No solution

► Example:


$$y + x = 5$$

$$y + x = -1$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:

The solution set correspond to intersection points between the lines determined by the equations. Three cases:




► One solution

► Example:

$$y - x = 2$$

$$y + x = 1$$




► No solution

► Example:

$$y + x = 5$$


$$y + x = -1$$



§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:

The solution set correspond to intersection points between the lines determined by the equations. Three cases:




► One solution

► Example:

$$y - x = 2$$

$$y + x = 1$$




► No solution

► Example:

$$y + x = 5$$

$$y + x = -1$$




► ∞ solutions

§1.1 SYSTEMS OF LINEAR EQUATIONS

2 LINEAR EQUATIONS IN 2 VARIABLES:

The solution set correspond to intersection points between the lines determined by the equations. Three cases:




► One solution

► Example:

$$y - x = 2$$

$$y + x = 1$$




► No solution

► Example:

$$y + x = 5$$

$$y + x = -1$$



► ∞ solutions

► Example:

$$2y + 2x = -2$$

$$y + x = -1$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

THEOREM

A system of linear equations has either

§1.1 SYSTEMS OF LINEAR EQUATIONS

THEOREM

A system of linear equations has either

1. no solution,

§1.1 SYSTEMS OF LINEAR EQUATIONS

THEOREM

A system of linear equations has either

1. no solution,
2. exactly one solution, or

§1.1 SYSTEMS OF LINEAR EQUATIONS

THEOREM

A system of linear equations has either


1. no solution,
2. exactly one solution, or
3. infinitely many solutions.

§1.1 SYSTEMS OF LINEAR EQUATIONS

THEOREM

A system of linear equations has either

1. no solution,
2. exactly one solution, or
3. infinitely many solutions.





§1.1 SYSTEMS OF LINEAR EQUATIONS

THEOREM

A system of linear equations has either

1. no solution,
2. exactly one solution, or
3. infinitely many solutions.






§1.1 SYSTEMS OF LINEAR EQUATIONS

THEOREM

A system of linear equations has either

1. no solution,
2. exactly one solution, or
3. infinitely many solutions.






§1.1 SYSTEMS OF LINEAR EQUATIONS

THEOREM

A system of linear equations has either

1. no solution,
2. exactly one solution, or
3. infinitely many solutions.






A system with at least one solution is called **consistent**.

§1.1 SYSTEMS OF LINEAR EQUATIONS

THEOREM

A system of linear equations has either

1. no solution,
2. exactly one solution, or
3. infinitely many solutions.



A system with at least one solution is called **consistent**. A system with no solutions is called **inconsistent**.

§1.1 SYSTEMS OF LINEAR EQUATIONS

EXISTENCE AND UNIQUENESS QUESTIONS

§1.1 SYSTEMS OF LINEAR EQUATIONS

EXISTENCE AND UNIQUENESS QUESTIONS

For a given system of linear equations we can ask

§1.1 SYSTEMS OF LINEAR EQUATIONS

EXISTENCE AND UNIQUENESS QUESTIONS

For a given system of linear equations we can ask

1. does at least one solution exist (i.e. is the system consistent?)

§1.1 SYSTEMS OF LINEAR EQUATIONS

EXISTENCE AND UNIQUENESS QUESTIONS

For a given system of linear equations we can ask

1. does at least one solution exist (i.e. is the system consistent?)
2. if a solution exists, is there only one solution (i.e. is the solution unique?)

§1.1 SYSTEMS OF LINEAR EQUATIONS

For a given system of equations

$$\begin{array}{rcccccccl} x_1 & + & 3x_2 & + & 2x_3 & + & 3x_4 & = & -4 \\ & & x_2 & - & 2x_3 & - & 2x_4 & = & 3 \\ - & x_1 & - & 3x_2 & + & 2x_3 & + & x_4 & = & 4 \end{array}$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

For a given system of equations

$$\begin{array}{cccccccl} x_1 & + & 3x_2 & + & 2x_3 & + & 3x_4 & = & -4 \\ & & x_2 & - & 2x_3 & - & 2x_4 & = & 3 \\ - & x_1 & - & 3x_2 & + & 2x_3 & + & x_4 & = & 4 \end{array}$$

we can express it as a **matrix of coefficients**

$$\begin{bmatrix} 1 & 3 & 2 & 3 \\ 0 & 1 & -2 & -2 \\ -1 & -3 & 2 & 1 \end{bmatrix}$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

For a given system of equations

$$\begin{array}{cccccccl} x_1 & + & 3x_2 & + & 2x_3 & + & 3x_4 & = & -4 \\ & & x_2 & - & 2x_3 & - & 2x_4 & = & 3 \\ - & x_1 & - & 3x_2 & + & 2x_3 & + & x_4 & = & 4 \end{array}$$

we can express it as a **matrix of coefficients**

$$\begin{bmatrix} 1 & 3 & 2 & 3 \\ 0 & 1 & -2 & -2 \\ -1 & -3 & 2 & 1 \end{bmatrix}$$

or as an **augmented matrix**

$$\left[\begin{array}{cccc|c} 1 & 3 & 2 & 3 & -4 \\ 0 & 1 & -2 & -2 & 3 \\ -1 & -3 & 2 & 1 & 4 \end{array} \right].$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

The matrix

$$\begin{bmatrix} 1 & 3 & 2 & 3 \\ 0 & 1 & -2 & -2 \\ -1 & -3 & 2 & 1 \end{bmatrix}$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

The matrix

$$\begin{bmatrix} 1 & 3 & 2 & 3 \\ 0 & 1 & -2 & -2 \\ -1 & -3 & 2 & 1 \end{bmatrix}$$

is a 3×4 **matrix**, because it has **3 rows** (horizontal)

§1.1 SYSTEMS OF LINEAR EQUATIONS

The matrix

$$\begin{bmatrix} 1 & 3 & 2 & 3 \\ 0 & 1 & -2 & -2 \\ -1 & -3 & 2 & 1 \end{bmatrix}$$

is a 3×4 **matrix**, because it has 3 **rows** (horizontal), and 4 **columns** (vertical).

§1.1 SYSTEMS OF LINEAR EQUATIONS

The matrix

$$\begin{bmatrix} 1 & 3 & 2 & 3 \\ 0 & 1 & -2 & -2 \\ -1 & -3 & 2 & 1 \end{bmatrix}$$

is a 3×4 **matrix**, because it has 3 **rows** (horizontal), and 4 **columns** (vertical).

A matrix with m rows and n columns is called an $\mathbf{m} \times \mathbf{n}$ **matrix**.

§1.1 SYSTEMS OF LINEAR EQUATIONS

We can modify an augmented matrix without changing the solutions of the corresponding system:

§1.1 SYSTEMS OF LINEAR EQUATIONS

We can modify an augmented matrix without changing the solutions of the corresponding system:

ELEMENTARY ROW OPERATIONS

§1.1 SYSTEMS OF LINEAR EQUATIONS

We can modify an augmented matrix without changing the solutions of the corresponding system:

ELEMENTARY ROW OPERATIONS

1. Replace one row by the sum of itself and a multiple of another row.

§1.1 SYSTEMS OF LINEAR EQUATIONS

We can modify an augmented matrix without changing the solutions of the corresponding system:

ELEMENTARY ROW OPERATIONS

1. Replace one row by the sum of itself and a multiple of another row.
2. Interchange two rows.

§1.1 SYSTEMS OF LINEAR EQUATIONS

We can modify an augmented matrix without changing the solutions of the corresponding system:

ELEMENTARY ROW OPERATIONS

1. Replace one row by the sum of itself and a multiple of another row.
2. Interchange two rows.
3. Multiply all entries in a row by a nonzero number.

§1.1 SYSTEMS OF LINEAR EQUATIONS

We can modify an augmented matrix without changing the solutions of the corresponding system:

ELEMENTARY ROW OPERATIONS

1. Replace one row by the sum of itself and a multiple of another row.
2. Interchange two rows.
3. Multiply all entries in a row by a nonzero number.

Two matrices that can be joined by a sequence of elementary row operations are called **row equivalent**.

§1.1 SYSTEMS OF LINEAR EQUATIONS

We can modify an augmented matrix without changing the solutions of the corresponding system:

ELEMENTARY ROW OPERATIONS

1. Replace one row by the sum of itself and a multiple of another row.
2. Interchange two rows.
3. Multiply all entries in a row by a nonzero number.

Two matrices that can be joined by a sequence of elementary row operations are called **row equivalent**.

THEOREM

Elementary row operations do not change the solution set of the associated system of equations

§1.1 SYSTEMS OF LINEAR EQUATIONS

We can modify an augmented matrix without changing the solutions of the corresponding system:

ELEMENTARY ROW OPERATIONS

1. Replace one row by the sum of itself and a multiple of another row.
2. Interchange two rows.
3. Multiply all entries in a row by a nonzero number.

Two matrices that can be joined by a sequence of elementary row operations are called **row equivalent**.

THEOREM

Elementary row operations do not change the solution set of the associated system of equations (row equivalent augmented matrices have the same solution sets).

§1.1 SYSTEMS OF LINEAR EQUATIONS

Example: Is the following system consistent?

$$\begin{array}{cccccccl} x_1 & + & 3x_2 & + & 2x_3 & + & 3x_4 & = & -4 \\ & & x_2 & - & 2x_3 & - & 2x_4 & = & 3 \\ - & x_1 & - & 3x_2 & + & 2x_3 & + & x_4 & = & 4 \end{array}$$

§1.1 SYSTEMS OF LINEAR EQUATIONS

Example: Is the following system consistent?

$$\begin{array}{cccccccl} x_1 & + & 3x_2 & + & 2x_3 & + & 3x_4 & = & -4 \\ & & x_2 & - & 2x_3 & - & 2x_4 & = & 3 \\ - & x_1 & - & 3x_2 & + & 2x_3 & + & x_4 & = & 4 \end{array}$$

If so, find a solution.

§1.1 SYSTEMS OF LINEAR EQUATIONS

Example: Is the following system consistent?

$$\begin{array}{cccccccl} x_1 & + & 3x_2 & + & 2x_3 & + & 3x_4 & = & -4 \\ & & x_2 & - & 2x_3 & - & 2x_4 & = & 3 \\ - & x_1 & - & 3x_2 & + & 2x_3 & + & x_4 & = & 4 \end{array}$$

If so, find a solution. How many solutions does it have?

§1.2 ROW REDUCTION AND ECHELON FORMS

For any nonzero row of a matrix A , a **leading entry** is the first (left-most) nonzero entry.

§1.2 ROW REDUCTION AND ECHELON FORMS

For any nonzero row of a matrix A , a **leading entry** is the first (left-most) nonzero entry.

Example: For the matrix

$$\begin{bmatrix} -2 & 2 & 1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 4 \end{bmatrix},$$

§1.2 ROW REDUCTION AND ECHELON FORMS

For any nonzero row of a matrix A , a **leading entry** is the first (left-most) nonzero entry.

Example: For the matrix

$$\begin{bmatrix} -2 & 2 & 1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 4 \end{bmatrix},$$

the first row has leading entry -2 ,

§1.2 ROW REDUCTION AND ECHELON FORMS

For any nonzero row of a matrix A , a **leading entry** is the first (left-most) nonzero entry.

Example: For the matrix

$$\begin{bmatrix} -2 & 2 & 1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 4 \end{bmatrix},$$

the first row has leading entry -2 , the second row doesn't have a leading entry (it's all zeros),

§1.2 ROW REDUCTION AND ECHELON FORMS

For any nonzero row of a matrix A , a **leading entry** is the first (left-most) nonzero entry.

Example: For the matrix

$$\begin{bmatrix} -2 & 2 & 1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 4 \end{bmatrix},$$

the first row has leading entry -2 , the second row doesn't have a leading entry (it's all zeros), while the leading entry of the third row is 3 .

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION ((REDUCED ROW) ECHELON FORM)

A matrix is said to be in **echelon form** if the following three conditions hold:

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION ((REDUCED ROW) ECHELON FORM)

A matrix is said to be in **echelon form** if the following three conditions hold:

1. All nonzero rows are above any rows with all zeros.

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION ((REDUCED ROW) ECHELON FORM)

A matrix is said to be in **echelon form** if the following three conditions hold:

1. All nonzero rows are above any rows with all zeros.
2. Each leading entry of a row is in a column to the right of the leading entry of the row above it.

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION ((REDUCED ROW) ECHELON FORM)

A matrix is said to be in **echelon form** if the following three conditions hold:

1. All nonzero rows are above any rows with all zeros.
2. Each leading entry of a row is in a column to the right of the leading entry of the row above it.
3. All entries in a column below a leading entry are zeros.

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION ((REDUCED ROW) ECHELON FORM)

A matrix is said to be in **echelon form** if the following three conditions hold:

1. All nonzero rows are above any rows with all zeros.
2. Each leading entry of a row is in a column to the right of the leading entry of the row above it.
3. All entries in a column below a leading entry are zeros.

Furthermore, it is in **reduced row echelon form (RREF)** if

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION ((REDUCED ROW) ECHELON FORM)

A matrix is said to be in **echelon form** if the following three conditions hold:

1. All nonzero rows are above any rows with all zeros.
2. Each leading entry of a row is in a column to the right of the leading entry of the row above it.
3. All entries in a column below a leading entry are zeros.

Furthermore, it is in **reduced row echelon form (RREF)** if

4. The leading entry in each nonzero row is 1.

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION ((REDUCED ROW) ECHELON FORM)

A matrix is said to be in **echelon form** if the following three conditions hold:

1. All nonzero rows are above any rows with all zeros.
2. Each leading entry of a row is in a column to the right of the leading entry of the row above it.
3. All entries in a column below a leading entry are zeros.

Furthermore, it is in **reduced row echelon form (RREF)** if

4. The leading entry in each nonzero row is 1.
5. Each leading 1 is the only nonzero entry in its column.

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Which of the following matrices are in echelon form, and which are in RREF?

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Which of the following matrices are in echelon form, and which are in RREF?

$$\begin{bmatrix} -2 & 2 & 1 & 3 \\ 0 & 0 & 3 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Which of the following matrices are in echelon form, and which are in RREF?

$$\begin{bmatrix} -2 & 2 & 1 & 3 \\ 0 & 0 & 3 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \text{ is in echelon form.}$$

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Which of the following matrices are in echelon form, and which are in RREF?

$$\begin{bmatrix} -2 & 2 & 1 & 3 \\ 0 & 0 & 3 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \text{ is in echelon form.}$$

$$\begin{bmatrix} -2 & 2 & 1 \\ 0 & 2 & 3 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Which of the following matrices are in echelon form, and which are in RREF?

$$\begin{bmatrix} -2 & 2 & 1 & 3 \\ 0 & 0 & 3 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \text{ is in echelon form.}$$

$$\begin{bmatrix} -2 & 2 & 1 \\ 0 & 2 & 3 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{ is not in echelon form (or RREF).}$$

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Which of the following matrices are in echelon form, and which are in RREF?

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Which of the following matrices are in echelon form, and which are in RREF?

$$\begin{bmatrix} 1 & 2 & 0 & 0 & 5 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix},$$

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Which of the following matrices are in echelon form, and which are in RREF?

$$\begin{bmatrix} 1 & 2 & 0 & 0 & 5 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix}, \text{ is in RREF.}$$

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Which of the following matrices are in echelon form, and which are in RREF?

$$\begin{bmatrix} 1 & 2 & 0 & 0 & 5 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix}, \text{ is in RREF.}$$

$$\begin{bmatrix} -2 & 2 & 1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 4 \end{bmatrix}$$

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Which of the following matrices are in echelon form, and which are in RREF?

$$\begin{bmatrix} 1 & 2 & 0 & 0 & 5 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix}, \text{ is in RREF.}$$

$$\begin{bmatrix} -2 & 2 & 1 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 4 \end{bmatrix} \text{ is not in echelon form (or RREF).}$$

§1.2 ROW REDUCTION AND ECHELON FORMS

THEOREM (EXISTENCE AND UNIQUENESS OF RREF)

Each matrix is row equivalent to one and only one matrix in RREF.

§1.2 ROW REDUCTION AND ECHELON FORMS

THEOREM (EXISTENCE AND UNIQUENESS OF RREF)

Each matrix is row equivalent to one and only one matrix in RREF.

DEFINITION

A **pivot position** in a matrix A is an entry of a matrix which corresponds to a leading 1 in the RREF of A .

§1.2 ROW REDUCTION AND ECHELON FORMS

THEOREM (EXISTENCE AND UNIQUENESS OF RREF)

Each matrix is row equivalent to one and only one matrix in RREF.

DEFINITION

A **pivot position** in a matrix A is an entry of a matrix which corresponds to a leading 1 in the RREF of A .

A **pivot column** of A is a column of A which contains a pivot position.

§1.2 ROW REDUCTION AND ECHELON FORMS

THEOREM (EXISTENCE AND UNIQUENESS OF RREF)

Each matrix is row equivalent to one and only one matrix in RREF.

DEFINITION

A **pivot position** in a matrix A is an entry of a matrix which corresponds to a leading 1 in the RREF of A .

A **pivot column** of A is a column of A which contains a pivot position.

NOTE: To locate the pivot positions of a matrix A , we need to use elementary row operations to put A in RREF.

§1.2 ROW REDUCTION AND ECHELON FORMS

THEOREM (EXISTENCE AND UNIQUENESS OF RREF)

Each matrix is row equivalent to one and only one matrix in RREF.

DEFINITION

A **pivot position** in a matrix A is an entry of a matrix which corresponds to a leading 1 in the RREF of A .

A **pivot column** of A is a column of A which contains a pivot position.

NOTE: To locate the pivot positions of a matrix A , we need to use elementary row operations to put A in RREF. The positions (columns) of the leading ones in the RREF tells us the locations of the pivots (pivot columns) in the original matrix A .

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Find the pivot positions and columns of the matrix

$$\begin{bmatrix} 3 & 2 & 1 & 0 \\ -6 & -4 & 3 & 1 \\ 9 & 6 & -2 & -1 \end{bmatrix}.$$

§1.2 ROW REDUCTION AND ECHELON FORMS

ROW REDUCTION ALGORITHM

§1.2 ROW REDUCTION AND ECHELON FORMS

ROW REDUCTION ALGORITHM

1. Begin with the leftmost nonzero column. This is a pivot column. The pivot position is at the top.

§1.2 ROW REDUCTION AND ECHELON FORMS

ROW REDUCTION ALGORITHM

1. Begin with the leftmost nonzero column. This is a pivot column. The pivot position is at the top.
2. Select a nonzero entry in the pivot column as a pivot. If necessary, interchange rows to move this entry into the pivot position.

§1.2 ROW REDUCTION AND ECHELON FORMS

ROW REDUCTION ALGORITHM

1. Begin with the leftmost nonzero column. This is a pivot column. The pivot position is at the top.
2. Select a nonzero entry in the pivot column as a pivot. If necessary, interchange rows to move this entry into the pivot position.
3. Use row replacement operations to create zeros in all positions below the pivot.

§1.2 ROW REDUCTION AND ECHELON FORMS

ROW REDUCTION ALGORITHM

1. Begin with the leftmost nonzero column. This is a pivot column. The pivot position is at the top.
2. Select a nonzero entry in the pivot column as a pivot. If necessary, interchange rows to move this entry into the pivot position.
3. Use row replacement operations to create zeros in all positions below the pivot.
4. Ignore the row containing the pivot position and all rows above it. Repeat the process until there are no more nonzero rows to modify.

§1.2 ROW REDUCTION AND ECHELON FORMS

ROW REDUCTION ALGORITHM

1. Begin with the leftmost nonzero column. This is a pivot column. The pivot position is at the top.
2. Select a nonzero entry in the pivot column as a pivot. If necessary, interchange rows to move this entry into the pivot position.
3. Use row replacement operations to create zeros in all positions below the pivot.
4. Ignore the row containing the pivot position and all rows above it. Repeat the process until there are no more nonzero rows to modify.
5. Beginning with the rightmost pivot and working upward and to the left, create zeros above each pivot. If a pivot is not 1, make it 1 by a scaling operation.

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Use the row reduction algorithm to find the RREF matrix which is row equivalent to

$$\begin{bmatrix} 0 & 3 & -6 & 4 & -5 \\ 3 & -7 & 8 & 8 & 9 \\ 3 & -9 & 12 & 6 & 15 \end{bmatrix}.$$

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Find solutions to the following system

$$\begin{array}{ccccccccc} & & 3x_2 & - & 6x_3 & + & 4x_4 & = & -5 \\ 3x_1 & - & 7x_2 & + & 8x_3 & + & 8x_4 & = & 9 \ . \\ 3x_1 & - & 9x_2 & + & 12x_3 & + & 6x_4 & = & 15 \end{array}$$

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Find solutions to the following system

$$\begin{array}{rrcrcl} & 3x_2 & - & 6x_3 & + & 4x_4 & = & -5 \\ 3x_1 & - & 7x_2 & + & 8x_3 & + & 8x_4 & = & 9 \\ 3x_1 & - & 9x_2 & + & 12x_3 & + & 6x_4 & = & 15 \end{array} .$$

Hint: The augmented matrix $\left[\begin{array}{cccc|c} 0 & 3 & -6 & 4 & -5 \\ 3 & -7 & 8 & 8 & 9 \\ 3 & -9 & 12 & 6 & 15 \end{array} \right]$ row

reduces to

$$\left[\begin{array}{cccc|c} 1 & 0 & -2 & 0 & -24 \\ 0 & 1 & -2 & 0 & -7 \\ 0 & 0 & 0 & 1 & 4 \end{array} \right] .$$

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION (BASIC AND FREE VARIABLES)

Variables corresponding to pivot columns are called **basic variables**,

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION (BASIC AND FREE VARIABLES)

Variables corresponding to pivot columns are called **basic variables**, other variables are called **free variables**.

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION (BASIC AND FREE VARIABLES)

Variables corresponding to pivot columns are called **basic variables**, other variables are called **free variables**.

In the above example, x_1, x_2 , and x_4 are basic variables, while x_3 is a free variable.

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION (BASIC AND FREE VARIABLES)

Variables corresponding to pivot columns are called **basic variables**, other variables are called **free variables**.

In the above example, x_1, x_2 , and x_4 are basic variables, while x_3 is a free variable.

After putting the matrix in RREF, we can solve for all basic variables in terms of the free variables.

§1.2 ROW REDUCTION AND ECHELON FORMS

DEFINITION (BASIC AND FREE VARIABLES)

Variables corresponding to pivot columns are called **basic variables**, other variables are called **free variables**.

In the above example, x_1, x_2 , and x_4 are basic variables, while x_3 is a free variable.

After putting the matrix in RREF, we can solve for all basic variables in terms of the free variables.

Different choices of values for the free variables give different solutions.

§1.2 ROW REDUCTION AND ECHELON FORMS

Example: Find the solutions to the following system

$$\begin{array}{rrcrcl} x & + & y & + & 3z & = & -1 \\ 2x & + & 2y & + & 6z & = & 4 \\ 3x & + & 2y & + & 1z & = & 1 \quad . \end{array}$$

§1.2 ROW REDUCTION AND ECHELON FORMS

THEOREM (EXISTENCE AND UNIQUENESS THEOREM)

A linear system is consistent if and only if the right-most column of the augmented matrix is *not* a pivot column

§1.2 ROW REDUCTION AND ECHELON FORMS

THEOREM (EXISTENCE AND UNIQUENESS THEOREM)

A linear system is consistent if and only if the right-most column of the augmented matrix is *not* a pivot column i.e. if and only if an echelon form of the augmented matrix has no row of the form

$$\left[\begin{array}{cccc|c} 0 & 0 & \cdots & 0 & b \end{array} \right], \quad \text{where } b \text{ is nonzero.}$$

§1.2 ROW REDUCTION AND ECHELON FORMS

THEOREM (EXISTENCE AND UNIQUENESS THEOREM)

A linear system is consistent if and only if the right-most column of the augmented matrix is *not* a pivot column i.e. if and only if an echelon form of the augmented matrix has no row of the form

$$\left[\begin{array}{cccc|c} 0 & 0 & \cdots & 0 & b \end{array} \right], \quad \text{where } b \text{ is nonzero.}$$

If the system is consistent, then it either has a unique solution if it has no free variables, or an infinite number of solutions if it has at least one free variable.

§1.3 VECTOR EQUATIONS

DEFINITION

A **vector** is an $n \times 1$ matrix (matrix with only one column).

§1.3 VECTOR EQUATIONS

DEFINITION

A **vector** is an $n \times 1$ matrix (matrix with only one column).

The set of all vectors with n rows is denoted \mathbb{R}^n .

§1.3 VECTOR EQUATIONS

DEFINITION

A **vector** is an $n \times 1$ matrix (matrix with only one column).

The set of all vectors with n rows is denoted \mathbb{R}^n .

Example: $\begin{bmatrix} 1 \\ -2 \end{bmatrix}$ and $\begin{bmatrix} -4 \\ 11 \end{bmatrix}$ are both vectors in \mathbb{R}^2 .

§1.3 VECTOR EQUATIONS

DEFINITION

A **vector** is an $n \times 1$ matrix (matrix with only one column).

The set of all vectors with n rows is denoted \mathbb{R}^n .

Example: $\begin{bmatrix} 1 \\ -2 \end{bmatrix}$ and $\begin{bmatrix} -4 \\ 11 \end{bmatrix}$ are both vectors in \mathbb{R}^2 .

Example: $\begin{bmatrix} 3 \\ 5 \\ 2 \\ 1 \end{bmatrix}$ is a vector in \mathbb{R}^4 ,

§1.3 VECTOR EQUATIONS

DEFINITION

A **vector** is an $n \times 1$ matrix (matrix with only one column).

The set of all vectors with n rows is denoted \mathbb{R}^n .

Example: $\begin{bmatrix} 1 \\ -2 \end{bmatrix}$ and $\begin{bmatrix} -4 \\ 11 \end{bmatrix}$ are both vectors in \mathbb{R}^2 .

Example: $\begin{bmatrix} 3 \\ 5 \\ 2 \\ 1 \end{bmatrix}$ is a vector in \mathbb{R}^4 , and we write $\begin{bmatrix} 3 \\ 5 \\ 2 \\ 1 \end{bmatrix} \in \mathbb{R}^4$.

§1.3 VECTOR EQUATIONS

Vectors in \mathbb{R}^2 can be identified with points in the plane:

$$\begin{bmatrix} a \\ b \end{bmatrix} \in \mathbb{R}^2$$

§1.3 VECTOR EQUATIONS

Vectors in \mathbb{R}^2 can be identified with points in the plane:

$$\begin{bmatrix} a \\ b \end{bmatrix} \in \mathbb{R}^2 \quad \longleftrightarrow \quad \text{the point } (a, b) \text{ in the } xy\text{-plane.}$$

§1.3 VECTOR EQUATIONS

Vectors in \mathbb{R}^2 can be identified with points in the plane:

$$\begin{bmatrix} a \\ b \end{bmatrix} \in \mathbb{R}^2 \quad \longleftrightarrow \quad \text{the point } (a, b) \text{ in the } xy\text{-plane.}$$

Vectors can be added and multiplied together by real scalars:

§1.3 VECTOR EQUATIONS

Vectors in \mathbb{R}^2 can be identified with points in the plane:

$$\begin{bmatrix} a \\ b \end{bmatrix} \in \mathbb{R}^2 \quad \longleftrightarrow \quad \text{the point } (a, b) \text{ in the } xy\text{-plane.}$$

Vectors can be added and multiplied together by real scalars:

$$\begin{bmatrix} 1 \\ -2 \end{bmatrix} + \begin{bmatrix} -4 \\ 11 \end{bmatrix} = \begin{bmatrix} -3 \\ 9 \end{bmatrix}$$

§1.3 VECTOR EQUATIONS

Vectors in \mathbb{R}^2 can be identified with points in the plane:

$$\begin{bmatrix} a \\ b \end{bmatrix} \in \mathbb{R}^2 \quad \longleftrightarrow \quad \text{the point } (a, b) \text{ in the } xy\text{-plane.}$$

Vectors can be added and multiplied together by real scalars:

$$\begin{bmatrix} 1 \\ -2 \end{bmatrix} + \begin{bmatrix} -4 \\ 11 \end{bmatrix} = \begin{bmatrix} -3 \\ 9 \end{bmatrix} \quad \text{and} \quad 5 \cdot \begin{bmatrix} -4 \\ 11 \end{bmatrix} = \begin{bmatrix} -20 \\ 55 \end{bmatrix}.$$

§1.3 VECTOR EQUATIONS

Vectors in \mathbb{R}^2 can be identified with points in the plane:

$$\begin{bmatrix} a \\ b \end{bmatrix} \in \mathbb{R}^2 \quad \longleftrightarrow \quad \text{the point } (a, b) \text{ in the } xy\text{-plane.}$$

Vectors can be added and multiplied together by real scalars:

$$\begin{bmatrix} 1 \\ -2 \end{bmatrix} + \begin{bmatrix} -4 \\ 11 \end{bmatrix} = \begin{bmatrix} -3 \\ 9 \end{bmatrix} \quad \text{and} \quad 5 \cdot \begin{bmatrix} -4 \\ 11 \end{bmatrix} = \begin{bmatrix} -20 \\ 55 \end{bmatrix}.$$

Warning: We don't multiply vectors together (though we will in the future).

§1.3 VECTOR EQUATIONS

Formally we have

$$\begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} + \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{bmatrix} =$$

§1.3 VECTOR EQUATIONS

Formally we have

$$\begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} + \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{bmatrix} = \begin{bmatrix} u_1 + w_1 \\ u_2 + w_2 \\ \vdots \\ u_n + w_n \end{bmatrix}$$

§1.3 VECTOR EQUATIONS

Formally we have

$$\begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} + \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{bmatrix} = \begin{bmatrix} u_1 + w_1 \\ u_2 + w_2 \\ \vdots \\ u_n + w_n \end{bmatrix}$$

and

$$c \cdot \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} =$$

§1.3 VECTOR EQUATIONS

Formally we have


$$\begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} + \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{bmatrix} = \begin{bmatrix} u_1 + w_1 \\ u_2 + w_2 \\ \vdots \\ u_n + w_n \end{bmatrix}$$

and

$$c \cdot \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} cu_1 \\ cu_2 \\ \vdots \\ cu_n \end{bmatrix}.$$


§1.3 VECTOR EQUATIONS

Geometric view of vector addition:




§1.3 VECTOR EQUATIONS

Geometric view of vector addition:




§1.3 VECTOR EQUATIONS

Geometric view of vector addition:




§1.3 VECTOR EQUATIONS

Geometric view of scalar multiplication:




§1.3 VECTOR EQUATIONS

Geometric view of scalar multiplication:



§1.3 VECTOR EQUATIONS

Geometric view of scalar multiplication:



§1.3 VECTOR EQUATIONS

A vector in \mathbb{R}^n is an ordered list of n real numbers, i.e.

$$\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}.$$

Let

$$\mathbf{0} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

denote the zero vector in \mathbb{R}^n .

§1.3 VECTOR EQUATIONS

THEOREM (ALGEBRAIC PROPERTIES OF \mathbb{R}^n)

For all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c, d in \mathbb{R} :

§1.3 VECTOR EQUATIONS

THEOREM (ALGEBRAIC PROPERTIES OF \mathbb{R}^n)

For all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c, d in \mathbb{R} :

1. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$

§1.3 VECTOR EQUATIONS

THEOREM (ALGEBRAIC PROPERTIES OF \mathbb{R}^n)

For all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c, d in \mathbb{R} :

1. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
2. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$

§1.3 VECTOR EQUATIONS

THEOREM (ALGEBRAIC PROPERTIES OF \mathbb{R}^n)

For all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c, d in \mathbb{R} :

1. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
2. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
3. $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$

§1.3 VECTOR EQUATIONS

THEOREM (ALGEBRAIC PROPERTIES OF \mathbb{R}^n)

For all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c, d in \mathbb{R} :

1. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
2. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
3. $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$
4. $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$, where $-\mathbf{u}$ denotes $-1 \cdot \mathbf{u}$

§1.3 VECTOR EQUATIONS

THEOREM (ALGEBRAIC PROPERTIES OF \mathbb{R}^n)

For all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c, d in \mathbb{R} :

1. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
2. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
3. $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$
4. $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$, where $-\mathbf{u}$ denotes $-1 \cdot \mathbf{u}$
5. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$

§1.3 VECTOR EQUATIONS

THEOREM (ALGEBRAIC PROPERTIES OF \mathbb{R}^n)

For all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c, d in \mathbb{R} :

1. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
2. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
3. $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$
4. $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$, where $-\mathbf{u}$ denotes $-1 \cdot \mathbf{u}$
5. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
6. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$

§1.3 VECTOR EQUATIONS

THEOREM (ALGEBRAIC PROPERTIES OF \mathbb{R}^n)

For all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c, d in \mathbb{R} :

1. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
2. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
3. $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$
4. $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$, where $-\mathbf{u}$ denotes $-1 \cdot \mathbf{u}$
5. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
6. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
7. $c(d\mathbf{u}) = (cd)\mathbf{u}$

§1.3 VECTOR EQUATIONS

THEOREM (ALGEBRAIC PROPERTIES OF \mathbb{R}^n)

For all vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c, d in \mathbb{R} :

1. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
2. $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
3. $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$
4. $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$, where $-\mathbf{u}$ denotes $-1 \cdot \mathbf{u}$
5. $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
6. $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
7. $c(d\mathbf{u}) = (cd)\mathbf{u}$
8. $1 \cdot \mathbf{u} = \mathbf{u}$.

§1.3 VECTOR EQUATIONS

DEFINITION (LINEAR COMBINATIONS)

Given vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ in \mathbb{R}^n , and scalars c_1, \dots, c_p , a vector of the form

§1.3 VECTOR EQUATIONS

DEFINITION (LINEAR COMBINATIONS)

Given vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ in \mathbb{R}^n , and scalars c_1, \dots, c_p , a vector of the form

$$\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p$$

§1.3 VECTOR EQUATIONS

DEFINITION (LINEAR COMBINATIONS)

Given vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ in \mathbb{R}^n , and scalars c_1, \dots, c_p , a vector of the form

$$\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p$$

is called a **linear combination** of the vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ with weights c_1, \dots, c_p .

§1.3 VECTOR EQUATIONS

DEFINITION (LINEAR COMBINATIONS)

Given vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ in \mathbb{R}^n , and scalars c_1, \dots, c_p , a vector of the form

$$\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p$$

is called a **linear combination** of the vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ with weights c_1, \dots, c_p .

Examples: The vectors

$$\sqrt{2}\cdot\mathbf{v}_1 + 3\cdot\mathbf{v}_2 - \mathbf{v}_3$$

§1.3 VECTOR EQUATIONS

DEFINITION (LINEAR COMBINATIONS)

Given vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ in \mathbb{R}^n , and scalars c_1, \dots, c_p , a vector of the form

$$\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p$$

is called a **linear combination** of the vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ with weights c_1, \dots, c_p .

Examples: The vectors

$$\sqrt{2}\cdot\mathbf{v}_1+3\cdot\mathbf{v}_2-\mathbf{v}_3 \qquad \mathbf{v}_1 = 1\cdot\mathbf{v}_1+0\cdot\mathbf{v}_2+0\cdot\mathbf{v}_3$$

§1.3 VECTOR EQUATIONS

DEFINITION (LINEAR COMBINATIONS)

Given vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ in \mathbb{R}^n , and scalars c_1, \dots, c_p , a vector of the form

$$\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p$$

is called a **linear combination** of the vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ with weights c_1, \dots, c_p .

Examples: The vectors

$$\sqrt{2}\cdot\mathbf{v}_1+3\cdot\mathbf{v}_2-\mathbf{v}_3 \qquad \mathbf{v}_1 = 1\cdot\mathbf{v}_1+0\cdot\mathbf{v}_2+0\cdot\mathbf{v}_3 \qquad \mathbf{0} = 0\cdot\mathbf{v}_1+0\cdot\mathbf{v}_2+0\cdot\mathbf{v}_3,$$

§1.3 VECTOR EQUATIONS

DEFINITION (LINEAR COMBINATIONS)

Given vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ in \mathbb{R}^n , and scalars c_1, \dots, c_p , a vector of the form

$$\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p$$

is called a **linear combination** of the vectors $\mathbf{v}_1, \dots, \mathbf{v}_p$ with weights c_1, \dots, c_p .

Examples: The vectors

$$\sqrt{2}\cdot\mathbf{v}_1+3\cdot\mathbf{v}_2-\mathbf{v}_3 \qquad \mathbf{v}_1 = 1\cdot\mathbf{v}_1+0\cdot\mathbf{v}_2+0\cdot\mathbf{v}_3 \qquad \mathbf{0} = 0\cdot\mathbf{v}_1+0\cdot\mathbf{v}_2+0\cdot\mathbf{v}_3,$$

are all linear combinations of the vectors $\mathbf{v}_1, \mathbf{v}_2$, and \mathbf{v}_3 .

§1.3 VECTOR EQUATIONS

Example: Can the vector $\mathbf{y} = \begin{bmatrix} 8 \\ -9 \\ 2 \end{bmatrix}$ be written as a linear combination of the vectors

$$\mathbf{v} = \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix}, \quad \text{and} \quad \mathbf{w} = \begin{bmatrix} 2 \\ -5 \\ 2 \end{bmatrix}?$$

§1.3 VECTOR EQUATIONS

A vector equation

$$x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \cdots + x_p\mathbf{v}_p = \mathbf{b}$$

has the same solution set as the linear system corresponding to the augmented matrix

$$\left[\begin{array}{cccc|c} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_p & \mathbf{b} \end{array} \right].$$

§1.3 VECTOR EQUATIONS

A vector equation

$$x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \cdots + x_p\mathbf{v}_p = \mathbf{b}$$

has the same solution set as the linear system corresponding to the augmented matrix

$$\left[\begin{array}{cccc|c} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_p & \mathbf{b} \end{array} \right].$$

Thus \mathbf{b} can be written as a linear combination of the \mathbf{v}_j if and only if the system of linear equations corresponding to the above augmented matrix is consistent.

§1.3 VECTOR EQUATIONS

DEFINITION (SPAN OF VECTORS)

Let $\mathbf{v}_1, \dots, \mathbf{v}_p$ be vectors in \mathbb{R}^n .

§1.3 VECTOR EQUATIONS

DEFINITION (SPAN OF VECTORS)

Let $\mathbf{v}_1, \dots, \mathbf{v}_p$ be vectors in \mathbb{R}^n . Then the **set spanned by the vectors** $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$, which we denote by $\text{Span } \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$, is the set of all vectors which can be written as linear combinations of the \mathbf{v}_j .

§1.3 VECTOR EQUATIONS

DEFINITION (SPAN OF VECTORS)

Let $\mathbf{v}_1, \dots, \mathbf{v}_p$ be vectors in \mathbb{R}^n . Then the **set spanned by the vectors** $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$, which we denote by $\text{Span } \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$, is the set of all vectors which can be written as linear combinations of the \mathbf{v}_j .

In other words $\text{Span } \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is the set of all vectors that can be written as

$$c_1\mathbf{v}_1 + \cdots + c_p\mathbf{v}_p$$

for some weights c_1, \dots, c_p .

§1.3 VECTOR EQUATIONS

Example: Let $\mathbf{v}_1, \mathbf{v}_2$ be vectors in \mathbb{R}^n .

§1.3 VECTOR EQUATIONS

Example: Let $\mathbf{v}_1, \mathbf{v}_2$ be vectors in \mathbb{R}^n . Then since

$$\mathbf{v}_1 = 1 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2$$

§1.3 VECTOR EQUATIONS

Example: Let $\mathbf{v}_1, \mathbf{v}_2$ be vectors in \mathbb{R}^n . Then since

$$\mathbf{v}_1 = 1 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2 \quad \text{and} \quad \mathbf{v}_2 = 0 \cdot \mathbf{v}_1 + 1 \cdot \mathbf{v}_2$$

§1.3 VECTOR EQUATIONS

Example: Let $\mathbf{v}_1, \mathbf{v}_2$ be vectors in \mathbb{R}^n . Then since

$$\mathbf{v}_1 = 1 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2 \quad \text{and} \quad \mathbf{v}_2 = 0 \cdot \mathbf{v}_1 + 1 \cdot \mathbf{v}_2$$

both \mathbf{v}_1 and \mathbf{v}_2 are in $\text{Span} \{ \mathbf{v}_1, \mathbf{v}_2 \}$.

§1.3 VECTOR EQUATIONS

Example: Let $\mathbf{v}_1, \mathbf{v}_2$ be vectors in \mathbb{R}^n . Then since

$$\mathbf{v}_1 = 1 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2 \quad \text{and} \quad \mathbf{v}_2 = 0 \cdot \mathbf{v}_1 + 1 \cdot \mathbf{v}_2$$

both \mathbf{v}_1 and \mathbf{v}_2 are in $\text{Span} \{ \mathbf{v}_1, \mathbf{v}_2 \}$.

Also, since $\mathbf{0} = 0 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2$ we also have that $\mathbf{0}$ is in $\text{Span} \{ \mathbf{v}_1, \mathbf{v}_2 \}$.

§1.3 VECTOR EQUATIONS

Example: Let $\mathbf{v}_1, \mathbf{v}_2$ be vectors in \mathbb{R}^n . Then since

$$\mathbf{v}_1 = 1 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2 \quad \text{and} \quad \mathbf{v}_2 = 0 \cdot \mathbf{v}_1 + 1 \cdot \mathbf{v}_2$$

both \mathbf{v}_1 and \mathbf{v}_2 are in $\text{Span} \{\mathbf{v}_1, \mathbf{v}_2\}$.

Also, since $\mathbf{0} = 0 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2$ we also have that $\mathbf{0}$ is in $\text{Span} \{\mathbf{v}_1, \mathbf{v}_2\}$.

Example: Let $\mathbf{v}_1 = \begin{bmatrix} 3 \\ 1 \\ 5 \end{bmatrix}$, $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \\ 3 \end{bmatrix}$, and $\mathbf{v}_3 = \begin{bmatrix} -3 \\ 2 \\ 2 \end{bmatrix}$.

§1.3 VECTOR EQUATIONS

Example: Let $\mathbf{v}_1, \mathbf{v}_2$ be vectors in \mathbb{R}^n . Then since

$$\mathbf{v}_1 = 1 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2 \quad \text{and} \quad \mathbf{v}_2 = 0 \cdot \mathbf{v}_1 + 1 \cdot \mathbf{v}_2$$

both \mathbf{v}_1 and \mathbf{v}_2 are in $\text{Span} \{\mathbf{v}_1, \mathbf{v}_2\}$.

Also, since $\mathbf{0} = 0 \cdot \mathbf{v}_1 + 0 \cdot \mathbf{v}_2$ we also have that $\mathbf{0}$ is in $\text{Span} \{\mathbf{v}_1, \mathbf{v}_2\}$.

Example: Let $\mathbf{v}_1 = \begin{bmatrix} 3 \\ 1 \\ 5 \end{bmatrix}$, $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \\ 3 \end{bmatrix}$, and $\mathbf{v}_3 = \begin{bmatrix} -3 \\ 2 \\ 2 \end{bmatrix}$.


Is $\mathbf{y} = \begin{bmatrix} 1 \\ 1 \\ 9 \end{bmatrix}$ in $\text{Span} \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$?

§1.3 VECTOR EQUATIONS

For a nonzero vector $\mathbf{v} \in \mathbb{R}^n$, $\text{Span } \{\mathbf{v}\}$ is the set of all vectors of the form $c\mathbf{v}$, for some scalar c .


§1.3 VECTOR EQUATIONS

For a nonzero vector $\mathbf{v} \in \mathbb{R}^n$, $\text{Span}\{\mathbf{v}\}$ is the set of all vectors of the form $c\mathbf{v}$, for some scalar c .




§1.3 VECTOR EQUATIONS

For a nonzero vector $\mathbf{v} \in \mathbb{R}^n$, $\text{Span}\{\mathbf{v}\}$ is the set of all vectors of the form $c\mathbf{v}$, for some scalar c .



§1.3 VECTOR EQUATIONS

For a nonzero vector $\mathbf{v} \in \mathbb{R}^n$, $\text{Span}\{\mathbf{v}\}$ is the set of all vectors of the form $c\mathbf{v}$, for some scalar c .




This is the line passing through the origin $\mathbf{0}$ and the vector \mathbf{v} .

§1.3 VECTOR EQUATIONS

For nonzero vectors $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$, with \mathbf{v} not a multiple of \mathbf{w} , $\text{Span} \{\mathbf{v}, \mathbf{w}\}$ is the unique plane in \mathbb{R}^3 containing the vectors \mathbf{v} , \mathbf{w} , and $\mathbf{0}$.


§1.3 VECTOR EQUATIONS

For nonzero vectors $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$, with \mathbf{v} not a multiple of \mathbf{w} , $\text{Span} \{\mathbf{v}, \mathbf{w}\}$ is the unique plane in \mathbb{R}^3 containing the vectors \mathbf{v}, \mathbf{w} , and $\mathbf{0}$.



§1.3 VECTOR EQUATIONS

For nonzero vectors $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$, with \mathbf{v} not a multiple of \mathbf{w} , $\text{Span}\{\mathbf{v}, \mathbf{w}\}$ is the unique plane in \mathbb{R}^3 containing the vectors \mathbf{v}, \mathbf{w} , and $\mathbf{0}$.



§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

DEFINITION

If A is an $m \times n$ matrix with columns $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$, and if \mathbf{x} is a vector in \mathbb{R}^n ,

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

DEFINITION

If A is an $m \times n$ matrix with columns $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$, and if \mathbf{x} is a vector in \mathbb{R}^n , then the product of A and \mathbf{x} , denoted $A\mathbf{x}$, is defined as

$$A\mathbf{x} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n.$$

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

DEFINITION

If A is an $m \times n$ matrix with columns $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$, and if \mathbf{x} is a vector in \mathbb{R}^n , then the product of A and \mathbf{x} , denoted $A\mathbf{x}$, is defined as

$$A\mathbf{x} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n.$$

In other words, $A\mathbf{x}$ is the linear combination of the columns of A with weights the corresponding entries of \mathbf{x} .

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

DEFINITION

If A is an $m \times n$ matrix with columns $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$, and if \mathbf{x} is a vector in \mathbb{R}^n , then the product of A and \mathbf{x} , denoted $A\mathbf{x}$, is defined as

$$A\mathbf{x} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n.$$

In other words, $A\mathbf{x}$ is the linear combination of the columns of A with weights the corresponding entries of \mathbf{x} .

Note: $m \times n$ matrices can only be multiplied with vectors in \mathbb{R}^n .

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

Example: Find the product of the matrix

$$A = \begin{bmatrix} 2 & -1 & 4 \\ 1 & 1 & 0 \\ 8 & 1 & -3 \\ 2 & -2 & 1 \end{bmatrix}$$

with the vector

$$\mathbf{v} = \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}.$$

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

Example: Write the system

$$\begin{array}{rcccccl} x_1 & + & 2x_2 & - & x_3 & = & 4 \\ 3x_1 & - & 4x_2 & - & 2x_3 & = & -8 \\ x_1 & + & x_2 & + & 5x_3 & = & 0 \end{array}$$

as a vector equation, and as a matrix equation, then solve.

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

Example: Write the system

$$\begin{array}{rcccccccl} x_1 & + & 2x_2 & - & x_3 & = & 4 \\ 3x_1 & - & 4x_2 & - & 2x_3 & = & -8 \\ x_1 & + & x_2 & + & 5x_3 & = & 0 \end{array}$$

as a vector equation, and as a matrix equation, then solve.

THEOREM

If A is an $m \times n$ matrix, \mathbf{u} and \mathbf{v} are vectors in \mathbb{R}^n and c is a scalar, then:

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

Example: Write the system

$$\begin{array}{rcccccccl} x_1 & + & 2x_2 & - & x_3 & = & 4 \\ 3x_1 & - & 4x_2 & - & 2x_3 & = & -8 \\ x_1 & + & x_2 & + & 5x_3 & = & 0 \end{array}$$

as a vector equation, and as a matrix equation, then solve.

THEOREM

If A is an $m \times n$ matrix, \mathbf{u} and \mathbf{v} are vectors in \mathbb{R}^n and c is a scalar, then:

1. $A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v}$

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

Example: Write the system

$$\begin{array}{rrcrcl} x_1 & + & 2x_2 & - & x_3 & = & 4 \\ 3x_1 & - & 4x_2 & - & 2x_3 & = & -8 \\ x_1 & + & x_2 & + & 5x_3 & = & 0 \end{array}$$

as a vector equation, and as a matrix equation, then solve.

THEOREM

If A is an $m \times n$ matrix, \mathbf{u} and \mathbf{v} are vectors in \mathbb{R}^n and c is a scalar, then:

1. $A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v}$
2. $A(c\mathbf{u}) = c(A\mathbf{u})$

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

THEOREM

If A is an $m \times n$ matrix with columns $\mathbf{a}_1, \dots, \mathbf{a}_n$, and if \mathbf{b} is in \mathbb{R}^m ,

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

THEOREM

If A is an $m \times n$ matrix with columns $\mathbf{a}_1, \dots, \mathbf{a}_n$, and if \mathbf{b} is in \mathbb{R}^m , the matrix equation

$$A\mathbf{x} = \mathbf{b}$$

has the same solution set as the vector equation

$$x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n = \mathbf{b},$$

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

THEOREM

If A is an $m \times n$ matrix with columns $\mathbf{a}_1, \dots, \mathbf{a}_n$, and if \mathbf{b} is in \mathbb{R}^m , the matrix equation

$$A\mathbf{x} = \mathbf{b}$$

has the same solution set as the vector equation

$$x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n = \mathbf{b},$$

which has the same solution set as the system of equations whose augmented matrix is

$$\left[\begin{array}{ccc|c} \mathbf{a}_1 & \cdots & \mathbf{a}_n & \mathbf{b} \end{array} \right].$$

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

Fact: The matrix equation $A\mathbf{x} = \mathbf{b}$ has a solution if and only if \mathbf{b} is a linear combination of the columns of A .

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

Fact: The matrix equation $A\mathbf{x} = \mathbf{b}$ has a solution if and only if \mathbf{b} is a linear combination of the columns of A .

This is the same as saying that \mathbf{b} is in $\text{Span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$.

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

Fact: The matrix equation $A\mathbf{x} = \mathbf{b}$ has a solution if and only if \mathbf{b} is a linear combination of the columns of A .

This is the same as saying that \mathbf{b} is in $\text{Span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$.

Example: Let

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}.$$

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

Fact: The matrix equation $A\mathbf{x} = \mathbf{b}$ has a solution if and only if \mathbf{b} is a linear combination of the columns of A .

This is the same as saying that \mathbf{b} is in $\text{Span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$.

Example: Let

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}.$$

Is the equation $A\mathbf{x} = \mathbf{b}$ consistent for all possible b_1, b_2, b_3 ?

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

THEOREM

Let A be an $m \times n$ matrix. Then the following statements are logically equivalent.

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

THEOREM

Let A be an $m \times n$ matrix. Then the following statements are logically equivalent. That is, for a particular matrix A , they are either all true or they are all false.

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

THEOREM

Let A be an $m \times n$ matrix. Then the following statements are logically equivalent. That is, for a particular matrix A , they are either all true or they are all false.

1. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

THEOREM

Let A be an $m \times n$ matrix. Then the following statements are logically equivalent. That is, for a particular matrix A , they are either all true or they are all false.

1. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
2. Each \mathbf{b} in \mathbb{R}^m is a linear combination of the columns of A .

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

THEOREM

Let A be an $m \times n$ matrix. Then the following statements are logically equivalent. That is, for a particular matrix A , they are either all true or they are all false.

1. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
2. Each \mathbf{b} in \mathbb{R}^m is a linear combination of the columns of A .
3. The columns of A span \mathbb{R}^m .

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

THEOREM

Let A be an $m \times n$ matrix. Then the following statements are logically equivalent. That is, for a particular matrix A , they are either all true or they are all false.

1. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
2. Each \mathbf{b} in \mathbb{R}^m is a linear combination of the columns of A .
3. The columns of A span \mathbb{R}^m .
4. A has a pivot position in every row.

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

THEOREM

Let A be an $m \times n$ matrix. Then the following statements are logically equivalent. That is, for a particular matrix A , they are either all true or they are all false.

1. For each \mathbf{b} in \mathbb{R}^m , the equation $A\mathbf{x} = \mathbf{b}$ has a solution.
2. Each \mathbf{b} in \mathbb{R}^m is a linear combination of the columns of A .
3. The columns of A span \mathbb{R}^m .
4. A has a pivot position in every row.

Warning! This theorem is about a *coefficient matrix*, not an *augmented matrix*. If an augmented matrix $[A \mid \mathbf{b}]$ has a pivot position in every row, then the equation $A\mathbf{x} = \mathbf{b}$ may or may not be consistent.

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

ROW-VECTOR RULE FOR COMPUTING $A\mathbf{x}$

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

ROW-VECTOR RULE FOR COMPUTING $A\mathbf{x}$

Example: Find the product of

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 0 & 5 & -3 \\ 1 & 1 & -3 \end{bmatrix} \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix},$$

§1.4 THE MATRIX EQUATION $A\mathbf{x} = \mathbf{b}$

ROW-VECTOR RULE FOR COMPUTING $A\mathbf{x}$

Example: Find the product of

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 0 & 5 & -3 \\ 1 & 1 & -3 \end{bmatrix} \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix},$$

Note: The j th entry of $A\mathbf{x}$ is just the sum of the products of the entries of the j th row of A with the corresponding entries of \mathbf{x} .

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

DEFINITION (HOMOGENEOUS SYSTEM)

A system of linear equations is said to be **homogeneous** if it can be written in the form $A\mathbf{x} = \mathbf{0}$, where A is an $m \times n$ matrix and $\mathbf{0}$ is the zero vector in \mathbb{R}^m .

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

DEFINITION (HOMOGENEOUS SYSTEM)

A system of linear equations is said to be **homogeneous** if it can be written in the form $A\mathbf{x} = \mathbf{0}$, where A is an $m \times n$ matrix and $\mathbf{0}$ is the zero vector in \mathbb{R}^m .

FACT

A homogeneous system ($A\mathbf{x} = \mathbf{0}$) always has at least one solution :

$$\mathbf{x} = \mathbf{0}.$$

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

DEFINITION (HOMOGENEOUS SYSTEM)

A system of linear equations is said to be **homogeneous** if it can be written in the form $A\mathbf{x} = \mathbf{0}$, where A is an $m \times n$ matrix and $\mathbf{0}$ is the zero vector in \mathbb{R}^m .

FACT

A homogeneous system ($A\mathbf{x} = \mathbf{0}$) always has at least one solution :

$$\mathbf{x} = \mathbf{0}.$$

This is called the **trivial solution**.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

QUESTION

For a matrix A , is there a **nontrivial solution** to $A\mathbf{x} = \mathbf{0}$?

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

QUESTION

For a matrix A , is there a **nontrivial solution** to $A\mathbf{x} = \mathbf{0}$?

FACT

$A\mathbf{x} = \mathbf{0}$ has a nontrivial solution if and only if the equation has at least one free variable.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

QUESTION

For a matrix A , is there a **nontrivial solution** to $A\mathbf{x} = \mathbf{0}$?

FACT

$A\mathbf{x} = \mathbf{0}$ has a nontrivial solution if and only if the equation has at least one free variable.

Example: Determine if the following homogeneous system has a nontrivial solution. Then describe the solution set.

$$\begin{array}{rrcrcl} 3x_1 & + & 5x_2 & - & 4x_3 & = & 0 \\ -3x_1 & - & 2x_2 & + & 4x_3 & = & 0 \\ 6x_1 & + & x_2 & - & 8x_3 & = & 0 \end{array}$$

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

Example: Find a description of the solution set of

$$2x_1 + x_2 - x_3 = 0.$$

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

Example: Find a description of the solution set of

$$2x_1 + x_2 - x_3 = 0.$$

Note: Any solution set (of a homogeneous equation) can be expressed as $\text{Span } \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ for some set of vectors $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

Example: Find a description of the solution set of

$$2x_1 + x_2 - x_3 = 0.$$

Note: Any solution set (of a homogeneous equation) can be expressed as $\text{Span } \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ for some set of vectors $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.

DEFINITION

Expressing a solution set as $t\mathbf{v}_1 + s\mathbf{v}_2$, where t and s can be any real number (as above), is called **parametric vector form**.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

THEOREM

Suppose that $A\mathbf{x} = \mathbf{b}$ is consistent for some fixed \mathbf{b} , and let \mathbf{p} be the solution, i.e. $A\mathbf{p} = \mathbf{b}$.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

THEOREM

Suppose that $A\mathbf{x} = \mathbf{b}$ is consistent for some fixed \mathbf{b} , and let \mathbf{p} be the solution, i.e. $A\mathbf{p} = \mathbf{b}$.

Then the solution set of $A\mathbf{x} = \mathbf{b}$ is the set of all vectors of the form $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$ where \mathbf{v}_h is any solution of the homogeneous equation $A\mathbf{x} = \mathbf{0}$.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

THEOREM

Suppose that $A\mathbf{x} = \mathbf{b}$ is consistent for some fixed \mathbf{b} , and let \mathbf{p} be the solution, i.e. $A\mathbf{p} = \mathbf{b}$.

Then the solution set of $A\mathbf{x} = \mathbf{b}$ is the set of all vectors of the form $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$ where \mathbf{v}_h is any solution of the homogeneous equation $A\mathbf{x} = \mathbf{0}$.

Warning! This theorem only applies to systems $A\mathbf{x} = \mathbf{b}$ that are consistent.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

THEOREM

Suppose that $A\mathbf{x} = \mathbf{b}$ is consistent for some fixed \mathbf{b} , and let \mathbf{p} be the solution, i.e. $A\mathbf{p} = \mathbf{b}$.

Then the solution set of $A\mathbf{x} = \mathbf{b}$ is the set of all vectors of the form $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$ where \mathbf{v}_h is any solution of the homogeneous equation $A\mathbf{x} = \mathbf{0}$.

Warning! This theorem only applies to systems $A\mathbf{x} = \mathbf{b}$ that are consistent. If the system is inconsistent then the solution set is empty.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS


All solutions to $A\mathbf{x} = \mathbf{0}$




§1.5 SOLUTION SETS OF LINEAR SYSTEMS

All solutions to $A\mathbf{x} = \mathbf{0}$


Trivial solution $\mathbf{0}$ •




§1.5 SOLUTION SETS OF LINEAR SYSTEMS




§1.5 SOLUTION SETS OF LINEAR SYSTEMS




§1.5 SOLUTION SETS OF LINEAR SYSTEMS




§1.5 SOLUTION SETS OF LINEAR SYSTEMS




§1.5 SOLUTION SETS OF LINEAR SYSTEMS



§1.5 SOLUTION SETS OF LINEAR SYSTEMS



§1.5 SOLUTION SETS OF LINEAR SYSTEMS



§1.5 SOLUTION SETS OF LINEAR SYSTEMS

Example: Find all solutions to the system

$$\begin{array}{rrcrcl} 3x_1 & + & 5x_2 & - & 4x_3 & = & 7 \\ -3x_1 & - & 2x_2 & + & 4x_3 & = & -1 \\ 6x_1 & + & x_2 & - & 8x_3 & = & -4. \end{array}$$

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

WRITING SOLUTION SETS IN PARAMETRIC VECTOR FORM

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

WRITING SOLUTION SETS IN PARAMETRIC VECTOR FORM

1. Row reduce the augmented matrix to reduced echelon form.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

WRITING SOLUTION SETS IN PARAMETRIC VECTOR FORM

1. Row reduce the augmented matrix to reduced echelon form.
2. Express each basic variable in terms of any free variables appearing in an equation.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

WRITING SOLUTION SETS IN PARAMETRIC VECTOR FORM

1. Row reduce the augmented matrix to reduced echelon form.
2. Express each basic variable in terms of any free variables appearing in an equation.
3. Write a typical solution \mathbf{x} as a vector whose entries depend on the free variables, if any.

§1.5 SOLUTION SETS OF LINEAR SYSTEMS

WRITING SOLUTION SETS IN PARAMETRIC VECTOR FORM

1. Row reduce the augmented matrix to reduced echelon form.
2. Express each basic variable in terms of any free variables appearing in an equation.
3. Write a typical solution \mathbf{x} as a vector whose entries depend on the free variables, if any.
4. Decompose \mathbf{x} into a linear combination of vectors (with numeric entries) using the free variables as parameters.