

CHEMICAL KINETICS

for Class XII
BOARD EXAMS

These notes
have been verified
by CBSE Science
Toppers &
reputated faculties


We have integrated
Previous Year Q's
in the notes

APNI
KAKSHA



Raman Dhattarwal

Flow Chart





Introduction

Rate of Reaction

Average Rate

Instantaneous Rate

Factors affecting Rate




Chemical Kinetics

Order and molecularity of a reaction



Integrated Rate Equation

Zero order Reaction

$A \rightarrow \text{Products}$




Half life of a Reaction



first order Reaction

Pseudo first order Reaction



Chemical Kinetics

Apni Kaksha :-


- For a chemical reaction $R \rightarrow P$: Three questions are important.
- i) Reaction will happen or not? : Thermodynamics : Reaction is feasible only if $\Delta H < 0$.
- ii) To which extent reaction will happen? : Chemical Equilibrium
- iii) What is the speed of chemical reaction? : Chemical kinetics : Concept of rate of a reaction.

Rate of reaction :- [velocity of reaction] [CBSE 2010/2015 / Delhi 2010] (1M)

The rate of change of concentration of any species taking part in a chemical reaction with time is known as rate of reaction.

$$\rightarrow \text{For a reaction } R \rightarrow P \quad \text{rate} = \frac{\text{Change in Concentration}}{\text{time}} = \frac{\Delta C}{\Delta t}$$

→ Unit : $\frac{(\text{mol/l})}{\text{s}}$ or $\frac{(\text{atm})}{\text{s}}$ → For gaseous reactant or product.



Types of rate :- [CBSE 2010C] (2M)

i) Average Rate :- $r_{av} = \frac{\text{Total change in conc}^n}{\text{Total time taken}}$

$$\text{Rate} = -\frac{\Delta [\text{Reactant}]}{\Delta t} = \frac{\Delta [\text{Product}]}{\Delta t} = \frac{c_2 - c_1}{t_2 - t_1} = \frac{\Delta C}{\Delta t}$$


→ $R \rightarrow P$
at time t_1 : c_1 [concentration of product after time t_1]

at time t_2 : c_2

ii) Instantaneous Rate :- Rate at a given time

$$r_{inst} = \lim_{\Delta t \rightarrow 0} r_{av} = \lim_{\Delta t \rightarrow 0} \frac{\Delta C}{\Delta t} = \frac{dc}{dt}$$

= Slope of concentration and time graph.



Rate of appearance and rate of disappearance :-

Let us consider a reaction $R \rightarrow P$

At time t_1 : $R_1 \quad P_1$

t_2 : $R_2 \quad P_2$


$$\Delta R = R_2 - R_1$$

$$\Delta P = P_2 - P_1$$

$$\Delta t = t_2 - t_1$$

$$\text{Rate of disappearance of R} = \frac{\text{Decrease in [R]}}{\text{Time taken}} = -\frac{\Delta[R]}{\Delta t}$$

$$\text{Rate of appearance of P} = \frac{\text{Increase in [P]}}{\text{Time taken}} = \frac{\Delta[P]}{\Delta t}$$




→ Rate of reaction = Rate of disappearance of $\text{Hg} | \text{U}_2$ = Rate of appearance of HgCl_2

$$= -\frac{\Delta[\text{Hg}]}{\Delta t} = -\frac{\Delta[\text{U}_2]}{\Delta t} = \frac{\Delta[\text{HgU}_2]}{\Delta t}$$

Note :- If stoichiometric coefficients of reactants or products are not equal to one, then rate of disappearance of any of the reactants or rate of appearance of products is divided by their respective stoichiometric coefficients.




$$\text{Rate of reaction} = -\frac{1}{a} \frac{\Delta[A]}{\Delta t} = -\frac{1}{b} \frac{\Delta[B]}{\Delta t} = \frac{1}{c} \frac{\Delta[C]}{\Delta t} = \frac{1}{d} \frac{\Delta[D]}{\Delta t}$$



→ Rate of disappearance of $\text{HI} = 2 \times \text{Rate of appearance of H}_2 | \text{I}_2$

→ Rate of reaction = Rate of appearance of $\text{H}_2 | \text{I}_2 = \frac{1}{2} \text{Rate of disappearance of HI}$.


$$= \frac{\Delta[\text{H}_2]}{\Delta t} = \frac{\Delta[\text{I}_2]}{\Delta t} = -\frac{1}{2} \frac{\Delta[\text{HI}]}{\Delta t}$$



$$\text{Rate of reaction} = -\frac{1}{5} \frac{\Delta[\text{Br}^-]}{\Delta t} = -\frac{\Delta[\text{BrO}_3^-]}{\Delta t} = -\frac{1}{6} \frac{\Delta[\text{H}^+]}{\Delta t}$$

Question :- For the reaction $2 \text{N}_2\text{O}_5(\text{g}) \rightarrow 4 \text{NO}_2(\text{g}) + \text{O}_2(\text{g})$, the rate of formation of $\text{NO}_2(\text{g})$ is $2.8 \times 10^{-3} \text{ M s}^{-1}$. Calculate the rate of disappearance of $\text{N}_2\text{O}_5(\text{g})$!

[CBSE 2010] 2M



$$\text{Rate of Reaction} = -\frac{1}{2} \frac{\Delta[\text{N}_2\text{O}_5]}{\Delta t} = \frac{1}{4} \frac{\Delta[\text{NO}_2]}{\Delta t}$$


$$\rightarrow \text{Rate of disappearance of } \text{N}_2\text{O}_5 = -\frac{\Delta [\text{N}_2\text{O}_5]}{\Delta t} = 2 \times \frac{1}{4} \frac{\Delta [\text{NO}_2]}{\Delta t}$$

Given that rate of formation of $\text{NO}_2 = 2.8 \times 10^{-3} \text{ M s}^{-1}$

$$= \frac{\Delta [\text{NO}_2]}{\Delta t} = \frac{1}{2} \times 2.8 \times 10^{-3} \text{ M s}^{-1}$$

$$= 1.4 \times 10^{-3} \text{ M s}^{-1}$$

Question :- The decomposition of N_2O_5 in CCl_4 at 310 K has been studied by monitoring the concentration of N_2O_5 in the solution. Initially the concentration of N_2O_5 is 2.33 mol L^{-1} and after 184 minutes, it is reduced to 2.08 mol L^{-1} . Calculate the average rate of this reaction and what is the rate of production of NO_2 during this period?



Answer :- Average Rate $= \frac{-1}{2} \frac{\Delta [\text{N}_2\text{O}_5]}{\Delta t} = \frac{-1}{2} \left[\frac{(2.08 - 2.33) \text{ mol L}^{-1}}{184 \text{ min}} \right]$

$$= 6.79 \times 10^{-4} \text{ mol L}^{-1} \text{ min}^{-1} = \frac{6.79 \times 10^{-4} \text{ mol L}^{-1}}{60 \text{ s}}$$

$$\rightarrow \text{Rate} = \frac{1}{4} \left[\frac{\Delta [\text{NO}_2]}{\Delta t} \right] = 6.79 \times 10^{-4} \frac{\text{mol L}^{-1}}{\text{min}} = 1.13 \times 10^{-5} \text{ mol L}^{-1} \text{ s}^{-1}$$

$$\rightarrow \frac{\Delta [\text{NO}_2]}{\Delta t} = 4 \times 6.79 \times 10^{-4} = 2.79 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1} = \text{Rate of production of } \text{NO}_2$$

Factors influencing rate of a reaction :- [Delhi 2010C] [CBSE 2015]

2M

1M

i) Concentration :- Rate of reaction increases with concentration of reactant.
↳ Liquid / Aqueous Medium

→ In case of gases, pressure $\uparrow \Rightarrow$ Rate \uparrow

ii) Temperature :- Rate of a reaction increases with increase of temperature.

iii) Nature of reactants :- Different reactants require different amount of energies for breaking of old bonds and for formation of new bonds.

iv) Catalyst :- It changes the rate of reaction, without being used (consumed) in the reaction.

Rate Law | Rate Equation | Rate Expression :- [Dependence of rate on concⁿ]

→ Rate law is expression in which reaction rate is given in terms of molar concⁿ of reactants with each term raised to some power. [CBSE 2011 | Delhi 2011/2012]

1M

→ The power raised on concentration terms in rate expression may or may not be same as stoichiometric coefficient of the reacting species in a balanced chemical equation.

→ Consider a general reaction : $aA + bB \rightarrow cC + dD$

Rate equation : Rate $\propto [A]^x [B]^y$

a, b, c, d are stoichiometric coefficients.

$$\boxed{\text{Rate} = K [A]^x [B]^y}$$

↓
Rate Constant

→ Exponent x and y may or may not be equal to stoichiometric coefficient (a and b).

→ The equation which relates, rate of a reaction to concentration of the reactants is called **rate equation** or **rate expression**.

Order of a reaction :- The sum of powers of the concentration of the reactants in rate law expression is called order of that chemical reaction.

For a reaction : $aA + bB \rightarrow cC + dD$

$$\text{Rate} = K [A]^x [B]^y \quad \text{then order of reaction} = x+y$$

1M [CBSE 2011/2010 / Delhi 2010]

→ Order of a reaction can be determined only by experiment. We can not predict theoretically.

Apni Kaksha

→ Order of a reaction can be 0, 1, 2 and even a fraction.

<u>Reaction</u>	<u>Experimental rate expression</u>	<u>order of reaction</u>
$\text{CHCl}_3 + \text{Cl}_2 \rightarrow \text{CCl}_4 + \text{HCl}$	$\text{Rate} = K [\text{CHCl}_3]^1 [\text{Cl}_2]^{1/2}$	$1 + \frac{1}{2} = 1.5$
$\text{CH}_3\text{-}\overset{\text{O}}{\underset{\downarrow}{\text{C}_2\text{H}_5}} + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{COOH} + \text{C}_2\text{H}_5\text{OH}$	$\text{Rate} = K [\text{CH}_3\text{COOC}_2\text{H}_5]^1 [\text{H}_2\text{O}]^0$	$1+0=1$

Units of rate constant - : From above : $\text{Rate} = K [A]^x [B]^y$

$$\text{Then } K = \frac{\text{Rate}}{[A]^x [B]^y} ; \text{ unit of } K = \frac{\text{Conc}^1 \text{ time}}{(\text{Conc})^x (\text{Conc})^y} = \frac{\text{Conc}^n}{\text{time}} \cdot \frac{1}{[\text{Conc}]^{x+y}}$$

$$\rightarrow \text{Unit of } k = \frac{\text{mol L}^{-1}}{\text{s}} \times \frac{1}{(\text{mol L}^{-1})^n}$$

$n = \text{order of reaction} = x+y$

\rightarrow For zero order reaction : $n=0$ then unit of $k = \text{mol L}^{-1} \text{s}^{-1}$

\rightarrow For first order reaction : $n=1$: Unit of $k = \text{s}^{-1}$

\rightarrow For second order reaction : $n=2$: unit of $k = \text{mol}^{-1} \text{L s}^{-1}$

Question :- Identify order of following reaction :-

- (i) $k = 2.3 \times 10^{-5} \text{ L mol}^{-1} \text{s}^{-1}$
- (ii) $3 \times 10^{-4} \text{ s}^{-1}$

Answer :- (i) $\text{L mol}^{-1} \text{s}^{-1}$: Unit of second order

so, this represent second order reaction.

[Delhi 2011 C]

2M

(ii) s^{-1} : Unit of first order reaction.

Question :- Write the rate equation for reaction $2\text{NO}(g) + \text{O}_2(g) \rightarrow \text{NO}_2(g)$ by using following experimental data table?

Solution :- Let us write rate expression

for reaction : Rate = $k [\text{NO}]^x [\text{O}_2]^y$ - (i)

Put the data of experiment (i) in equation (i) :-

$$\rightarrow 0.096 \text{ mol L}^{-1} \text{s}^{-1} = k [0.30]^x [0.30]^y \quad \text{(ii)}$$

Experiment	Initial [NO]/ mol L ⁻¹	Initial [O ₂]/ mol L ⁻¹	Initial rate of formation of NO ₂ / mol L ⁻¹ s ⁻¹
1.	0.30	0.30	0.096
2.	0.60	0.30	0.384
3.	0.30	0.60	0.192
4.	0.60	0.60	0.768

similarly, experiment (ii) and equation (i) :- $0.384 \text{ mol L}^{-1} \text{s}^{-1} = k [0.60]^x [0.30]^y$ - (iii)

\rightarrow Experiment (iii) and equation (i) :- $0.192 \text{ mol L}^{-1} \text{s}^{-1} = k [0.30]^x [0.60]^y$ - (iv)

$$\text{Divide equation (ii) by (iii) :- } \frac{0.096}{0.384} = \left[\frac{0.30}{0.60} \right]^x \left[\frac{0.30}{0.60} \right]^y$$

$$\Rightarrow \frac{1}{4} = \frac{1}{2^x} \quad \text{then } x=2$$

Apni
Kaksha

$$\text{Divide (iv) by (i) :- } \frac{0.192}{0.096} = \left[\frac{0.30}{0.30} \right]^x \left[\frac{0.60}{0.30} \right]^y \Rightarrow 2 = 2^y \text{ then } y=1$$

put the value of x and y in equation (i) :-

$$\boxed{\text{Rate} = k [\text{NO}]^2 [\text{O}_2]^1} \quad \text{-(v)}$$

Order of reaction = $2+1=3$

value of rate constant :- Put the value of rate, [NO], [O₂] in equation (v) from any experiment.

Question :- For the reaction $2\text{NO}(g) + \text{Cl}_2(g) \rightarrow 2\text{NOCl}(g)$, the following data were collected. All the measurements were taken at 263K.

- i Write the expression for rate law.
- ii Calculate the value of rate constant and specify its unit.
- iii What is the initial rate of disappearance of Cl_2 in experiment 4? [Delhi 2012] 3M

Exp. No.	Initial [NO] (M)	Initial $[\text{Cl}_2]$ (M)	Initial rate of disappearance of Cl_2 (M/min)
1.	0.15	0.15	0.60
2.	0.15	0.30	1.20
3.	0.30	0.15	2.40
4.	0.25	0.25	?

Answer :- Rate law : $\text{Rate} = K [\text{NO}]^x [\text{Cl}_2]^y$

→ Initial rate becomes $(\text{Rate})_0 = K [\text{NO}]^x [\text{Cl}_2]^y$

→ Comparing experiments 1 and 2 : $(\text{Rate})_1 = K [0.15]^x [0.15]^y = 0.60$ — i

$$(\text{Rate})_2 = K [0.15]^x [0.30]^y = 1.20 \quad \text{ii}$$

→ Dividing equation (ii) by equation (i) →

$$\frac{(\text{Rate})_2}{(\text{Rate})_1} = \frac{1.20}{0.60} = \frac{K [0.15]^x [0.30]^y}{K [0.15]^x [0.15]^y} \quad \text{Then, } 2^y = 2^1$$

$$y=1$$

→ Comparing experiments 1 and 3 :

$$(\text{Rate})_1 = K [0.15]^x [0.15]^y = 0.60 \quad \text{iii}$$

$$(\text{Rate})_3 = K [0.30]^x [0.15]^y = 2.40 \quad \text{iv}$$

Dividing equation (iv) by equation (iii) :-

$$\frac{(\text{Rate})_3}{(\text{Rate})_1} = \frac{K [0.30]^x [0.15]^y}{K [0.15]^x [0.15]^y} = \frac{2.40}{0.60}$$

$$\Rightarrow 2^x = 4 \quad \text{then } x=2$$

Rate law : $\text{Rate} = K [\text{NO}]^2 [\text{Cl}_2]$

$$\# \text{ Rate constant } K = \frac{\text{Rate}}{[\text{NO}]^2 [\text{Cl}_2]} = \frac{0.60}{(0.15)^2 (0.15)} = 177.51 \text{ mol}^{-2} \text{ L}^2 \text{ min}^{-1}$$

Initial rate of disappearance of Cl_2 in experiment 4 is r_4 .




$$r_4 = K [\text{NO}]^2 [\text{Cl}_2] = 177.51 \times (0.25)^2 (0.25)$$

$$r_4 = 2.77 \frac{\text{M}}{\text{min}}$$

Molecularity of a reaction -:

→ The no. of reacting species [atoms, ions or molecules] taking part in a one step reaction [an elementary reaction], which must collide simultaneously in order to bring about a chemical reaction is called molecularity of a reaction.

[CBSE 2020/2011C/2014] 1M



**Apni
Kaksha**

Note:- Generally molecularity greater than 3 is not observed , because simultaneous collisions of more than 3 molecules are not possible .

Difference between order and molecularity

[CBSE 2014/2010/Delhi 2011C]

Molecularity


Order

- i) It is defined as the no. of molecules of reactant taking part in a reaction.
- i) It is defined as the sum of the power of concentration terms that appear in rate law.
- ii) Example -: $\text{NH}_4\text{NO}_2 \xrightarrow{\Delta} \text{N}_2 + 2\text{H}_2\text{O}$
- ii) $\text{NH}_4\text{NO}_2 \xrightarrow{\Delta} \text{N}_2 + 2\text{H}_2\text{O}$
- molecularity = 1
- Rate = $k[\text{NH}_4\text{NO}_2]$, order = 1
- iii) It is a theoretical value.
- iii) It is an experimental value.
- iv) It is always a whole no. It can neither be zero nor fractional.
- iv) It may be zero, fractional or integer.
- v) It is derived from rate determining step in the mechanism of a reaction.
- v) It is derived from rate expression


Note :- Let us consider a reaction $2\text{H}_2\text{O}_2 \xrightarrow[\text{Alkaline Medium}]{\text{I}^-} 2\text{H}_2\text{O} + \text{O}_2$.

Evidences suggest that this reaction takes place in two step.

[Delhi 2019]



2M



→ Slow step [first step] is the rate determining step [RDS].

→ Rate equation for reaction can be written on the basis of RDS.

$$\text{Rate} = -\frac{d[H_2O_2]}{dt} = k [H_2O_2]^1[I^-]^1$$

→ Overall order of reaction = $1+1=2$

Integrated Rate Equations

Zero Order Reaction :- Let us consider a zero order reaction $R \rightarrow P$

$$\text{Then rate} = -\frac{d[R]}{dt} = k[R]^0 = k \quad \text{Differential rate Equation.}$$

$$\Rightarrow d[R] = -k dt \implies \text{on integrating both side : } \int d[R] = -k \int dt + I \quad \text{constant of integration.}$$
$$\Rightarrow [R] = -kt + I \quad \text{(i)}$$

At $t=0$, the concentration of reactant $R = [R]_0$.


By equation (i) $\Rightarrow [R]_0 = -k \times 0 + I \text{ then } I = [R]_0$.

→ put the value of I in equation (i) :-

$$[R] = -kt + [R]_0$$

$y = mx + c$

Integrated rate Equation.




→ Straight line graph between $[R]$ and t

→ Slope = $-k$

Example of zero order reaction :- (i) Enzyme catalysed reactions.

(ii) Decomposition of NH_3 on a hot platinum surface at high pressure.


[Delhi 2016]



1M

$$\rightarrow \text{Rate} = k[NH_3]^0$$

Explanation :- At high pressure the metal surface gets saturated with gas molecules. So, a further change in reaction condition is unable to change the amount of ammonia on the surface of the catalyst, making rate of reaction independent of its concentration.



Integrated rate equation of first order reactions :-

Consider a first order reaction : $R \rightarrow P$

$$\text{Rate} = -\frac{d[R]}{dt} = k[R] \Rightarrow \frac{d[R]}{[R]} = -k dt$$

$$\text{on integration} \Rightarrow \int \frac{d[R]}{[R]} = -k \int dt + I \quad \text{--- (i)}$$

→ When $t=0$ then $[R]= [R]_0$.

$$\rightarrow \text{Put this value in equation (i) } \therefore \ln[R]_0 = -k \cdot 0 + I \quad \ln[R] = -k t + I$$


$$I = \ln[R]_0 \quad \text{--- (ii)} \quad \begin{matrix} \text{constant} \\ \text{of integration.} \end{matrix}$$

→ By equation (i) and (ii) :-

$$\ln[R] = -k \cdot t + \ln[R]_0 \quad \text{--- (iii)}$$

$$\downarrow \quad \downarrow \quad \downarrow$$

$$y = m \cdot x + c$$




$$\rightarrow 2.303 \log \frac{[R]_0}{[R]} = k t$$

$$\log \frac{[R]_0}{[R]} = \frac{k}{2.303} t$$

$$\downarrow \quad \downarrow \quad \downarrow$$

$$y = m \cdot x$$



$$\text{Slope} = \frac{k}{2.303}$$


$$\text{On rearranging equation (iii) : } \ln[R] - \ln[R]_0 = -k t$$

$$\rightarrow \ln \frac{[R]}{[R]_0} = -k t$$

$$k = \frac{1}{t} \ln \frac{[R]_0}{[R]}$$

$$\rightarrow \frac{[R]}{[R]_0} = e^{-kt} \quad \text{then } [R] = [R]_0 e^{-kt}$$

Graph between $[R]$ and time $t \rightarrow$



At t_1 , $[R]_1$ Then By equation (iii) →

$$\ln[R]_1 = -k t_1 + \ln[R]_0$$

At t_2 , $[R]_2$

$$\ln[R]_2 = -k t_2 + \ln[R]_0$$

$$K = \frac{1}{(t_2 - t_1)} \ln \frac{[R]_1}{[R]_2}$$

Example -: i) All natural and artificial radioactive decay of unstable nuclei take place by 1st order kinetics. $^{226}_{88}\text{Ra} \longrightarrow {}^+_2\text{He} + {}^{222}_{86}\text{Rn}$

ii) Hydrogenation of ethene



$$\hookrightarrow \text{Rate} = k[R_a]$$

Question -: The initial concentration of N_2O_5 in the following 1st order reaction $\text{N}_2\text{O}_5(g) \longrightarrow 2\text{NO}_2(g) + \frac{1}{2}\text{O}_2(g)$ was $1.24 \times 10^{-2} \text{ mol L}^{-1}$ at 318 K. The concentration of N_2O_5 after 60 minutes was $0.20 \times 10^{-2} \text{ mol L}^{-1}$. Calculate the rate constant of the reaction at 318 K?

Answer -: For 1st order reaction : $k = \frac{2.303}{(t_2 - t_1)} \log \frac{[R]_1}{[R]_2}$

$$\rightarrow k = \frac{2.303}{(60 \text{ min} - 0 \text{ min})} \log \frac{1.24 \times 10^{-2} \text{ mol L}^{-1}}{0.20 \times 10^{-2} \text{ mol L}^{-1}} = \frac{2.303}{60} \log 6.2 \text{ min}^{-1}$$

First order gas phase reaction :-

Let us consider a 1st order reaction : $A(g) \longrightarrow B(g) + C(g)$

p_i : Initial pressure of A.

At $t=0$: p_i atm 0 0

p_t : Total pressure at time t.

At time t : $(p_i - x)$ x x

p_A, p_B, p_C : Partial pressure of A, B, C respectively.

Total pressure at time t

x : Decrease in pressure after time t.

$$p_t = p_A + p_B + p_C$$

$$p_t = (p_i - x) + x + x$$

$$\text{then } x = p_t - p_i$$

Partial pressure] at time t
of A

$$\rightarrow \text{Rate constant } k = \frac{2.303}{t} \log \left(\frac{p_i}{p_t} \right)$$


$$K = \frac{2.303}{t} \log \frac{p_i}{2p_i - p_t}$$

$$p_A = p_i - x = p_i - (p_t - p_i)$$

$$p_A = 2p_i - p_t$$

Question :- Experimental data for the first order thermal decomposition of SO_2Cl_2 at constant volume.

[CBSE 2014 | Delhi 2014] (3M)



Exp. Time Total pressure

1. 0 s $P_i = 0.4$

2. 100 s $P_f = 0.7$

Answer :- $k = \frac{2.303}{t} \log \frac{P_i}{2P_i - P_f}$

$$= \frac{2.303}{100 \text{ s}} \log \frac{0.4}{2 \times 0.4 - 0.7} = \frac{2.303}{100} (2 \log 2) = 1.307 \times 10^{-2} \text{ s}^{-1}$$

Half life of a reaction :- "The time in which the concentration of a reactant is reduced to one half of its initial concentration." [Delhi 2015 C] (1M)

→ For reaction $R \rightarrow P$

$$\text{At } t=0 \quad [R]_0$$

$$\text{At } t=t_{1/2} \quad \frac{[R]_0}{2}$$

→ Half life for zero order reaction

$$\hookrightarrow \text{Rate constant } k = \frac{[R]_0 - [R]}{t}$$

$$\text{At } t = t_{1/2} \text{ (Half life)} \quad [R] = \frac{[R]_0}{2}$$

$$\hookrightarrow k = \frac{[R]_0 - \frac{[R]_0}{2}}{t_{1/2}}$$

$$k = \frac{[R]_0}{2 t_{1/2}}$$

$$t_{1/2} = \frac{[R]_0}{2k}$$

→ Half life $t_{1/2}$ for 1st order reaction

$$\hookrightarrow \text{Rate constant } k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$$

$$\hookrightarrow k = \frac{2.303}{t_{1/2}} \log \frac{[R]_0}{[R]_0/2}$$

$$k = \frac{2.303}{t_{1/2}} \log 2$$

$$k = \frac{0.693}{t_{1/2}}$$

$$t_{1/2} = \frac{0.693}{k}$$

→ Half life of zero order reaction depends on initial concentration of reactant.


a.) $t_{1/2} \propto [R]_0$

b.) $t_{1/2} \propto \frac{1}{k}$


→ Half life ($t_{1/2}$) is independent of initial concentration. [Delhi 2017] (1M)

$A \rightarrow \text{Products}$

Reactant Concentration versus Time




Rate versus Reactant Concentration




Order	Reaction type	Differential rate law	Integrated rate law	Straight line plot	Half-life	Units of k
0	$R \rightarrow P$	$d[R]/dt = -k$	$kt = [R]_0 - [R]$	$[R]$ vs t	$[R]_0/2k$	conc time $^{-1}$ or mol L $^{-1}$ s $^{-1}$
1	$R \rightarrow P$	$d[R]/dt = -k[R]$	$[R] = [R]_0 e^{-kt}$ or $kt = \ln\{[R]_0/[R]\}$	$\ln[R]$ vs t	$\ln 2/k$	time $^{-1}$ or s $^{-1}$

Apni Kaksha :-)

[A] vs time for a 0 order reaction



[A] vs time for a 1st order reaction



Question :- A first order reaction is found to have a rate constant $K = 5.5 \times 10^{-14} \text{ s}^{-1}$. Find the half life of the reaction.

Answer :- For first order, Half life $t_{1/2} = \frac{0.693}{K} = \frac{0.693}{5.5 \times 10^{-14}}$

$$t_{1/2} = 1.26 \times 10^{14} \text{ s} \quad [\text{CBSE 2013}]$$

1M

Question :- A first order reaction takes place in 30 min for 50% completion. Calculate the time required for 90% completion of this reaction. Given $\log 2 = 0.3010$

[CBSE 2015 | Delhi 2015]

3M

Answer :- 50% completion time = $t_{1/2} = 30 \text{ min}$

$$K = \frac{0.693}{t_{1/2}} = \frac{0.693}{30 \text{ min}} = 0.023 \text{ min}^{-1}$$

→ For the same reaction, time required for 90% completion :-

$$K = 0.023 \text{ min}^{-1}; [R]_0 = 100 \quad [R] = 100 - 90 = 10$$

$$\rightarrow t = \frac{2.303}{K} \log \frac{[R]_0}{[R]} = \frac{2.303}{0.023 \text{ min}^{-1}} \log \frac{100}{10} = 100.13 \text{ min.}$$

Pseudo first order reaction :- A chemical reaction between two substances when one reactant is present in excess quantity. [CBSE 2011] 2M

i) Hydrolysis of ethyl acetate :- $\text{CH}_3\text{C}(=\text{O})\text{OC}_2\text{H}_5 + \text{H}_2\text{O} \xrightarrow{\text{H}^+} \text{CH}_3\text{C}(=\text{O})\text{OH} + \text{C}_2\text{H}_5\text{OH}$

If At time $t=0$: 0.01 mol 10 mol 0 mol 0 mol


and, at time t : 0 mol 9.99 mol 0.01 0.01

→ $[H_2O]_{t=0} \approx [H_2O]_t$ means that concentration of H_2O does not change. So it can not change the rate. $[H_2O] = \text{constant}$

→ Rate of reaction = $k' [H_2O] [CH_3COOC_2H_5]$ $k' [H_2O] = \text{constant}$
 $= K$

Rate of reaction = $K [CH_3COOC_2H_5] \rightarrow$ It indicates 1st order kinetics.

ii) Hydrolysis of cane sugar :-



$$\boxed{\text{Rate} = K [C_{12}H_{22}O_{11}]}$$

Question :- For the hydrolysis of methyl acetate in aqueous solution, the following results were obtained.

- i) If concentration of water remains constant then show that it follows pseudo first order reaction.

- ii) Calculate average rate of reaction between time interval 10 to 20 second.


Time (second)	0	10	20
$[\text{CH}_3-\overset{\overset{\bullet}{\text{O}}}{\underset{\underset{\text{H}_2\text{O}}{\text{C}}}{}-\text{OCH}_3]$ [mol L ⁻¹]	0.10	0.05	0.025

[CBSE 2015]

5M

Answer :- i) $[\text{H}_2\text{O}] = \text{constant}$ (During reaction)

$$\text{So, rate} = k [\text{Methyl acetate}]$$



$$\text{First order, then } k = \frac{2.303}{t} \log \frac{[\text{CH}_3-\overset{\overset{\bullet}{\text{O}}}{\underset{\underset{\text{H}_2\text{O}}{\text{C}}}{}-\text{OCH}_3]_0}{[\text{CH}_3-\overset{\overset{\bullet}{\text{O}}}{\underset{\underset{\text{H}_2\text{O}}{\text{C}}}{}-\text{OCH}_3]}$$

$$\text{At time } t_1 = 10 \text{ second} \Rightarrow k_1 = \frac{2.303}{10} \log \frac{0.10}{0.05} = 6.93 \times 10^{-2} \text{ s}^{-1}$$

$$\text{At time } t_2 = 20 \text{ second} \Rightarrow k_2 = \frac{2.303}{20} \log \frac{0.10}{0.025} = 6.93 \times 10^{-2} \text{ s}^{-1}$$

Rate constant for the reaction has a constant value under any given time interval. Hence the given reaction follows pseudo first order kinetics.

$$\text{ii) Average rate} = - \frac{\Delta [\text{CH}_3-\overset{\overset{\bullet}{\text{O}}}{\underset{\underset{\text{H}_2\text{O}}{\text{C}}}{}-\text{OCH}_3]}{\Delta t} = - \frac{[0.025 - 0.05]}{20 - 10} = 0.0025 \text{ mol L}^{-1} \text{ s}^{-1}$$

Important Previous Year Questions

Apni Kaksha :-

- 1) For a first order reaction $\text{H}_2\text{O}_2 \rightarrow \text{H}_2\text{O(l)} + \text{O}_2(g)$, rate constant $k = 1.06 \times 10^{-3} \text{ min}^{-1}$.

2M [Delhi 2014 C]

- Q) How long will it take for 15% of a sample of H_2O_2 to decompose?

Answer :- Let us consider $[\text{H}_2\text{O}_2]_0 = 100 \text{ mol/L}$, then $[\text{H}_2\text{O}_2]_t = \frac{100 - 15}{85} \text{ mol/L}$

$$k = \frac{2.303}{t} \log \frac{[\text{H}_2\text{O}_2]_0}{[\text{H}_2\text{O}_2]} \Rightarrow t = \frac{2.303}{1.06 \times 10^{-3}} \log \frac{100}{85} = 153.3 \text{ min.}$$

(b) → How long time it will take for 95% of the sample to decompose?

$$\rightarrow \text{Now: } [\text{H}_2\text{O}_2]_0 = 100 \text{ mol/L} \quad [\text{H}_2\text{O}_2]_t = 100 - 95 = 15 \text{ mol/L}$$

$$t = \frac{2.303}{k} \log \frac{[\text{H}_2\text{O}_2]_0}{[\text{H}_2\text{O}_2]_t} = \frac{2.303}{1.06 \times 10^{-3}} \log 6.667 = \underline{1790.05 \text{ min}}$$

[2.] The rate constant for a zero order reaction is $0.0030 \text{ mol L}^{-1} \text{ s}^{-1}$. How long will it take for the initial concentration to fall from 0.10 M to 0.075 M ?

[Delhi 2010]

Answer -: For zero order reaction $\rightarrow k = \frac{[R]_0 - [R]}{t}$

$$[R]_0 = 0.10 \text{ M}$$

$$[R] = 0.075 \text{ M}$$

$$k = 0.0030 \text{ mol L}^{-1} \text{ s}^{-1}$$

$$\text{Time } t = \frac{[R]_0 - [R]}{k} = \frac{0.10 - 0.075}{0.0030}$$

$$\underline{t = 0.33 \text{ second}}$$

[3.] For a reaction $\text{NaO}_5 \rightarrow 2\text{NO}_2 + \frac{1}{2}\text{O}_2$

i) Show that it follows first order reaction?

ii) calculate the half life?

time (s)	0	300	600
$[\text{NaO}_5] (\text{mol L}^{-1})$	1.6×10^{-2}	0.8×10^{-2}	0.4×10^{-2}

[Delhi 2017]

3M

Answer -: For first order reaction $k = \frac{2.303}{t} \log \frac{[\text{NaO}_5]_0}{[\text{NaO}_5]}$

a) At $t = 300 \text{ second}$, $[\text{NaO}_5]_0 = 1.6 \times 10^{-2} \text{ mol L}^{-1} \text{ s}^{-1}$ and $[\text{NaO}_5] = 0.8 \times 10^{-2} \text{ mol L}^{-1} \text{ s}^{-1}$

$$k = \frac{2.303}{300} \log \frac{1.6 \times 10^{-2}}{0.8 \times 10^{-2}} = 2.31 \times 10^{-3} \text{ s}^{-1}$$

b) At $t = 600$, $[\text{NaO}_5] = 0.4 \times 10^{-2}$

$$k = \frac{2.303}{600} \log \frac{1.6 \times 10^{-2}}{0.4 \times 10^{-2}} = 2.31 \times 10^{-3} \text{ s}^{-1}$$

→ Rate constant remains same in both condition (a & b). So we can say above reaction is first order.

ii) $K = 2.31 \times 10^{-3} \text{ s}^{-1}$ then $t_{\frac{1}{2}} = \frac{0.693}{K} = \frac{0.693}{2.31 \times 10^{-3}} = 300 \text{ s}$.

[4] → first order reaction $2 \text{N}_2\text{O}_5(g) \rightarrow 4 \text{NO}_2(g) + \text{O}_2(g)$ at 40°C , data is given below —

3M

[Delhi 2011]

i) Calculate the rate constant (with its unit).

$$\text{At time } t = 20 \text{ min}, [\text{N}_2\text{O}_5] = 0.209 \frac{\text{mol}}{\text{L}}$$

$$\& [\text{N}_2\text{O}_5]_0 = 0.4 \frac{\text{mol}}{\text{L}}$$

$$K = \frac{2.303}{20} \log \frac{0.4}{0.209} = \frac{2.303}{20} \log 1.904$$

$[\text{N}_2\text{O}_5](\text{mol/L})$	Time (min)
0.400	0.00
0.209	20.0
0.209	40.0
0.151	60.0
0.109	80.0

$$K = 0.01625 \text{ min}^{-1} = 1.62 \times 10^{-2} \text{ min}^{-1}$$

ii) → What will be concentration of N_2O_5 after 100 min?

$$K = \frac{2.303}{100} \log \frac{0.4}{[\text{N}_2\text{O}_5]} = 1.62 \times 10^{-2} \text{ then } \log \frac{0.4}{[\text{N}_2\text{O}_5]} = \frac{1.62}{2.303} = 0.705$$

$$\rightarrow \frac{0.4}{[\text{N}_2\text{O}_5]} = \text{antilog } 0.705 = 5.07 \text{ then } [\text{N}_2\text{O}_5] = \frac{0.4}{5.07} = 0.078 \frac{\text{mol}}{\text{L}}$$

iii) → Initial rate of reaction? All the Best ✓

$$\text{Initial rate} = K [\text{N}_2\text{O}_5] = 0.4 \times 1.62 \times 10^{-2} = 6.4 \times 10^{-3} \frac{\text{mol}}{\text{L min}}$$

5. :- for a reaction $A + B \rightarrow P$ the rate is given by Rate = $K[A][B]^2$

[CBSE 2015 | Delhi 2015]

a) → How is the rate affected if $[B]$ is doubled?

2M

$$\text{Rate of reaction} = K [A] [2B]^2 = 4 [A] [B]^2 = 4 \times \text{original rate}$$

b) → What is overall order of reaction if A is present in large excess?

→ If A is present in large excess, then the reaction will be independent of the concentration of A.

$$\text{Rate} = \underbrace{K[A]}_{\text{constant}} [B]^2 = K' [B]^2$$

Thus order of reaction = 2