

Cold response in *Saccharomyces cerevisiae*: new functions for old mechanisms

Jaime Aguilera, Francisca Randez-Gil & Jose Antonio Prieto

Department of Biotechnology, Instituto de Agroquímica y Tecnología de los Alimentos, Consejo Superior de Investigaciones Científicas, Burjassot, Valencia, Spain

Correspondence: Francisca Randez-Gil, Department of Biotechnology, Instituto de Agroquímica y Tecnología de los Alimentos (CSIC), PO Box 73, 46100 Burjassot (Valencia), Spain. Tel.: +34 963900022; fax: +34 963636301; e-mail: randez@iata.csic.es

Received 4 July 2006; revised 15 December 2006; accepted 22 December 2006.
First published online 9 February 2007.

DOI:10.1111/j.1574-6976.2007.00066.x

Editor: Karl-Dieter Entian

Keywords

yeast; cold-shock; stress; freezing; signalling; signal transduction pathways.

Abstract

The response of yeast cells to sudden temperature downshifts has received little attention compared with other stress conditions. Like other organisms, both prokaryotes and eukaryotes, in *Saccharomyces cerevisiae* a decrease in temperature induces the expression of many genes involved in transcription and translation, some of which display a cold-sensitivity phenotype. However, little is known about the role played by many cold-responsive genes, the sensing and regulatory mechanisms that control this response or the biochemical adaptations at or near 0 °C. This review focuses on the physiological significance of cold-shock responses, emphasizing the molecular mechanisms that generate and transmit cold signals. There is now enough experimental evidence to conclude that exposure to low temperature protects yeast cells against freeze injury through the cold-induced accumulation of trehalose, glycerol and heat-shock proteins. Recent results also show that changes in membrane fluidity are the primary signal triggering the cold-shock response. Notably, this signal is transduced and regulated through classical stress pathways and transcriptional factors, the high-osmolarity glycerol mitogen-activated protein kinase pathway and Msn2/4p. Alternative cold-stress generators and transducers will also be presented and discussed.

Introduction

Decreases in ambient temperature are common in almost every ecological niche. A drop in temperature may take place seasonally, daily or just unexpectedly, depending on the region, climate and environment. The effects of low temperature on life have been studied thoroughly, as cold influences the structural and functional properties of cellular components negatively, both physically and chemically. A decrease in membrane fluidity and diffusion rates, alterations in molecular topology or modifications in enzyme kinetics reportedly occur as a consequence of low temperatures (Gast *et al.*, 1993; Thieringer *et al.*, 1998; Inouye, 1999). To withstand such temperature downshifts, organisms have developed mechanisms that enable them to adapt and survive. These mechanisms (commonly referred to as the 'cold-shock response') have been extensively studied in bacteria and plants. In *Escherichia coli*, *Bacillus subtilis* and other prokaryotes, a certain group of proteins is specifically (but not uniquely) induced upon a cold-shock (Thieringer *et al.*, 1998; Gualerzi *et al.*, 2003). These cold-shock proteins are involved in transcription, translation and other funda-

mental functions that play a role in maintaining nucleic acid structure. However, little is known about the mechanisms underlying the transcriptional activation of cold-responding genes. In plants, cold-inducible genes cover a wider range of functions, and some of them also respond to drought or high salinity (Shinozaki & Yamaguchi-Shinozaki, 1996; Seki *et al.*, 2002b). Moreover, it is thought that there is some cross-communication between the signal transduction pathways responding to these stresses (Seki *et al.*, 2002a, 2004), although an exclusive pathway for cold-induced gene expression has also been proposed (Shinozaki & Yamaguchi-Shinozaki, 1996).

Although the cold-shock response has been widely studied in bacteria and plants, relatively little attention has been paid to fungi, and specifically to yeast. The budding yeast *Saccharomyces cerevisiae* has colonized different natural and artificial environments, all of which can be subject to low temperatures. Yeast growing on the ground or on the surface of grapes and other fruits are exposed to temperature downshifts like those described above. On the other hand, when these yeasts are used in certain brewing and wine fermentations, they are exposed to temperatures of around

10–12 °C, far below their natural physiological temperature (25–30 °C). Moreover, yeasts, as well as other microorganisms, are stored at very low temperature (4 °C) in industrial and laboratory conditions. This temperature is chosen because it is generally accepted that although it is growth-restrictive, viability is fully maintained over long periods of time.

All these features indicate that *S. cerevisiae* must have the necessary molecular machinery to survive and adapt to cold, and that this is probably one of the reasons why this species is so widespread. Therefore, the study of the cold-shock response in *S. cerevisiae* is of crucial interest for both basic and technological reasons (Randez-Gil *et al.*, 2003). The growth in high-throughput analytical techniques has led to a number of publications on this subject in recent years. Our knowledge of the cold response in yeast, although limited, is expanding and we are gaining new insight into other molecular mechanisms, signalling pathways and regulators that, until now, were apparently unrelated to low temperature. The state of the art and some possible models for cold-signal generation, signal transduction and function of cold-responding genes, as well as the prospects for future research, are discussed in this review.

Time and temperature-dependent and -independent genetic responses to cold

On analysing the different gene expression patterns in *S. cerevisiae* after a decrease in temperature, it becomes clear that there is a common response involving certain groups of genes (Sahara *et al.*, 2002; Homma *et al.*, 2003; Kandror *et al.*, 2004; Schade *et al.*, 2004; Murata *et al.*, 2006). Among these are the genes encoding members of the DAN/TIR family of putative cell-wall mannoproteins, temperature-shock inducible protein (*TIP1*), *TIR1/SRP1*, *TIR2* and *TIR4*, which are induced at 10, 4 °C (Sahara *et al.*, 2002; Homma *et al.*, 2003; Schade *et al.*, 2004; Murata *et al.*, 2006; Panadero *et al.*, 2006) and even at 0 °C (Kandror *et al.*, 2004). Similarly, *TIR*-related DNA sequences, seripauperin (PAU) family proteins (*PAU1*, *PAU2*, *PAU4*, *PAU5*, *PAU6* and *PAU7*), which have been shown to display phospholipid-interacting activity (Zhu *et al.*, 2001), are also up-regulated after cold-shock (Homma *et al.*, 2003; Murata *et al.*, 2006).

Apart from these characteristic cold-stress markers, part of the genetic response to low temperature seems to be time-dependent. Examination of different microarray-based studies reveals sequential changes in the transcriptional profile during the time-course of exposure to 10 °C. Researchers have noted two (Schade *et al.*, 2004) or even three (Sahara *et al.*, 2002) stages, depending on the stress duration. Initial responses (0–2 h) include the enhanced expression of key genes involved in phospholipid synthesis, such as *INO1* and *OPI3*, and fatty-acid desaturation (*OLE1*) (Sahara *et al.*, 2002;

Schade *et al.*, 2004). Other genes that are induced within this period are those related to transcription (RNA helicases such as *DBP2*, RNA polymerase subunits such as *RPA49* and RNA processing proteins such as *NSR1*, among others), and an important number of ribosomal protein genes. Thus, 94 of the 323 genes found to be up-regulated after yeast was incubated for 2 h at 10 °C encode ribosomal proteins (Sahara *et al.*, 2002).

Remarkably, most of these transcription-related and ribosomal genes are abruptly repressed after longer incubation periods (4–24 h) at 10 °C (Sahara *et al.*, 2002; Schade *et al.*, 2004) and also at 4 °C (Homma *et al.*, 2003; Murata *et al.*, 2006). This phase is characterized to some degree by the transcriptional activation of typical stress-marker genes. Indeed, some members of the main gene family involved in cellular protection, the heat shock protein (*HSP*) genes, are induced at this stage. Genes encoding HSPs, namely *HSP12*, *HSP26*, *HSP42*, *HSP104*, *SSA4*, *SSE2* and *YRO2*, are up-regulated after 4–12 h of transfer to 10 °C (Sahara *et al.*, 2002; Schade *et al.*, 2004), as well as at similar times at 4 °C (Homma *et al.*, 2003; Murata *et al.*, 2006), 0 °C (Kandror *et al.*, 2004) and –80 °C (Odani *et al.*, 2003). Other members of the HSP family (i.e. *CIS3*, *HMS2*, *HSC82*, *HSP30*, *HSP60*, *HSP78*, *HSP82*, *HSP150*, *SSA1*, *SSA2*) are repressed at 10 °C (Sahara *et al.*, 2002), whereas at lower temperatures their mRNA content increases (Murata *et al.*, 2006). This discrepancy indicates that different members of the HSP gene family are differentially regulated in both a time- and a temperature-dependent manner. The inventory of genes that are induced after 4–6 h of cold incubation and beyond also includes genes involved in the metabolism of reserve carbohydrate glycogen (*GLG1*, *GSY1*, *GLC3*, *GAC1*, *GPH1* and *GDB1*), and trehalose (*TPS1*, *TPS2* and *TSL1*), and genes for detoxifying reactive oxygen species (ROS) and defence against oxidative stress, such as catalase (*CTT1*), glutaredoxin (*TTR1*), thioredoxin (*PRX1*) and glutathione transferase (*GTT2*).

Cold-sensitive mutants and the cold ribosome adaptation hypothesis

In view of the available data, it appears that yeasts adapt to temperature decreases in terms of the duration and the extent of the stress. This is especially evident if we consider the opposite regulation displayed by ribosomal and transcription-related genes between short and long incubation periods at 10 or 4 °C (Sahara *et al.*, 2002; Schade *et al.*, 2004; Murata *et al.*, 2006), which is consistent with previous reports about cold growth defective mutants. Several of the 106 mutants displaying growth defects at low temperatures (Hampsey, 1997) (catalogued in <http://mips.gsf.de/proj/yeast/CYGD/db/index.html>) are affected in ribosomal proteins and translation (Winzeler *et al.*, 1999; Zhang *et al.*, 2001),

or in proteins involved in pre-rRNA processing for ribosome biogenesis or assembly (Lee & Baserga, 1997), protein folding (Craig & Jacobsen, 1985), exocytosis (Lehman *et al.*, 1999) and nucleus–cytosol exchange (Noguchi *et al.*, 1997, 1999). These observations indicate the need for remodelling the translational machinery and secondary structure of nucleic acids compromised by cold-shock; moreover, this could support a model for cold ribosome adaptation in *S. cerevisiae* that is similar to that proposed for *E. coli* (Jones & Inouye, 1996). At temperatures that are low, but still permit growth, there is an up-regulation of genes that are essential for cold growth (Fig. 1). By contrast, such genes are expressed minimally or repressed after prolonged periods of stress (when all the transcriptional and translational machinery has already been remodelled) or under severe cold conditions as at 4 °C (generally considered to be a growth-restrictive temperature, although minimal growth still occurs), where low or no *de novo* protein synthesis is required (Fig. 1).

Even so, results from transcriptomic experiments reveal that several genes respond to cold conditions regardless of exposure time and temperature. This finding suggests that not all the cold-regulated genes are involved in functions related to growth.

The freeze-protective function of the cold response

Although some of the cold-responsive proteins might be essential for the cells to adapt and resume growth under the new unfavourable environmental conditions, some evidence suggests that the main aim of other common cold-shock responses is to protect cells against freeze injury (Fig. 1). For example, the stability of Ole1p, the only fatty-acid desatur-

ase known in *S. cerevisiae* (Stukey *et al.*, 1989, 1990), seems to be important for cold growth (Loertscher *et al.*, 2006); however, overexpression of *OLE1* does not confer growth advantages at low temperatures (Kajiwar *et al.*, 2000). By contrast, production of recombinant desaturases increased the unsaturation index and fluidity of the yeast membrane, and positively influenced freeze tolerance of baker's yeast cells (Rodríguez-Vargas *et al.*, 2007). These observations suggest that changes in membrane composition after a decrease in temperature might be important in influencing cell survival upon freezing. Membrane organization and dynamic properties are the main targets of freeze injury (Wolfe & Bryant, 1999). Hence, the cold-instigated induction of fatty acid desaturases, cell-wall mannoproteins and seripauperins are consistent with this idea. Similarly, it has been shown that, in addition to temperature downshifts (Zhang *et al.*, 2003), freezing and thawing generate superoxide anions and free radicals (Park *et al.*, 1997; Du & Takagi, 2005), which is consistent with the cold-provoked up-regulation of genes for ROS detoxification and defence against oxidative stress, and with genes encoding HSPs. In this sense, HSPs are molecular chaperones involved in the response to several kinds of stress, preventing protein denaturation and misfolding (Papp *et al.*, 2003; Young *et al.*, 2003).

A downshift in temperature also activates the metabolism of reserve carbohydrates. It has been suggested that glycogen turnover occurs in response to different kinds of stress, such as heat-shock or oxidative damage, because the genes for both its synthesis and its degradation are simultaneously induced, with no net accumulation of this polysaccharide (Parrou *et al.*, 1997). In spite of the similar transcriptional profile observed at low temperatures (Sahara *et al.*, 2002; Schade *et al.*, 2004; Murata *et al.*, 2006), such recycling does




Fig. 1. Main responses of *Saccharomyces cerevisiae* to decreasing temperatures. At mid- to low temperatures, the specific transcriptional and translational machinery is up-regulated in order to enable cold-adapted growth. These mechanisms are switched off when the temperature is low enough to be growth-restrictive. By contrast, freeze-protective mechanisms, such as fatty acid desaturation and synthesis of osmolytes, are triggered even at permissive temperatures, and intensified at near-freezing conditions, a situation in which further protection systems, such as HSPs, are also activated.

not appear to be the case for cold stress, because glycogen is significantly accumulated in yeast after 12 h of treatment at 10 °C (Schade *et al.*, 2004). Like glycogen, trehalose starts to accumulate after 12 h of incubation at 10, 4 or 0 °C regardless of whether there is growth or not (Kandror *et al.*, 2004; Schade *et al.*, 2004). Consistent with this, the genes involved in the biosynthesis of this disaccharide are cold-induced after 4–8 h of incubation at 10 °C. The same activation occurs at lower temperatures, showing maximum induction levels at 0 °C (Sahara *et al.*, 2002; Kandror *et al.*, 2004; Schade *et al.*, 2004; Murata *et al.*, 2006). Trehalose is known to accumulate at high levels in response to different stress conditions (Blomberg, 2000; Estruch, 2000), and several protecting roles have been proposed for this disaccharide, such as membrane and protein stabilization (Singer & Lindquist, 1998; Elbein *et al.*, 2003; Gancedo & Flores, 2004). Moreover, high levels of trehalose have also been correlated with freezing resistance (Kim *et al.*, 1996). Although trehalose is not needed for growth at 10 °C (Schade *et al.*, 2004), the results obtained by Kandror *et al.* (2004) revealed that this compound protects cells against very cold conditions. In fact, viability of cells incubated at 0 °C for 5–20 days correlated with the intracellular trehalose content, and cold-instigated accumulation of the disaccharide protected the cells from viability losses due to freezing. Similar behaviour has been observed with glycerol. *GPD1*, the gene encoding the main enzyme involved in glycerol synthesis (Albertyn *et al.*, 1994), is activated upon a shift to low temperature, and cells begin to accumulate this osmolyte, displaying higher values at 4 °C than at 12 °C (Panadero *et al.*, 2006). Like trehalose, glycerol accumulation provides freeze-protection, although it is not needed for growth at 12 °C (Panadero *et al.*, 2006). Glycerol is the only osmoprotectant solute accumulated by yeast upon hyperosmotic stress, and is also a by-product of redox homeostasis. The protective role of glycerol against freezing is well known and has been explained in terms of the osmotic shrinkage resulting from freezing/thawing processes (Wolfe & Bryant, 1999; Panadero *et al.*, 2006).

In view of the above results, there is a clear link between exposure to low temperatures and resistance to freezing. In natural environments, frosts are often preceded by periods of very low temperatures, so this hypothesis is meaningful from an evolutionary point of view. However, there are still some aspects that need to be clarified. For example, there is no evidence of the possible function of glycogen in cold response, and a better explanation is needed for the protective role of trehalose at 0 °C, a temperature at which cells are not frozen. On the other hand, the function of DAN/TIR and PAU family proteins remains unclear. Future work will contribute to clarifying the dual aspects of cold-shock responses, as a way to survive, adapt and grow at low temperatures, as well as a protective mechanism for subsequent freezing.

The cold signal and cold-sensing mechanism

For transcriptional activation to be triggered, yeast cells must sense temperature downshifts. Evidence indicates that changes in the physical state of the membrane could be perceived as a primary signal of a variation in temperature (Vigh *et al.*, 1998). This concept was first put forward by Carratù *et al.* (1996), who demonstrated that a membrane lipid perturbation causes a signal to be triggered, inducing the transcription of heat-shock genes in yeast. Based on this finding and additional evidence, the group of Murata demonstrated that, in the cyanobacteria *Synechocystis*, a drop in temperature causes a reduction of the membrane fluidity, and this plays a role in the perception of low temperatures and the subsequent signal transduction (Inaba *et al.*, 2003; Los & Murata, 2004). Genetically engineered *Synechocystis* cells that lacked genes encoding fatty acid desaturases displayed higher membrane rigidity than the wild-type at both low and physiological temperatures, showing enhanced cold inducibility of gene activation (Inaba *et al.*, 2003). In *Synechocystis*, expression of these desaturase genes upon cold-shock is controlled by the histidine kinase Hik33 (Suzuki *et al.*, 2000, 2001), an integral membrane protein that functions as a cold sensor. Similarly, DesK in *B. subtilis* (Aguilar *et al.*, 2001) and TRP-channels in the mammalian nervous system (McKemy *et al.*, 2002; Peier *et al.*, 2002), which are integral membrane proteins, act as cold-sensors. Furthermore, the extent to which the DesK protein senses temperature is regulated by the rigidity of the membrane lipid bilayer (Mansilla & de Mendoza, 2005). Overall, these observations indicate that changes in the physical state of the membrane caused by a temperature downshift are recognized by cold-sensors anchored in the membrane, generating the cold-signal.

Recently, Hik33 has been shown to regulate the expression of osmostress-inducible genes in *Synechocystis* (Mikami *et al.*, 2002), suggesting that both osmotic- and cold-stress could be perceived in this organism by common mechanisms and sensing elements. Like Hik33, Sln1p, the only known yeast histidine kinase sensor, has recently been reported in the cold-sensing mechanism (Panadero *et al.*, 2006). Sln1p, together with Ypd1p and Ssk1p, forms a phosphorelay system, which transmits the osmostress signal through different elements of the high-osmolarity glycerol (HOG) pathway, the most important osmotic stress-responding cascade in *S. cerevisiae* (Hohmann, 2002; Westfall *et al.*, 2004). Activation of the HOG pathway was also found to be induced by the membrane-rigidifier agent dimethylsulfoxide (DMSO) (Panadero *et al.*, 2006). Several lines of research in plants and cyanobacteria have shown that this chemical indeed mimics the changes in membrane fluidity caused by a sharp decrease in temperature, and triggers the

activation of several cold-induced mitogen-activated protein kinases (MAPKs) (Örvar *et al.*, 2000; Sangwan *et al.*, 2002). Thus, Sln1p appears to function under either of these stressful conditions (Hayashi & Maeda, 2006; Panadero *et al.*, 2006) and it is likely, therefore, that the basic mechanism of its activation by low temperature and osmotic pressure could be similar or even identical.

The fluidity state of the cell membrane might be a key factor in integrating the sensing mechanism of cold and hyperosmolarity. There is evidence that different osmosensors, such as EnvZ in *E. coli* or OpuA, a transmembrane transporter of *Lactobacillus lactis* with osmosensor and osmoregulator properties, are stimulated by changes in the fluidity of membrane lipids (Tokishita & Mizuno, 1994; van der Heide & Poolman, 2000; van der Heide *et al.*, 2001). Moreover, studies in *B. subtilis* and *S. cerevisiae* indicate that osmotic stress reduces cell-membrane fluidity (López *et al.*, 2000; Laroche *et al.*, 2001). Altogether, these observations suggest a model in which Sln1p monitors the changes in membrane fluidity caused by different stressors. Moreover, the fact that Sln1p responds to DMSO supports this view. However, more work is required to confirm this idea and to understand the exact molecular mechanism of cold-signal perception.

Cold-induced transduction pathways and transcription factors

Signal transduction pathways are the link between the sensing mechanism and the genetic response. As is the case for the cold signalling mechanism, to date there is no signal transduction pathway or transcription factor that is known to respond exclusively to low temperatures. Although the existence of such a pathway cannot be ruled out, available data point to known signal transduction mechanisms, operating under other kinds of stimuli, which may also be triggered by cold stress. We shall now go on to describe putative and known regulatory mechanisms controlling cold-instigated gene expression.

The MOX factors

The coordinated expression of the *DAN/TIR* genes under anaerobic conditions reportedly depends on the transcriptional activator *MOX4*, whereas their repression, in conditions of oxygen availability, is mediated by the two repression factors *MOX1* and *MOX2* in a heme-dependent fashion (Abramova *et al.*, 2001a). All of these factors act through the consensus sequence AR1 (Cohen *et al.*, 2001). Mox4p, Mox1p and Mox2p (or some factors dependent upon them) are believed to form a heme-sensitive complex similar to the galactose-sensitive Gal4p-Gal80p

(Zenke *et al.*, 1996; Platt & Reece, 1998). Like *GAL4* regulation, *MOX4* expression is autoregulated in anaerobic cells via its AR1 sequence. Interestingly, expression of *TIR* genes under cold-shock was eliminated in a *mox4* mutant strain (Abramova *et al.*, 2001b). Furthermore, *MOX4* mRNA was also more abundant in cold-shocked cells (Abramova *et al.*, 2001a), suggesting that induction of the activator contributes to up-regulation of *TIR* genes under this condition (Fig. 2).

Regulation of *OLE1* expression and RUP processing

Unlike *DAN/TIR* genes, the promoter of *OLE1* does not contain AR1 sequences. Unsaturated fatty acids (UFAs) mediate *OLE1* repression via the *cis* transactivation element FAR (fatty acid regulated). Two genes for fatty acid synthesis (*FAA1* and *FAA4*), the transcriptional activator *HAP1* and the acyl-coenzyme A (CoA)-binding protein Acb1p have also been reported to regulate *OLE1* expression (Choi *et al.*, 1996). In addition, Spt23p and Mga2p, two functionally and structurally related factors, are also necessary for *OLE1* transcription (Zhang *et al.*, 1999) (Fig. 2). Spt23p appears to play a major role in fatty acid regulation, whereas Mga2p is known to be essential for the hypoxic induction of *OLE1* through a consensus sequence named LORE (for low-oxygen response element) (Jiang *et al.*, 2001). This sequence is also found in several genes induced by hypoxia, including *TIR1*. All these genes could, therefore, share regulation via LORE (Vasconcelles *et al.*, 2001). The transcriptional induction of *OLE1* under cold-shock is also dependent on Mga2p (Nakagawa *et al.*, 2002), suggesting that the LORE-binding complex could drive the expression of a subset, at least, of cold-responsive genes (Fig. 2).

Moreover, Mga2p could also play a role as a cold-sensor, as previously postulated (Nakagawa *et al.*, 2002). This hypothesis was constructed on the basis that a novel pathway has been described in *S. cerevisiae*, the so-called 'RUP (regulated ubiquitin/proteasome-dependent) processing' pathway, which regulates the activation of Spt23p and Mga2p by UFAs (Hoppe *et al.*, 2000, 2001). Spt23p and Mga2p are initially created as dormant precursors, anchored in the endoplasmic reticulum or nuclear envelope membranes by their C-terminal tails. The shortage of UFAs leads to the ubiquitination of Spt23p and Mga2p, releasing the N-terminal transcription factor domain into the cytosol by the action of the 26S proteasome in a dual way: first, processing the transcriptional factor into its active form and second, destroying it after the activation of genes such as *OLE1* (Auld *et al.*, 2006). In the processing of Mga2p, UFAs exert a modest influence, with hypoxia being the main signal that triggers its processing (Jiang *et al.*, 2002). In this scenario, a change in the physical state of the membrane




Fig. 2. Cold-instigated activation of the MOX factors and RUP processing. (1) When the repressor complex Mox1p–Mox2p is released, the Mox4p factor promotes the transcription of *TIP/TIR* genes. The cold-provoked induction of these genes is eliminated in a *mox4* mutant, suggesting that low temperatures are able to release the repressor complex. (2) The transcriptional induction of *OLE1* under cold-shock is dependent on the transcription factor Mga2p, which is released from the nuclear envelope by proteasome-dependent deubiquitination provoked by the action of the Rsp5p factor. This action could therefore be triggered by cold stress (see text for details).

due to a downshift in temperature or to hypoxia conditions (Nakagawa *et al.*, 2003) could activate Mga2p by starting off the RUP processing (Fig. 2).

The cAMP-PKA and general stress response

Thirty-two protein phosphatase genes have been identified in the *S. cerevisiae* genome (Sakumoto *et al.*, 1999). From all of them, only *YVH1* was found to be induced by cold-shock (Hampsey, 1997) and to play a role in growth at low temperatures (Sakumoto *et al.*, 1999). *YVH1* encodes a dual-specificity protein phosphatase (DSP), which exhibits a catalytic domain displaying protein-tyrosine phosphatase activity and a C-terminal cysteine-rich motif (Fauman & Saper, 1996; Muda *et al.*, 1999). Yvh1p shares the control of some cellular processes such as sporulation, growth and glycogen accumulation with protein kinase A (PKA). The cAMP-PKA pathway in *S. cerevisiae* plays a major role in the control of the genetic response to a wide variety of stresses (Thevelein & de Winde, 1999; Estruch, 2000; Wilson & Roach, 2002; Santangelo, 2006), through the phosphorylation of the Msn2p and Msn4p transcription factors (Griffioen & Thevelein, 2002) (Fig. 3, pathway 1). Upon stressful conditions, Msn2/4p migrates to the nucleus and promotes the transcription of an important number of stress-respond-

ing genes, which have the recognition sequence called STRE (stress response element) in their promoter (Martínez-Pastor *et al.*, 1996). It has been suggested that Yvh1p could negatively modulate the cAMP level or a downstream component of the PKA signalling cascade. In fact, deletion of *YVH1* resulted in decreased expression of a multi-STRE reporter construction (Beeser & Cooper, 2000). Like Yvh1p, Mck1p, a Ser/Thr/Tyr protein kinase, might also be involved in the cold-shock signalling via PKA. Genetic data indicate that Yvh1p acts upstream of Mck1p, at least in sporulation signalling (Beeser & Cooper, 1999). Moreover, Mck1p is involved in the phosphorylation of Bcy1p (Griffioen *et al.*, 2003), the negative regulator of PKA. This modification occurs under environmental stress conditions such as heat-shock, salt or oxidative stress and determines the distribution of Bcy1p between the nucleus and cytoplasm. The specific relocation of Bcy1p could help to drive the PKA activity towards certain substrates under these conditions. In addition, Mck1p is necessary for the correct stress response, owing to its ability to promote indirectly a DNA-Msn2p complex (Hirata *et al.*, 2003), without the need of Msn2p phosphorylation. Deletion of *MCK1* leads to a growth defect at low temperature (de Jesus Ferreira *et al.*, 2001) and inhibition of STRE-dependent transcription (Hirata *et al.*, 2003), similar to that found in the *yvh1* mutant (Sakumoto *et al.*, 1999).




Fig. 3. cAMP-PKA and TOR pathways. (1) Mutants in the Tpk1/2/3p (PKA)-negative regulators Yvh1p and Mck1p show growth defects at low temperatures, indicating the involvement of the cAMP-PKA pathway in the cold response. This pathway starts with the formation of the GDP-binding complex Ras1/2p/Ira1/2p/Cdc25p, which activates Cyr1p for cAMP synthesis. Inhibition of this synthesis by cold, in addition to Yvh1p and Mck1p activation, would prevent Tpk1/2/3p-mediated phosphorylation of the transcription factors Msn2/4p, avoiding their nuclear export. (2) Disruption of either *TPD3* or *CDC55* causes cold sensitivity. Inhibition of Tap42p phosphorylation caused by cold would prevent the formation of the complex Tap42p/Php21/22p/TPD3p/Cdc55p, also blocking the exportation of Msn2/4p from the nucleus, with the subsequent induction of genes encoding trehalose, glycerol and other STRE genes. By contrast, localization of Msn2/4p is regulated by PKA in a TORC1-dependent manner, whereas the TORC2 complex acts on Rom2p activation (see text for details).

The cold signal transduction via the cAMP-PKA signaling pathway would, as anticipated, be essential in determining the genetic response to cold. Indeed, genes induced at the late phase of a cold-shock, like those encoding trehalose

and glycogen metabolism enzymes, as well as *HSP12*, *HSP26*, *HSP42*, *SSA4* and the ROS-defensive genes *CTT1*, *PRX1* and *TTR1*, are known to be strictly dependent on Msn2/4p (Kandror *et al.*, 2004; Schade *et al.*, 2004).

Expression of early responding genes such as *DPB2*, *NSR1*, *OPI3* and *RPA49*, as well as *PAU* genes, whose cold-provoked induction is maintained over time, has also been reported to depend on a functional PKA pathway (Wang *et al.*, 2004). Moreover, signalling through the cAMP-PKA pathway would play a major role in providing cold adaptation and tolerance to freezing. Mutants lacking *MSN2/MSN4* die more rapidly at 0 °C or upon freezing (Kandror *et al.*, 2004). Mutants in *RAS2*, the gene for a regulatory protein in the synthesis of cAMP, clearly showed a better freeze-thaw tolerance than the wild-type, while mutants in *BCY1* were sensitive to this stress (Park *et al.*, 1997). In the same regard, mutants from a commercial baker's yeast strain, showing partial inactivation in adenylate cyclase, were better at maintaining freeze resistance (van Dijck *et al.*, 2000). These results indicate that any event involving a decrease in the level of PKA activity improves cell tolerance to freeze stress. Consistent with this, deletion of *IRA2* encoding the Ras GTPase (an activator of the PKA pathway) led to a freeze-sensitivity phenotype. Similarly, mutants in *PDE2*, the high-affinity phosphodiesterase encoding gene, also showed impaired freezing tolerance (Park *et al.*, 2005b).

The PKC pathway

Signalling through the so-called PKC pathway (for a review, see Levin, 2005) is triggered under conditions that jeopardize cell-wall stability, including heat-shock (Kamada *et al.*, 1995) and hypo-osmotic stress (Davenport *et al.*, 1995). Activation of the G-protein Rho1, the master regulator of the pathway, triggers a linear MAPK signalling cascade that starts with Pkc1p, a homologue to the mammalian protein kinase C (PKC), and finish with the Slt2p/Mpk1p kinase. The PKC pathway controls important cell functions, such as cell-wall maintenance and actin polarization. In addition, Pkc1p regulates additional targets that are separated from the MAPK cascade (Levin, 2005). Activation of Rho1p is stimulated by Rom2p, which has GDP/GTP exchanging activity. Disruption of *ROM2* leads to temperature-sensitive growth defects at both high (37 °C) and low (11 °C) temperatures (Manning *et al.*, 1997). These defects include failure to form normal bud polarization and sensitivity to the microtubule depolymerizing drug benomyl. Although the activation of the PKC pathway by heat stress is well documented (Kamada *et al.*, 1995; Heinisch *et al.*, 1999), the phenotype of the *rom2* null mutant suggests a general function of PKC in perceiving and signalling changes in environmental temperature. Some collateral observations support this idea. For example, mutants in several genes encoding proteins related to actin polarization are cold sensitive. Among them, Num1p (Revardel & Aigle, 1993) interacts with Bni1p, a formin that nucleates the assembly of linear actin filaments in response to activation by the PKC

pathway (Levin, 2005). In addition, some mutants in Rho3p, a homologue of Rho1p that is involved in Bni1p activation during bud growth, also confer sensitivity to low temperatures (Imai *et al.*, 1996).

PKC may also play a role in connection with the cAMP-PKA pathway. Evidence of this interaction has been provided by Park *et al.* (2005a). These authors demonstrated that Rom2p can negatively regulate the Ras-cAMP pathway by controlling cAMP levels. Thus, a *rom2* null mutant showed sensitivity to freeze stress, a phenotype that was suppressed by additional mutation of *RAS2* or in a *tpk1 tpk2* background. By contrast, the absence of either *IRA2* or *PDE2* exacerbated the freeze-sensitivity phenotype of the single *rom2* mutant.

The TOR pathway

In plants, cold acclimatization is associated with rapid and reversible changes in the phosphorylation status of specific pre-existing proteins that allow the expression of cold-acclimatized specific (CAS) genes (Monroy *et al.*, 1993). The type 2A Ser/Thr protein phosphatase (PP2A) is involved in basic cellular processes, such as metabolism, transcription and signal transduction (Ariño *et al.*, 1993). One of the signals regulated by PP2A is a downshift in temperature (Monroy *et al.*, 1998). The composition of PP2A subunits determines its specificity, activity and sub-cellular localization. In plants, *TAP46* codes for a PP2A regulatory subunit, and is induced by cold-shock, suggesting that Tap46p targets PP2A under this condition (Harris *et al.*, 1999). In *S. cerevisiae*, the homologue to plant *TAP46* is *TAP42*. As in plants, Tap42p associates to PP2A catalytic subunits, Sit4p or Pph21/22p (di Como & Arndt, 1996; Jiang & Broach, 1999). This association depends on the Tor proteins, phosphatidylinositol kinase-related kinases, which are members of the target of rapamycin (TOR) signalling cascade, a pathway that controls cellular functions necessary for cell growth and metabolism in response to environmental cues (Schmelzle & Hall, 2000; Wullschlegel *et al.*, 2006). By contrast, inhibition of yeast TOR activity leads to transcriptional activation of genes so they can adapt to nutrient depletion (di Como & Arndt, 1996; Jiang & Broach, 1999; Rohde *et al.*, 2004). Recently, it has been shown that inactivation of Tap42p prolongs the up-regulation of several Msn2/4p-dependent genes after heat-shock (Düvel *et al.*, 2003). Therefore, a functional TOR pathway would be involved in nuclear export of the transcription factors under nonstressing conditions, a fact that has already been demonstrated for Msn2p (Santhanam *et al.*, 2004). Phosphorylated Tap42p competes with Cdc55p/Tpd3p for binding to the catalytic subunit Pph21/22p, increasing protein synthesis under these conditions (Jiang & Broach, 1999). Interestingly, disruption of either *TPD3* or *CDC55* causes cold

sensitivity (Healy *et al.*, 1991). These data strongly suggest that, as in plants, PP2A via Tap42p, Cdc55p and Tpd3p may play a role in cold response, and that the transcription factors Msn2/4p mediate the TOR signalling (Fig. 3, pathway 2). Hence, the cold-provoked induction of Msn2/4p-dependent genes would be signalled through TOR, as well as via the PKA pathway. In this respect, it has recently been proposed that the cAMP-PKA pathway is a TOR effector branch (Schmelzle *et al.*, 2004) involved in Msn2p localization and other read-outs that are not controlled by Tap42p/Sit4p.

Further evidence that the TOR pathway is involved in cold response comes from the functional analysis of distinct TOR complexes (Loewith *et al.*, 2002; Wullschlegel *et al.*, 2006). As well as its shared role with Tor1p in cell growth, Tor2p displays a unique function in the organization of cytoskeleton. Thus, *tor2* mutants exhibit abnormal polarized distribution of the actin cytoskeleton, a phenotype that is rescued by overexpression of the actin-specific TCP-1 chaperone (Schmidt *et al.*, 1996). Interestingly, mutants in genes encoding different subunits of this chaperone (Stoldt *et al.*, 1996), e.g. *CCT2*, *CCT3* and *TCP1*, or in all the components of its cochaperone, the GimC complex (Siegers *et al.*, 1999), showed clear cold-sensitive phenotypes (Chen *et al.*, 1994; Geissler *et al.*, 1998). This suggests a Tor2p-mediated activation of protein complexes involved in maintaining the cytoskeleton under cold stress. Moreover, the abnormal actin polarization displayed by *tor2* mutants was reverted by enhanced expression of *PKC1*, *ROM2* and other effectors of the PKC pathway (Helliwell *et al.*, 1998). Hence, it seems clear that maintenance of an adequate cell polarization upon cold-shock requires interplay between TOR and PKC pathways.

However, we cannot rule out the possibility of an indirect effect of low temperatures in the activity of the TOR pathway. Changes in the physical state of the plasma membrane due to abrupt shifts in temperature could significantly alter the activity of membrane-associated enzymes and transporters, lowering nutrient uptake, and leading to TORC1 complex inactivation. For example, stress conditions, such as heat-shock, have been reported to cause a severe decrease in amino acid import (Schmidt *et al.*, 1998; Chung *et al.*, 2001). This decrease is a consequence of the vacuolar degradation of high-affinity amino acid permeases such as Tat2p or Fur4p (Skrzypek *et al.*, 1998; Bultynck *et al.*, 2006), an event that is also stimulated by a downshift in environmental temperature (Abe & Horikoshi, 2000).

Role of the HOG pathway

As mentioned above, the histidine kinase Sln1p induces signalling through the HOG pathway in response to a downshift in temperature (Hayashi & Maeda, 2006; Panadero *et al.*, 2006). The HOG pathway, one of the five MAP

kinase pathways discovered in *S. cerevisiae* (Gustin *et al.*, 1998; Schwartz & Madhani, 2004), has traditionally been thought to be involved in the genetic response to hyper-osmotic stress alone (Hohmann, 2002; Westfall *et al.*, 2004). However, recent findings have established novel roles for this pathway, such as adaptation to citric acid stress (Lawrence *et al.*, 2004), heat-shock (Winkler *et al.*, 2002), methylglyoxal resistance (Aguilera *et al.*, 2005), distribution of proteins within the Golgi complex (Reynolds *et al.*, 1998) or cell-wall maintenance (García-Rodríguez *et al.*, 2000). Recently, results obtained by our group (Panadero *et al.*, 2006) have revealed that downstream elements of the HOG pathway are also involved in cold-signal transduction. The transcription factor Hog1p, which is the last step of the MAPK cascade, is phosphorylated after a downshift in temperature, a fact that is prevented in mutants lacking upstream elements of the pathway. Signalling through HOG causes the activation of *GPD1*, *GLO1* and other Hog1p-dependent osmotically activated genes in response to cold-shock (Panadero *et al.*, 2006) (Fig. 4). These results are in good agreement with previously reported data from other organisms. The *Arabidopsis* MKK2 pathway mediates signal transduction upon both saline and




Fig. 4. The HOG pathway. Sensitivity of the Sln1p/Ypd1p/Ssk1p histidine kinase complex to membrane rigidification caused by low temperature generates a signal that is transduced through the MAPK phosphorylation cascade. Hog1p is phosphorylated and migrates into the nucleus, promoting the transcription of genes for glycerol synthesis, among others (see text for details).

cold stress (Teige *et al.*, 2004). In the fission yeast *Schizosaccharomyces pombe*, the homologue pathway to HOG (named SAPK) is also involved in the cold-shock response (Soto *et al.*, 2002). Similarly, activation by hypothermic stress of mammalian MAPK family members, including JNK (c-Jun N-terminal kinase) and p38, the functional homologues to Sty1p and Hog1p (de Nadal *et al.*, 2002), has been reported in human cells (Gon *et al.*, 1998; Ohsaka *et al.*, 2002; Roberts *et al.*, 2002). Both JNK and p38 are also activated during natural freezing and thawing of the wood frog (Greenway & Storey, 2000). This result strongly suggests that cold-stress responses by MAPKs, particularly p38 and relatives, may be adaptive in dealing with freeze stress, as recently reported (Panadero *et al.*, 2006). In fact, a yeast strain lacking *HOG1* showed impaired viability after freezing and frozen storage as compared with the wild-type (Panadero *et al.*, 2006).

Concluding remarks

Although *S. cerevisiae* cold-sensitive mutants were first described over 10 years ago, only very recent studies have started to reveal the existence of cold-shock and freezing-protective responses in this organism, giving some clues to their physiological significance. Many genes with up- or down-regulated expression at low temperatures are known to participate in cellular adaptation to other kinds of stressful situations, such as thermal stress, high osmolarity, oxidation or presence of certain poisons. This phenomenon is very well known in yeast and has been called the general stress response. Hence, it is not surprising that exposure to low or subfreezing temperatures are also included in the scope of this systemic mechanism. Similarly, known sensing molecules, signal transduction machinery and transcription factors that mediate the genetic response to other stressful conditions, appear to be involved in cold response. However, cold activation of several signalling pathways might be indirect. In this way, the loss in membrane fluidity, as a physical consequence of exposure to low temperatures, would diminish the protein mobility of integral membrane proteins, such as amino acid transporters, and be interpreted by the cells as a situation of nutrient starvation.

Diversification of the functional roles of transcription pathways, which were initially believed to respond exclusively to one kind of stress, is an idea supported by an increasing number of new studies, especially in the case of the HOG pathway. Moreover, links between hyperosmotic stress, ratio of unsaturated to saturated fatty acids, and membrane rigidification lead us to speculate about a scenario in which the level of membrane fluidity is the physical perturbation sensed by the cell that triggers the response to different kinds of stresses. However, not all of the cold-induced response is referred to multistress-related genes. This is the case for the *TIP/TIR* genes, whose function and

activation mechanism have yet to be clarified. Hence, the existence of an exclusive mechanism for the cold response in yeast cannot be discarded. Given the increasing interest shown by several groups in this field, we can expect that related works in the near future will shed light on the currently speculative aspects of this response.

Acknowledgements

We thank in particular all members of the IATA Baker's Yeast Laboratory for their contributions towards several aspects of this work. Our research described in this review was supported by the Comisión Interministerial de Ciencia y Tecnología (projects AGL2000-0524, AGL2001-1203 and AGL2004-462). J.A. is supported by an I3P grant from the Spanish Ministry of Education and Science. The English text was revised by F. Barraclough.

References

- Abe F & Horikoshi K (2000) Tryptophan permease gene *TAT2* confers high-pressure growth in *Saccharomyces cerevisiae*. *Mol Cell Biol* **20**: 8093–8102.
- Abramova N, Cohen BD, Sertil O, Kapoor R, Davis KJA & Lowry CV (2001a) Regulatory mechanisms controlling expression of the *DAN/TIR* mannoprotein genes during anaerobic remodelling of the cell wall in *Saccharomyces cerevisiae*. *Genetics* **157**: 1169–1177.
- Abramova N, Sertil O, Mehta S & Lowry CV (2001b) Reciprocal regulation of anaerobic and aerobic cell wall mannoprotein gene expression in *Saccharomyces cerevisiae*. *J Bacteriol* **183**: 2881–2887.
- Aguilar PS, Hernandez-Arriaga AM, Cybulski LE, Erazo AC & de Mendoza D (2001) Molecular basis of thermosensing: a two-component signal transduction thermometer in *Bacillus subtilis*. *EMBO J* **20**: 1681–1691.
- Aguilera J, Rodríguez-Vargas S & Prieto JA (2005) The HOG MAP kinase pathway is required for the induction of methylglyoxal-responsive genes and determines methylglyoxal resistance in *Saccharomyces cerevisiae*. *Mol Microbiol* **56**: 228–239.
- Albertyn J, Hohmann S, Thevelein JM & Prior BA (1994) *GPD1*, which encodes glycerol-3-phosphate dehydrogenase, is essential for growth under osmotic stress in *Saccharomyces cerevisiae*, and its expression is regulated by the high-osmolarity glycerol response pathway. *Mol Cell Biol* **14**: 4135–4144.
- Ariño J, Pérez-Callejón E, Cunillera N, Camps M, Posas F & Ferrer A (1993) Protein phosphatases in higher plants: multiplicity of type 2A phosphatases in *Arabidopsis thaliana*. *Plant Mol Biol* **21**: 475–485.
- Auld KL, Brown CR, Casolari JM, Komili S & Silver PA (2006) Genomic association of the proteasome demonstrates

- overlapping gene regulatory activity with transcription factor substrates. *Mol Cell* **21**: 861–871.
- Beeser AE & Cooper TG (1999) The dual-specificity protein phosphatase Yvh1p acts upstream of the protein kinase Mck1p in promoting spore development in *Saccharomyces cerevisiae*. *J Bacteriol* **181**: 5219–5224.
- Beeser AE & Cooper TG (2000) The dual-specificity protein phosphatase Yvh1p regulates sporulation, growth, and glycogen accumulation independently of catalytic activity in *Saccharomyces cerevisiae* via the cyclic AMP-dependent protein kinase cascade. *J Bacteriol* **182**: 3517–3528.
- Blomberg A (2000) Metabolic surprises in *Saccharomyces cerevisiae* during adaptation to saline conditions: questions, some answers and a model. *FEMS Microbiol Lett* **182**: 1–8.
- Bultynck G, Heath VL, Majeed AP, Galan JM, Haguenaer-Tsapir R & Cyert MS (2006) Slm1 and slm2 are novel substrates of the calcineurin phosphatase required for heat stress-induced endocytosis of the yeast uracil permease. *Mol Cell Biol* **26**: 4729–4745.
- Carratù L, Franceschelli S, Pardini CL, Kobayashi GS, Horvath I, Vigh L & Maresca B (1996) Membrane lipid perturbation modifies the set point of the temperature of heat shock response in yeast. *Proc Natl Acad Sci USA* **93**: 3870–3875.
- Chen X, Sullivan DS & Huffaker TC (1994) Two yeast genes with similarity to TCP-1 are required for microtubule and actin function in vivo. *Proc Natl Acad Sci USA* **91**: 9111–9115.
- Choi J-Y, Stukay J, Hwang S-Y & Martin CE (1996) Regulatory elements that control transcription activation and unsaturated fatty acid-mediated repression of the *Saccharomyces cerevisiae* *OLE1* gene. *J Biol Chem* **271**: 3581–3589.
- Chung N, Mao C, Heitman J, Hannun YA & Obeid LM (2001) Phytosphingosine as a specific inhibitor of growth and nutrient import in *Saccharomyces cerevisiae*. *J Biol Chem* **276**: 35614–35621.
- Cohen BD, Sertil O, Abramova NE, Davies KJ & Lowry CV (2001) Induction and repression of *DAN1* and the family of anaerobic mannoprotein genes in *Saccharomyces cerevisiae* occurs through a complex array of regulatory sites. *Nucleic Acids Res* **29**: 799–808.
- Craig EA & Jacobsen K (1985) Mutations in cognate genes of *Saccharomyces cerevisiae* hsp70 result in reduced growth rates at low temperatures. *Mol Cell Biol* **5**: 3517–3524.
- Davenport KR, Sohaskey M, Kamada Y, Levin DE & Gustin MC (1995) A second osmosensing signal transduction pathway in yeast. Hypotonic shock activates the PKC1 protein kinase-regulated cell integrity pathway. *J Biol Chem* **270**: 30157–30161.
- de Jesus Ferreira MC, Bao X, Laizé V & Hohmann S (2001) Transposon mutagenesis reveals novel loci affecting tolerance to salt stress and growth at low temperature. *Curr Genet* **40**: 27–39.
- de Nadal E, Alepuz PM & Posas F (2002) Dealing with osmotic stress through MAP kinase activation. *EMBO Rep* **3**: 735–740.
- di Como CJ & Arndt KT (1996) Nutrients, via the Tor proteins, stimulate the association of Tap42 with type 2A phosphatases. *Genes Dev* **10**: 1904–1916.
- Du X & Takagi H (2005) N-Acetyltransferase Mpr1 confers freeze tolerance on *Saccharomyces cerevisiae* by reducing reactive oxygen species. *J Biochem* **138**: 391–397.
- Düvel K, Santhanam A, Garrett S, Schnepfer L & Broach JR (2003) Multiple roles of Tap42 in mediating rapamycin-induced transcriptional changes in yeast. *Mol Cell* **11**: 1467–1478.
- Elbein AD, Pan YT, Pastuszak I & Carroll D (2003) New insights on trehalose: a multifunctional molecule. *Glycobiology* **13**: 17R–27R.
- Estruch F (2000) Stress-controlled transcription factors, stress-induced genes and stress tolerance in budding yeast. *FEMS Microbiol Rev* **24**: 469–486.
- Fauman EB & Saper MA (1996) Structure and function of protein tyrosine phosphatases. *Trends Biochem Sci* **21**: 413–417.
- Gancedo C & Flores CL (2004) The importance of a functional trehalose biosynthetic pathway for the life of yeasts and fungi. *FEMS Yeast Res* **4**: 351–359.
- García-Rodríguez LJ, Durán A & Roncero C (2000) Calcofluor antifungal action depends on chitin and a functional high-osmolarity glycerol response (HOG) pathway: evidence for a physiological role of the *Saccharomyces cerevisiae* HOG pathway under noninducing conditions. *J Bacteriol* **182**: 2428–2437.
- Gast K, Damaschun G, Damaschun H, Misselwitz R & Zirwer D (1993) Cold denaturation of yeast phosphoglycerate kinase: kinetics of changes in secondary structure and compactness on unfolding and refolding. *Biochemistry* **32**: 7747–7752.
- Geissler S, Siegers K & Schiebel E (1998) A novel protein complex promoting formation of functional α - and γ -tubulin. *EMBO J* **17**: 952–966.
- Gon Y, Hashimoto S, Matsumoto K, Nakayama T, Takeshita I & Horie T (1998) Cooling and rewarming-induced IL-8 expression in human bronchial epithelial cells through p38 MAP kinase-dependent pathway. *Biochem Biophys Res Commun* **249**: 156–160.
- Greenway SC & Storey KB (2000) Activation of mitogen-activated protein kinases during natural freezing and thawing in the wood frog. *Mol Cell Biochem* **209**: 29–37.
- Griffioen G & Thevelein JM (2002) Molecular mechanisms controlling the localisation of protein kinase A. *Curr Genet* **41**: 199–207.
- Griffioen G, Swinnen S & Thevelein JM (2003) Feedback inhibition on cell wall integrity signaling by Zds1 involves Gsk3 phosphorylation of a cAMP-dependent protein kinase regulatory subunit. *J Biol Chem* **278**: 23460–23471.
- Gualerzi CO, Giuliodori AM & Pon CL (2003) Transcriptional and post-transcriptional control of cold-shock genes. *J Mol Biol* **331**: 527–539.
- Gustin MC, Albertyn J, Alexander M & Davenport K (1998) MAP kinase pathways in the yeast *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* **62**: 1264–1300.

- Hampsey M (1997) A review of phenotypes in *Saccharomyces cerevisiae*. *Yeast* **13**: 1099–1133.
- Harris DM, Myrick TL & Rundle SJ (1999) The *Arabidopsis* homolog of yeast *TAP42* and mammalian $\alpha 4$ binds to the catalytic subunit of protein phosphatase 2A and is induced by chilling. *Plant Physiol* **121**: 609–617.
- Hayashi M & Maeda T (2006) Activation of the HOG pathway upon cold stress in *Saccharomyces cerevisiae*. *J Biochem (Tokyo)* **139**: 797–803.
- Healy AM, Zolnierowick S, Stapleton AE, Goebel M, DePaoli-Roach AA & Pringle JR (1991) *CDC55*, a *Saccharomyces cerevisiae* gene involved in cellular morphogenesis: identification, characterization, and homology to the β subunit of mammalian type 2A protein phosphatase. *Mol Cell Biol* **11**: 5767–5780.
- Heinisch JJ, Lorberg A, Schmitz HP & Jacoby JJ (1999) The protein kinase C-mediated MAP kinase pathway involved in the maintenance of cellular integrity in *Saccharomyces cerevisiae*. *Mol Microbiol* **32**: 671–680.
- Helliwell SB, Howald I, Barbet N & Hall MN (1998) TOR2 is part of two related signaling pathways coordinating cell growth in *Saccharomyces cerevisiae*. *Genetics* **148**: 99–112.
- Hirata Y, Andoh T, Asahara T & Kikuchi A (2003) Yeast glycogen synthase kinase-3 activates Msn2p-dependent transcription of stress responsive genes. *Mol Biol Cell* **14**: 302–312.
- Hohmann S (2002) Osmotic stress signalling and osmoadaptation in yeasts. *Microbiol Mol Biol Rev* **66**: 300–372.
- Homma T, Iwahashi H & Komatsu Y (2003) Yeast gene expression during growth at low temperature. *Cryobiology* **46**: 230–237.
- Hoppe T, Matuschewski K, Rape M, Schlenker S, Ulrich HD & Jentsch S (2000) Activation of a membrane-bound transcription factor by regulated ubiquitin/proteasome-dependent processing. *Cell* **102**: 577–586.
- Hoppe T, Rape M & Jentsch S (2001) Membrane-bound transcription factors: regulated release by RIP or RUP. *Curr Opin Cell Biol* **13**: 344–348.
- Imai J, Toh-e A & Matsui Y (1996) Genetic analysis of the *Saccharomyces cerevisiae* *RHO3* gene, encoding a Rho-type small GTPase, provides evidence for a role in bud formation. *Genetics* **142**: 359–369.
- Inaba M, Suzuki I, Szalontai B, Kanesaki Y, Los DA, Hayashi H & Murata N (2003) Gene-engineered rigidification of membrane lipids enhances the cold inducibility of gene expression in *Synechocystis*. *J Biol Chem* **278**: 12191–12198.
- Inouye M (1999) Cold-shock response and adaptation. *J Mol Microbiol Biotechnol* **1**: 191.
- Jiang Y & Broach JR (1999) Tor proteins and protein phosphatase 2A reciprocally regulate Tap42 in controlling cell growth in yeast. *EMBO J* **18**: 2782–2792.
- Jiang Y, Vasconcelles MJ, Wretzel S, Light A, Martin CE & Goldberg MA (2001) *MGA2* is involved in the low-oxygen response element-dependent hypoxic induction of genes in *Saccharomyces cerevisiae*. *Mol Cell Biol* **21**: 6161–6169.
- Jiang Y, Vasconcelles MJ, Wretzel S, Light A, Gilooly L, McDaid K, Oh C-S, Martin CE & Goldberg MA (2002) Mga2p processing by hypoxia and unsaturated fatty acids in *Saccharomyces cerevisiae*: impact on LORE-dependent gene expression. *Eukaryot Cell* **1**: 481–490.
- Jones PG & Inouye M (1996) RbfA, a 30S ribosomal binding factor, is a cold-shock protein whose absence triggers the cold-shock response. *Mol Microbiol* **21**: 1207–1218.
- Kajiwarra S, Aritomi T, Suga K, Ohtaguchi K & Kobayashi O (2000) Overexpression of the *OLE1* gene enhances ethanol fermentation by *Saccharomyces cerevisiae*. *Appl Microbiol Biotechnol* **53**: 568–574.
- Kamada Y, Jung US, Piotrowski J & Levin DE (1995) The protein kinase C-activated MAP kinase pathway of *Saccharomyces cerevisiae* mediates a novel aspect of the heat shock response. *Genes Dev* **9**: 1559–1571.
- Kandror O, Bretschneider N, Kreydin E, Cavalieri D & Goldberg AL (2004) Yeast adapt to near-freezing temperatures by STRE/Msn2,4-dependent induction of trehalose synthesis and certain molecular chaperones. *Mol Cell* **13**: 771–781.
- Kim J, Alizadeh P, Harding T, Hefner-Gravink A & Klionsky DJ (1996) Disruption of the yeast *ATH1* gene confers better survival after dehydration, freezing and ethanol shock: potential commercial applications. *Appl Environ Microbiol* **62**: 1563–1569.
- Laroche C, Beney L, Marechal PA & Gervais P (2001) The effect of osmotic pressure on the membrane fluidity of *Saccharomyces cerevisiae* at different physiological temperatures. *Appl Microbiol Biotechnol* **56**: 249–254.
- Lawrence CL, Botting CH, Antrobus R & Coote PJ (2004) Evidence of a new role for the high-osmolarity glycerol mitogen-activated protein kinase pathway in yeast: regulating adaptation to citric acid stress. *Mol Cell Biol* **24**: 3307–3323.
- Lee SJ & Baserga SJ (1997) Functional separation of pre-rRNA processing steps revealed by truncation of the U3 small nucleolar ribonucleoprotein component, Mpp10. *Proc Natl Acad Sci USA* **94**: 13536–13541.
- Lehman K, Rossi G, Adamo JE & Brennwald P (1999) Yeast homologues of tomosyn and lethal giant larvae function in exocytosis and are associated with the plasma membrane SNARE, Sec9. *J Cell Biol* **146**: 125–140.
- Levin DE (2005) Cell wall integrity signaling in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* **69**: 262–291.
- Loertscher J, Larson LL, Matson CK, Parrish ML, Felthausen A, Sturm A, Tachibana C, Bard M & Wright R (2006) Endoplasmic reticulum-associated degradation is required for cold adaptation and regulation of sterol biosynthesis in the yeast *Saccharomyces cerevisiae*. *Eukaryot Cell* **5**: 712–722.
- Loewith R, Jacinto E, Wulfschleger S, Lorberg A, Crespo JL, Bonenfant D, Oppliger W, Jenoe P & Hall MN (2002) Two TOR complexes, only one of which is rapamycin sensitive, have distinct roles in cell growth control. *Mol Cell* **10**: 457–468.
- López CS, Heras H, Garda H, Ruzal S, Sánchez-Rivas C & Rivas E (2000) Biochemical and biophysical studies of *Bacillus subtilis*

- envelopes under hyperosmotic stress. *Int J Food Microbiol* **55**: 137–142.
- Los DA & Murata N (2004) Membrane fluidity and its roles in the perception of environmental signals. *Biochim Biophys Acta* **1666**: 142–157.
- Manning BD, Padhmanabha R & Snyder M (1997) The Rho-GFP Rom2p localizes to sites of polarized cell growth and participates in cytoskeletal functions in *Saccharomyces cerevisiae*. *Mol Biol Cell* **8**: 1829–1844.
- Mansilla MC & de Mendoza D (2005) The *Bacillus subtilis* desaturase: a model to understand phospholipid modification and temperature sensing. *Arch Microbiol* **183**: 229–235.
- Martínez-Pastor MT, Marchler G, Schüller C, Marchler-Bauer A, Ruis H & Estruch F (1996) The *Saccharomyces cerevisiae* zinc finger proteins Msn2p and Msn4p are required for transcriptional induction through the stress response element (STRE). *EMBO J* **15**: 2227–2235.
- McKemy DD, Neuhauser WM & Julius D (2002) Identification of a cold receptor reveals a general role for TRP channels in thermosensation. *Nature* **416**: 52–58.
- Mikami K, Kanesaki Y, Suzuki I & Murata N (2002) The histidine kinase Hik33 perceives osmotic stress and cold stress in *Synechocystis* sp. PCC 6803. *Mol Microbiol* **46**: 905–915.
- Monroy AF, Sarhan F & Dhindsa RS (1993) Cold-induced changes in freezing tolerance, protein phosphorylation, and gene expression. Evidence for a role of calcium. *Plant Physiol* **102**: 1227–1235.
- Monroy AF, Sangwan V & Dhindsa RS (1998) Low temperature signal transduction during cold acclimation: protein phosphatase 2A as an early target for cold-inactivation. *Plant J* **13**: 653–660.
- Muda M, Manning ER, Orth K & Dixon JE (1999) Identification of the human YVH1 protein-tyrosine phosphatase orthologue reveals a novel zinc binding domain essential for in vivo function. *J Biol Chem* **274**: 23991–23995.
- Murata Y, Homma T, Kitagawa E *et al.* (2006) Genome-wide expression analysis of yeast response during exposure to 4 °C. *Extremophiles* **10**: 117–128.
- Nakagawa Y, Sakumoto N, Kaneko Y & Harashima S (2002) Mga2p is a putative sensor for low temperature and oxygen to induce *OLE1* transcription in *Saccharomyces cerevisiae*. *Biochem Biophys Res Commun* **291**: 707–713.
- Nakagawa Y, Ueda A, Kaneko Y & Harashima S (2003) Merging of multiple signals regulating $\Delta 9$ fatty acid desaturase gene transcription in *Saccharomyces cerevisiae*. *Mol Gen Genomics* **269**: 370–380.
- Noguchi E, Kayashi N, Nakashima N & Nishimoto T (1997) Yrb2p, a Nup2p-related yeast protein, has a functional overlap with Rna1p, a yeast Ran-GTPase-activating protein. *Mol Cell Biol* **17**: 2235–2246.
- Noguchi E, Sayito Y, Saber S & Nishimoto T (1999) Disruption of the *YRB2* gene retards nuclear protein export, causing a profound mitotic delay, and can be rescued by overexpression of *XPO1/CRM1*. *J Biochem* **125**: 574–585.
- Odani M, Komatsu Y, Oka S & Iwahashi H (2003) Screening of genes that respond to cryopreservation stress using yeast DNA microarray. *Cryobiology* **47**: 155–164.
- Ohsaka Y, Ohgiya S, Hoshino T & Ishizaki K (2002) Phosphorylation of c-Jun N-terminal kinase in human hepatoblastoma cells is transiently increased by cold exposure and further enhanced by subsequent warm incubation of the cells. *Cell Physiol Biochem* **12**: 111–118.
- Örvar BL, Sangwan V, Omann F & Dhindsa RS (2000) Early steps in cold sensing by plant cells: the role of actin cytoskeleton and membrane fluidity. *Plant J* **23**: 785–794.
- Panadero J, Pallotti C, Rodríguez-Vargas S, Randez-Gil F & Prieto JA (2006) A downshift in temperature activates the high osmolarity glycerol (HOG) pathway, which determines freeze tolerance in *Saccharomyces cerevisiae*. *J Biol Chem* **281**: 4638–4645.
- Papp E, Nardai G, Soti C & Csermely P (2003) Molecular chaperones, stress proteins and redox homeostasis. *Biofactors* **17**: 249–257.
- Park JI, Grant CM, Attfield PV & Dawes IW (1997) The freeze-thaw stress response of the yeast *Saccharomyces cerevisiae* is growth phase specific and is controlled by nutritional state via the RAS-cyclic AMP signal transduction pathway. *Appl Environ Microbiol* **63**: 3818–3824.
- Park JI, Collinson EJ, Grant CM & Dawes IW (2005a) Rom2p, the Rho1 GTP/GDP exchange factor of *Saccharomyces cerevisiae*, can mediate stress responses via the Ras-cAMP pathway. *J Biol Chem* **280**: 2529–2535.
- Park JI, Grant CM & Dawes IW (2005b) The high-affinity cAMP phosphodiesterase of *Saccharomyces cerevisiae* is the major determinant of cAMP levels in stationary phase: involvement of different branches of the Ras-cyclic AMP pathway in stress responses. *Biochem Biophys Res Commun* **327**: 311–319.
- Parrou JL, Teste MA & François J (1997) Effects of various types of stress on the metabolism of reserve carbohydrates in *Saccharomyces cerevisiae*: genetic evidence for a stress-induced recycling of glycogen and trehalose. *Microbiology* **143**: 1891–1900.
- Peier AM, Moqrich A, Hergarden AC *et al.* (2002) A TRP channel that senses cold stimuli and menthol. *Cell* **108**: 705–715.
- Platt A & Reece RJ (1998) The yeast galactose genetic switch is mediated by the formation of a Gal4p-Gal80p-Gal3p complex. *EMBO J* **17**: 4086–4091.
- Randez-Gil F, Aguilera J, Codón A, Rincón AM, Estruch F & Prieto JA (2003) Baker's yeast: challenges and future aspects. *Functional Genetics of Industrial Yeasts* (de Winder JH, ed), pp. 57–97. Springer-Verlag, Heidelberg.
- Revardel E & Aigle M (1993) The *NUM1* yeast gene: length polymorphism and physiological aspects of mutant phenotype. *Yeast* **9**: 495–506.
- Reynolds TB, Hopkins BD, Lyons MR & Graham TR (1998) The high osmolarity glycerol response (HOG) MAP kinase pathway controls localization of a yeast golgi glycosyltransferase. *J Cell Biol* **143**: 935–946.

- Roberts JR, Rowe PA & Demaine AG (2002) Activation of NF- κ B and MAP kinase cascades by hypothermic stress in endothelial cells. *Cryobiology* **44**: 161–169.
- Rodríguez-Vargas S, Sánchez-García A, Martínez-Rivas JM, Prieto JA & Randez-Gil F (2007) Fluidization of membrane lipids enhances the tolerance of *Saccharomyces cerevisiae* to freezing and salt stress. *Appl Environ Microbiol* **73**: 110–116.
- Rohde JR, Campbell S, Zurita-Martínez SA, Cutler NS, Ashe M & Cardenas ME (2004) TOR controls transcriptional and translational programs via Sap-Sit4 protein phosphatase signaling effectors. *Mol Cell Biol* **24**: 8332–8341.
- Sahara T, Goda T & Ohgiya S (2002) Comprehensive expression analysis of time-dependent genetic responses in yeast cells to low temperature. *J Biol Chem* **277**: 50015–50021.
- Sakumoto N, Mukai Y, Uchida K *et al.* (1999) A series of protein phosphatase gene disruptants in *Saccharomyces cerevisiae*. *Yeast* **15**: 1669–1679.
- Sangwan V, Örvar BL, Beyerly J, Hirt H & Dhindsa RS (2002) Opposite changes in membrane fluidity mimic cold and heat stress activation of distinct plant MAP kinase pathways. *Plant J* **31**: 629–638.
- Santangelo GM (2006) Glucose signaling in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* **70**: 253–282.
- Santhanam A, Hartley A, Düvel K, Broach JR & Garrett S (2004) PP2A phosphatase activity is required for stress and Tor kinase regulation of yeast stress response factor Msn2p. *Eukaryot Cell* **3**: 1261–1271.
- Schade B, Jansen G, Whiteway M, Entian KD & Thomas DY (2004) Cold adaptation in budding yeast. *Mol Biol Cell* **15**: 5492–5502.
- Schmelzle T & Hall MN (2000) TOR, a central controller of cell growth. *Cell* **103**: 253–262.
- Schmelzle T, Beck T, Martin DE & Hall MN (2004) Activation of the RAS/cyclic AMP pathway suppresses a TOR deficiency in yeast. *Mol Cell Biol* **24**: 338–351.
- Schmidt A, Kunz J & Hall MN (1996) TOR2 is required for organization of the actin cytoskeleton in yeast. *Proc Natl Acad Sci USA* **93**: 13780–13785.
- Schmidt A, Beck T, Koller A, Kunz J & Hall MN (1998) The TOR nutrient signalling pathway phosphorylates NPR1 and inhibits turnover of the tryptophan permease. *EMBO J* **17**: 6924–6931.
- Schwartz MA & Madhani HD (2004) Principles of MAP kinase signaling specificity in *Saccharomyces cerevisiae*. *Annu Rev Genet* **38**: 725–748.
- Seki M, Ishida J, Narusaka M *et al.* (2002a) Abstract monitoring the expression pattern of around 7,000 *Arabidopsis* genes under ABA treatments using a full-length cDNA microarray. *Funct Integr Genomics* **2**: 282–291.
- Seki M, Narusaka M, Ishida J *et al.* (2002b) Monitoring the expression profiles of 7000 *Arabidopsis* genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant J* **31**: 279–292.
- Seki M, Satou M, Sakurai T *et al.* (2004) RIKEN *Arabidopsis* full-length (RAFL) cDNA and its applications for expression profiling under abiotic stress conditions. *J Exp Bot* **55**: 213–223.
- Shinozaki K & Yamaguchi-Shinozaki K (1996) Molecular responses to drought and cold stress. *Curr Opin Biotechnol* **7**: 161–167.
- Siegers K, Waldmann T, Leroux MR, Grein K, Shevchenko A, Schiebel E & Hartl FU (1999) Compartmentation of protein folding in vivo: sequestration of non-native polypeptide by the chaperonin-GimC system. *EMBO J* **8**: 75–84.
- Singer MA & Lindquist S (1998) Thermotolerance in *Saccharomyces cerevisiae*: the Yin and Yang of trehalose. *Trends Biotechnol* **16**: 460–468.
- Skrzypek MS, Nagiec MM, Lester RL & Dickson RC (1998) Inhibition of amino acid transport by sphingoid long chain bases in *Saccharomyces cerevisiae*. *J Biol Chem* **273**: 2829–2834.
- Soto T, Beltrán FF, Paredes V, Madrid M, Millar JBA, Vicente-Soler J, Cansado J & Gacto M (2002) Cold induces stress-activated protein kinase-mediated response in the fission yeast *Schizosaccharomyces pombe*. *Eur J Biochem* **269**: 5056–5065.
- Stoldt V, Rademacher F, Kehren V, Ernst JF, Pearce DA & Sherman F (1996) Review: the Cct eukaryotic chaperonin subunits of *Saccharomyces cerevisiae* and other yeasts. *Yeast* **12**: 523–529.
- Stukey JE, McDonough VM & Martin CE (1989) Isolation and characterization of *OLE1*, a gene affecting fatty acid desaturation from *Saccharomyces cerevisiae*. *J Biol Chem* **264**: 16537–16544.
- Stukey JE, McDonough VM & Martin CE (1990) The *OLE1* gene of *Saccharomyces cerevisiae* encodes the $\Delta 9$ fatty acid desaturase and can be functionally replaced by the rat stearyl-CoA desaturase gene. *J Biol Chem* **265**: 20144–20149.
- Suzuki I, Los DA, Kanesaki Y, Mikami K & Murata N (2000) The pathway for perception and transduction of low-temperature signals in *Synechocystis*. *EMBO J* **19**: 1327–1334.
- Suzuki I, Kanesaki Y, Mikami K, Kanehisa M & Murata N (2001) Cold-regulated genes under control of the cold sensor Hik33 in *Synechocystis*. *Mol Microbiol* **40**: 235–244.
- Teige M, Scheikl E, Eulgem T, Doczi R, Ichimura K, Shinozaki K, Dangl JL & Hirt H (2004) The MKK2 pathway mediates cold and salt stress signaling in *Arabidopsis*. *Mol Cell* **15**: 141–152.
- Thevelein JM & de Winder JH (1999) Novel sensing mechanisms and targets for the cAMP-protein kinase A pathway in the yeast *Saccharomyces cerevisiae*. *Mol Microbiol* **33**: 904–918.
- Thieringer HA, Jones PG & Inouye M (1998) Cold shock and adaptation. *BioEssays* **20**: 49–57.
- Tokishita S & Mizuno T (1994) Transmembrane signal transduction by the *Escherichia coli* osmotic sensor, EnvZ: intermolecular complementation of transmembrane signalling. *Mol Microbiol* **13**: 435–444.
- van der Heide T & Poolman B (2000) Osmoregulated ABC-transport system of *Lactococcus lactis* senses water stress via changes in the physical state of the membrane. *Proc Natl Acad Sci USA* **97**: 7102–7106.
- van der Heide T, Stuart MC & Poolman B (2001) On the osmotic signal and osmosensing mechanism of an ABC transport system for glycine betaine. *EMBO J* **20**: 7022–7032.

- van Dijck P, Ma P, Versele M, Gorwa M-F, Colombo S, Lemaire K, Bossi D, Loïez A & Thevelein JM (2000) A baker's yeast mutant (*fil1*) with a specific, partially inactivating mutation in adenylate cyclase maintains a high stress resistance during active fermentation and growth. *J Mol Microbiol Biotechnol* **2**: 521–530.
- Vasconcelles MJ, Jiang Y, McDaid K, Gilooley L, Wretzel S, Porter DL, Martin CE & Goldberg MA (2001) Identification and characterization of a low oxygen response element involved in the hypoxic induction of a family of *Saccharomyces cerevisiae* genes. *J Biol Chem* **276**: 14374–14384.
- Vigh L, Maresca B & Harwood JL (1998) Does the membrane's physical state control the expression of heat shock and other genes? *Trends Biochem Sci* **23**: 369–374.
- Wang Y, Pierce M, Schnepel L, Guldal CG, Zhang X, Tavazoie S & Broach JR (2004) Ras and Gpa2 mediate one branch of a redundant glucose signaling pathway in yeast. *PLoS Biol* **2**: 610–622.
- Westfall PJ, Ballon DR & Thorner J (2004) When the stress of your environment makes you go HOG wild. *Science* **306**: 1511–1512.
- Wilson WA & Roach PJ (2002) Nutrient-regulated protein kinases in budding yeast. *Cell* **111**: 155–158.
- Winkler A, Arkind C, Mattison CP, Burkholder A, Knoche K & Ota I (2002) Heat stress activates the yeast high-osmolarity glycerol mitogen-activated protein kinase pathway, and protein tyrosine phosphatases are essential under heat stress. *Eukaryot Cell* **1**: 163–173.
- Winzler EA, Shoemaker DD, Astromoff A *et al.* (1999) Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* **285**: 901–906.
- Wolfe J & Bryant G (1999) Freezing, crying, and/or vitrification of membrane-solute-water systems. *Cryobiology* **39**: 103–129.
- Wullschlegel S, Loewith R & Hall MN (2006) TOR signalling in growth and metabolism. *Cell* **124**: 471–484.
- Young JC, Barral JM & Hartl FU (2003) More than folding: localized functions of cytosolic chaperones. *Trends Biochem Sci* **28**: 541–547.
- Zenke FT, Engles R, Vollenbroich V, Meyer J, Hollenberg CP & Breunig KD (1996) Activation of Gal4p by galactose-dependent interaction of galactokinase and Gal80p. *Science* **272**: 1662–1665.
- Zhang L, Ohta A, Horiuchi H, Takagi M & Imai R (2001) Multiple mechanisms regulate expression of low temperature responsive (*LOT*) genes in *Saccharomyces cerevisiae*. *Biochem Biophys Res Commun* **283**: 531–535.
- Zhang L, Onda K, Imai R, Fukuda R, Horiuchi H & Ohta A (2003) Growth temperature downshift induces antioxidant response in *Saccharomyces cerevisiae*. *Biochem Biophys Res Commun* **307**: 308–314.
- Zhang S, Skalsky Y & Garfinkel DJ (1999) *MGA2* or *SPT23* is required for transcription of the 9 fatty acid desaturase gene, *OLE1*, and nuclear membrane integrity in *Saccharomyces cerevisiae*. *Genetics* **151**: 473–483.
- Zhu H, Bilgin M, Bangham R *et al.* (2001) Global analysis of protein activities using proteome chips. *Science* **293**: 2101–2105.