

Math H185 Lecture 3

Keshav Balwant Deoskar

March 2, 2024

These are notes taken from lectures on Complex Analysis delivered by Professor Tony Feng for UC Berkeley's Math H185 class in the Spring 2024 semester.

Contents

1	January 22 - Holomorphic functions	2
1.1	Sequences and Series	2
1.2	Complex Differentiability	2
1.3	Stability Properties	3

1 January 22 - Holomorphic functions

1.1 Sequences and Series

Recall that a sequence of complex numbers $\{z_n \in \mathbb{C}\}$ is said to *converge* to $z \in \mathbb{C}$ if for all $\epsilon > 0$ there exists a natural number $N \geq 1$ such that

$$|z_n - z| < \epsilon$$

for all $n \geq N$.

Equivalently,

$$\lim_{n \rightarrow \infty} |z_n - z| = 0$$

In HW1, we show that if $z_n = x_n + iy_n$ and $z = x + iy$ where $x, y, x_n, y_n \in \mathbb{R}$ then

$$\lim_{n \rightarrow \infty} |z_n - z| = 0 \iff \begin{cases} \lim_{n \rightarrow \infty} |x_n - x| = 0 \\ \lim_{n \rightarrow \infty} |y_n - y| = 0 \end{cases}$$

1.2 Complex Differentiability

Let $f : U \subseteq_{\text{open}} \mathbb{C} \rightarrow \mathbb{C}$.

Holomorphic Functions

We say f is Holomorphic at $z_0 \in U$ if

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists in \mathbb{C} . If so, its value is denoted as $f'(z_0)$.

Note: Keep in mind that h is a complex number.

- This means that for any sequence $h_n \rightarrow 0$ or $\forall \epsilon > 0 \exists \delta > 0$ such that

$$|h| < \delta \implies \left| \frac{f(z_0 + h) - f(z_0)}{h} - f'(z_0) \right| < \epsilon$$

- This is the most important definition of the course.

Remark: Although \mathbb{C} is the same as \mathbb{R}^2 as a metric space, Holomorphicity is much stronger than differentiability of a function $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ because the limits along every path to a point are required to be equal. In contrast to this, differentiability in \mathbb{R}^2 requires that *a* limit exists along each path, but not that all limits are equal.

Example: Consider the function $f(z) = \bar{z}$.

We observe that

$$\begin{aligned}\frac{f(z+h) - f(z)}{h} &= \frac{\overline{z+h} - \bar{z}}{h} \\ &= \frac{\bar{h}}{h}\end{aligned}$$

Now, if we take the limit as $h \rightarrow 0$ along the real line, we get $\frac{\bar{h}}{h} = \frac{h}{h} = 1$, however if we take the limit along the imaginary line we get $\frac{\bar{h}}{h} = \frac{-h}{h} = -1$

On the other hand if we consider the counterpart of this function in \mathbb{R}^2 as $f(x, y) = (x, -y)$, this function is *smooth everywhere*. In contrast to this, the complex function $f(z) = \bar{z}$ is *not holomorphic at any* $z_0 \in \mathbb{C}$.

We will see that holomorphic functions have strong rigidity properties not shared by real differentiable functions. For instance,

- If f, g are holomorphic on a connected open set $U \subseteq \mathbb{C}$ and $f = g$ on a line segment in U , then in fact they agree at *all* points in U : $f(z) = g(z) \forall z \in U$. This is the **Principle of Analytic Continuation**.
- Another example of surprising rigidity is that if f is holomorphic on U i.e. it is once differentiable on U , then in fact it is *infinitely* differentiable on U .

Examples:

1. $f(z) = z^n$

Calculate:

$$\begin{aligned}\frac{f(z+h) - f(z)}{h} &= \frac{(z+h)^n - z^n}{h} \\ &= \text{binom thm.} \left[\frac{1}{h} (z^n + nz^{n-1}h + \dots + nh^{n-1} + h^n) - z^n \right] \\ &= nz^{n-1} + h(\dots)\end{aligned}$$

So, just like in Real Analysis, we have

$$\boxed{\lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h} = nz^{n-1}}$$

1.3 Stability Properties

While holomorphicity is different from real differentiability, there are a number of properties which are justified by the same $\epsilon - \delta$ proofs as those from \mathbb{R}^2 analysis.

- If $f, g : U \rightarrow \mathbb{C}$ are holomorphic at $z_0 \in U$ then

– $(f + g)$ is holomorphic at z_0 and

$$(f + g)'(z_0) = f'(z_0) + g'(z_0)$$

– fg is holomorphic at z_0 and

$$(fg)'(z_0) = f'(z_0)g(z_0) + f(z_0)g'(z_0)$$

- Chain Rule: $(f \circ g)$ is holomorphic at z_0 and

$$(f \circ g)'(z_0) = f'(g(z_0)) \cdot g'(z_0)$$

- Division: (f/g) is holomorphic at z_0 if $g(z_0) \neq 0$ and

$$\left(\frac{f}{g}\right)' = \frac{f'(z_0)g(z_0) + f(z_0)g'(z_0)}{g(z_0)^2}$$

Polynomials: Finite sum of monomials.

$$f(z) = a_n z^n + \cdots + a_0$$

By linearity (Stability property 1), *all* Polynomials are holomorphic on \mathbb{C} .

Rational Functions: Ratios of Polynomials.

$$h(z) = \frac{f(z)}{g(z)}$$

By Stability property 3, all rational functions are holomorphic on $\{z \in \mathbb{C} : g(z) \neq 0\} \subseteq_{\text{open}} \mathbb{C}$.

Warm-down examples: Where are the following functions holomorphic, and what are their derivatives in those regions?

1. $f(z) = \frac{1}{z}$
2. $f(z) = z^2 + 3z + \frac{1}{2}$
3. $f(z) = \operatorname{Re}(z)$
4. $f(z) = i \cdot \operatorname{Im}(z)$
5. $f(z) = \operatorname{Re}(z) + i \cdot \operatorname{Im}(z)$

Answers:

1. Holomorphic on $\mathbb{C} \setminus \{0\}$, and derivative in the region is

$$-\frac{1}{z^2}$$

2. Holomorphic on \mathbb{C} , and derivative in the region is

$$2z + 3$$

3. Not holomorphic *anywhere*, since limit vertically is always zero but limit horizontally will be non-zero.
4. Not holomorphic *anywhere*, since limit horizontally is always zero but limit vertically will be non-zero.
5. Holomorphic on \mathbb{C} , and derivative in the region is 1 ($f(z) = z$, so $f'(z) = 1$ at all points).