

# Math H185 Lecture64

Keshav Balwant Deoskar

January 31, 2024

These are notes taken from lectures on Complex Analysis delivered by Professor Tony Feng for UC Berkeley's Math H185 class in the Sprng 2024 semester.

## Contents

<b>1</b>	<b>January 28 - Integrating over curves</b>	<b>2</b>
1.1	What is a curve? . . . . .	2
1.2	So, how do we actually integrate? . . . . .	2
1.3	Most fundamental example: $f(z) = z^n, n \in \mathbb{Z}$ . . . . .	3

# 1 January 28 - Integrating over curves

In calculus, we learned how to integrate over intervals. Today, we'll learn how to integrate complex valued functions over curves in the complex plane.

## Integral over a curve

We define the integral of a function  $f(z)$  over a curve  $\gamma$

$$\int_{\gamma} f(z)dz := \int_{\gamma} \operatorname{Re}(f(z))dz + i \int_{\gamma} \operatorname{Im}(f(z))dz$$

### 1.1 What is a curve?

- A **parameterized curve** is a continuous function  $\gamma : \underbrace{[a, b]}_{\subset \mathbb{R}} \rightarrow \mathbb{C}$
- We say  $\gamma$  is **piece-wise smooth** if there exist finite subdivisions of  $[a, b]$  on which  $\gamma$  is smooth (in the Math 104 sense i.e. the real and imaginary parts are separately infinitely differentiable).  
[include graphs of piece-wise smooth paths]
- Example:  $\gamma(t) = z_0 + re^{i\theta}$  where  $z_0 \in \mathbb{C}$ ,  $r \in \mathbb{R}_{\geq 0}$ ,  $t \in [0, 2\pi]$ . This path traces out a circle of radius  $r$  centered around the point  $z_0$ .  
[include graph]
- Example: Given a path  $\gamma : [a, b] \rightarrow \mathbb{C}$ . Let  $\gamma^- : [a, b] \rightarrow \mathbb{C}$  be  $\gamma^-(t) = \gamma(a + b - t)$ . Then  $\gamma^-$  is the same curve but traversed in reverse orientation.

### 1.2 So, how do we actually integrate?

Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a "nice" curve, where "nice" means piece-wise smooth parameterized. Then,

**Theorem:** The integral of  $f(z)$  over  $\gamma$  is

$$\int_{\gamma} f(z)dz = \int_a^b f(\gamma(t))\gamma'(t)dt$$

and these integrals have some nice properties:

1. Linearity: Say  $\lambda, \mu \in \mathbb{C}$  Then

$$\int_{\gamma} \lambda f(z) + \mu g(z)dz = \lambda \int_{\gamma} f(z)dz + \mu \int_{\gamma} g(z)dz$$

2. orientation:

$$\int_{\gamma} f(z)dz = - \int_{\gamma^-} f(z)dz$$

#### Examples:

- (a)  $\gamma : [a, b] \rightarrow \mathbb{C}$  given by  $\gamma(t) = t$ :

$$\int_{\gamma} f(z)dz = \int_a^b f(t)dt$$

(b)  $\gamma : [a, b] \rightarrow \mathbb{C}$  given by  $\gamma(t) = it$ :

$$\int_{\gamma} f(z) dz = \int f(it) i dt = i \int f(it) dt$$

Recall that a **primitive** (i.e. antiderivative) of  $f$  is  $F$  such that  $F'(z) = f(z)$ . Then, the fundamental theorem of calculus is

**Fundamental Theorem of Calculus:** For a "nice" curve  $\gamma : [a, b] \rightarrow \mathbb{C}$  and function  $f$  with primitive  $F$ ,

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a))$$

**Corollary:** If  $\gamma$  is a **closed** curve i.e.  $\gamma(a) = \gamma(b)$  and  $f$  has a primitive on (an open neighborhood of)  $\gamma$ , then

$$\int_{\gamma} f(z) dz = 0$$

### 1.3 Most fundamental example: $f(z) = z^n, n \in \mathbb{Z}$

Consider the function  $f(z) = z^n, n \in \mathbb{Z}$  and the curve  $\gamma : [0, 2\pi] \rightarrow \mathbb{C}$  where  $\gamma(t) = re^{it}$ . What is  $\int_{\gamma} f(z) dz$ ?

The integral is

$$\begin{aligned} \int_{\gamma} f(z) dz &= \int_0^{2\pi} re^{int} (rie^{it}) dt \\ &= r^{n+1} i \int_0^{2\pi} e^{i(n+1)t} dt \end{aligned}$$

If  $n \neq -1$ :  $e^{i(n+1)t} dt$  has primitive

$$\frac{1}{i(n+1)} e^{i(n+1)t}$$

So, the integral is

$$\int_0^{2\pi} e^{i(n+1)t} dt = 0$$

If  $n = -1$ : Then, we have

$$\begin{aligned} \int_{\gamma} f(z) dz &= i \int_0^{2\pi} e^{i(n+1)t} dt \\ &= \int_0^{2\pi} 1 dt \\ &= 2\pi i \end{aligned}$$

Precisely why does the primitive not work for  $n = -1$ ? The issue lies with the fact that the primitive of  $\frac{1}{z}$  is the **logarithm**.

The complex logarithm isn't defined along a full circle around the origin. We'll revisit this when studying branch cuts soon.

#### Conclusion

We find that

$$\int_{\partial B_r(0)} z^n dz = \begin{cases} 0 & n \neq -1 \\ 2\pi i & n = -1 \end{cases}$$

**Note:** A very interesting observation is that this is *independent of  $r$* . This is surprising, and foreshadows some incredible results we'll see soon.