

Math H185 Lecture 3

Keshav Balwant Deoskar

January 24, 2024

These are notes taken from lectures on Complex Analysis delivered by Professor Tony Feng for UC Berkeley's Math H185 class in the Spring 2024 semester.

Contents

1	January 24 - Power Series	2
1.1	Convergence	2
1.2	Differentiation of Power Series	5

1 January 24 - Power Series

Power Series

A **Power Series** is a formal expression

$$\sum_{n \geq 0} a_n z^n, z_n \in \mathbb{C}$$

for which operations are defined as:

- Addition:

$$\sum_{n \geq 0} a_n z^n$$

- multiplication

- "Formal" here means we temporarily ignore whether it makes sense to plug in complex numbers into such formulae.

1.1 Convergence

Defining these formal expressions is cool, but when does a power series actually define a function? It does so when the series **converges**.

Example: Geometric series:

Let $a \in \mathbb{C}$, then for the geometric series we have $a_n = a^n$. So,

$$\sum_{n \geq 0} a^n z^n = 1 + az + a^2 z^2 + \dots$$

converges if $S_n = \sum_{n \geq 0}^{N-1} a^n z^n$ has a limit.

By the same argument as in the reals, we can get a closed form expression for S_N :

$$S_N = \frac{1 - (az)^N}{1 - (az)}$$

To deal with convergence, we break into cases and take the limit.

- $|az| < 1$ **case:**

$$|az| < 1 \implies |az|^N \xrightarrow{N \rightarrow \infty} 0$$

So,

$$\lim_{N \rightarrow \infty} S_N = \frac{1}{1 - az}$$

- $|az| > 1$ **case:**

$$|az| > 1 \implies |az|^N \xrightarrow{N \rightarrow \infty} \infty$$

so

$$\lim_{N \rightarrow \infty} S_N \text{ diverges}$$

- $|az| = 1$ **case:**

Diverges if $|az| \neq 1$ and if $|az| = 1$. In this case, both diverge, but in general there may be more complicated behavior.

Conclusion: The geometric series converges (absolutely) for when $|z| < |a|$.

Recall that

$$\sum_{n \geq 0} z_n, z_n \in \mathbb{C}$$

converges absolutely if

$$\sum_{n \geq 0} |z_n|$$

converges.

So, we notice that the series converges for any z such that $|z| < 1/|a|$. This region is just an open disk of radius $|a|$. In general, power series has radii of convergence.

Radius of Convergence

Def: A complex Power Series

$$\sum_{n \geq 0} a_n z^n$$

has Radius of Convergence

$$r = \left(\lim_{n \rightarrow \infty} \sup |a_n|^{1/n} \right)^{-1} \in \mathbb{R}$$

Example: For $a_n = a^n$, we have

$$r = \left(\lim_{n \rightarrow \infty} \sup |a_n|^{1/n} \right)^{-1} = \frac{1}{|a|}$$

which matches with the result obtained earlier.

Theorem:

1. If $|z| < r$, then $f(z)$ converges absolutely.
2. If $|z| > r$, then it diverges.
3. At $|z| = r$, more care is needed.

Proof Sketch:

1. Consider z such that $|z| < (1 - \epsilon)r$ for some $\epsilon > 0$.

$$\begin{aligned}
\Rightarrow |a_n z^n| &< |a_n|(1 - \epsilon)^n r^n \\
&\leq |a_n|(1 - \epsilon)^n \left(\frac{1}{|a_n|^{1/n}} \right)^n \quad \text{Assume } a_n \neq 0 \\
&\leq (1 - \epsilon)^n \quad (\text{If } a_n = 0, \text{ this inequality is true trivially}) \\
&\Rightarrow \text{Convergence by Geometric series}
\end{aligned}$$

Term by term, the series is smaller than the geometric series (which converges), thus it also converges (Dominated Convergence Theorem).

2. If $|z| > r$, then $|z| > r/(1 - \epsilon)$ for some $\epsilon > 0$ while

$$|a_n|^{1/n} > \left(\limsup_{k \rightarrow \infty} |a_k|^{1/k} \right) (1 - \epsilon)$$

for infinitely many n .

$$\begin{aligned}
\Rightarrow |a_n z^n| &> \frac{1}{r^n} (1 - \epsilon)^n \cdot \frac{r^n}{(1 - \epsilon)^n} > 1 \quad \text{for inf. many } n \\
\Rightarrow \text{the sum diverges}
\end{aligned}$$

Example: Consider a polynomial

$$f(z) = \sum_{n=0}^N a_n z^n$$

$$\begin{aligned}
\Rightarrow \lim_{n \rightarrow \infty} \sup |a_n|^{1/n} &= 0 \\
\Rightarrow r &= \infty
\end{aligned}$$

Example: Consider the exponential function

$$\exp(z) = e^z = \sum_{n \geq 0} \frac{z^n}{n!}$$

$$\Rightarrow r = \lim_{n \rightarrow \infty} (n!)^{1/n} =_{\text{claim}} \infty$$

Proof of claim: We will show that for any b , $\frac{1}{2}b < (n!)^{1/n}$ for all $n \gg 0$.

Note that we can shift the center of the series from 0 to some point z_0 .
So, instead of

$$\sum_{n \geq 0} a_n z^n$$

We have

$$\sum_{n \geq 0} a_n (z - z_0)^n$$

The radius of convergence is given by the same expression.

1.2 Differentiation of Power Series

Knowing convergence properties of a series is also useful for the purposes of *differentiation*.

Theorem: Let r be the radius of convergence of $f(z) = \sum_{n \geq 0} a_n (z - z_0)^n$. Then, f is (complex) differentiable on $B_r(z_0)$ and

$$f'(z_0) = \sum_{n \geq 0} n a_n (z - z_0)^{n-1}$$

Proof: The proof follows the same argument as that in real analysis. Write the proof later.
One proof can be found [here](#).

We will break the proof into two parts:

- (a) First, we show that if $\sum_{n \geq 0} a_n (z - z_0)^n$ has radius of convergence r , then so does $\sum_{n \geq 0} n a_n (z - z_0)^{n-1}$.
- (b) Second, we show that if $f(z) = \sum_{n \geq 0} a_n (z - z_0)^n$ then the derivative at z_0 is indeed

$$f'(z_0) = \sum_{n \geq 0} n a_n (z - z_0)^{n-1}$$

Proof: (Check if this is correct in Office Hours)

- (a) Earlier, we proved that for any complex power series there exists a number r called the radius of convergence such that the series converges for any $z \in \mathbb{C}$ such that $|z| < r$. We also showed that r is given by

$$r = \limsup_{n \rightarrow \infty} |a_n|^{1/n}$$

So, what we want to do here is show that the radii of convergence for the two series R, R' are equal.

Now,

$$\begin{aligned} R' &= \limsup_{n \rightarrow \infty} |n a_n|^{1/n} \\ &= \limsup_{n \rightarrow \infty} |n|^{1/n} \cdot |a_n|^{1/n} \\ &= \limsup_{n \rightarrow \infty} |n|^{1/n} \cdot |a_n|^{1/n} \\ &= \limsup_{n \rightarrow \infty} |1| \cdot |a_n|^{1/n} \\ &= R \end{aligned}$$

So the two series have equal radii of convergence ($= R$).

(b) Next, we want to show that if

$$f(z) = \sum_{n \geq 0} a_n (z - z_0)^n$$

and

$$g(z) = \sum_{n \geq 0} n a_n (z - z_0)^{n-1}$$

Then, in fact, $f'(z) = g(z)$.