

# Math 214 Notes

Keshav Balwant Deoskar

February 6, 2024

These are notes taken from lectures on Differential Topology delivered by Eric C. Chen for UC Berkeley's Math 214 class in the Spring 2024 semester. Any errors that may have crept in are solely my fault.

## Contents

<b>1</b>	<b>February 6 - Tangent Vectors, Differentials of smooth maps</b>	<b>2</b>
1.1	Properties of Derivations . . . . .	2
1.2	Tangent Space to $\mathbb{R}^n$ . . . . .	3
1.3	The Differential of a Smooth Map . . . . .	4
1.4	Tangent Space of Smooth Manifolds (with or without boundary) . . . . .	5
1.5	Coordinates . . . . .	5

# 1 February 6 - Tangent Vectors, Differentials of smooth maps

## Recap

- Intuitively, we can visualize the tangent plane to a point on a sphere embedded in  $\mathbb{R}^3$ , however this doesn't generalize very well to higher dimensions.
- For abstract manifolds, recall that we defined a map  $v : C^\infty(M) \rightarrow \mathbb{R}$  to be a **derivation at**  $p \in M$  if it satisfies two properties:

– R-linearity:

$$v(af + bg) = av(f) + bv(g)$$

– "Product Rule":

$$v(fg) = f(p)v(g) + v(f)g(p)$$

- The space of derivations at  $p$  is defined to be the Tangent Plane  $T_pM$ , and it has a natural vector space structure. Intuitively, each element of the tangent space is a direction in which we can take a directional derivative.
- Example: Take  $M = \mathbb{R}^n$  and  $\vec{p} \in \mathbb{R}^n$ ,  $\vec{v}_0 = (v^1, \dots, v^n) \in \mathbb{R}^n$

Let

$$v(f) = \sum_{i=1}^n v^i \left( \frac{\partial}{\partial x^i} f \right) (p)$$

## 1.1 Properties of Derivations

Lemma: If  $v \in T_pM$ , then  $v(c) = 0$  where  $c \in \mathbb{R}$  is constant.

Proof:  $v(c) = v(c1) = c \cdot v(1)$ , so it suffices to show that  $v(1) = 0$ . Now,

$$\begin{aligned} v(1) &= v(1 \cdot 1) \\ &= 1 \cdot v(1) + v(1) \cdot 1 \end{aligned}$$

$$\implies v(1) = 0$$

Lemma: If  $f, g \in C^\infty(M)$  agree on a neighborhood  $U \subseteq M$  containing  $p$ , and  $v \in T_p M$ , then  $v(f) = v(g)$ .

Proof: Find a precompact neighborhood  $B \ni p$  such that  $p \in \overline{B} \subseteq U$ . There exists a bump function  $\psi \in C^\infty$  such that

$$\begin{cases} \psi \equiv 1 \text{ on } \overline{B} \\ \text{supp}(\psi) \subseteq U \end{cases}$$

Note that  $\psi \cdot (f - g) \equiv 0$  on all of  $M$ , so

$$\begin{aligned} 0 &= v(\psi \cdot (f - g)) \\ &= \psi(p) \cdot v(f - g) + \underbrace{v(\psi) \cdot (f - g)(p)}_{=0} \\ &= v(f - g) \\ &= v(f) - v(g) \end{aligned}$$

Now that we've proved some properties of derivations, we can consider other perspectives from which we can understand the tangent plane  $T_p M$ .

## 1.2 Tangent Space to $\mathbb{R}^n$

Lemma: At  $\vec{a} = (a^1, \dots, a^n) \in \mathbb{R}^n$ , for each  $\vec{v} = (v^1, \dots, v^n) \in \mathbb{R}^n$  we can define the map

$$D_{\vec{v}}|_{\vec{a}} : C^\infty(\mathbb{R}^n) \rightarrow \mathbb{R}^n$$

as

$$f \mapsto (\partial_{\vec{v}} f)(\vec{a}) = \sum_{i=1}^n v^i \frac{\partial f}{\partial x^i} \Big|_{\vec{a}}$$

which is a derivation at  $\vec{a}$ , so  $D_{\vec{v}}|_{\vec{a}} \in T_{\vec{a}} \mathbb{R}^n$   
and we will prove that

$$D : \mathbb{R}^n \rightarrow T_{\vec{a}} \mathbb{R}^n$$

defined by

$$\vec{v} \mapsto D_{\vec{v}}|_{\vec{a}}$$

is an isomorphism.

Proof:

Leaving linearity and injectivity as exercises, we show surjectivity. Take a derivation  $v \in T_{\vec{a}} \mathbb{R}^n$ , and let  $u^i = v(x^i) \in \mathbb{R}, i = 1, \dots, n$  (we're going to essentially do the reverse of  $D$ ).

c We will show that

$$v = D_{\vec{u}}|_{\vec{a}} = \sum_{i=1}^n u^i \frac{\partial}{\partial x^i} \Big|_{\vec{a}}$$

First note, for  $f \in C^\infty(M)$ , we have

$$\begin{aligned} f(\vec{x}) &= f(\vec{a}) + \int_0^1 \frac{d}{dt} f(\vec{a} + t(\vec{x} - \vec{a})) dt \\ &= f(\vec{a}) + \sum_{i=1}^n (x^i - a^i) \int_0^1 \underbrace{\frac{\partial f}{\partial x^i}(\vec{a} + t(\vec{x} - \vec{a}))}_{h^i(\vec{x})} dt \end{aligned}$$

Note that

$$h^i(\vec{a}) = \frac{\partial f}{\partial x^i}(\vec{a})$$

Apply the derivation  $v$  to  $f(\vec{x})$ :

$$\begin{aligned} v(f) &= v(f(\vec{a})) + \sum_{i=1}^n v(x^i - a^i) h^i(\vec{a}) + \sum_{i=1}^n v(a^i - a^i) v(h^i) \\ &= \sum_{i=1}^n u^i \cdot \frac{\partial f}{\partial x^i}(\vec{a}) \end{aligned}$$

### 1.3 The Differential of a Smooth Map

Give smooth manifolds  $M^m, N^n$  and a smooth map  $F : M \rightarrow N$ , the **differential of  $F$**  at  $p \in M$  is defined as

$$\begin{aligned} dF_p : T_p M &\rightarrow T_{F(p)} N \\ v &\mapsto dF_p(v) \end{aligned}$$

where  $(dF_p(v))(f) = v(f \circ F)$ , for  $f \in C^\infty(N)$ .

[Insert image]

One way to interpret this is as follows: If we think of a derivative (derivation at point  $p \in M$ ) to be a tangent/velocity curve then  $dF_p$  tells us how this velocity curve changes under the map  $F : M \rightarrow N$ .

Example: Take  $M = \mathbb{R}^m$ ,  $N = \mathbb{R}^n$  and

$$F(\vec{x}) = (F^1(x^1, \dots, x^m), \dots, F^n(x^1, \dots, x^m))$$

Then, for  $f \in C^\infty(N)$ ,

$$\begin{aligned} \left( dF_p \left( \frac{\partial}{\partial x_i} \Big|_p \right) \right) (p) &= \left( \frac{\partial}{\partial x_i} \Big|_p \right) (f \circ F) \\ &= [finishafterlec] \end{aligned}$$

Read: More properties of differential (Maybe prove later).

- Composition of diffeomorphisms, differential of identity is isomorphism, etc.

Now that we've studied the differential of a map between euclidean spaces, let's work with more general manifolds.

## 1.4 Tangent Space of Smooth Manifolds (with or without boundary)

Proposition: Let  $M^n$  be a smooth manifold,  $p \in U \subseteq_{\text{open}} M$  and consider the inclusion map  $\iota : U \rightarrow M$ . Then,

$$\begin{aligned} d\iota_p : T_p U &\rightarrow T_p U \\ v &\mapsto d\iota_p(v) \end{aligned}$$

is an isomorphism [Recall that by definition  $(d\iota_p(v))(f) = v(f \circ \iota)$ ].

This is essentially an application of the property we proved earlier that the derivation only depends on a small neighborhood at each point.

Proof:

*Injectivity:* Since it's a linear map, it is injective if

$$\begin{aligned} d\iota_p(v) = 0 &\iff d\iota_p(v)(f) = 0 \text{ for all } f \in C^\infty(M) \\ &\iff v(f \circ \iota) = 0 \\ &\iff v(f|_U) = 0 \end{aligned}$$

We want to show  $f(\tilde{f}) = 0$  for all  $\tilde{f} \in C^\infty(U)$ . Choose  $f \in C^\infty(M)$  such that  $\tilde{f} = f$  near  $p$ . Then,

$$\begin{aligned} 0 &= v(f|_U) \\ &= v(\tilde{f}) \text{ (by locality of derivations)} \end{aligned}$$

This shows injectivity.

*Surjectivity:* Given a derivation  $\tilde{v} \in T_p M$ , we want to find  $v \in T_p U$  such that  $\tilde{v}(f) = v(\tilde{f}|_U)$  for all  $\tilde{f} \in C^\infty$ . Given a function  $f \in C^\infty(U)$ , define  $v \in T_p U$  by

- Choose some (doesn't matter which) extension  $\tilde{f} \in C^\infty(M)$  such that  $f = \tilde{f}$  on a neighborhood of  $p$ .
- Set  $v(f) \equiv \tilde{v}(\tilde{f})$  and check that this is well defined i.e. independent of  $\tilde{f}$  choice (results from locality of derivations)

Corollary:  $\dim(T_p M^n) = n$

Proof: Given  $p \in M^n$ , choose a smooth chart  $(U, \phi)$  with  $p \in U$ . Then, from the above result, we know

$$T_p M \xleftarrow[\text{by prop}]{\cong} T_p U \xrightarrow[\text{isom.}]{\cong} T_{\phi(p)} \phi(U) \xrightarrow[\text{by incl. map}]{\cong} T_{\phi(p)} \mathbb{R}^n = \mathbb{R}^n$$

## 1.5 Coordinates

[Write from image and include graphic]

Note that

$$\left. \frac{\partial}{\partial x^1} \right|_p, \dots, \left. \frac{\partial}{\partial x^m} \right|_p$$

is a basis for  $T_p M$ .

Read: Differential of a smooth map in coordinates.