

Math H185 Lecture 3

Keshav Balwant Deoskar

March 2, 2024

These are notes taken from lectures on Complex Analysis delivered by Professor Tony Feng for UC Berkeley's Math H185 class in the Spring 2024 semester.

Contents

1	January 24 - Power Series	2
1.1	Convergence	2
1.2	Differentiation of Power Series	5

1 January 24 - Power Series

Power Series

A **Power Series** is a formal expression

$$\sum_{n \geq 0} a_n z^n, z_n \in \mathbb{C}$$

for which operations are defined as:

- Addition:

$$\sum_{n \geq 0} a_n z^n$$

- multiplication

- "Formal" here means we temporarily ignore whether or not it makes sense to plug in complex numbers into such formulae.

1.1 Convergence

Defining these formal expressions is cool, but when does a power series actually define a function? It does so when the series **converges**.

Example: Geometric series:

Let $a \in \mathbb{C}$, then for the geometric series we have $a_n = a^n$. So,

$$\sum_{n \geq 0} a^n z^n = 1 + az + a^2 z^2 + \dots$$

converges if $S_n = \sum_{n \geq 0}^{N-1} a^n z^n$ has a limit.

By the same argument as in the reals, we can get a closed form expression for S_N :

$$S_N = \frac{1 - (az)^N}{1 - (az)}$$

To deal with convergence, we break into cases and take the limit.

- $|az| < 1$ **case:**

$$|az| < 1 \implies |az|^N \xrightarrow{N \rightarrow \infty} 0$$

So,

$$\lim_{N \rightarrow \infty} S_N = \frac{1}{1 - az}$$

- $|az| > 1$ **case:**

$$|az| > 1 \implies |az|^N \xrightarrow{N \rightarrow \infty} \infty$$

so

$$\lim_{N \rightarrow \infty} S_N \text{ diverges}$$

- $|az| = 1$ **case:**

Now, if $|az| = 1$ but $az \neq 1$, then $(az)^N = \underbrace{(az) \times \cdots \times (az)}_{N \text{ times}}$ just means we rotate around on the unit circle without converging to any point in particular. So, the sum diverges.

If instead we have $az = 1$, then the denominator $1 - az$ vanishes and the sum diverges again.

Note: This time it happened to be the case that both case falling under $|az| = 1$ diverged, but in general we can have more complicated behavior.

Conclusion: The geometric series converges (absolutely) for when $|z| < |a|$.

Recall that

$$\sum_{n \geq 0} z_n, z_n \in \mathbb{C}$$

converges absolutely if

$$\sum_{n \geq 0} |z_n|$$

converges.

So, we notice that the series converges for any z such that $|z| < 1/|a|$. This region is just an open disk of radius $|a|$. In general, power series have radii of convergence.

Radius of Convergence

Def: A complex Power Series

$$\sum_{n \geq 0} a_n z^n$$

has Radius of Convergence

$$r = \left(\lim_{n \rightarrow \infty} \sup |a_n|^{1/n} \right)^{-1} \in \mathbb{R}$$

Example: For $a_n = a^n$, we have

$$r = \left(\lim_{n \rightarrow \infty} \sup |a_n|^{1/n} \right)^{-1} = \frac{1}{|a|}$$

which matches with the result obtained earlier.

Theorem:

1. If $|z| < r$, then $f(z)$ converges absolutely.
2. If $|z| > r$, then it diverges.
3. At $|z| = r$, more care is needed.

Proof Sketch:

1. Consider z such that $|z| < (1 - \epsilon)r$ for some $\epsilon > 0$.

$$\begin{aligned}
\Rightarrow |a_n z^n| &< |a_n|(1 - \epsilon)^n r^n \\
&\leq |a_n|(1 - \epsilon)^n \left(\frac{1}{|a_n|^{1/n}} \right)^n \quad \text{Assume } a_n \neq 0 \\
&\leq (1 - \epsilon)^n \quad (\text{If } a_n = 0, \text{ this inequality is true trivially}) \\
&\Rightarrow \text{Convergence by Geometric series}
\end{aligned}$$

Term by term, the series is smaller than the geometric series (which converges), thus it also converges (Dominated Convergence Theorem).

2. If $|z| > r$, then $|z| > r/(1 - \epsilon)$ for some $\epsilon > 0$ while

$$|a_n|^{1/n} > \left(\limsup_{k \rightarrow \infty} |a_k|^{1/k} \right) (1 - \epsilon)$$

for infinitely many n .

$$\begin{aligned}
\Rightarrow |a_n z^n| &> \frac{1}{r^n} (1 - \epsilon)^n \cdot \frac{r^n}{(1 - \epsilon)^n} > 1 \quad \text{for inf. many } n \\
\Rightarrow \text{the sum diverges}
\end{aligned}$$

Example: Consider a polynomial

$$f(z) = \sum_{n=0}^N a_n z^n$$

$$\begin{aligned}
\Rightarrow \lim_{n \rightarrow \infty} \sup |a_n|^{1/n} &= 0 \\
\Rightarrow r &= \infty
\end{aligned}$$

Example: Consider the exponential function

$$\exp(z) = e^z = \sum_{n \geq 0} \frac{z^n}{n!}$$

$$\Rightarrow r = \lim_{n \rightarrow \infty} (n!)^{1/n} =_{\text{claim}} \infty$$

Proof of claim: We will show that for any b , $b < (n!)^{1/n}$ for all $n \gg 0$.

For very large n , we have $n! = \underbrace{1 \cdot 2 \cdots b}_{\geq 1} \cdot \underbrace{(b+1)}_{\geq b} \cdot \underbrace{(b+2)}_{\geq b} \cdots \underbrace{n}_{\geq b}$. So,

$$(n!)^{1/n} > (b^{n-m})^{1/n} = b^{1-m/n} = b \cdot b^{-m/n}$$

and

$$\lim_{n \rightarrow \infty} b^{-m/n} = 1$$

So, in the $n \rightarrow \infty$ limit,

$$(n!)^{1/n} > b$$

Since this holds for *any* constant b , it must be the case that

$$\lim_{n \rightarrow \infty} (n!)^{1/n} = \infty \implies \boxed{r = \infty}$$

Note: So far we've only considered series centered at the origin but we can shift the center of the series from 0 to some point z_0 .

So, instead of

$$\sum_{n \geq 0} a_n z^n$$

We have

$$\sum_{n \geq 0} a_n (z - z_0)^n$$

The radius of convergence is given by the same expression.

1.2 Differentiation of Power Series

Knowing convergence properties of a series is also useful for the purposes of *differentiation*.

Theorem: Let r be the radius of convergence of $f(z) = \sum_{n \geq 0} a_n (z - z_0)^n$. Then, f is (complex) differentiable on $B_r(z_0)$ and

$$f'(z_0) = \sum_{n \geq 0} n a_n (z - z_0)^{n-1}$$

Proof: The proof follows the same argument as that in real analysis. Write the proof later. One proof can be found [here](#). Otherwise see Stein-Shakarchi Theorem 2.6.

We will break the proof into two parts:

- (a) First, we show that if $\sum_{n \geq 0} a_n (z - z_0)^n$ has radius of convergence r , then so does $\sum_{n \geq 0} n a_n (z - z_0)^{n-1}$.
- (b) Second, we show that if $f(z) = \sum_{n \geq 0} a_n (z - z_0)^n$ then the derivative at z_0 is indeed

$$f'(z_0) = \sum_{n \geq 0} n a_n (z - z_0)^{n-1}$$