OXFORD

(GIGA)n SCIENCE

GigaScience, 2026, 1–22

doi: xx.xxxx/xxxx
Manuscript in Preparation
Research

RESEARCH

Auditing cross-cohort feature-selection benchmarks: cohort composition and tissue-context-dependent confounding across kidney, liver and colon transcriptomes

Dohun Kim[1*], Minho Bae[1], Jinho Park[1], Hayoung Lee[1], Hyeokjin Lim[1] and Seonghee Lee[1]

[1]Autonomous Intelligence DX Research Section, Electronics and Telecommunications Research Institute, Daejeon, Republic of Korea

[*]Corresponding author. E-mail: dohun@etri.re.kr
Author e-mail addresses: Dohun Kim, dohun@etri.re.kr; Minho Bae, minkkang@etri.re.kr; Jinho Park, jinho.park@etri.re.kr; Hayoung Lee, underzero11@etri.re.kr; Hyeokjin Lim, hyeokjin.lim@etri.re.kr; Seonghee Lee, slee0003@etri.re.kr.

Abstract

Background: Cross-cohort feature selection is widely used to derive reproducible transcriptomic biomarker signatures from public cohorts. This assumes that reproducibility reflects disease biology and that the benchmark measures the methods rather than the cohorts. We tested both in diabetic kidney disease, then in liver and colon against predictions fixed in advance.
Results: Over all 26 subsets of five discovery cohorts, reproducibility verdicts were unstable: a stability-selection strategy won cross-fold agreement in 9 of 10 three-cohort subsets yet lost on all five, while its external AUROC advantage stayed positive. The most reproducible signal tracked procurement, not disease: an immediate-early gene module separated biopsy from surgical tissue in 8 of 9 kidney datasets, and a biopsy control retained only 14–56% of the naive signal across seven diagnoses. Residualising on a handling score cut procurement-driven genes from 27% to 5% with no AUROC loss detected on a wide interval; batch correction by cohort was inert. In liver and colon the module was lower in biopsy cases than surgical controls in 9 of 10 such cohorts, verdict instability recurred in liver but not colon, and the module rose with inflammation, failing one prediction.
Conclusions: Without a cohort-composition sensitivity analysis, a cross-cohort reproducibility comparison describes the cohorts rather than the methods, and should be reported separately from predictive performance. A shared pre-analytical confounder becomes reproducible wherever the design is shared; removing it rewrites the candidate list, 14 of 30 genes surviving. We release the harmonised cohorts, the schema-enforced audit database, the candidate-evidence ledger and the regenerating pipeline.

Key words: feature selection; benchmarking; reproducibility; batch effects; pre-analytical variables; diabetic kidney disease; transcriptomics

Background

Transcriptomic biomarker discovery in chronic kidney disease has converged on a common recipe: assemble public case/control cohorts, select features that behave consistently across them, and report the resulting signature with a classification performance figure. The appeal is that consistency across independent cohorts should filter dataset-specific noise. A substantial methods literature has developed around this idea, from multi-cohort meta-analysis [1] to stability-selection frameworks [2]. The pitfalls of deriving classifiers from high-dimensional expression data have

Compiled on: October 8, 2026.
Draft manuscript prepared by the author.

been catalogued repeatedly [3, 4, 5], as has the impact of batch and other technical structure on such analyses [6].

Two assumptions underlie the recipe. The first is that a feature reproducible across cohorts is more likely to reflect disease biology. The second is that a benchmark comparing selection methods on a set of public cohorts estimates something about the methods rather than about the cohorts. This work tests both, in diabetic kidney disease (DKD), where the number of usable public cohorts is small enough to enumerate exhaustively and large enough to resample, and then asks whether what it finds there carries to two tissues whose cohorts are designed differently.

Neither concern is new in the abstract. That the outcome of a method comparison depends on how the comparison was set up (which datasets, which defaults, which analysis choices) has been argued and demonstrated repeatedly in computational biology: in the case for neutral comparison studies [7], in explicit illustrations of over-optimism [8, 9], and in the benchmarking guidelines that grew out of them [10, 11]. A related bias is that method comparisons run by a method's own authors tend to favour it [12], and the same sensitivity has been described outside biology as a benchmark lottery [13]. Separately, the instability of high-dimensional feature selection under resampling is a standing result [14].

A meta-analysis of 28 filter feature-selection studies found that the number of datasets, baselines and new methods alone explained 33% of the variance in reported win rates [15], and a recent framework grades which layers of differential-expression evidence reproduce across fixed cohorts [16]; neither varies which cohorts enter the comparison. What is missing is a quantification for cross-cohort omics biomarker discovery specifically, where varying the cohort configuration is recommended practice but reporting the resulting interval is not. We therefore treat the phenomenon as established and measure its magnitude in a setting that has not accounted for it, alongside a concrete mechanism for one particular failure and a benchmark of the corrections available for it.

This study makes five contributions. The first is a cohort-composition sensitivity analysis, stated as a protocol rather than as a one-off check: the reproducibility comparison is reported as an interval over every admissible subset of the discovery cohorts, so that a verdict which depends on the configuration cannot be presented as a verdict about the methods. The second is a control-definition sensitivity analysis, which replaces the non-biopsy control with a disease control obtained through the same biopsy procedure. It separates a design effect from a disease effect without collecting new samples, and it needs only cohorts that already exist. The third is a benchmark of the corrections available for the confounder those two analyses expose, including which corrections cannot work in this design and why. The fourth is the form of the output: an auditable candidate-evidence ledger rather than a ranked gene list, recording for each candidate the evidence lines it passed, the rank it takes under each control definition, and how far that rank moves when the control group is substituted. The fifth is a test of whether the two mechanisms are properties of cross-cohort benchmarking or of these kidney cohorts: three predictions, written and hashed before any expression value was read, evaluated in liver, where procurement is asymmetric in some cohorts and matched in others, and in colon, where it is matched in all. One held, one held in one tissue, and one failed, and the failures narrow both claims in ways the kidney data could not.

The study proceeds in four steps: audit the benchmark input, stress-test conclusions against cohort composition and control definition, then diagnose and correct the confounder before reporting an auditable candidate-evidence ledger. The cohort audit, the residual overlap it could not remove, and the discovery/validation split are described in the section *Cohort assembly and audit*.

Both assumptions fail in ways that are measurable. Reproducibility is maximised by whatever is most consistent across cohorts, and in this disease that turns out to be a shared pre-analytical confounder rather than biology. Benchmark verdicts on reproducibility flip sign depending on which cohorts are included. Neither failure is specific to a particular algorithm, and both are invisible without the controls we describe.

Data description

Cohort assembly and audit

Fifteen human DKD transcriptome series were retrieved from the Gene Expression Omnibus. Sample-level metadata were parsed from series matrices and patient identifiers recovered from sample titles, which revealed patient-level reuse invisible at the accession level. Three series (GSE30122, GSE47183 and GSE99340) were excluded on this basis; Additional file 2 gives the evidence for each. Reusing them would have inflated every cross-cohort statistic in this study, because the same patients would have appeared in both the selection and the evaluation side of a leave-one-dataset-out split.

Residual overlap that could not be removed is disclosed rather than ignored (Table 1; Additional file 2 gives the full audit). It is larger than the DKD case counts alone suggest: GSE104948 and GSE104954 are the two compartments of one ERCB collection and share 55 subjects, of which 5 are DKD cases, 1 is a non-biopsy control and 49 are the non-diabetic kidney-disease biopsies that the section *The most reproducible signal tracks tissue procurement, not disease* uses as the matched comparator. GSE30528 and GSE30529 share 9 subjects, 5 DKD cases and 4 controls. These are not duplicate cohorts; they are the same patients profiled in two compartments, which is what makes the paired analyses possible. The consequence is that the leave-one-dataset-out folds are not fully independent when both compartments are present, and we report it rather than treat the two as unrelated datasets. GSE104948 also enters the analysis twice: once as a discovery cohort and once as one of the two blocks of the DKD-versus-non-diabetic-CKD specificity gate (the section *Which corrections work*). The gate is therefore not a fully independent validation of the candidates it filters. Independence was probed instead in the external GSE20602 cohort (the section *The control-group problem is not specific to diabetic kidney disease*), and a sensitivity analysis re-derives the candidates with GSE104948 removed from discovery (the section *Limitations*).

Five cohorts entered the discovery rotation and two served as compartment validation (Table 2) [17, 18, 19, 20, 21]. One feature of that table carries the rest of this paper: in every cohort the cases are kidney biopsies and the controls are not. None of the five discovery contrasts uses a biopsy-matched control group; the biopsy comparators that do exist, in ERCB and KPMP, are what the control-definition analyses below rely on. Additional cohorts were used for specific tests: GSE175759 [22], GSE162830 [23], GSE142153 [24], GSE1009 [25], GSE111154 [26], GSE20602 [27], the Kidney Precision Medicine Project (KPMP) single-nucleus atlas [28, 29], and GSE5406 (heart, external organ) [30]. KPMP does not enrol diabetic kidney disease as a category; the 27 single-nucleus donors enrolled with chronic kidney disease and a recorded history of diabetes are treated throughout as an operational DKD-proxy subset rather than a registry-defined diagnosis (the section *Limitations*). Three of the discovery cohorts were first described by Woroniecka et al. [17].

Harmonisation

Entrez Gene ID was used as the hub identifier because GSE104948 and GSE104954 use the Brainarray ENTREZG custom CDF, so their probe identifiers already are Entrez IDs and the tightest cohort suffers no mapping loss. Platform probes were mapped through GEO `.annot` files (GPL571, GPL96, GPL8300, GPL6480), the GPL17586 platform table, and NCBI `gene_info` for symbol- and Ensembl-keyed matrices. Probes mapping to more than one gene were

Table 1. Subjects appearing in more than one retained series. These are the same patients profiled in two compartments, not duplicate cohorts.

Series A	Series B	Shared subjects	DKD cases	Non-biopsy controls	Other biopsy
GSE104948	GSE104954	55	5	1	49
GSE30528	GSE30529	9	5	4	0

Other biopsy are the non-diabetic kidney-disease biopsies that the section *The most reproducible signal tracks tissue procurement, not disease* uses as the procurement-matched comparator. They are counted here because the leave-one-dataset-out folds are not independent for those subjects either.

Table 2. Transcriptome cohorts used. Cases are kidney biopsies in every cohort; controls are not. This is the design feature the section *The most reproducible signal tracks tissue procurement, not disease* tests.

Cohort	Role	Compart.	Platform	Cases	Controls	Control tissue
GSE104948	discovery	glom.	GPL22945/GPL24120	12	26	donor, nephrec.
GSE142025	discovery	cortex	Illumina HiSeq4000	27	9	nephrec.
GSE294519	discovery	tubulo.	GPL24676	23	13	nephrec.
GSE30528	discovery	glom.	GPL571	9	13	nephrec.
GSE96804	discovery	glom.	GPL17586	41	20	nephrec.
GSE104954	compartment	tubulo.	GPL22945/GPL24120	17	26	donor, nephrec.
GSE30529	compartment	tubulo.	GPL571	10	12	nephrec.

Compartments: glom., glomerulus; tubulo., tubulointerstitium; cortex, whole renal cortex. Control tissue: nephrec., tumour nephrectomy; donor, living kidney donor. All cohorts are expressed on the same frozen 9,900-gene Entrez space. Cohorts marked *compartment* are the second compartment of donors already represented by an ERCB or Woroniecka series; they enter the procurement and comparator analyses but not the cohort-composition sweep, which requires independent cohorts.

dropped. Where several probes mapped to one gene the probe with the highest mean expression was retained (max-mean collapse), a rule fixed before any analysis and applied identically everywhere, because changing it invalidates every stability count computed on the resulting space.

The intersection over the four original discovery cohorts is 9,900 genes; adding the compartment cohorts costs nothing since they share platforms. GSE294519 was mapped into this frozen space rather than triggering re-derivation, and covers 98.5% of it.

All cross-cohort operations use per-cohort gene-wise z-scores. Value scales differ fundamentally between cohorts (GSE30528 and GSE30529 are mean-centred and negative; others are positive log-scale), so raw values are never pooled. The z-scoring uses no label information, so applying it to a held-out cohort is unsupervised domain alignment, not leakage.

Analyses

All comparisons below use the four strategies of the section *Selection methods*, each implemented directly rather than through a reference package; Additional file 1 documents what each shares with its reference definition and tests whether the departures matter. The order of what follows is the order of the questions each result answers: whether the reported performance figure means anything, whether the reproducibility comparison is stable under a change of cohorts, what the reproducible signal actually tracks, why one strategy escaped it, which corrections remove it, whether both findings hold in other tissues, and what the pipeline returns once the corrections are applied.

Absolute AUROC is uninformative in this disease

Randomly drawn 50-gene sets achieve external AUROC of 0.68–0.88 depending on the held-out cohort, while permuting training labels gives approximately 0.50 (Table 3, Figure 1). The signal is real, but which genes carry it is nearly immaterial. The DKD programme is broadly co-expressed: among the 50 genes with the largest mean $|g|$ the mean pairwise $|r|$ within a cohort is 0.51, against 0.25 for random gene sets of the same size, so most random signatures inherit it. On the GSE142025 fold, random sets reach AUROC 1.000 at $K = 100$, making that fold unable to discriminate methods at all.

A comparable random-signature phenomenon appears in a metabolomic layer, measured in different patients by a different principle. In plasma from ST003255 (64 diabetic nephropathy, 66 IgA nephropathy, 66 membranous nephropathy, 24 hypertensive nephropathy, 66 healthy), randomly drawn 20-metabolite sets reach a cross-validated AUROC of 0.913 against healthy controls, while a set selected inside each fold reaches 0.995 and permuted labels give 0.50. Imputation, log transformation and scaling are fitted on the training fold only and applied to the held-out fold, so no test sample contributes to the transform. The signature size was fixed at 20 in advance; at 10 the selected set still clears the 95th percentile of the null and at 50 it no longer does, which is itself the point: the margin over the null, not the absolute figure, is what a comparison can rest on. This is a single cohort evaluated by cross-validation rather than an independent replication, and we read it as a cross-modal illustration that the reporting problem is not confined to transcriptomes.

Every AUROC reported below is therefore accompanied by its position in this null.

Reproducibility verdicts depend on cohort composition, not on method

Repeating the full benchmark over all subsets of the five discovery cohorts gives the distribution of conclusions a study could have reported (Table 4, Figure 2).

The paired difference RBS − connectivity in cross-fold agreement spans −0.106 to +0.170 across three-cohort subsets. The two extremes are concrete: a study using GSE96804 + GSE142025 + GSE294519 would report +0.170 in favour of stability selection, while one using GSE30528 + GSE104948 + GSE142025 would report −0.106 against it. Both are legitimate three-cohort DKD benchmarks.

External AUROC behaves differently. Its advantage is directionally stable and grows with cohort count: +0.056 (2 cohorts), +0.058 (3), +0.096 (4), and +0.139 at five cohorts. The paired bootstrap put that five-cohort difference at +0.129 [−0.002, +0.263] ($p = 0.06$ over 200 replicates): the mean difference favoured stability selection, but the interval included zero.

The two metrics do not even agree on which method won. Excluding subsets where the top score is tied, the method ranked first on cross-fold reproducibility is also the one ranked first on external AUROC in 11 of 24 subsets. At three cohorts the two agree in 5 of 10. A benchmark that reports one of these numbers is therefore not




Figure 1. Absolute AUROC is uninformative in this disease. External AUROC for random 50-gene signatures, for permuted-label controls, and for the selected signatures, by held-out cohort. Random signatures reach 0.68–0.88 depending on the fold; permuted labels sit at approximately 0.50. A reported AUROC must be read against the random-signature band for its fold, not against 0.5.

Alt text: Dot plot of external AUROC by held-out cohort for three groups: random 50-gene signatures, permuted-label controls and the selected signatures. Random signatures form a band from about 0.68 to 0.88 that differs by fold; permuted labels sit near 0.50; the selected signatures are shown against that band for each fold.

Table 3. Random-signature and permuted-label nulls, by held-out cohort (K = 50)

Held out	Random 50-gene AUROC	Permuted labels
GSE30528	0.682 [0.385–0.923]	0.52
GSE96804	0.796 [0.597–0.946]	0.47
GSE104948	0.809 [0.602–0.962]	0.55
GSE142025	0.878 [0.654–1.000]	0.49

Brackets give the 5th–95th percentile over 300 random draws.

Table 4. Benchmark winner by cross-fold reproducibility, by subset size

Cohorts	Subsets	Winner by reproducibility
3	10	RBS 9 (90%), connectivity 1 (10%)
4	5	RBS 3 (60%), connectivity 1 (20%), RBS_orth 1 (20%)
5	1	connectivity 1 (100%)

reporting a weaker version of the other.

The instability is not a property of one selector pair. Repeating the 26-subset sweep with each of six further inner selectors of stability selection, against the same connectivity comparator, reproduced it in every case (Table 5): the three-cohort reproducibility difference changed sign under all seven selectors, in 1 to 5 of the 10 subsets, and with all five cohorts the connectivity comparator won on reproducibility while stability selection won on external AUROC under every selector. Random forest was the least stable, reversing in half of the three-cohort subsets. What the seven share is the comparator and the cohorts, so the instability belongs to the cohort set rather than to any selector, within the classical families tested.

The comparator side was checked the same way. Replacing the reimplementation by the WGCNA package itself, run with its tutorial defaults on every fold (settings in the note to Table 5), reproduced the reversal: the three-cohort difference spanned −0.081 to +0.261, and with all five cohorts the package won on reproducibility (−0.133) while stability selection won on external AUROC (+0.080). The package was the weaker comparator on both axes (mean Jaccard 0.064 and AUROC 0.711 over the 26 subsets, against 0.107 and 0.783 for the reimplementation), so the reimplementation did not flatter stability selection.

The practical consequence is that cross-fold agreement, the metric most often used to argue that a selection method produces reproducible signatures, is the metric least able to support that argument from a single cohort configuration. We therefore state what a benchmark on public omics cohorts should report alongside its headline figure: the range of the method difference over all admissible cohort subsets, whether that difference changes sign, how often each method wins at a fixed subset size with ties excluded, and the two metrics side by side rather than one standing for both. The patient-overlap audit and the control-procurement table are inputs to that protocol, not preliminaries to it. Without the overlap audit, the independence of admissible subsets cannot be assessed; without the control-procurement table, their sensitivity cannot be interpreted as a design finding. Table 6 collects these items, together with the controls the later sections add, as the checklist we recommend.

The most reproducible signal tracks tissue procurement, not disease

A note on what is and is not measured. No DKD cohort we examined reports warm or cold ischaemia time, fixation, or time to stabilisation. Procurement route is read from the sample source annotation (needle biopsy versus nephrectomy or donor tissue),




Figure 2. The reproducibility verdict depends on which cohorts are benchmarked; the predictive advantage stays positive. The benchmark was repeated over all 26 non-trivial subsets of the five discovery cohorts; the panels show the 16 of size three or more. Two-cohort subsets are excluded because they leave only two folds, so cross-fold agreement collapses towards zero and ties between methods cannot be broken. **a** Percentage of subsets in which each method wins on cross-fold reproducibility, by subset size, counting only subsets with a single top scorer. **b** The same comparison shown for every individual subset: stability selection minus the connectivity comparator in cross-fold agreement. The difference spans −0.106 to +0.170 and changes sign at every subset size except five. Custom co-expression hub comparator; not the reference WGCNA implementation. **c** The same subsets scored by external AUROC instead. Here the advantage is positive at every size and grows monotonically (+0.058, +0.096, +0.139 for sizes three through five). Reproducibility and predictive performance are therefore not interchangeable summaries of the same quantity.

Alt text: Three panels. Left, stacked bars by benchmark size (three, four, five cohorts) of the percentage of subsets in which each selection method wins on cross-fold reproducibility. Middle, one point per subset of stability selection minus the connectivity comparator in cross-fold agreement, spanning minus 0.106 to plus 0.170 with points on both sides of zero at every size except five. Right, the mean external AUROC difference with its range, positive at every size.

Table 5. Verdict instability under each inner selector of stability selection

Inner selector	Jaccard, 3 cohorts	Neg.	AUROC, 3 cohorts	Neg.	Jaccard, 5	AUROC, 5
ReliefF (headline)	−0.106 to +0.170	1	−0.027 to +0.153	2	−0.053	+0.139
Univariate *F*	−0.111 to +0.150	1	−0.098 to +0.150	2	−0.063	+0.101
LASSO	−0.127 to +0.119	2	−0.112 to +0.146	1	−0.090	+0.078
Elastic net	−0.163 to +0.119	2	−0.033 to +0.119	1	−0.106	+0.091
mRMR	−0.105 to +0.154	1	−0.096 to +0.129	2	−0.037	+0.088
Random forest	−0.211 to +0.074	5	−0.039 to +0.127	3	−0.144	+0.125
Boruta	−0.157 to +0.120	2	−0.106 to +0.106	3	−0.145	+0.111
Comparator replaced by the WGCNA package (ReliefF inner selector)						
WGCNA package	−0.081 to +0.261	1	−0.007 to +0.283	1	−0.133	+0.080

Stability selection minus the connectivity comparator, the inner selector of stability selection varied; the comparator is unchanged. Ten three-cohort subsets per row; Neg., subsets in which the difference is negative. Five, the single five-cohort benchmark. Rows above the rule use the custom co-expression hub comparator, not the reference WGCNA implementation. The last row replaces the custom comparator by the WGCNA package run with its tutorial defaults on every fold (soft power by scale-free fit, 4–20 here; unsigned TOM; dynamic tree cut; module most correlated with the trait; hubs by |kME|).

so the attribution to pre-analytical handling is an inference, not a measurement. It rests on three things: the association between procurement route and module level holds across nine datasets and four platforms; the module does not separate two groups that share a procurement route, even when one is diabetic and the other is not; and warm ischaemia is independently documented to induce exactly these transcripts within minutes, including in kidney [31], and ischaemia time is a documented driver of transcriptome variation across human tissues at scale [32]. Since this analysis was completed, an independent single-cell comparison of reference tissue sources within KPMP has reported the same dependence: stress-response genes are higher in nephrectomy and living-donor tissue than in percutaneous biopsies from healthy volunteers, and the choice of reference changes which genes are called differentially expressed [33]. We use "procurement-associated" throughout and reserve "artifact" for the interpretation, which the data support but do not prove.

Across nine kidney datasets the IEG module tracks how tissue was obtained, not disease (Table 7, Figure 3). The Spearman correlation between procurement (biopsy versus non-biopsy) and module level across all groups is $\rho = -0.752$, $p < 0.001$.

The KPMP single-nucleus data provide the decisive control because they contain both contrasts with donor-level replication (27 DKD-proxy donors as defined in the section *Cohort assembly and audit*, 10 non-diabetic CKD, 12 reference, 6 nephrectomy donors; statistics on donor × cell-type pseudobulk, Figure 4; the three contrasts are tabulated in Additional file 4).

GSE162830, the single dataset where all 32 samples were processed identically (OCT, −80 °C, laser microdissection), shows no procurement effect at all (−0.05), which is what the interpretation

Table 6. What a cross-cohort feature-selection benchmark on public cohorts should report, and where each item is applied here

Report	Because	Here
The method difference as a range over all admissible cohort subsets	a single configuration is one draw from that range	the section *Reproducibility verdicts depend on cohort composition, not on method*
Whether that difference changes sign across subsets	a sign change makes the verdict a statement about the cohorts	the section *Reproducibility verdicts depend on cohort composition, not on method*
Winner frequency at a fixed subset size, ties excluded	with few folds, ties inflate apparent wins	the section *Reproducibility verdicts depend on cohort composition, not on method*
Reproducibility and predictive performance side by side	the two agreed on the winner in only 11 of 24 subsets	the section *Reproducibility verdicts depend on cohort composition, not on method*
Every AUROC with its position in a random-signature null	random 50-gene sets reached 0.68–0.88	the section *Absolute AUROC is uninformative in this disease*
A patient-overlap audit across accessions	reused patients make subsets and folds non-independent	the section *Cohort assembly and audit*
Procurement route of cases and of controls, per cohort	collinearity with diagnosis is a design confounder no selector can remove	the section *The most reproducible signal tracks tissue procurement, not disease*
A confounder-matched contrast wherever one exists	it did more work here than any algorithm	the sections *Which corrections work*, *The control-group problem is not specific to diabetic kidney disease*
Pre-analytical variables to the SPREC or BRISQ standard	they turn the inference into a measurable covariate	unavailable in every cohort used

Table 7. Immediate-early module level by procurement route, across nine kidney datasets

Dataset	Biopsy	Non-biopsy	Difference
GSE142025	−0.63	+1.26	−1.90
GSE104954	−0.23	+1.37	−1.60
GSE175759	−0.21	+1.10	−1.31
GSE96804	−0.39	+0.80	−1.19
GSE104948	−0.15	+0.88	−1.03
KPMP snRNA	−0.33	+0.64	−0.97
GSE30529	−0.26	+0.22	−0.48
GSE30528	−0.19	+0.13	−0.32
GSE162830	+0.13	+0.18	−0.05

Values are the module mean z-score computed within each dataset.

predicts.

Because cases are biopsies and controls surgical tissue in every discovery contrast, this confounder is shared across cohorts and therefore maximally reproducible. A criterion that rewards cross-cohort consistency will rank it first: uncorrected RBS places 10 immediate-early genes in its top 50.

This is organ- and protocol-specific, not universal, and we report it in the main text rather than as supplementary material because it bounds the claim. In heart failure (GSE5406, 194 explanted ventricles versus 16 donor hearts) the module separates explant from donor only weakly ($g = -0.23$) and instead separates ischaemic from idiopathic cardiomyopathy, both explants ($p = 4 \times 10^{-8}$). Explanted and donor hearts undergo similar cardioplegia and cold storage, whereas needle biopsies and nephrectomies differ substantially in warm ischaemia. The confounder is a property of the procurement contrast, not of the gene module.

Why one method escaped the confounder, and why that is not robustness

This section exists to close a loophole in the section *The most reproducible signal tracks tissue procurement, not disease* rather than to open a second topic. If a widely used criterion simply avoids the confounder, the problem is less serious than we claim. It does avoid it here, but for a reason that does not generalise.

The connectivity comparator placed no immediate-early genes in its top 50 while stability selection placed 10. This exclusion alone does not establish robustness. In the real data 12 of 19 immediate-early genes are inside the disease-correlated module and have high gene–trait correlation (*DUSP1* $|gs| = 0.710$, among the highest in the module), yet rank 790th of 990 by connectivity, because intramodular connectivity is a *sum* of adjacencies and therefore rewards membership of the largest coherent block: the median block size (module genes with $|r| > 0.5$) is 19 for the immediate-early genes against 705 for the top 50 by connectivity, and 395 for




Figure 3. The most reproducible signal tracks tissue procurement, and it is organ-specific. a The immediate-early gene module, z-scored within dataset, for every diagnosis group in nine kidney datasets, ordered by effect. Biopsy-procured groups sit below nephrectomy- and donor-procured groups in eight of the nine (Spearman $\rho = -0.75$, $p < 0.001$); GSE162830, the one histology-matched design, is the exception. **b** The same module as Hedges' g for four contrasts. In KPMP, DKD against non-biopsy reference tissue gives −0.84 while DKD against other biopsies gives +0.08. In the heart-failure cohort GSE5406, disease against donor reaches only −0.23 and the within-procurement contrast (ischaemic against idiopathic) is +0.33. The kidney effect therefore appears when procurement differs and vanishes when it does not, and it does not transfer to heart, which bounds the claim to kidney procurement rather than to tissue injury in general.

Alt text: Two panels. Left, a dumbbell chart of nine kidney datasets in which biopsy-procured groups (orange) sit below nephrectomy- or donor-procured groups (blue) on the immediate-early module score in eight of nine; GSE162830 shows no gap. Right, horizontal bars of Hedges g for four contrasts: in KPMP, diabetic kidney disease against non-biopsy reference tissue minus 0.84 and against other biopsies plus 0.08; in heart failure, disease against donor minus 0.23 and ischaemic against idiopathic plus 0.33.




Figure 4. A biopsy-versus-biopsy contrast removes the effect. KPMP single-nucleus data reduced to donor-level pseudobulk. Comparing DKD with non-diabetic CKD, both procured by needle biopsy, gives $p = 0.21$; comparing DKD with non-biopsy reference tissue in the same resource gives $p = 6.0 \times 10^{-76}$. The contrast is the negative control for Figure 3: when procurement is held constant the signal disappears, and it is not a property of DKD.

Alt text: Box plots of the immediate-early module score in KPMP single-nucleus pseudobulk by donor. Diabetic kidney disease against non-diabetic chronic kidney disease, both needle biopsies, shows overlapping distributions (p equals 0.21); diabetic kidney disease against non-biopsy reference tissue shows a clear separation.

the whole module.

This is not an artifact of how the comparator was implemented. Rebuilding it across every combination of module definition (gene–trait correlation or clustering on the topological overlap matrix),




Figure 5. Connectivity criteria are biased by module size, which is why one method escaped. Simulation sweeping the ratio between the size of the disease block and that of the confounder block. **a** Percentage of the top 50 features that come from the confounder block. **b** The difference between a connectivity criterion and a stability criterion, which changes sign as the ratio crosses one. In the real data the immediate-early block contains 19 genes against 705 in the top-connectivity block, placing it in the regime where connectivity misses the confounder for reasons unrelated to robustness. All seven configurations are significant by Wilcoxon signed-rank on paired replicates.

Alt text: Two line panels over the ratio of disease-block size to confounder-block size in a simulation. Left, the percentage of the top 50 features drawn from the confounder block for each criterion as the ratio changes. Right, the difference between the connectivity and stability criteria, crossing zero where the ratio equals one. A marker shows the real-data regime, 19 immediate-early genes against 705 top-connectivity genes.

hub criterion (intramodular connectivity or module membership), adjacency (signed or unsigned) and soft power ($\beta \in \{4, 6, 8, 12\}$, spanning both our value and the $\beta \geq 10$ the scale-free criterion would select on these data) gives 32 variants, and *all 32 place no immediate-early gene in the top 50*. The WGCNA package itself, run with its tutorial defaults on every fold of the cohort sweep, does not: in 9 of 75 folds (3 of the 30 distinct training combinations the folds evaluate) it placed immediate-early genes in its top 50, up to 18 of 50, always where dynamic tree cut had produced a small trait module (61 to 199 genes against a median of 588). Because the package ranks by |kME| within the chosen module, this is consistent with module-dependent exclusion rather than a direct test of the summed-connectivity account; the median rank of an immediate-early gene within its module ranges from 664 to 3,531. Under a signed network the immediate-early genes are not module members at all, so the confounder is missed earlier still. Raising β improves their rank without approaching the top 50, so our lower β is conservative against our own argument. Additional file 1 reports the full grid.

Simulation confirms the mechanism and shows that it reverses (Figure 5; the paired differences are tabulated in Additional file 1).

The sign tracks the block-size ratio; effect-size and stability criteria show no such dependence (differences −0.025 to +0.022 across the same configurations). The connectivity comparator avoided the kidney confounder because that confounder happened to form a small block. Where a confounder forms a large programme, connectivity is the worst of the three.

Which corrections work

Corrections are compared on three axes at once: how much of the confounder survives, whether the surviving signal is DKD-specific, and what the correction costs in external AUROC (Table 8, Figure 6).

ComBat returns the same values as no correction in this implementation and evaluation setting. Cohort-level batch adjustment alone cannot disentangle procurement from disease when the two are confounded *within* cohorts. RUVg with empirically chosen controls also fails; the control set, not the framework, does the work. Residualisation and surrogate-variable adjustment both succeed. Table 8 reports identical fold-mean AUROCs to three decimal places for no correction and for residualisation. In a separate patient-stratified paired bootstrap analysis with 200 replicates, which resamples patients within each cohort and class and re-runs the whole leave-one-dataset-out evaluation in every replicate at a reduced inner bootstrap ($B = 30$ rather than the 150 of the fold means), the mean uncorrected minus residualised difference was +0.079 (95% percentile interval −0.048 to +0.229). This exploratory uncertainty estimate does not establish performance preservation: a substantial loss remains compatible with the interval, whose tails are imprecisely estimated.

The evaluation is not circular: the procurement flag derives from ERCB effect sizes, not from module membership, and correlation with the handling score separates flagged from unflagged genes at AUC = 0.480 (0.5 indicates independence).

The correction is not specific to the selector it was benchmarked with. Repeating the comparison with each of the seven inner selectors of the section *Selection methods*, at $B = 40$ rather than 150 because the question is the direction of the change rather than its third decimal, the procurement-flagged fraction of the top 50 falls for every one of them (Table 9), from 17.0–35.5% to 3.5–12.5%. The cost is not uniform. External AUROC rises for ReliefF (+0.021), falls by less than 0.05 for Boruta and mRMR, and falls by 0.05 to 0.11 for Elastic net, LASSO, Random forest and Univariate *F*. The removal is therefore a property of the correction, its cost a property of the selector, and a study adopting the correction should report that cost for the selector it uses.




Figure 6. What each correction removes, and what it costs. a Percentage of the top 50 selected features flagged as procurement-driven, lower being better. **b** Specificity pass-rate expressed as fold-enrichment over the 8.3% background rate at which a random gene passes the gate, so that no bar can be read as an absolute success rate. **c** External AUROC, showing the cost of correcting. Handling-score residualisation and surrogate-variable analysis reduce procurement-driven features from 27% to 5–6%; residualisation does so at a fold-mean AUROC identical to no correction to three decimal places, surrogate-variable analysis at 0.915 against 0.926. The separate paired bootstrap estimate of the residualisation cost is uncertain and is given in the text. ComBat returned the same values as no correction in this evaluation setting; the confounder is within-cohort rather than between-cohort.

Alt text: Three panels of horizontal bars for six correction strategies. Left, the percentage of the top 50 selected features flagged as procurement-driven: 27 percent with no correction or ComBat, 5 to 6 percent with handling-score residualisation or surrogate-variable analysis. Middle, specificity pass rate as fold enrichment over the 8.3 percent background. Right, external AUROC, 0.926 for no correction and residualisation and 0.915 for surrogate-variable analysis.

Table 8. Correction benchmark at $K = 50$

Correction	IEG in top 50	Procurement-flagged	Specificity pass	AUROC	Cross-fold
None	8.5	27%	33%	0.926	0.284
ComBat (batch = cohort)	8.5	27%	33%	0.926	0.284
RUVg, empirical controls	7.5	28%	26%	0.924	0.154
RUVg, module as controls	0.8	18%	40%	0.905	0.215
Surrogate-variable adjustment	2.0	6%	44%	0.915	0.251
Residualise on handling score	1.0	5%	40%	0.926	0.260

Specificity pass is the fraction of selected genes clearing the DKD-versus-other-CKD gate. Two rates are involved and they have different denominators. The *background pass rate* is 8.3%: 820 of the 9,900 genes in the frozen space clear the gate, so a random set clears it at that rate and the 40–44% figures are approximately 5× background. The gate's *permutation-based false-discovery estimate* is 16.7%: of those 820 passers, 137 are expected when the labels are permuted. The two ERCB blocks profile the same patients in two compartments, so the labels are permuted at the patient level; shuffling each block independently returns 13.7%, which understates the gate because it removes a dependence the real data has.

Table 9. The handling-score correction under each inner selector ($K = 50$, $B = 40$): procurement-flagged share of the selected features and external AUROC, uncorrected and corrected

Selector	Flagged, uncorrected	Flagged, corrected	AUROC, uncorrected	AUROC, corrected	Change in AUROC
Univariate *F*	24.5%	6.0%	0.921	0.807	−0.114
LASSO	33.0%	12.5%	0.956	0.879	−0.077
Elastic net	35.5%	3.5%	0.909	0.857	−0.052
mRMR	28.0%	6.5%	0.941	0.896	−0.046
ReliefF	29.5%	5.0%	0.928	0.949	+0.021
Random forest	17.0%	7.5%	0.895	0.815	−0.079
Boruta	22.0%	6.5%	0.918	0.901	−0.017

Each row is the mean over leave-one-dataset-out folds on the four headline cohorts. The ReliefF row repeats the comparison of Table 8 at the lower B; the small differences between the two rows are the bootstrap resolution, not a change in method.

The control-group problem is not specific to diabetic kidney disease

Everything above concerns one disease, which leaves open whether the effect is a property of diabetic kidney disease or of the design that every public kidney biopsy study shares. The ERCB compartments answer this inside a single resource, because they contain

Table 10. Genes at $|g| \geq 0.5$ before and after matching the control group, by diagnosis

Diagnosis	Patients	vs non-biopsy	vs other biopsy	Median $\lvert g \rvert$
IgA nephropathy	52	3,800	0	14%
Systemic lupus erythematosus	65	4,506	166	30%
Hypertensive nephropathy	35	3,251	9	30%
Diabetic nephropathy	29	5,107	833	37%
ANCA-associated vasculitis	43	5,878	1,981	42%
Minimal change disease	27	3,351	690	54%
Membranous glomerulonephropathy	38	3,282	702	56%

The last column is the ratio of median $|g|$ under the matched contrast to median $|g|$ under the naive one, a retained effect size rather than a retained count of genes, so it is not sensitive to the differing group sizes. All seven Wilcoxon tests on the paired effect vectors give $p < 10^{-300}$.

eight biopsy diagnoses alongside the same non-biopsy controls. We call the comparison that follows a *control- definition sensitivity analysis*: holding the case set fixed, it asks how far the ranking and the effect-size conclusions move when the control group is replaced by an alternative that is clinically defensible. It does not separate clinical from pre-analytical causes, and we do not claim it does; procurement is our reading of why the two definitions differ here, supported by the evidence in the section *The most reproducible signal tracks tissue procurement, not disease* rather than established by this comparison. For each diagnosis we formed the two contrasts the argument turns on, namely diagnosis versus non-biopsy control and diagnosis versus the remaining biopsy diagnoses, and counted genes reaching $|g| \geq 0.5$ (Table 10). Seven of the eight are reported; thin membrane disease was left out because its 9 cases give neither contrast usable power.

Shrinking is not the whole of it. A signal that fell uniformly would leave the ranking, and therefore the candidate list, unchanged; what happens instead is that the top of the ranking is rebuilt (Figure 7). Taking the 50 largest effects under each contrast, the two lists share 1 gene in diabetic nephropathy and between 0 and 18 across the seven diagnoses. The comparison holds the case set fixed and varies the control group; it is not a within-patient pairing, for which no public data exist. Rank correlation over all 9,900 genes is 0.69 for diabetic nephropathy, so the bulk of the transcriptome moves together while the extreme that feeds selection does not. The movement has direction: myeloid and inflammatory genes fall (*TYROBP* 31 to 678, *CASP1* 16 to 603) while matrix genes rise (*FMOD* 1264 to 4, *OLFML3* 5530 to 377). *MOXD1* moves from 204 to 40; that is a change in priority under a contrast with less procurement asymmetry, not evidence that it is a marker.

The consequence for a candidate list is direct. Running the same pipeline with and without the handling-score correction returns 30 candidates each, sharing only 14 (Additional file 3). Which genes a study nominates is therefore set by how its control tissue was obtained as much as by the disease.

Every diagnosis loses most of its signal, and diabetic nephropathy is unremarkable among them: it ranks fourth of seven, and the median across the other six is 36% against its 37%. In IgA nephropathy the 3,800 genes reaching $|g| \geq 0.5$ against non-biopsy controls fall to none. The problem therefore belongs to the design rather than to the disease, and the argument of The sections *The most reproducible signal tracks tissue procurement, not disease* and *Which corrections work* applies to kidney biopsy studies generally. One qualification carries over from the section *The most reproducible signal tracks tissue procurement, not disease*: the matched comparator here pools the remaining diagnoses, and pooling can itself produce a result, so these figures should be read as the pooled-comparator case.

Three datasets never used elsewhere in this work provide external checks of the candidate list. In GSE20602, which compares nephrosclerosis glomeruli against tumour nephrectomy, the same control type as GSE30528, the candidates move further in the diabetic contrast than in the nephrosclerosis one (median $|g_{\mathrm{DKD}}| - |g_{\mathrm{NSC}}|$ of +1.02 against −0.17 for the remaining genes, Mann–Whitney $p = 5.1 \times 10^{-18}$), which replicates the specificity gate outside ERCB. In GSE111154 the candidates are already altered in early diabetic nephropathy (permutation $p = 0.0015$), though with four patients per group this is weak. In GSE142153, blood from diabetic nephropathy patients, the candidates are indistinguishable from random genes ($p = 0.80$): the signature is tissue-restricted, which supports its specificity and limits its clinical accessibility at the same time.

The comparator behaves the same way in that metabolomic layer, which is the clearest evidence that the recommendation is not specific to transcriptomics. Replacing the healthy controls of ST003255 with the non-diabetic kidney disease patients of the same study drops the random-set AUROC from 0.913 to 0.724, and the selected set clears its null by a wider margin: 0.854 against a 95th percentile of 0.800, where against healthy controls it was 0.995 against 0.977. The reason is visible in the effect sizes. Across the 187 named metabolites the diabetic-versus-healthy and other-kidney-disease-versus-healthy effect vectors correlate at $r = 0.851$, so most of what separates these patients from healthy people is shared with kidney disease in general. Twenty-nine metabolites survive the matched contrast at $q < 0.05$. A disease-relevant comparator distinguishes diabetic-kidney-disease-associated changes from metabolic changes shared across kidney diseases; it does not establish the procurement mechanism seen in the transcriptome cohorts, which is a property of how those tissues were obtained and has no counterpart in plasma.

Both mechanisms outside kidney: predictions fixed before the data

The section *The control-group problem is not specific to diabetic kidney disease* showed that the control-group problem is not specific to diabetic nephropathy, but every diagnosis it could test was a kidney biopsy diagnosis. Whether verdict instability and a reproducible procurement confounder belong to cross-cohort benchmarking or to kidney cohorts needs tissues designed differently. Three predictions were fixed before any liver or colon expression value was read (the section *Predictions fixed before the liver and colon data were read*), and each verdict below is judged against its written wording (Table 11, Figure 8). The analyses that followed the verdicts were stated before they ran and are reported as post hoc.

Verdict instability (P1). In liver the difference in cross-fold agreement between stability selection and the connectivity comparator ranged from −0.008 to +0.174 across the ten three-cohort subsets, crossing zero in one of the ten subsets tested and narrowly, while the external AUROC difference stayed positive in all ten. That one reversal is small, so we asked whether it is an artefact of the resampling seed. Repeating the ten three-cohort comparisons under five bootstrap seeds reproduced it in the same subset (GSE89632 + GSE48452 + GSE83452) every time, at −0.017 to −0.008, while the other nine subsets stayed positive under all five seeds. The fifty runs are repeated computations on one dataset rather than fifty independent tests, so this supports robustness to computational




Figure 7. Changing the control-group definition rebuilds the top of the ranking. a Each line is one gene, drawn between its rank by $|g|$ when diabetic nephropathy cases are compared with non-biopsy controls (left) and its rank when the same cases are compared with biopsies of other diagnoses (right). Blue rises under the matched control, orange falls. The 46 genes drawn are those reaching either candidate list; 23 rise and 23 fall. Rank is on a log scale because the difference between first and twentieth matters for selection and the difference between five thousandth and five thousand five hundredth does not. **b** Genes shared by the two top-50 lists, computed over all 9,900 genes rather than the subset drawn in **a**, for each of the seven ERCB biopsy diagnoses. This is the same case set compared against two control groups, not a within-patient pairing; no public resource provides the latter. *MOXD1* rising from 204 to 40 is a change in priority under a contrast with less procurement asymmetry, not evidence that it is a marker.

Alt text: Two panels. Left, a slope chart of 46 genes drawn between their rank against non-biopsy controls and their rank against other-diagnosis biopsies in diabetic nephropathy, on a logarithmic rank axis; blue lines rise under the matched control and orange lines fall, 23 each. Right, horizontal bars of the number of genes shared by the two top-50 lists for seven ERCB biopsy diagnoses.

randomness and says nothing about patient-sample uncertainty. P1 therefore holds in liver with that scope: a small reversal, repeatable, in one configuration of ten. In colon the agreement difference never crossed zero (+0.006 to +0.192), and the prediction failed there. Post hoc, the tissues differed most in how far their cohorts agree about the disease: the median pairwise correlation of gene-wise effect sizes across the five core cohorts was 0.66 in colon against 0.22 in kidney and 0.15 in liver. Cohort size does not account for the difference. With each colon cohort subsampled to the case and control counts of a kidney cohort, concordance stayed between 0.60 and 0.65 in five replicates, and the agreement difference crossed zero in two of them, by at most 0.013, against a stated threshold of three (Figure 8c). The sign changes of The section *Reproducibility verdicts depend on cohort composition, not on method* were therefore associated with cohorts that disagree about the disease signal rather than with cohort size, which three tissues cannot raise to a necessary condition. Concordance can be computed before any selection method is run, and a low value warns that a single-configuration verdict is unlikely to transfer; three tissues are too few to treat it as a threshold.

The module at the cohort level (P2). Where cases were needle biopsies and controls were surgical or donor tissue, the immediate-early score was lower in cases in 9 of 10 cohorts across the two tissues, with liver GSE89632 at $g = -4.60$ [−6.11, −3.82] and GSE24807 at −2.83, both above the 99th percentile of random modules. The prediction nonetheless failed, because matched cohorts were not near zero. In colon, where both arms are endoscopic biopsies, the score was higher in inflamed cases in 9 of 10 cohorts (+0.51 to +3.43). In matched liver cohorts it was small in three of four (−0.28 to +0.40; GSE162694 +0.67), and in GSE48452, whose case arm mixes needle and intraoperative biopsies, it was −0.37. The module is therefore not a pure procurement readout: it also responds to active inflammation. Adjusting for the inflammation score shrank the colon case effect by a median of 49% and left every biopsy-versus-surgical cohort negative, all ten after adjustment against nine before (median standardised coefficient −0.94 before and −1.14 after; matched cohorts +0.89 and +0.30; Mann–Whitney $p = 1.6 \times 10^{-4}$ between the groups after adjustment). The kidney attribution survives this and is, if anything, strengthened: the one kidney cohort with a positive raw effect, GSE294519, turns negative once inflammation is accounted for.

Reproducibility of the module (P3). Over leave-one-dataset-out folds on each tissue's five core cohorts, the uncorrected stability top 50 contained a mean of 8.0 immediate-early genes in kidney,




Figure 8. Outside kidney, the module follows procurement where asymmetric and inflammation where matched. **a** Hedges' g of the handling score, case minus control, with 95% bootstrap intervals, for every cohort whose procurement could be coded. Above the dashed line, biopsy cases against surgical or donor controls; below, both arms obtained by the same procedure. Hollow points are not a second g: they are that g multiplied by the ratio of the adjusted to the unadjusted standardised case coefficient from the inflammation regression, which carries the adjustment onto the same axis. The two coefficients on their own scale are in Additional file 7. **b** Mean number of immediate-early genes in the stability-selection top 50 over leave-one-dataset-out folds on each tissue's five core cohorts, uncorrected and after residualising on the handling score. **c** Stability selection minus the connectivity comparator in cross-fold agreement for every three-cohort subset; orange points are subsets in which the verdict reversed. Custom co-expression hub comparator; not the reference WGCNA implementation. ρ is the median pairwise Spearman correlation of gene-wise effect sizes across the five core cohorts. The rightmost column repeats colon with each cohort subsampled to kidney case and control counts (five replicates of ten subsets). Filled points in **a**, panel **b** and the first three columns of **c** test the registered predictions; the inflammation adjustment, concordance and subsampling are post hoc.

Alt text: Three panels. Left, a dot plot of Hedges g for the handling score, case minus control, with 95 percent bootstrap intervals, one row per cohort, split into cohorts with biopsy cases against surgical or donor controls (above the dashed line) and cohorts with both arms obtained by the same procedure (below); hollow points show the inflammation-adjusted value. Middle, bars of the mean number of immediate-early genes in the stability-selection top 50 per tissue, with and without correction. Right, per-subset reproducibility differences for kidney, liver and colon with reversed verdicts highlighted.

0.6 in liver and 0.0 in colon, and residualising on the handling score removed 7.2 of them in kidney (Figure 8b). The two new tissues fail to put the module among the most reproducible genes for different reasons. Liver carries the largest procurement effects in this study, but only two of its five core cohorts are asymmetric, so the effect is not shared. Colon shares a direction of effect across cohorts, but there the module is one inflammation-responsive set among stronger ones. Relaxing the liver expression filter raised its coverage of the module from 11 to 15 genes and left the count at 0.6.

Control substitution in liver (post hoc). In GSE48452 the two control definitions rebuilt the top of the ranking as in ERCB: the 50 largest effects against surgical and against bariatric controls shared 2 genes, against 1 for diabetic nephropathy. The immediate-early effect moved in the expected direction (g = −0.38 [−1.45, +0.31] against surgical controls, −0.02 [−0.84, +0.61] against bariatric controls), but the intervals overlap and the overall signal barely shrank (median $|g|$ ratio 0.94, rank correlation 0.48 over all genes). With 12 and 16 controls, this comparison is recorded as unresolved.

Table 11. Predictions fixed before the liver and colon data were read, with the kidney reference computed by the same code

Prediction	Kidney, DKD	Liver, MASH	Colon, UC	Verdict
P1. The three-cohort RBS − connectivity difference in cross-fold agreement spans zero	−0.106 to +0.170 (1 of 10 negative)	−0.008 to +0.174 (1 of 10)	+0.006 to +0.192 (0 of 10)	held in liver, failed in colon
and the external AUROC difference changes sign in fewer subsets	−0.027 to +0.153 (2 of 10)	+0.040 to +0.377 (0 of 10)	−0.005 to +0.067 (1 of 10)	
P2. $\lvert g_{IEG} \rvert < 0.5$ in procurement-matched cohorts; negative, and larger than in every matched cohort, where cases are biopsies and controls surgical	7 asymmetric, 6 negative	3 asymmetric, all negative; 3 of 4 matched below 0.5	10 matched, 1 below 0.5, 9 positive	failed in both
P3. Immediate-early genes in the uncorrected stability top 50: kidney > liver > colon, colon at most 1; correction reduces the count substantially only in kidney	8.0 (7.2 removed)	0.6 (0.6 removed)	0.0	held

Wording abbreviated from the registered file (Additional file 7). The single negative liver subset recurs under five bootstrap seeds (−0.017 to −0.008), and the other nine stay positive in all five. Kidney values are a reference, not part of any prediction. By the same per-subset count the second clause of P1 would not hold in kidney either (2 AUROC sign changes against 1); the stability of the predictive verdict in the section *Reproducibility verdicts depend on cohort composition, not on method* rests on the mean advantage per subset size, which is positive at every size in all three tissues. The kidney P3 count is a mean over folds at the sweep's inner *B*, and so differs from the 10 of the section *Why one method escaped the confounder, and why that is not robustness*.

What the test changes is the scope of both claims rather than either claim. Verdict instability is not a property of five particular kidney cohorts, since it recurred in liver, but it is conditional on cohorts that disagree about effects. The procurement confounder appears wherever procurement is asymmetric and becomes reproducible where the asymmetry is shared; but the score used to detect and remove it also carries inflammation, so correcting on that score is tissue-conditional, and a control obtained by the same procedure is the design alternative to prefer where clinically defensible and available.

What the pipeline returned, and what happened to it

Applied without correction, the pipeline produced 30 specificity-gated candidates, of which two (*OLFML3*, *PRSS23*) had no DKD literature. Three independent tests removed them: the direction reversed in GSE175759, they were indistinguishable from random genes in the KPMP confounder-matched contrast, and the corrected selector did not rank them. The corrected selector instead recovered established DKD genes at a higher rate (20 of 30 already proposed as DKD markers versus 13 of 30 uncorrected), including *FMOD*, *LUM*, *MMP2* and *THBS2*; *FMOD* and *LUM* had previously been reported as DKD hub genes [34]. A germline test against a Korean DKD genome-wide association study (2,532 cases) with SNP-count-matched permutation nulls was a clean null (1 of 30 at empirical $p < 0.05$, expected 1.5; Kolmogorov–Smirnov $p = 0.74$).

Additional file 5 shows what stands behind each of the 30, and why none of them clears the bar. Two axes decide a biomarker claim: the gene must be strong in the data and it must not already be in the DKD literature. Six candidates are strong but reported, six are unreported but weak, and the cell requiring both is empty.

We report no novel biomarker. The corrected pipeline declining to promote its own uncorrected output is the intended behaviour.

Protein-level corroboration in two patient-independent datasets

The candidate list is transcriptomic. Whether any of it survives a change of molecule is a different question, and one the confounder analysis cannot answer. Two public cohorts measure protein abundance in human kidney tissue under a diabetic-versus-non-diabetic contrast, in patients unrelated to each other and to any cohort used here. Their control arms are not the same: one enrolled healthy participants, the other non-diabetic pathology cases (Additional file 6 records what each source states and what it leaves unstated). A SOMAscan aptamer panel of cortex biopsies (23 diabetic kidney disease, 10 healthy; Mendeley 83k89shdx5 [35]) carries 841 of our 9,900 genes and 7 of the 30 candidates. Mass spectrometry of archival formalin-fixed cortex (5 diabetic, 7 non-diabetic; PRIDE PXD041884 [36]) quantifies 1,142 and 7. The two overlap in only two candidates, so together they measure 12 of the 30.

The second cohort matters for a reason beyond sample size. Both of its arms are archival explant cases from one institution, so the biopsy-versus-nephrectomy contrast that dominates the transcriptome cohorts (the section *The most reproducible signal tracks tissue procurement, not disease*) is absent from its design. That is a weaker statement than procurement being matched: neither source reports ischaemia, fixation or storage time, so pre-analytical exposure is unmeasured in both, exactly as it is in the transcriptome cohorts.

Of the 12 candidates measured, 5 reach $q < 0.05$ in at least one cohort: *MMP2*, *MMP7*, *THBS2* and *CA10* on the aptamer panel, *MMP7* and *ACTN1* by mass spectrometry (Table 12). Only *MMP7* is significant in both, and only there does the direction match the transcriptome in two measurement principles at once. The other four rest on a single cohort each and should be read as such.

Two qualifications belong here. The first is about *MMP7*: it is the headline finding of the publication that produced the aptamer cohort [35], so recovering it there is not an independent observation: it is the same data reaching the same conclusion. What is independent is the mass-spectrometry cohort, where the same protein moves the same way in different patients on a different instrument.

The second is about what the mass spectrometer actually measured. Peptides are assigned to proteins, and some peptides belong to more than one, so a quantified value is not automatically specific to the gene it is labelled with. Of the 25 distinct peptides behind *ACTN1*, only 8 are unique to it; the remaining 17 are shared with *ACTN4*, *ACTN2* and *ACTN3*, and *ACTN4* is itself a podocyte gene, so that signal cannot be attributed to *ACTN1* alone. *MMP7* carries an

Table 12. Candidates measured at the protein level in two patient-independent datasets. Neither shares patients with the other or with any transcriptome cohort used here, and the two use different measurement principles. Transcriptome effect is the mean across the harmonised cohorts; protein effect and *q* are from the proteomic cohort named in each block.

Gene	Transcriptome *g*	Protein *g*	Protein *q*	Direction
SOMAscan aptamer panel; 23 DKD, 10 healthy (Mendeley 83k89shdx5)				
MMP2	+1.58	+1.71	0.008	agrees
MMP7	+1.82	+2.04	0.008	agrees
THBS2	+1.78	+1.31	0.013	agrees
CA10	−1.85	−1.05	0.046	agrees
LUM	+1.72	+1.00	0.057	agrees
TNFRSF12A	+0.95	+0.32	0.587	agrees
TNNT2	−1.68	+0.19	0.952	differs
LC-MS/MS of FFPE cortex; 5 DKD, 7 non-diabetic (PRIDE PXD041884)				
MMP7	+1.82	+3.34	0.045	agrees
ACTN1	+1.81	+1.78	0.046	agrees
CTNNBIP1	−1.42	+0.75	0.391	differs
RNASE6	+1.71	−0.59	0.542	differs
LUM	+1.72	+0.43	0.587	agrees
VCAN	+1.66	+0.42	0.589	agrees
COL1A2	+2.01	+0.39	0.614	agrees

Neither cohort is genome-wide. The aptamer panel carries 841 of the 9,900 genes in the frozen space and 7 of the 30 candidates; the mass-spectrometry cohort quantifies 1,142 and 7. Twelve distinct candidates are measured in total, two (*LUM*, *MMP7*) in both. Absence means the protein was not measured, not that it failed. Pooled across the two cohorts, candidates reach $q < 0.05$ more often than the proteins that are measured but not candidates (Cochran–Mantel–Haenszel $p = 0.030$; permutation without replacement $p = 0.035$; permutation with the background matched on abundance and peptide count $p = 0.039$).

unambiguous accession but rests on 2 distinct peptides. Additional file 6 reports aptamer identifiers, accessions and peptide counts for every candidate in the table.

Neither cohort is significant on its own (Fisher $p = 0.11$ and $p = 0.18$): seven candidates cannot separate a real enrichment from chance. Because their background rates differ widely (28% of measured non-candidates reach $q < 0.05$ on the panel, 11% by mass spectrometry), they must be pooled within strata rather than concatenated. Stratified, candidates reach $q < 0.05$ more often than the proteins measured alongside them: over the 14 candidate measurement events, the Mantel–Haenszel common odds ratio against the stratum-specific backgrounds is 3.3 (95% CI 1.1–10.2), an interval wide enough that the size of the effect is not established even though it excludes 1 (Cochran–Mantel–Haenszel [37] $p = 0.030$; permutation over 200,000 draws without replacement within each stratum $p = 0.035$, against a null expectation of 2.7 significant candidates rather than the six observed). A candidate set could reach that simply by being made of abundant, readily detected proteins, so the permutation was repeated drawing each null protein from the background matched to that candidate on abundance rank and, for the mass-spectrometry cohort, on peptide count: $p = 0.039$ (Additional file 6 reports the achieved balance). Direction agreement alone does not separate them ($p = 0.19$), which is expected: 60–67% of background proteins already agree in direction.

The analysis plan (candidate versus measured non-candidate, Benjamini–Hochberg [38] $q < 0.05$, direction as a separate test) was fixed on the first cohort and applied unchanged to the second, which was located by a systematic search of PRIDE rather than chosen for its result. Both cohorts are reported in full, including the two non-significant single-cohort tests and the null direction test.

None of the five is novel. *MMP2* and *THBS2* are tier 2; *MMP7*, *CA10* and *ACTN1* tier 4. *MOXD1*, the tier-2 candidate with the thinnest diabetic kidney disease literature (1 paper, none proposing it as a marker, against 2 to 147 papers for the other five in that tier), is on neither dataset. KPMP does quantify it in kidney tissue by mass spectrometry, so it is measurable; it is simply not measured anywhere the contrast is diabetic versus non-diabetic. The layer that could have promoted it does not do so.

Discussion

One boundary should be stated before anything else in this section. What follows is a warning about cross-cohort omics benchmarking in general, but the quantification behind it is concentrated in one disease, five discovery cohorts, with the two mechanisms then tested against written predictions in two further tissues (the section *Both mechanisms outside kidney: predictions fixed before the data*). That test turned both from kidney observations into conditional statements. Verdict instability recurred in liver but not in colon, whose cohorts agree about gene-wise effects far more than kidney or liver cohorts do; the procurement confounder appeared wherever procurement was asymmetric and was prominent among the most reproducible genes only where the asymmetry was shared by every cohort. The connectivity blind spot does not depend on the disease either, as the simulation in the section *Why one method escaped the confounder, and why that is not robustness* shows. The magnitudes do. A reader should take the numbers below as measurements in three tissues, and the two conditions, low effect concordance and a design shared across cohorts, as hypotheses testable by the same procedure wherever more than two cohorts exist.

Cross-cohort consistency is a proxy for validity only when the cohorts differ in the ways that matter. In public DKD transcriptomics they do not: every discovery contrast compares biopsies against surgical or donor tissue, so the strongest shared signal is one associated with procurement, a dependence on reference tissue source that has since been observed independently in single-cell data [33]. This is a batch effect in the sense of Leek et al. [6], with the aggravating feature that it is confounded with the outcome by design rather than merely correlated with it. Any criterion that rewards consistency will find it, and no amount of additional cohorts of the same design will reveal the problem: more cohorts make the confounder look more reproducible, not less.

The instability of benchmark verdicts has a simpler cause and a broader reach. With five cohorts there are ten three-cohort configurations, and the sign of the reproducibility comparison differs between them. That benchmark rankings shift with benchmark composition is not itself a new observation [7, 8, 9, 10, 12]; what these data add is the magnitude in a setting where the practice of reporting a single configuration is universal, and the observation that

the two metrics behave differently: reproducibility flips sign, while the mean external AUROC advantage does not. The cross-tissue test adds an association. The flips went with cohorts that disagree about the disease signal: in colon, at a median effect correlation of 0.66, reversals were rare and small even after subsampling to kidney cohort sizes (two of five replicates, at most 0.013). Three tissues cannot establish this as a necessary condition. Cross-cohort comparisons of reproducibility should be treated as descriptions of a particular cohort collection rather than of the methods.

Practically, three things follow. Report reproducibility and validity as separate axes, since they came apart measurably here. Include a confounder-matched contrast in the design: the 310 non-diabetic CKD biopsies in ERCB did more work than any algorithm in this study, and only two of the public resources we used, ERCB and KPMP, contain such a comparator. And report pre-analytical variables (warm and cold ischaemia time, fixation, and storage) in the SPREC [39, 40] or BRISQ [41] format. No DKD cohort we examined reports ischaemia time, which is why the attribution in the section *The most reproducible signal tracks tissue procurement, not disease* has to remain an inference; had these variables been recorded, the confounder would be a covariate rather than a conjecture.

The handling score has an incidental use as a quality-control metric. It is computable per sample from expression alone, requires no labels, and in these data tracks procurement route closely enough to flag heterogeneous batches.

The cross-tissue test qualifies that use and the correction built on it. The module responds to active inflammation as well as to handling: in colon, where both arms are endoscopic biopsies, it rose in inflamed cases in 9 of 10 cohorts. As a quality-control flag it is interpretable only alongside an inflammation score, and residualising on it in an inflamed tissue would remove disease biology together with the artefact. In kidney that risk is bounded, since adjusting for inflammation made the biopsy-versus-surgical effect larger rather than smaller. It is nonetheless why we regard the score-based correction as tissue-conditional, and a control obtained by the same procedure as the design alternative to prefer where clinically defensible and available.

Limitations

The empirical comparisons cover a limited set of established selection strategies, including a custom co-expression hub comparator, rather than an exhaustive benchmark of contemporary methods. The reversals recur under seven classical inner selectors and when the custom comparator is replaced by the WGCNA package (Table 5), so they are not an artefact of the reimplementation; the exclusion of the immediate-early module, by contrast, held for the reimplementation and its 32 variants but not for the package on every fold, and appears to depend on module detection rather than to be a property of connectivity criteria as a class. The magnitudes reported apply to the implementations and cohort configurations examined here and should not be generalised to entire method families. The audit procedure can be applied to additional selectors, but their sensitivity to cohort composition remains to be evaluated.

The confounder-matched evidence rests on ERCB and KPMP; no other public DKD resource provides non-diabetic kidney disease biopsies procured identically to the cases, so the specificity argument rests on two resources. A partial external check is available and reported in The section *The control-group problem is not specific to diabetic kidney disease*. The gate's background pass rate (8.3%) and its permutation-based false-discovery estimate (16.7%) are different quantities, as the footnote to Table 8 sets out.

GSE104948 contributed both to discovery and to the matched-control specificity gate, so that gate is not a fully independent validation. Re-deriving the candidates with GSE104948 removed from discovery changes which genes are returned: 17 of the 30 survive, among them *FMOD*, *LUM*, *MMP7* and *MOXD1*. It does not change the conclusions drawn from the gate. The rate at which ranked genes clear the gate is no lower once the gate no longer shares a cohort with discovery (pipeline stage `sens_no_ercb_check`), and the external replication holds (candidate median $|g_{\mathrm{DKD}}| - |g_{\mathrm{NSC}}|$ of +1.00 against +1.02, Mann–Whitney $p = 4.8 \times 10^{-16}$ against 5.1×10^{-18}). The overlap is real and partial; it is not what produces the specificity result.

The procurement effect did not replicate in heart, where explanted and donor hearts are handled similarly, but did appear in liver cohorts with biopsy cases and surgical or donor controls. Our simulation model is deliberately simple and does not capture platform effects or realistic correlation structure beyond block correlation. The five discovery cohorts remain few for LODO-based claims; the cohort-sensitivity analysis quantifies rather than removes that limitation.

The cross-tissue test has limits of its own. Only three liver cohorts had asymmetric procurement, one of them with a mixed case arm, and the liver core gene space retained 11 of the 19 module genes. Procurement outside kidney was coded from series text and source publications, not from recorded sample handling. Two predictions failed as written, and the explanations offered for the failures, inflammation and effect concordance, were examined only post hoc and in three tissues, too few to establish concordance as the condition for instability rather than one correlate of it. The liver control substitution rests on 12 and 16 controls and is unresolved.

The central design claim rests on single-modality transcriptomics; the cross-modal and germline analyses support or bound it and are not its basis. Three things the phrase multi-omics can mean are supported to different degrees by public data, and only one is claimed here. Linking layers measured in the same participant could not be constructed from the public resources analysed. Cross-modal corroboration, which checks a candidate list in another layer in unmatched patients under a similar rather than identical comparison, is what the section *Protein-level corroboration in two patient-independent datasets* does, against healthy participants in one cohort and non-diabetic pathology cases in the other. Within-layer discovery in the protein layer is not attempted, and Additional file 6 shows why it cannot yet be.

The obstacle to the first is worth stating precisely, because it is not that measurements are never paired. Of the 80 KPMP single-nucleus donors, the 27 in the operational DKD-proxy subset of the section *Cohort assembly and audit* include 3 who also carry regional proteomics and 7 who carry spatial metabolomics. What is missing differs by layer. For regional proteomics it is the control arm: no participant in any control category carries it (0 of 12 healthy reference), so the contrast cannot be formed however many cases are paired. For spatial metabolomics both arms exist, 7 proxy and 6 healthy reference participants alongside transcriptomics, and the obstacle is the released format: imaging mass spectrometry distributed as image data rather than a per-participant matrix, behind a public API that exposes file counts without a retrieval path. No public resource therefore carries a multi-layer design complete enough to form a diabetic-versus-control contrast, and counting enrolment categories rather than recorded histories would have reached that conclusion for the wrong reason.

Potential implications

A cross-cohort reproducibility comparison in omics biomarker discovery is a statement about a particular collection of cohorts, not about the methods compared, unless it is accompanied by a cohort-composition sensitivity analysis. In diabetic kidney disease the reproducibility verdict changes sign across legitimate subsets of the same five cohorts, while the predictive verdict does not, so the two should be reported as separate axes rather than as interchangeable summaries of robustness. In a test against predictions fixed in advance, the same instability recurred in liver but not in colon,

whose cohorts agree closely about effects, so cross-cohort effect concordance, computable before any method is run, indicates when a single configuration is least trustworthy.

The signal that survives such a comparison most reliably in this disease is associated with tissue procurement rather than with disease, because every discovery contrast compares biopsies with surgical tissue. That is not a peculiarity of diabetic kidney disease: substituting a matched comparator for the non-biopsy control removes most of the signal in all seven ERCB biopsy diagnoses we could test, and diabetic nephropathy is unremarkable among them. The same module was lower in biopsy cases than in surgical controls in liver, and was prominent among the most reproducible genes only in kidney, where every cohort shares the asymmetry (8.0 genes against 0.6 in liver and 0.0 in colon). Whether a given method filters that confounder depends on the size of the block it forms, not on the method's robustness. Residualisation on a specified control set and surrogate-variable adjustment both remove most of it, at no significant cost in predictive performance for the selector benchmarked here and at a cost that varies from none to about 0.1 AUROC across seven selectors, whereas correcting for batch by cohort cannot, because the confounder lies within cohorts rather than between them. Because the module also rises with inflammation, correcting on it is tissue-conditional, and a control obtained by the same procedure should be preferred where clinically defensible and available.

What the correction buys is worth stating plainly, because it is not what a reader might expect. It changes which genes are reported, not whether a biomarker can be reported at all. Only 14 of the 30 candidates are the same genes before and after, and the corrected list moves towards established biology, recovering 20 rather than 13 genes already proposed as diabetic kidney disease markers. What it does not do is produce a defensible novel candidate: the cell requiring a gene to be both strong in the data and absent from the literature is empty in both lists. The value of removing this confounder is therefore diagnostic rather than generative. It tells a reader that a list assembled without the correction is substantially a different list, and that the difference does not run in the direction of novelty, which is the more useful thing to know before any candidate set is carried forward.

We therefore recommend that studies of this kind report a cohort-composition interval alongside any reproducibility claim, include a confounder-matched contrast in the design, and report pre-analytical variables to the SPREC or BRISQ standard, which would convert the confounder described here from an inference into a measurable covariate.

Methods

The study has two layers. The first layer is a conventional cross-cohort benchmark: retrieve public cohorts, audit them for patient reuse, harmonise them into one frozen gene space, and compare four selection strategies under leave-one-dataset-out evaluation. The second layer is the set of controls placed around that benchmark, namely the random-signature and permuted-label nulls, the sweep over cohort subsets, the procurement contrast, and the substitution of the control group. Cohort assembly and harmonisation are described under Data description (the sections *Cohort assembly and audit* and *Harmonisation*). The subsections below cover the four strategies (the section *Selection methods*), the nulls and the sensitivity sweeps (the section *Null controls and uncertainty*), the confounder score and the corrections benchmarked against it (the section *Confounder score, corrections and simulation*), the test of both mechanisms in liver and colon (the section *Predictions fixed before the liver and colon data were read*), and how each reported number is regenerated from source data (the section *Implementation*).

Selection methods

Four strategies were compared, all on the same gene space and the same folds.

DEG_meta. Per-cohort Welch t-statistic combined across cohorts by Stouffer's method weighted by $\sqrt{n}$ [42]. The plain effect-size approach.

WGCNA_hub. Soft-thresholded correlation network, module defined by gene–trait correlation, genes ranked by intramodular connectivity: a connectivity-based hub ranking in the manner of WGCNA [43], of the kind used by prior DKD work [34]. It is implemented directly rather than called from the reference package, and it departs from the canonical pipeline in three respects: the soft power is fixed at $\beta = 6$ rather than chosen by the scale-free topology criterion, modules are defined by gene–trait correlation rather than by dynamic tree cut on the topological overlap matrix, and hubs are ranked by intramodular connectivity rather than by module membership. We do not claim this reproduces WGCNA. Table 13 states element by element what is shared and what is not; Additional file 1 rebuilds the comparator across those axes and shows that the ranking behaviour the section *Why one method escaped the confounder, and why that is not robustness* rests on is unchanged in every variant.

RBS. Cross-cohort stability selection (robust biomarker score), used here as an instrument rather than proposed as a contribution: it stands in for the family of stability-selection approaches [2] so that the benchmark contains a representative of that family, and no claim of novelty attaches to it. Bootstrap selection is run inside each cohort separately and aggregated across cohorts, never on the pooled matrix. In the headline benchmark the inner selector is ReliefF, run on $B = 200$ class-stratified bootstrap replicates per cohort at $K = 50$. Let π_{cj} be the fraction of those replicates of cohort c in which gene j is selected. The four components are importance I_j, the selector score of gene j divided by the largest score in the same replicate, averaged over replicates and cohorts, so that a gene the selector never touches scores zero; stability $S_j = M^{-1}\sum_c \pi_{cj}$; cross-cohort reproducibility R_j; and perturbation robustness P_j, the selection frequency of gene j over 40 refits per cohort under class-stratified 70% subsampling, Gaussian noise at half the gene-wise standard deviation, masking of 20% of genes, and a synthetic batch shift, computed only for the 300 genes ranked highest by $I_j S_j R_j$ so that perturbation stress-tests candidates rather than discovers them:

$$
\begin{aligned}
R_j &= \delta_j \exp\Big(\frac{1}{M}\sum_c \log(\pi_{cj} + \epsilon)\Big), \\
\mathrm{RBS}_j &= \delta_j \exp\Big(\frac{\sum_k w_k \log(V_{kj} + \epsilon)}{\sum_k w_k}\Big),
\end{aligned}
\tag{1}
$$

where $V_{\cdot j} = (I_j, S_j, R_j, P_j)$, $w = (1, 1, 2, 1)$, $\epsilon = 10^{-3}$ inside R_j and 10^{-6} in the combination, and $\delta_j \in \{0, 1\}$ is the sign-concordance gate, equal to 1 only when g_{cj} carries the same non-zero sign in every cohort. The geometric mean requires a gene to be adequate on all four components rather than to buy its way in on one, and the gate is hard rather than a soft penalty. Baseline selectors within the bootstrap were compared (univariate F, LASSO [44], elastic net [45], mRMR [46], ReliefF [47], random forest [48], Boruta [49]), all implemented directly rather than through packages that pin older `numpy`/`scikit-learn` versions.

RBS_orth. RBS after residualising every gene on a per-sample handling score (the section *Confounder score, corrections and simulation*).

Two cohort sets are used and the distinction matters when reading the tables below. The headline benchmark and its nulls run on the four cohorts fixed before the analysis began (GSE30528, GSE96804, GSE104948, GSE142025). The cohort-composition

Table 13. What the correlation-module comparator shares with the canonical WGCNA pipeline, and where it departs. Additional file 1 tests whether any departure changes a conclusion.

Element	Canonical WGCNA	Used here	Status
Similarity measure	Pearson correlation	Pearson correlation	same
Adjacency	signed or unsigned power	unsigned power	restricted
Soft-threshold power	chosen by scale-free fit	fixed at $\beta = 6$	differs
Module detection	TOM distance, dynamic tree cut	clustering on correlation distance	differs
Hub score	kME, eigengene correlation	intramodular connectivity kIN	differs
Module eigengene	first principal component	not computed	absent

The departures are why the manuscript calls this a correlation-module comparator rather than WGCNA. We do not claim it reproduces the reference package; the package itself was also run as a second comparator (the section *Reproducibility verdicts depend on cohort composition, not on method*).

sweep of the section *Reproducibility verdicts depend on cohort composition, not on method* adds GSE294519 as a fifth, because that analysis exists precisely to vary the cohort set and a wider set of compositions makes it stronger.

Evaluation is leave-one-dataset-out (LODO). Write the harmonised cohorts as $D_1, \ldots, D_M$. For held-out cohort m the training set is $T_m = \bigcup_{j \neq m} D_j$; a selection strategy S returns the top-K gene set $G_m^{(K)} = S(T_m, K)$; a plain L_2 logistic model f_m is fitted on T_m restricted to those genes and assigns each sample $i \in D_m$ a score $s_i = f_m(x_i)$. The classifier is deliberately simple because the object of study is the features, not the classifier.

Let P_m and N_m be the case and control samples of D_m, with $n_1 = |P_m|$ and $n_0 = |N_m|$. External AUROC is the Mann–Whitney statistic, ties counted as half:

$$\mathrm{AUROC}_m = \frac{1}{n_1 n_0} \sum_{i \in P_m} \sum_{j \in N_m} \left[\mathbf{1}(s_i > s_j) + \tfrac{1}{2}\mathbf{1}(s_i = s_j) \right], \tag{2}$$

and the reported value is the mean over folds, $\overline{\mathrm{AUROC}} = M^{-1} \sum_{m=1}^{M} \mathrm{AUROC}_m$. The tie term matters here because several cohorts are small enough for scores to tie exactly.

The second axis is reproducibility, the agreement between the gene sets selected on different training sets:

$$J = \binom{M}{2}^{-1} \sum_{m<m'} \frac{|G_m^{(K)} \cap G_{m'}^{(K)}|}{|G_m^{(K)} \cup G_{m'}^{(K)}|}. \tag{3}$$

The two axes are reported separately throughout and are not interchangeable: the section *Reproducibility verdicts depend on cohort composition, not on method* shows they disagree.

Effect size. Every effect size reported here is Hedges' g [50], the standardised mean difference with the small-sample correction. For gene j with case and control groups of size n_1 and n_0,

$$\begin{aligned} g_j &= \frac{\bar{x}_{1j} - \bar{x}_{0j}}{s_{p,j}} \left(1 - \frac{3}{4(n_1 + n_0) - 9}\right), \\ s_{p,j}^2 &= \frac{(n_1 - 1)s_{1j}^2 + (n_0 - 1)s_{0j}^2}{n_1 + n_0 - 2}. \end{aligned} \tag{4}$$

The specificity gate keeps genes with $|g_j| \geq 0.5$ in the diabetic-versus-other- kidney-disease contrast, averaged over the two ERCB compartments.

Null controls and uncertainty

Two nulls are computed per fold and per signature size. For the random-signature null, $B = 300$ gene sets $R_1, \ldots, R_B$ of the same size K are drawn uniformly from the same frozen gene space, and the model is refitted and rescored on the same fold with real labels. An observed AUROC is reported as its position in that null,

$$\pi_m = \frac{1}{B} \sum_{b=1}^{B} \mathbf{1}\left(\mathrm{AUROC}_m^{(R_b)} < \mathrm{AUROC}_m\right), \tag{5}$$

never as the raw value. The second null permutes the case/control labels within each cohort and leaves everything else unchanged, so it measures what the evaluation returns when no signal exists at all. This control follows Venet et al. [51], who showed that most random signatures are significantly associated with outcome in breast cancer.

All method comparisons use a paired patient-level bootstrap of $R = 40$ replicates, stratified by cohort and class, with the entire LODO evaluation, selection included, re-run inside each replicate. The interval is formed on the paired difference computed within a replicate,

$$\begin{aligned} \Delta^{(r)} &= \overline{\mathrm{AUROC}}_A^{(r)} - \overline{\mathrm{AUROC}}_B^{(r)}, \\ r &= 1, \ldots, R, \end{aligned} \tag{6}$$

and quoted as the 2.5th and 97.5th percentiles of $\Delta^{(1)}, \ldots, \Delta^{(R)}$. Pairing within the replicate is what makes the interval informative: the two methods see identical patients, so the cohort-to-cohort variance that dominates the marginal AUROC cancels. Resampling folds (3–5 units) or the pairwise Jaccard values (which are correlated) does not support an interval. R is small for a percentile interval because each replicate repeats the whole evaluation, selection included; it is set by compute, and the intervals should be read as bootstrap ranges whose finest resolvable two-sided p is $2/R = 0.05$.

Cohort-composition sensitivity is assessed exhaustively: every subset of the five discovery cohorts of size 2–5 (26 subsets) receives a complete benchmark, with 30 inner bootstrap replicates per cohort rather than the 200 of the headline run, since the sweep asks which method wins rather than for its score to the third decimal.

Confounder score, corrections and simulation

The immediate-early gene (IEG) module (*FOS*, *FOSB*, *JUN*, *JUNB*, *EGR1*, *ATF3*, *DUSP1*, *ZFP36*, *NR4A1*, *NR4A2*, *BTG2*, *EGR2*, *EGR3*, *IER2*, *KLF2*, *KLF4*, *SOCS3*, *DUSP2*, *JUND*) is a per-sample, label-free proxy for tissue handling; its mean z-score over that module is the handling score h_i of sample i.

Residualisation removes that axis from every gene. The slope is estimated *within* class, not over all samples: because cases are biopsies and controls are not, h is nearly collinear with case status, and a total slope would absorb the disease difference itself. Writing $\tilde{h}$ and $\tilde{x}_j$ for h and gene j centred inside each class,

$$\beta_j = \frac{\tilde{h}^\top \tilde{x}_j}{\tilde{h}^\top \tilde{h}}, \qquad x_{ij}^{\mathrm{corr}} = x_{ij} - \lambda(h_i - \bar{h})\beta_j, \tag{7}$$

with $\lambda = 1$ for full removal. Corrections compared: no correction; residualisation on the handling score by Eq. (7); RUVg with the module as negative controls; RUVg with empirically chosen controls (the least associated genes) [52]; a surrogate-variable adjustment protecting the design [53]; and ComBat with cohort as batch [54].

The simulation uses a generative model with a disease block and a confounder block, each with within-block correlation, cohort-specific effect heterogeneity (σ_d, σ_c drawn once per cohort), and a confounder correlated with case status by ρ. Block sizes, ρ, heterogeneity and cohort count are swept. The stability arm was run both with a simplified stand-in and with the actual RBS implementation; conclusions were identical.

Predictions fixed before the liver and colon data were read

The two findings make different predictions outside kidney. Instability of reproducibility verdicts under a change of cohorts is a statistical claim and should appear in any tissue; a procurement confounder should become reproducible only where every cohort shares the same asymmetry between arms. Two tissues separate these. In liver with metabolic dysfunction-associated steatohepatitis (MASH), procurement is asymmetric in some cohorts (needle biopsy cases, living-donor or surgical controls) and matched in others (both arms needle biopsies, or both intraoperative at bariatric surgery); in ulcerative colitis (UC), both arms are endoscopic biopsies in every cohort. Three predictions (the section *Both mechanisms outside kidney: predictions fixed before the data*) were written after reading only series- and sample-level metadata and before any expression value was downloaded, and the file's SHA-256 digest and time stamp were recorded when it was written. A second file stating post hoc analyses was written and hashed after the verdicts were known and before those analyses ran. A pipeline stage verifies both digests on every run, and both files are reproduced in Additional file 7. The digests are local records: they establish that neither file changed after it was written, not an externally certified date, since no third-party registry entry exists. We therefore call these predictions stated and hash-fixed before the analysis.

Cases were MASH against histologically normal liver and actively inflamed UC colon against normal control colon, one sample per subject, with ileal and Crohn's disease samples excluded. Procurement was coded per arm from the series text, the sample fields and the source publication, with its evidence recorded, before any effect was computed. In each tissue five core cohorts, matching the five kidney discovery cohorts, entered the composition sweep; the remaining eligible cohorts entered only the per-cohort module test (Additional file 7). Patient reuse across series from the same group or platform was audited on the expression values themselves: a sample whose best-correlated partner in another series reached $r \geq 0.9999$ was called identical, and a pair of series whose median gap between the best and second-best partner correlation exceeded five times that of an unrelated same-platform pair was called reprocessed from shared individuals. The audit confirmed GSE59071 as a subset of GSE75214 (116 identical samples) and GSE17470 as overlapping GSE24807 (9 identical), and both were excluded; it also found GSE48452 and GSE61260 likely to share individuals (median gap 0.0109 against 0.0016), and GSE61260, whose deposited matrix could not be mapped to samples, was not analysed.

Microarrays were mapped to Entrez identifiers through the platform annotation, RNA-seq counts were transformed to $\log_2(\mathrm{CPM} + 1)$ after keeping genes with CPM ≥ 1 in at least 20% of samples, and probes were collapsed by maximum mean, as for kidney. Selection used the genes shared by each tissue's five core cohorts: 7,805 in liver, including 11 of the 19 immediate-early genes, and 15,789 in colon, including all 19. The four strategies, $K = 50$, the inner selector, the soft-thresholding power and the inner $B = 30$ of the kidney composition sweep were reused by calling the same code, and the kidney reference was recomputed with it. The per-cohort effect g_{IEG} is Hedges' g of the handling score, case minus control, with a 95% interval from 1,000 bootstrap resamples within arms and a percentile against 2,000 random gene modules of the same size.

Four post hoc analyses followed the plan. Control substitution was repeated in GSE48452, which contains MASH cases with two control groups: liver from oncological surgery and histologically normal liver from obese individuals at the same bariatric operations as most cases. The immediate-early effect was adjusted for inflammation by regressing the standardised handling score on case status within each cohort, with and without an inflammation score fixed in advance from 13 genes sharing none with the module (*PTPRC*, *CD14*, *CD68*, *FCGR3A*, *S100A8*, *S100A9*, *LCN2*, *CXCL8*, *IL1B*, *TNF*, *SAA1*, *MMP3*, *CHI3L1*). Cross-cohort effect concordance was defined as the median pairwise Spearman correlation of gene-wise g across the five core cohorts, and colon was re-run with each core cohort subsampled without replacement to the case and control counts of a kidney discovery cohort, in five replicates. Finally, the liver RNA-seq filter was relaxed to a raw count of at least 1 in 20% of samples to test whether the liver module coverage affected the result.

Implementation

All analyses are implemented in Python 3.14 against `numpy`, `scipy`, `pandas`, `scikit-learn` and `matplotlib`, with pinned versions. The pipeline runs from a single entry point covering 133 declared stages, of which 122 reproduce the results reported here; every stage that resamples takes a fixed seed, and reruns reproduce the reported numbers, with the run-time dependence on external reference resources recorded under Regeneration below.

Every comparator is implemented directly rather than called from a reference package, for reasons of speed and single-environment reproducibility set out in Additional file 1. That file also states, for each comparator, what it shares with its reference definition and where it departs, and reports the variant grid establishing that no conclusion here depends on those departures. Readers assessing the comparisons in the sections *Reproducibility verdicts depend on cohort composition, not on method* and *Why one method escaped the confounder, and why that is not robustness* should read it alongside them.

Availability of source code and requirements

- **Project name:** dkd-cross-cohort-audit
- **Project home page:** https://github.com/kdh6126/dkd-cross-cohort-audit
- **Operating system(s):** Platform independent (developed and run on Windows 11)
- **Programming language:** Python 3.14; R 4.6.1 for the reference WGCNA comparator only
- **Other requirements:** Python packages pinned in requirements.txt (numpy 2.5.2, scipy 1.18.1, pandas 3.0.5, scikit-learn 1.9.0, matplotlib 3.11.1, pyarrow 25.0.1, h5py 3.16.0); R package WGCNA 1.74; a LaTeX distribution to rebuild the manuscript
- **License:** MIT
- **Restrictions to use by non-academics:** None
- **Archived version:** https://doi.org/10.5281/zenodo.23051697 (version 1.0.0)

Data availability

The data sets supporting the results of this article are available in the Zenodo repository, https://doi.org/10.5281/zenodo.23051697 [55], with the primary public records listed below.

All data supporting the conclusions of this article are publicly available, and all code required to regenerate every figure and table is openly released.

Primary data. All transcriptome cohorts analysed here are available from the Gene Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/) under accessions GSE30528, GSE30529, GSE96804, GSE104948, GSE104954, GSE142025, GSE294519, GSE162830, GSE175759, GSE142153, GSE1009, GSE111154, GSE20602 and GSE5406. The three series excluded by the cohort audit for patient-level reuse are GSE30122, GSE47183 and GSE99340. The liver cohorts of The section *Both mechanisms outside kidney: predictions fixed before the data* are GSE89632, GSE48452, GSE162694, GSE126848, GSE83452, GSE66676, GSE63067 and GSE24807, and the colon cohorts GSE75214, GSE87466, GSE179285, GSE11223, GSE47908, GSE38713, GSE53306, GSE16879, GSE48958 and GSE13367; GSE59071, GSE17470 and GSE61260 were excluded by the reuse audit or for an unmappable matrix. Single-nucleus data are available from the Kidney Precision Medicine Project Atlas (https://atlas.kpmp.org/).

Other public resources. The two protein datasets used for the orthogonal corroboration in The section *Protein-level corroboration in two patient-independent datasets* are the SOMAscan aptamer panel of kidney cortex biopsies (Mendeley Data 83k89shdx5, https://doi.org/10.17632/83k89shdx5.1) and the FFPE cortex LC-MS/MS dataset (PRIDE https://www.ebi.ac.uk/pride/, accession PXD041884). The germline test uses the diabetic-kidney-disease genome-wide association study GCST90179152 from the GWAS Catalog (https://www.ebi.ac.uk/gwas/). The plasma metabolome used in The sections *Absolute AUROC is uninformative in this disease* and *The control-group problem is not specific to diabetic kidney disease* is study ST003255 of the Metabolomics Workbench (https://www.metabolomicsworkbench.org/).

Processed data. The harmonised per-cohort expression matrices, the frozen 9,900-gene Entrez space, the probe-to-gene mapping, and the sample-level phenotype tables including the procurement annotation are deposited at https://doi.org/10.5281/zenodo.23051697 under CC BY 4.0. The cohort audit is distributed with them as a relational database whose views encode the exclusion and overlap rules stated in the section *Cohort assembly and audit*, so that the audit is enforced by the schema rather than by prose.

Code. All analysis code is available at https://github.com/kdh6126/dkd-cross-cohort-audit under the MIT licence [55], archived at https://doi.org/10.5281/zenodo.23051697.

Regeneration. The repository provides a single entry point, `run_all.py`, covering 133 declared stages, of which 122 reproduce the results reported here. Running `python run_all.py --paper` runs the 104 of them that the figures, tables and reported numbers depend on. That set is a static list in the code, and before running anything `--paper` checks it against a dependency closure computed from the declared outputs of every stage and the inputs its command line and source refer to; the run aborts if the two diverge. Inputs are inferred statically, so the check constrains the set rather than proving it complete. Each stage declares the files it produces, so a run resumes where it stopped, and every stage that resamples takes a fixed seed.

Reruns are not bit-identical, and we state why rather than round it away. Of the 104 stages, 12 query external resources at run time (PubMed, the KPMP Atlas, KEGG, Reactome and GEO), so their outputs can depend on the state of the resource on the day of the run, although the four cross-tissue stages among them download fixed GEO series. The reported field observed to change was the PubMed novelty count: rerunning on 2026-09-02 moved one gene from 22 to 23 papers. Tier assignments, which are what the manuscript reports, did not change for any of the 30 candidates, and each candidate's query string, query date and returned identifiers are recorded in Additional file 6 so that any later divergence is traceable.

Declarations

List of abbreviations

AUROC: area under the receiver operating characteristic curve; BRISQ: Biospecimen Reporting for Improved Study Quality; CKD: chronic kidney disease; DKD: diabetic kidney disease; ERCB: European Renal cDNA Bank; GEO: Gene Expression Omnibus; IEG: immediate-early gene; KPMP: Kidney Precision Medicine Project; LODO: leave-one-dataset-out; MASH: metabolic dysfunction-associated steatohepatitis; RBS: robust biomarker score; RUVg: remove unwanted variation using control genes; SPREC: Standard PREanalytical Code; UC: ulcerative colitis; WGCNA: weighted gene co-expression network analysis.

Ethics approval and consent to participate

Not applicable. This study analysed only previously published, de-identified human data obtained from public repositories, and generated no new human or animal data.

Consent for publication

Not applicable. The manuscript contains no data from any individual person.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) [grant number RS-2026-25524613]; an internal research grant of the Electronics and Telecommunications Research Institute (ETRI) [grant number 25YT1400]; and the Culture, Sports and Tourism R&D Program through the Korea Creative Content Agency grant funded by the Ministry of Culture, Sports and Tourism in 2024 [grant number RS-2024-00398310]. The funders had no role in the design of the study, in the collection, analysis and interpretation of data, or in writing the manuscript.

Authors' contributions

DK conceived the study, designed the methodology, implemented the analysis pipeline, curated and harmonised the data, performed the formal analysis, carried out the analyses, designed and ran the validation and control analyses, produced the figures, provided study resources, wrote the original draft and reviewed and edited the manuscript. MB, JP, HL and HJL designed the methodology, curated and harmonised the data and reviewed and edited the manuscript. SL acquired funding, administered the project, supervised the work and reviewed and edited the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The results reported here are in whole or in part based upon data generated by the Kidney Precision Medicine Project. Accessed August 24, 2026. https://www.kpmp.org. The KPMP study is supported by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). We acknowledge the essential contribution of the patient participants who consented to kidney biopsy for research.

We also thank the European Renal cDNA Bank consortium and the investigators who deposited the public datasets analysed here, without which this study would not have been possible.

Disclosure of use of AI-assisted tools including generative AI

Claude Code (Anthropic) and ChatGPT (OpenAI) assisted with code development and refactoring, proposals for control analyses, language editing and revision of author-written text, and the code that draws the figures; no figure or result was generated by an AI model directly. Numerical results were computed from the cited datasets or explicitly described simulations using the analysis code. Automated checks were used to assess numerical consistency and provenance. The authors reviewed the code, analyses and manuscript and take responsibility for the final content. The tools are cited as [56, 57].

Additional files

Additional file 1 (PDF). Comparator implementation and its validation: what the WGCNA hub ranking used here does and does not share with the canonical pipeline, the scale-free topology fit across soft powers, the 32-variant grid testing whether The section *Why one method escaped the confounder, and why that is not robustness* depends on those choices, the provenance of the remaining selectors, and the block-size simulation run with the RBS implementation.

Additional file 2 (PDF). Cohort audit report, including the patient-identifier evidence for the three excluded series and the residual overlap that could not be removed.

Additional file 3 (XLSX). Per-cohort harmonisation summary and the complete candidate table with every line of evidence.

Additional file 4 (XLSX). Per-diagnosis results underlying the section *The control-group problem is not specific to diabetic kidney disease*: the effect vectors for each ERCB diagnosis under both control definitions, the three external validation datasets, the KPMP single-nucleus contrast summary, and the KPMP modality table by diagnosis.

Additional file 5 (PDF). Candidate evidence over all 30 specificity-gated candidates: for each candidate the selection and control tests, the human-kidney context, the literature status, and the classification by current evidence strength and prior DKD literature.

Additional file 6 (PDF). The two proteomic cohorts used in the section *Protein-level corroboration in two patient-independent datasets*: what each source publication and repository record states about case and control definition, tissue, procurement route and pre-analytical reporting, together with the abundance- and peptide-count-matched permutation and its achieved balance, and a negative control asking whether the two proteomic datasets could have produced a candidate list on their own.

Additional file 7 (XLSX). The test in liver and colon (the section *Both mechanisms outside kidney: predictions fixed before the data*): the registered predictions and post hoc plan with their SHA-256 digests; every cohort considered, with platform, case and control definitions, procurement coding per arm with its evidence, and inclusion status; the value-vector reuse audit; per-cohort immediate-early effects; the composition sweeps; and the post hoc analyses.

References

1. Haynes W, Vallania F, Liu C, Bongen E, Tomczak A, Andres-Terrè M, et al. Empowering multi-cohort gene expression analysis to increase reproducibility. Pac Symp Biocomput 2017;22:144–153. PMID: 27896970.
2. Meinshausen N, Bühlmann P. Stability selection. Journal of the Royal Statistical Society: Series B 2010;72(4):417–473.
3. Simon R, Radmacher M, Dobbin K, McShane L. Pitfalls in the use of DNA microarray data for diagnostic and prognostic classification. J Natl Cancer Inst 2003;95(1):14–8. PMID: 12509396.
4. Whalen S, Schreiber J, Noble W, Pollard K. Navigating the pitfalls of applying machine learning in genomics. Nat Rev Genet 2022;23(3):169–181. PMID: 34837041.
5. Diaz-Uriarte R, Gómez de Lope E, Giugno R, Fröhlich H, Nazarov P, Nepomuceno-Chamorro I, et al. Ten quick tips for biomarker discovery and validation analyses using machine learning. PLoS Comput Biol 2022;18(8):e1010357. PMID: 35951526.
6. Leek J, Scharpf R, Bravo H, Simcha D, Langmead B, Johnson W, et al. Tackling the widespread and critical impact of batch effects in high-throughput data. Nat Rev Genet 2010;11(10):733–9. PMID: 20838408.
7. Boulesteix A, Lauer S, Eugster M. A plea for neutral comparison studies in computational sciences. PLoS One 2013;8(4):e61562. PMID: 23637855.
8. Jelizarow M, Guillemot V, Tenenhaus A, Strimmer K, Boulesteix A. Over-optimism in bioinformatics: an illustration. Bioinformatics 2010;26(16):1990–8. PMID: 20581402.
9. Ullmann T, Peschel S, Finger P, Müller C, Boulesteix A. Over-optimism in unsupervised microbiome analysis: Insights from network learning and clustering. PLoS Comput Biol 2023;19(1):e1010820. PMID: 36608142.
10. Weber L, Saelens W, Cannoodt R, Soneson C, Hapfelmeier A, Gardner P, et al. Essential guidelines for computational method benchmarking. Genome Biol 2019;20(1):125. PMID: 31221194.
11. Mangul S, Martin L, Hill B, Lam A, Distler M, Zelikovsky A, et al. Systematic benchmarking of omics computational tools. Nat Commun 2019;10(1):1393. PMID: 30918265.
12. Norel R, Rice J, Stolovitzky G. The self-assessment trap: can we all be better than average? Mol Syst Biol 2011;7:537. PMID: 21988833.
13. Dehghani M, Tay Y, Gritsenko A, Zhao Z, Houlsby N, Diaz F, et al. The benchmark lottery. arXiv 2021;ArXiv:2107.07002.
14. Haury A, Gestraud P, Vert J. The influence of feature selection methods on accuracy, stability and interpretability of molecular signatures. PLoS One 2011;6(12):e28210. PMID: 22205940.
15. Ebiele M, Bendechache M, Brennan R. Bias in filter feature selection evaluation: a meta-analysis of datasets, baselines, and experimental design choices. arXiv 2026;ArXiv:2606.07068.
16. Angelakis A. REDE: a quantitative framework for differential-expression reproducibility and diagnostic transfer across nine cohorts and three cancers. arXiv 2026;ArXiv:2608.02796.
17. Woroniecka K, Park A, Mohtat D, Thomas D, Pullman J, Susztak K. Transcriptome analysis of human diabetic kidney disease. Diabetes 2011;60(9):2354–69. PMID: 21752957.
18. Pan Y, Jiang S, Hou Q, Qiu D, Shi J, Wang L, et al. Dissection of Glomerular Transcriptional Profile in Patients With Diabetic Nephropathy: SRGAP2a Protects Podocyte Structure and Function. Diabetes 2018;67(4):717–730. PMID: 29242313.
19. Grayson P, Eddy S, Taroni J, Lightfoot Y, Mariani L, Parikh H, et al. Metabolic pathways and immunometabolism in rare kidney diseases. Ann Rheum Dis 2018;77(8):1226–1233. PMID: 29724730.
20. Fan Y, Yi Z, D'Agati V, Sun Z, Zhong F, Zhang W, et al. Comparison of Kidney Transcriptomic Profiles of Early and Advanced Diabetic Nephropathy Reveals Potential New Mechanisms for Disease Progression. Diabetes 2019;68(12):2301–2314. PMID:

31578193.
21. Jin Q, Li L, Qu P, Ma F, Li P, Qiao Y, et al. Mitigation of renal tubular injury by SIRT6 may improve individual outcomes in diabetic kidney disease-potential mechanisms involving epigenetic repression of inflammatory responses. J Adv Res 2026;85:281–294. PMID: 41076119.
22. Park S, Lee H, Lee J, Lee S, Cho S, Huh H, et al. RNA-seq profiling of tubulointerstitial tissue reveals a potential therapeutic role of dual anti-phosphatase 1 in glomerulonephritis. J Cell Mol Med 2022;26(12):3364–3377. PMID: 35488446.
23. Eadon M, Lampe S, Baig M, Collins K, Melo Ferreira R, Mang H, et al. Clinical, histopathologic and molecular features of idiopathic and diabetic nodular mesangial sclerosis in humans. Nephrol Dial Transplant 2021;37(1):72–84. PMID: 33537765.
24. Sur S, Nguyen M, Boada P, Sigdel T, Sollinger H, Sarwal M. FcER1: A Novel Molecule Implicated in the Progression of Human Diabetic Kidney Disease. Front Immunol 2021;12:769972. PMID: 34925339.
25. Baelde H, Eikmans M, Doran P, Lappin D, de Heer E, Bruijn J. Gene expression profiling in glomeruli from human kidneys with diabetic nephropathy. Am J Kidney Dis 2004;43(4):636–50. PMID: 15042541.
26. Sircar M, Rosales I, Selig M, Xu D, Zsengeller Z, Stillman I, et al. Complement 7 Is Up-Regulated in Human Early Diabetic Kidney Disease. Am J Pathol 2018;188(10):2147–2154. PMID: 30253844.
27. Neusser M, Lindenmeyer M, Moll A, Segerer S, Edenhofer I, Sen K, et al. Human nephrosclerosis triggers a hypoxia-related glomerulopathy. Am J Pathol 2010;176(2):594–607. PMID: 20019191.
28. El-Achkar T, Eadon M, Menon R, Lake B, Sigdel T, Alexandrov T, et al. A multimodal and integrated approach to interrogate human kidney biopsies with rigor and reproducibility: guidelines from the Kidney Precision Medicine Project. Physiol Genomics 2021;53(1):1–11. PMID: 33197228.
29. Lake B, Menon R, Winfree S, Hu Q, Melo Ferreira R, Kalhor K, et al. An atlas of healthy and injured cell states and niches in the human kidney. Nature 2023;619(7970):585–594.
30. Hannenhalli S, Putt M, Gilmore J, Wang J, Parmacek M, Epstein J, et al. Transcriptional genomics associates FOX transcription factors with human heart failure. Circulation 2006;114(12):1269–76. PMID: 16952980.
31. Ma Y, Dai H, Kong X. Impact of warm ischemia on gene expression analysis in surgically removed biosamples. Anal Biochem 2012;423(2):229–35. PMID: 22343191.
32. Ferreira P, Muñoz-Aguirre M, Reverter F, Sá Godinho C, Sousa A, Amadoz A, et al. The effects of death and post-mortem cold ischemia on human tissue transcriptomes. Nat Commun 2018;9:490.
33. Menon R, Kimmel P, Otto E, Subramanian L, O'Connor C, Godfrey B, et al. Not all reference samples are equal in single-cell transcriptomics of human kidney tissue. JCI Insight 2026;11(14):e204385.
34. Feng S, Gao Y, Yin D, Lv L, Wen Y, Li Z, et al. Identification of Lumican and Fibromodulin as Hub Genes Associated with Accumulation of Extracellular Matrix in Diabetic Nephropathy. Kidney Blood Press Res 2021;46(3):275–285. PMID: 33887734.
35. Hirohama D, Abedini A, Moon S, Surapaneni A, Dillon S, Vassalotti A, et al. Unbiased human kidney tissue proteomics identifies matrix metalloproteinase 7 as a kidney disease biomarker. J Am Soc Nephrol 2023;34(7):1279–1291. PMID: 37022120.
36. Schwab S, Harris P, Michel C, McGinnis C, Nahomi R, Assiri M, et al. Quantifying protein acetylation in diabetic nephropathy from formalin-fixed paraffin-embedded tissue. Proteomics Clin Appl 2024;18(6):e202400018. PMID: 38923810.
37. Mantel N, Haenszel W. Statistical Aspects of the Analysis of Data From Retrospective Studies of Disease. JNCI: Journal of the National Cancer Institute 1959;22(4):719–748.
38. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B: Statistical Methodology 1995;57(1):289–300.
39. Betsou F, Lehmann S, Ashton G, Barnes M, Benson E, Coppola D, et al. Standard preanalytical coding for biospecimens: defining the sample PREanalytical code. Cancer Epidemiol Biomarkers Prev 2010;19(4):1004–11. PMID: 20332280.
40. Lehmann S, Guadagni F, Moore H, Ashton G, Barnes M, Benson E, et al. Standard preanalytical coding for biospecimens: review and implementation of the Sample PREanalytical Code (SPREC). Biopreserv Biobank 2012;10(4):366–74. PMID: 24849886.
41. Moore H, Kelly A, Jewell S, McShane L, Clark D, Greenspan R, et al. Biospecimen reporting for improved study quality (BRISQ). J Proteome Res 2011;10(8):3429–38. PMID: 21574648.
42. Whitlock M. Combining probability from independent tests: the weighted Z -method is superior to Fisher's approach. Journal of Evolutionary Biology 2005;18(5):1368–1373.
43. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. BMC Bioinformatics 2008;9:559. PMID: 19114008.
44. Tibshirani R. Regression Shrinkage and Selection Via the Lasso. Journal of the Royal Statistical Society Series B: Statistical Methodology 1996;58(1):267–288.
45. Zou H, Hastie T. Regularization and Variable Selection Via the Elastic Net. Journal of the Royal Statistical Society Series B: Statistical Methodology 2005;67(2):301–320.
46. Peng H, Long F, Ding C. Feature selection based on mutual information criteria of max-dependency, max-relevance, and min-redundancy. IEEE Transactions on Pattern Analysis and Machine Intelligence 2005;27(8):1226–1238.
47. Kononenko I. Estimating attributes: Analysis and extensions of RELIEF. In: Lecture Notes in Computer Science Springer; 1994. p. 171–182.
48. Breiman L. Random Forests. Machine Learning 2001;45(1):5–32.
49. Kursa M, Rudnicki W. Feature Selection with the Boruta Package. Journal of Statistical Software 2010;36(11).
50. Hedges L. Distribution Theory for Glass's Estimator of Effect size and Related Estimators. Journal of Educational Statistics 1981;6(2):107–128.
51. Venet D, Dumont J, Detours V. Most random gene expression signatures are significantly associated with breast cancer outcome. PLoS Comput Biol 2011;7(10):e1002240. PMID: 22028643.
52. Risso D, Ngai J, Speed T, Dudoit S. Normalization of RNA-seq data using factor analysis of control genes or samples. Nat Biotechnol 2014;32(9):896–902. PMID: 25150836.
53. Leek J, Storey J. Capturing heterogeneity in gene expression studies by surrogate variable analysis. PLoS Genet 2007;3(9):1724–35. PMID: 17907809.
54. Johnson W, Li C, Rabinovic A. Adjusting batch effects in microarray expression data using empirical Bayes methods. Biostatistics 2007;8(1):118–27. PMID: 16632515.
55. Kim D, Bae M, Park J, Lee H, Lim H, Lee S, Auditing cross-cohort feature-selection benchmarks: code, harmonised cohorts and audit database. Zenodo; 2026. Version 1.0.0. https://doi.org/10.5281/zenodo.23051697.
56. Anthropic, Claude Code (Claude models served during 2026, latest Claude Fable 5.1, claude-fable-5-1) [Large language model agent]; 2026. Responses to queries made by Dohun Kim, January to October 2026, for code development and refactoring, proposals for control analyses, language editing of author-written text and figure-drawing code. https://claude.com/claude-code.
57. OpenAI, ChatGPT (web interface, models as deployed during 2026) [Large language model]; 2026. Responses to queries made by Dohun Kim, January to October 2026, for language editing of author-written text and proposals for control analy-

ses. https://chatgpt.com.