

Molecular analysis of primary melanoma T cells identifies patients at risk for metastatic recurrence

Pruessmann *et al.* 2020 Nat Cancer. doi: [10.1038/s43018-019-0019-5](https://doi.org/10.1038/s43018-019-0019-5)

Presented by Kim Dill-McFarland


2021.06.07 @kdillmcfarland

Molecular analysis of primary melanoma T cells identifies patients at risk for metastatic recurrence

Pruessmann *et al.* 2020 Nat Cancer. doi: [10.1038/s43018-019-0019-5](https://doi.org/10.1038/s43018-019-0019-5)

Presented by Kim Dill-McFarland
2021.06.07 @kdillmcfarland

2020 Melanoma cumulative risk of incidence



All rights reserved. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization / International Agency for Research on Cancer concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate borderlines for which there may not yet be full agreement.

Data source: GLOBOCAN 2020
Graph production: IARC
(<http://gco.iarc.fr/today>)
World Health Organization

Melanoma risk factors ¹

- Ultraviolet (UV) light exposure
 - Especially sunburn
- Age
- Skin irregularities like freckles and moles
 - < 30% of skin cancers originate in a mole
- Skin color
 - Rates of melanoma are lower in non-Hispanic Black compared to White populations (1:25)
 - Mortality due to melanoma is disproportionately high in non-Hispanic Black populations (1:10) ²

Recurrence

- Surgical removal often effective but...
- Metastatic recurrence common at > 20% of primary melanomas within 5 years ³
- Prediction of recurrence can aid monitoring and treatment plans


T cells as predictors of recurrence

- Melanomas cause an adaptive immune response ⁵
- Tumor-infiltrating lymphocytes associated with less recurrence ⁶
- T-cell receptor clonality predicts response to some therapies ⁷


Primary hypothesis

T-cell fraction (TCFr) and / or T-cell clonality
improves prediction of primary melanoma
recurrence compared to current clinical
measures alone

Design



TCRB sequencing targets CDR3



Genomic segments:

Leading peptide

Variable

Diversity

Joining

Constant

T-cell metrics

- T-cell clonality
 - > 100 T cells per sample
 - Simpson measure = $\sqrt{\sum p_i^2}$ p_i = abundance of clone i / total clones
 - 0 = highly polyclonal, high-diversity


- T-cell fraction (TCFr)
 - > 500 nucleated cells per sample
 - Total T cells / total nucleated cells

T-cell fraction (TCFr) as a
predictor of recurrence

TCFr associated with recurrence independently from other variables



	TCFr association P-value
Recurrence	< 1E4
Survival	< 1E4
Tumor thickness	0.52
Age	0.76
Sex	0.41
Nodal disease	0.15
Ulceration	0.89
Mitotic rate	0.41

Design




TCFr cutoff to best predict recurrence


- 80% subsample bootstrapping
- 0.05 incremental TCFr cutoffs
- Minimize change in true positive rate (TPR) and F-score = $2 \times \frac{\text{precision} \times \text{recall}}{\text{precision} + \text{recall}}$




20% TCFr validated in testing set




20% TCFr improves recurrence prediction



20% TCFr among top predictors of recurrence




TCFr correlated with more traditional measures
but TCFr is more predictive



Analysis of deviance $P < 0.01$
TCFr model vs

- TIL MIA grade
- TIL briskness

TCFr correlated with some T-cell subsets
but TCFr is more predictive



None predict recurrence, $P > 0.05$

Conclusions

- T-cell fraction (TCFr) is a unique measure not correlated with a number of current clinical measures
- TCFr less than 20% is associated with higher primary melanoma recurrence
- 20% TCFr improves recurrence prediction in conjunction with current clinical measures, particularly tumor thickness
- Treg proportion within TCFr may further improve prediction

Primary hypothesis

T-cell fraction (TCFr) ~~and / or T-cell clonality~~
improves prediction of primary melanoma
recurrence compared to current clinical
measures alone

Next steps

Incorporate TCFr measures with


- anti-PD-1 blockade (or other treatment) trial
- metastatic melanoma
- < 1 mm melanoma

My next step proposed


Hypothesis

T-cell fraction (TCFr) improves prediction of high-mutation but not low-mutation burden cancer outcomes


Design



Cancer type selection



Sample size



- Expect high-burden hazard ratio near melanoma (0.8)
- Expect sample failure at mean 1 to 5-year-old samples (15%)
- Goal: 118 samples per cancer (after failure ≈ 100)

Potential variables

Outcome

- Recurrence
- Survival

Co-variate

- Common risks (thickness, location, etc)
- Demographics (age, sex, etc)
- Disease history

Explanatory

- TCFr (20%, continuous)
- Mutation burden (high/low, continuous)

Analyses

- Cox's proportional-hazard model
 - Separate cancers vs combined analysis
 - ± TCFr
 - ± mutation burden
- Sparse partial least-squares regression (sPLS) ¹⁰

References

1. American Cancer Society. www.cancer.org/cancer/melanoma-skin-cancer/about/key-statistics.html
2. American Cancer Society. 2019. *Cancer Facts & Figures for African Americans*.
3. Elsaesser *et al* 2012 PLoS One. doi: [10.1371/journal.pone.0029791](https://doi.org/10.1371/journal.pone.0029791)
4. Waldman *et al* 2020 Nat Rev Immunol. doi: [10.1038/s41577-020-0306-5](https://doi.org/10.1038/s41577-020-0306-5)
5. Snyder *et al* 2014 N Engl J Med doi: [10.1056/NEJMoa1406498](https://doi.org/10.1056/NEJMoa1406498)
6. Clemente *et al* 1996 Cancer doi: [10.1002/\(SICI\)1097-0142\(19960401\)77:7<1303::AID-CNCR12>3.0.CO;2-5](https://doi.org/10.1002/(SICI)1097-0142(19960401)77:7<1303::AID-CNCR12>3.0.CO;2-5)
7. Tumeh *et al* 2014 Nature doi: [10.1038/nature13954](https://doi.org/10.1038/nature13954)
8. Migalska *et al* 2018 Sci Rep doi: [10.1038/s41598-018-30037-0](https://doi.org/10.1038/s41598-018-30037-0)
9. Martincorena and Campbell 2015 Science doi: [10.1126/science.aab4082](https://doi.org/10.1126/science.aab4082)
10. Lee *et al* 2013 Stat Med doi: [10.1002/sim.5975](https://doi.org/10.1002/sim.5975)

Molecular analysis of primary melanoma T cells identifies patients at risk for metastatic recurrence

Wiebke Pruessmann^{1,8,10}, Julie Rytlewski ^{2,10}, James Wilmott ^{3,10}, Martin C. Mihm Jr¹, Grace H. Attrill³, Beatrice Dyring-Andersen^{1,9}, Paul Fields ², Qian Zhan¹, Andrew J. Colebatch^{3,4}, Peter M. Ferguson^{3,4}, John F. Thompson ^{3,5}, Klaus Kallenbach⁶, Erik Yusko², Rachael A. Clark¹, Harlan Robins^{2,7}, Richard A. Scolyer ^{3,4,10} and Thomas S. Kupper ^{1,10}★